

BMJ Open Performance of injury severity measures in trauma research: a literature review and validation analysis of studies from low-income and middle-income countries

Amber Mehmood,¹ Yuen W Hung,¹ Huan He,^{1,2} Shahmir Ali,^{3,4} Abdul M Bachani¹

To cite: Mehmood A, Hung YW, He H, *et al.* Performance of injury severity measures in trauma research: a literature review and validation analysis of studies from low-income and middle-income countries. *BMJ Open* 2019;**9**:e023161. doi:10.1136/bmjopen-2018-023161

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-023161>).

Received 23 March 2018

Revised 1 August 2018

Accepted 13 November 2018



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Amber Mehmood;
amehmoo2@jhu.edu

ABSTRACT

Introduction Characterisation of injury severity is an important pillar of scientific research to measure and compare the outcomes. Although majority of injury severity measures were developed in high-income countries, many have been studied in low-income and middle-income countries (LMICs). We conducted this study to identify and characterise all injury severity measures, describe how widely and frequently they are used in trauma research from LMICs, and summarise the evidence on their performance based on empirical and theoretical validation analysis.

Methods First, a list of injury measures was identified through PubMed search. Subsequently, a systematic search of PubMed, Global Health and EMBASE was undertaken on LMIC trauma literature published from January 2006 to June 2016, in order to assess the application and performance of injury severity measures to predict in-hospital mortality. Studies that applied one or more global injury severity measure(s) on all types of injuries were included, with the exception of war injuries and isolated organ injuries.

Results Over a span of 40 years, more than 55 injury severity measures were developed. Out of 3862 non-duplicate citations, 597 studies from 54 LMICs were listed as eligible studies. Full-text review revealed 37 studies describing performance of injury severity measures for outcome prediction. Twenty-five articles from 13 LMICs assessed the validity of at least one injury severity measure for in-hospital mortality. Injury severity score was the most commonly validated measure in LMICs, with a wide range of performance (area under the receiver operating characteristic curve (AUROC) between 0.9 and 0.65). Trauma and Injury Severity Score validation studies reported AUROC between 0.80 and 0.98.

Conclusion Empirical studies from LMICs frequently use injury severity measures, however, no single injury severity measure has shown a consistent result in all settings or populations and thus warrants validation studies for the diversity of LMIC population.

INTRODUCTION

Injury remains a major public health problem globally, causing significant death

Strengths and limitations of this study

- The study comprises three parts: summary of all injury severity measures, description of their use in low-income and middle-income countries (LMICs) and their performance to predict in-hospital mortality in LMIC settings.
- Injury severity measures, whether developed exclusively for characterising trauma and injuries, or non-injury severity measures incorporated in trauma research, are both included in this study.
- A systematic electronic search of PubMed, Global Health and EMBASE on literature published from January 2006 to June 2016.
- Validation studies conducted in LMICs are used to estimate the performance of injury severity measures.
- Performance of injury severity measures to predict other outcomes such as blood transfusion requirement, intensive care unit admission or hospital length of stay are not the focus of this study.

and disability across all the age and sex spectrum.¹ A disproportionate share, 90%, of all trauma deaths occur in low-income and middle-income countries (LMICs), where resources to deal with this crisis are inadequate. An efficient and effective trauma system has been found to be a key component. It is estimated that approximately two million lives could be saved annually if LMICs could implement trauma systems comparable with trauma care systems available in high-income countries (HICs).² However, this would require a careful assessment of the gaps and planning to ensure the most efficient use of available resources. Injury severity scoring systems can provide a foundation for benchmarking and performance improvement in the arena of trauma care.³ Characterisation of injury severity is a critical pillar in the provision, and improvement of trauma care for key

activities such as field triage, prognostication, prediction of risk-adjusted outcomes, quality improvement, evaluation of cost and effectiveness of trauma service delivery, planning of services and organisation of resources.⁴ Many injury measures have been formulated over time with a wide range of methodologies.⁵ While no single injury measure is considered the best or the most comprehensive, assessment of injuries in a patient has been aided by assigning numerical values to several indicators including physiological or biochemical parameters, anatomical descriptors, age and so on, and combining these values to an overall measure of injury severity.^{6,7} Although injury severity measures are most often used for the purpose as they were developed, such as triage or mortality prediction, it is not uncommon to validate and use them for other functions.^{8,9}

There has been a proliferation of injury severity measures over the past few decades.^{7,10} While a variety of injury severity measures have been developed exclusively for trauma and injuries, other non-injury severity measures have also been incorporated in trauma research on many occasions.^{11–14} These severity measures use a range of clinical, biochemical, demographic and physical attributes to create indicators for prognostic predictions and performance evaluation.^{4,15} However, both the utilisation and validation of injury scores in clinical care or outcome research has been sparse in LMICs.¹⁶ There are multiple reasons for this but in many cases, especially for those injury severity measures developed in high-income settings, the information needs are challenging for a low-resource environment.^{11,15,17–19} Many well-recognised injury measures were sometimes applied without being validated in the populations under study. Subsequently, studies have documented poor performance of injury severity measures such as Trauma and Injury Severity Score (TRISS), when applied to other populations using the coefficients derived from the Major Trauma Outcome Study.^{20–26} However, there is a dearth in the literature on utilisation of common injury severity measures, and whether they show acceptable performance in terms of validity and reliability to support their use in LMICs. This gap limits our ability to translate high-quality injury research methods developed in HICs into effective decision support and quality improvement systems for LMICs. The aim of this study was therefore to fill this gap in the literature through a thorough review of the literature; specifically we sought to: (1) identify all the measures and scoring systems that were ever developed to measure injury severity, and summarise their characteristics; (2) describe how widely and frequently the key measures are used in LMICs and (3) summarise the evidence on their measurement performance based on empirical validation analysis and theoretical analysis of their applicability.

METHODS

For our first aim, we conducted a literature search for terms ‘injury AND severity measures’ OR ‘injury AND

scores’ OR ‘Injury AND scales’, as well ‘Trauma AND severity measures’ to include those that are not exclusive to injuries but have been used in trauma and injury research. A list of injury measures was identified through PubMed search. Subsequently, using bibliographies of the results of the primary search, a secondary search was performed to find the original literature of the injury measure development. Full text of all publications was reviewed to understand and describe the initial purpose and scope of development of the injury measure, its main components, year of first publication and country of development.

For the specific aims 2 and 3, we conducted a detailed literature review to assess the application and performance of injury severity measures to predict in-hospital mortality, conducted in LMICs. We included studies of global trauma populations and specific injury pathologies and used World Bank’s classification for LMICs in the year 2016.

Eligibility criteria

For the purpose of determining the applications of different injury severity measures in LMICs, we included studies that applied one or more global injury severity measure(s) on any type of injury population, except for studies that focused only on poisoning, drowning and ocular trauma. We excluded studies that applied exclusively organ specific injury severity measure(s), population from low income country treated in a high-income setting, as well as studies describing only combat injuries or those from military trauma registries due to the environment and contexts largely different from general LMICs settings.

Information sources and search strategy

We conducted a systematic electronic search of PubMed, Global Health and EMBASE on literature published from January 2006 to June 2016. We used combinations of search terms including medical subject heading and keywords on two groups: ‘trauma or injury measures’, and a list of ‘LMICs’ (online supplementary file 1). We applied human subjects restrictions but language restrictions were not applied. All references were exported to Endnote V.7 and duplicated studies were excluded using Endnote before exporting them into an Excel spreadsheet.

Two authors (AM and SA) independently screened the titles and abstracts of all studies resulted from the above search strategy to identify the eligible studies for the applications of injury severity measures in LMICs. Full-text version of all the eligible articles were sought, and if full text was not available in English language, the abstracts were excluded from further analysis. All eligible full-text articles were reviewed for relevance and data collection.

Data abstraction

Data were extracted from the selected studies using a predesigned electronic data collection form. The studies were further categorised into validation studies




Figure 1 Flow diagram of search strategy and study selection according to Preferred Reporting Items for Systematic Reviews and Meta-Analysis guidelines.

or empirical/non-validation studies, or excluded if they did not match the inclusion criteria on full-text review (figure 1).

To assess the performance of injury severity measures and prediction of in-hospital mortality, we selected studies that estimated the Area Under the Receiver Operating Characteristic curve (AUROC) or correlation between specific injury severity measure and in-hospital mortality, based on the studies identified with applications of injury severity measures in LMICs. Studies that did not specify the outcome of assessment or did not include any estimates of AUROC, correlation or sensitivity and specificity were excluded. Three authors (AM, HH and YWH) screened these identified studies for the performance on predicting in-hospital mortality. Any disagreements were resolved by discussions among the three authors.

For the purpose of determining applications of different injury severity measures in LMICs, three authors (AM, HH and YWH) extracted information on the injury severity measures used in each study, whether performance was assessed on in-hospital mortality prediction, and the country in which the study was conducted. The studies and corresponding injury measures were assessed in detail for study population, type of injury and injury mechanism, injury severity measures, study methods, in-hospital mortality prediction and their corresponding performance in predicting in-hospital mortality. The performance of the injury severity measures is reported

as AUROC and calibration as Hosmer-Lemeshow (H-L) goodness of fit test.

Patient and public involvement

This study did not involve patients or human subjects directly or indirectly, and the results of the analysis were solely based on the previously published literature.

RESULTS

The results are described in order of specific objectives of the study. Our study demonstrates considerable growth in the science of injury severity measurement globally as well as in LMICs. Table 1 summarises the search results of different injury measures, categorised according to the primary purpose of their development and their core components. It shows clearly that the science of injury severity measures had essentially taken off in early 1970s, and it is still ongoing with similar enthusiasm. Almost 60 severity measures or scoring systems have been developed either exclusively for injury and trauma research, or have been used in measuring the severity of injuries. Many injury severity measures were developed to support epidemiological research and performance evaluation; examples include Abbreviated Injury Scale (AIS), Injury Severity Score (ISS) and New Injury Severity Scores (NISS), A Severity Categorization of Trauma and International Classification for Diseases-9 ISS (ICISS). Others,

Table 1 List of injury severity measures, their purpose and components

	Measures	Year, country	Components
Primary purpose: epidemiological research and evaluation			
1	Abbreviated Injury Scale (AIS) ³¹	1971, USA	Anatomical description of injuries.
2	Comprehensive Research Injury Scale ³²	1972, USA	Energy dissipation, threat-to-life, permanent impairment, treatment period, incidence.
3	Injury Severity Score (ISS) ³³	1974, USA	AIS.
4	Estimated Survival Probability (ESP) index ³⁴	1978, USA	International Classification for Diseases (ICDA) codes.
5	Penetrating and Blunt code ³⁵	1978, USA	Anatomical description of injuries with limited physiological responses.
6	Wisconsin Trauma Index ¹⁷	1980, USA	Involvement of different organ systems, burns, age, pre-existing condition.
7	Anatomic Index ³⁶	1980, USA	Hospital adaptation of ICDA Discharge Diagnosis.
8	Revised ESP score ³⁷	1982, USA	Hospital ICDA Discharge Diagnosis, age.
9	Probability of Death Score ³⁸	1983, Denmark	Recategorisation of AIS.
10	TRauma and Injury Severity Score (TRISS) ISS ²³	1987, USA	Revised Trauma Score (RTS), ISS, age, mechanism of trauma.
11	Organ Injury Scale ³⁹⁻⁴²	1989, USA	Anatomic description, blood loss.
12	Anatomic Profile ⁴³	1990, USA	AIS, summary scores for body regions A through D.
13	A Severity Categorization of Trauma ⁴³	1990, USA	Emergency department RTS, patient age, AIS-85.
14	Neural networks ⁴⁴	1993, USA	RTS, ISS, age.
15	ICD-9 Injury severity score (ICISS) ⁴⁵	1996, USA	ISS, ICD-9 injury descriptors.
16	New ISS (NISS) ⁴⁶	1997, USA	AIS.
17	MAX AIS ⁴⁷	2002, USA	Maximum AIS score.
18	Trauma Registry AIS Score ⁴⁸	2003, USA	AIS derived survival risk ratio.
19	Turkish Injury Scale ⁴⁹	2003, Turkey	Injury severity according to Turkish Penal Code.
20	Revised Injury Severity Classification Score ²¹	2009, Germany	AIS, age, sex, head injury, biochemical and physiological parameters, cardiopulmonary resuscitation.
Primary purpose: triage and decision support			
21	Trauma Index (TI) ⁵⁰	1971, USA	Region and type of injury, cardiovascular, central nervous system, respiratory status.
22	Glasgow Coma Scale (GCS) ⁵¹	1974, UK	Eye opening, motor and verbal response.
23	Illness-injury Severity index ⁵²	1979, USA	Physiological parameters, region and type of injury, pre-existing condition.
24	Trauma Score ⁵³	1981, USA	Respiratory effort, capillary refill, respiratory rate (RR), systolic blood pressure (SBP), GCS.
25	Circulation, Respiration, Abdomen, Motor and Speech (CRAMS) ⁵⁴	1982, USA	Capillary refill, respiration, abdominal injuries, motor and verbal response.
26	Prehospital Index (PHI) ⁸	1986, USA	SBP, pulse, RR and level of consciousness.
27	Rapid Acute Physiology Score ⁵⁵	1987, USA	Truncated version of Acute Physiology And Chronic Health Evaluation (APACHE) II Pulse, BP, GCS, RR.
28	RTS ⁵⁶	1989, USA	Sum of weighted values of GCS, BP, RR.
29	Kampala Trauma Score (KTS) ⁵⁷	1996, Uganda	Age, number of serious injury, SBP, RR, neurological status (AVPU).
30	Full Outline of UnResponsiveness (FOUR) score ⁵⁸	2005, USA	Physiological score consisting of eye, motor, brainstem and respiratory components.

Continued

Table 1 Continued

	Measures	Year, country	Components
31	Trauma Associated Severe Hemorrhage Score ⁵⁹	2006, Germany	SBP, haemoglobin, free peritoneal fluid, base excess, complex fractures, pulse and sex.
32	Prehospital paediatric trauma classification ⁶⁰	2006, Brazil	Physiological status, trauma mechanism and anatomic injuries.
33	Ganga Hospital Score ⁶¹	2006, India	Severity of injury to the skin, bones and muscles of the limb; presence of comorbidities.
34	Assessment of Blood Consumption (ABC) Score ⁶²	2008, USA	SBP, positive abdominal ultrasound, pulse and penetrating injury.
35	Emergency Trauma Score ⁶³	2009, Germany	Age, prehospital GCS, base excess, prothrombin time.
36	Acidosis, Blood loss, Cold, Damage (ABCD) ⁶⁴	2012, USA	Acidosis, blood loss, temperature, NISS.
Primary purpose: outcome prediction			
37	Glasgow Outcome Scale ⁶⁵	1975, UK	Assessment of disability from recovery to death.
38	APACHE I ¹⁸	1981, USA	Physiological variables, age, preadmission health status; all disease categories.
39	Penetrating Abdominal Trauma Index ⁶⁶	1981, USA	Anatomical injury severity for each organ involved in penetrating trauma.
40	Simplified Acute Physiology Score (SAPS or s-APACHE) ⁶⁷	1984, France	Abbreviated version of APACHE.
41	APACHE II ⁶⁸	1985, USA	Physiological variables, age, chronic health; all disease categories.
42	Pediatric Risk of Mortality score ⁶⁹	1988, USA	14 Physiological and biochemical parameters.
43	Mangled Extremity Score ⁷⁰	1990, USA	Composite score of tissue damage, ischaemia, shock and age.
44	APACHE III ⁷¹	1991, USA	Acute physiological abnormalities, age, pre-existing functional limitations.
45	Shock Index ⁷²	1992, USA	Ratio of pulse rate vs SBP.
46	Rixen Score ⁷³	1999, Germany	Age, GCS, ISS, base excess, prothrombin time.
47	GCS Extended ⁷⁴	2000, UK, South Africa	Eye, verbal and motor response PLUS Amnesia Scale.
48	KTS-II ⁷⁵	2002, Uganda	Age, SBP, RR on admission, neurological status (AVPU), number of serious injuries.
49	Rapid Emergency Medicine Score ⁷⁶	2004, Germany	Coma, respiratory frequency, oxygen saturation, blood pressure, pulse rate and age.
50	FLAMES Score ⁷⁷	2008, Canada	Age, APACHE II score, extent of burn and sex.
51	Trauma Mortality Prediction Model ICD-9 ⁷⁸	2009, USA	ICISS, ICD-9.
52	Mechanism, GCS, Age, Pressure (MGAP) ⁷⁹	2010, France	Mechanism (blunt vs penetrating), GCS, age, SBP.
53	Sequential Trauma Score ¹¹	2010, Germany	Age, mechanism, clinical interventions, biochemical and physiological parameters, AIS.
54	GCS, age, pressure (GAP) ⁸⁰	2011, Japan	GCS, age, SBP.
55	NORwegian survival prediction Model In Trauma ⁸¹	2014, Norway	NISS, RTS, age, preinjury comorbidity score.
56	Exponential ISS ⁸²	2014, China	AIS-derived injury score.
57	Tangent ISS ⁸³	2015, China	AIS-derived injury score.

AVPU, alert, voice, pain, unresponsive; FLAMES, fatality by Longevity, APACHE II score, Measured Extent of burn, and sex.

such as Revised Trauma Score (RTS); Circulation, Respiration, Abdomen, Motor and Speech (CRAMS); Acidosis, Blood loss, Cold, Damage (ABCD) and Kampala Trauma Scores (KTS) were developed to help in decision making,

for example, prehospital triage and in-hospital patient disposition. A number of injury measures were developed for the purpose of outcome prediction; Trauma Mortality Prediction Model (TMPM), Rapid Emergency Medicine




Figure 2 Low-income and middle-income countries' publications using trauma/injury severity measures: 2006–2016.

Score and Glasgow Coma Scale (GCS), Age, Pressure are some examples.

Table 1 highlights that a number of empirically developed anatomical, physiological and composite measures such as AIS, or GCS, later became the basis of more complex measures such as RTS, ISS and Revised Injury Severity Classification score, and some of them (RTS, ISS, NISS) in turn became components of a more complex scoring system such as TRISS, Sequential Trauma Score and so on. The use of injury measures in studies published by different LMICs is depicted in figure 2. A total of 597 studies from 54 LMICs were listed as eligible studies between 2006 and 2016 which were a combination of empirical, epidemiological, review and validation studies. China, Turkey, Iran, South Africa, Colombia and Brazil are some of the upper-middle-income countries that contributed to the majority of injury literature published in the last 10 years (figure 3), whereas India, Pakistan, Nigeria and Tanzania are some of the lower-middle-income and low-income countries that extensively used injury measures in a number of injury and trauma-related publications. Thirty-one publications described multi-country studies which may also include an HIC. Approximately 31% (n=186) of all studies were related to head or traumatic brain injuries (TBI).

Table 2 outlines different injury measures used in publications from 54 LMICs in injury-related research. GCS, ISS, TRISS and RTS are the most commonly

used injury measures; however, some attempts have been made to develop new injury measures. Examples include Exponential Injury Severity Score (EISS), Ganga Hospital Score for lower limb fractures, Tangent Injury severity score (TISS) and some novel biomarkers such as lactate and serum acetylcholinesterase. Other scores that were not traditionally used in injury or trauma research such as McLaughlin, Modified Rankin, South African Triage Score, Modified Early Warning System and Rwanda mortality prediction model have also been used for prediction of mortality in trauma populations. Glasgow Outcome Scale is widely used in documenting the outcomes of TBI, and Functional Independence Measure was used in some studies focusing on functional outcomes of injured patients. Some attempts have been made to modify existing injury measures; for example, in Simplified RTS, Glasgow Coma Scale was replaced by five graded levels of consciousness, or NISS was used instead of traditional ISS in TRISS method.

Full-text review of eligible articles was conducted to understand the validity of these new or existing injury measures and revealed that 37 studies examined the performance of injury severity measures for the prediction of hospital length of stay, in-hospital mortality and functional outcome of injured patients. Online supplementary file 2 details 25 of 37 validation studies, as the remaining 12 use different outcomes (eg, respiratory failure, intensive care unit (ICU) admission, etc) or use




Figure 3 Top 10 countries with trauma/injury publications.

a different algorithm. These 25 articles from 13 LMICs assessed the validity of at least one injury severity measure in hospital settings. ISS was the most commonly validated measure in LMICs in the past 10 years, assessed in 11 studies. TRISS was the second most commonly validated injury severity measure in LMICs, followed by GCS, APACHE II and NISS. GCS was more commonly assessed among head/TBI, while also validated among patients with general injuries. The majority of validation studies included all injury mechanisms, some studies included critically ill populations such as ICU patients, while others included patients admitted to the emergency room. The proportion of mortality also varied widely among different settings, ranging from 0.6% to 40%.

Among injury severity measures that were validated in multiple contexts, many presented a wide range of AUROC estimates. Out of the 11 validation studies on ISS, 5 estimated AUROC above 0.90, and 2 of the studies had AUROC below 0.70 with 95% CI overlapping 0.65. Similarly, as majority of the validation studies on TRISS reported AUROC between 0.80 and 0.98, three studies reported 95% CI of AUROC overlapping 0.70. More than a third of the validation studies did not present 95% CI estimates of AUROC, and more than half of the validation studies did not provide estimates on calibration (15 studies).

A majority of the validation studies included only adults and sometimes adolescents. A third of the validation studies included both adults and children, and one study included only paediatric injury population. Many of the validation studies also did not report proportion of missing data. Of those articles that mentioned about

missing data, all excluded records with missed information from analyses.

Besides using in-hospital death as outcome, other studies included morbidity outcomes such as length of hospitalisation, damage control resuscitation, severe trauma, life-threatening injury, respiratory failure and sepsis. These morbidity outcomes are less standardised and therefore limit the ability for comparison.

DISCUSSION

Our review points to an ongoing search for a comprehensive yet simple scoring system applicable to LMICs research and trauma care needs. While Glasgow Coma Scale, AIS and its derivatives, and TRISS methodology have established themselves as gold standards in injury research, there seems to be a need for injury severity measures that are reliable even in the light of the realities facing patient care systems in LMICs. Looking closely at the components of injury measures, it is evident that many complex measures require a host of information starting from prehospital phase until the discharge from the hospital. Henceforth, resources required to record the anatomical and biochemical evidence of injury severity are more readily available in high-income settings but may be difficult to obtain in resource-constrained environments.

Injuries and their physiological response are complex mechanisms, and the outcome of injuries is frequently affected by a number of factors ranging from age and pre-existing conditions of the patient to biochemical response of the body. It is difficult to account for all

Table 2 Injury measures used in last 10 years' published literature from LMICs

Country	Injury measures
Algeria	GCS, ISS.
Argentina	GCS, GOS-E, Modified Rankin Scale.
Bangladesh	GCS, GOS.
Benin	GCS.
Bosnia	ISS.
Brazil	AIS, RTS, ISS, NISS, APACHE II, SAPS II, RAPS, REMS, GCS, MAIS, TRISS, FIM, Abdominal Trauma Index, OIS, MESS.
Cambodia	GCS, GOS.
Cameroon	ISS, TRISS, KTS, RTS, GCS, KTS II.
China	AIS, ISS, TRISS, GCS, APACHE II, NISS, Trauma Index, Prehospital Index, GOS, serum acetylcholinesterase, Exponential Injury Severity Score, Tangent ISS, FOUR score, SAPS II.
Colombia	RTS, GCS, ISS, NISS, ABCD, ABC, McLaughlin, GOS, AIS.
Croatia	GCS, GOS.
Cuba	GCS, GOS.
Egypt	GCS, APACHE II, GOS.
Ethiopia	TRISS, GCS.
Ghana	KTS II.
Guinea	GCS.
India	AIS, ISS, TRISS, KTS, RTS, GAP, MGAP, GCS, OIS, PTS, SOFA, NISS, ICISS, Ganga Hospital Score.
Indonesia	ISS, AIS, TRISS, GCS, REMS.
Iran	AIS, ISS, TRISS, RTS, GCS, APACHE II, NISS, ASCOT, Modified ISS, APACHE III, GOS-E, Abdominal Trauma Index, Simplified RTS, MESS.
Iraq	TRISS, PATI, ISS, Simplified RTS.
Jamaica	ISS, GCS.
Jordan	GCS, FIM, GOS.
Kenya	GCS, GOS, ISS, TRISS.
Lebanon	ISS, NISS.
Malawi	KTS, RTS, MGAP, GCS.
Malaysia	AIS, GOS, GCS, RTS, ISS.
Mali	GCS.
Mexico	OIS, APACHE II, ISS, PATI, AIS.
Montenegro	ISS.
Morocco	APACHE II, SAPS II.
Mozambique	RTS, ISS.
Nepal	AIS, ISS, GCS, GOS.
Niger	GCS.
Nigeria	RTS, GCS, ISS, Facial Injury Severity, AIS, PTS, GOS, MESS.

Continued

Table 2 Continued

Country	Injury measures
Pakistan	ISS, RTS, TRISS, GCS, OIS, GOS, TI.
Papua New Guinea	GCS, GOS.
Paraguay	ISS.
Rwanda	GCS, ISS, TRISS, Rwanda Mortality Probability Model.
Senegal	GCS.
Serbia	GCS, ISS, APACHE II, SOFAS, SAPS II.
South Africa	AIS, ISS, RTS, GCS, NISS, MEWS, South African Triage Score, GOS, Lactate, s-APACHE, RAPS, REMS, APACHE II, OIS.
Sri Lanka	ISS, GCS, GOS.
Suriname	ISS.
Tanzania	ISS, GCS, KTS, PTS, RTS, KTS II, OIS.
Thailand	GCS, ISS, TRISS, APACHE II, ABCD, Modified TRISS, GOS.
Trinidad	TRISS.
Tunisia	GCS, ISS, PTS, PRISM, GOS, FIM.
Turkey	ISS, AIS, RTS, TRISS, GCS, Pediatric Trauma score, Organ Specific Scores, Lactate, GOS, PATI, NISS, Turkish Penal Code.
Uganda	KTS, GCS, Lactate, KTS II.
Ukraine	GCS.
Uruguay	ISS, APACHE II, SAPS II, SOFAS, GCS.
Uzbekistan	GCS.
Vietnam	SOFA score.
Zambia	KTS, KTS II.

ABC, Assessment of Blood Consumption; ABCD, Acidosis, Blood loss, Cold, Damage; AIS, Abbreviated Injury Scale; APACHE, Acute Physiology And Chronic Health Evaluation; ASCOT, A Severity Categorization of Trauma; FIM, Functional Independence Measure; GCS, Glasgow Coma Scale; GOS, Glasgow Outcome Scale; GOS-E, GOS-Extended; ICISS, International Classification for Diseases-9 Injury Severity Score; ISS, Injury Severity Score; KTS, Kampala Trauma Score; LMICs, low-income and middle-income countries; MAIS, Maximum AIS; MESS, Mangled Extremity Score; MGAP, Mechanism, GCS, Age, Pressure; NISS, New ISS; OIS, Organ Injury Score; PATI, Penetrating Abdominal Trauma Index; PRISM, Pediatric Risk of Mortality; PTS, Pediatric Trauma Score; RAPS, Rapid Acute Physiology Score; RTS, Revised Trauma Score; SAPS, Simplified Acute Physiology Score; SOFAS, Sequential Organ Failure Assessment Score; TI, Trauma Index; TRISS, Trauma and Injury Severity Score.

factors in a single model or severity measure; therefore, use of non-injury-specific-measures such as APACHE II, SOFAS and SAPS has gained traction in trauma research. Simple yet composite measures such as MGAP and KTS have become more popular which have been widely used and validated across the globe.^{9 25–27} Our review demonstrated that, although a number of injury severity measures were developed during the 1990s and early

2000s, there have been limited applications in LMICs. Furthermore, very few validation studies were conducted in low-income settings (online supplementary file 2). Over 70% of publications on injury research in LMICs have been published from only 11 countries (figure 3) which is obviously incomparable with their burden of injuries; moreover, the body of research comprises mostly of descriptive or epidemiological studies. Comparison of the most commonly applied injury measures aligns with the most commonly validated injury severity measures, including GCS, ISS, TRISS, APACHE and KTS scores. It is important to note that the majority validation studies have been conducted in upper-middle-income countries such as China, Turkey, Brazil and Thailand; involved single centres; or included specific study population such as head or abdominal injuries. New methods and models such as EISS, TISS and new TRISS have not been validated in other LMICs, outside of their origin.

A subset of studies found relatively low performance of injury severity measures which demonstrates large deviation from studies conducted in predominantly high-income settings (eg, TRISS, ISS). These differences may be due to a wide range of factors, such as delays in recording time sensitive injury data (such as blood pressure or GCS), training of personnel administering AIS codes, limited resources and equipment available for diagnosis, missed injuries and so on. Some recent studies confirm that commonly used injury severity measures that depend on in-depth information may not perform well in mortality prediction, especially with limited or incomplete data.^{25 26} Such differences underline the importance of assessing the performance and calibration of measures in specific contexts prior to their use in trauma registries or for outcome prediction. A review of publications on validation studies demonstrated that limited statistical analysis was performed in validation studies and the issue of missing data was not addressed. This may introduce bias in the estimates of performance of the injury severity measures. As mentioned before, many of the validation studies were limited with small sample size and single institutions, restricting to the specific setting and a lack of comparison among similar institutions within the country. Very often, the validation studies did not include statistical inference of the estimation, further restricting the ability to compare performance among injury severity measures inspected. Calibration is another feature of the measure that should be more commonly assessed.

Overall, our study has been able to highlight several important issues. First, the '10–90' funding and research gap are also quite evident for injury and trauma, and we have observed that the amount of injury research from LMICs is still far less than the burden of injuries faced by these countries.²⁸ The quality and depth of research is also not sufficient, being mostly limited to small empirical studies. The findings of validation studies focusing on mortality prediction highlight large variability in performance of commonly applied injury measures including GCS, ISS, RTS, TRISS and KTS. However, lack of large

multicentre databases restricts the generalisability of results in large populations, even within a country.

The results nevertheless corroborate the assumption that no single injury measure has shown a consistent result in all settings and thus underscores the importance of context specific validation studies. This has also been reported previously from systematic reviews for injury severity measures such as ISS, NISS, ICISS and TMPM, mainly featuring studies from high-income settings.^{29 30} Furthermore, application of injury measures in field triage or emergency room disposition is also heavily influenced by the system of trauma care delivery, and hence, their performance in terms of prediction of survival, hospital length of stay or complications has to be tested and validated in specific settings where they are being used.

Our study has a few limitations. First, we conducted this literature review between 2006 and 2016, covering a 10-year period, and studies that were published outside of this timeframe are not included. Second, we have limited our literature search to three databases; nonetheless, inclusion of the Global Health database enabled us to review several Latin/South American publications that would have been otherwise missed. Third, we limited our detailed analysis of validation studies to those that focused on mortality prediction; this was due to a very limited number of studies focusing on a specific non-fatal outcome. We also did not focus on studies that used alternative coefficients for some of the established measures, as they were not consistently tested across settings.

CONCLUSION

The science of injury severity measurement has been growing to predict injury outcomes, help in decision-making and support epidemiological research. Empirical studies from upper-income and lower-middle-income countries frequently use injury severity measures. However, there is still a lack of large multicentre validation studies. The evidence base from low-income countries is even less established, where most of the burden of injury and trauma lies. No single injury severity measure has shown a consistent result in all settings and thus underscores the importance of context specific validation studies.

Author affiliations

¹Johns Hopkins International Injury Research Unit, Health Systems Program, Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

²School of Public Administration, Southwestern University of Finance and Economics, Chengdu, Sichuan, China

³Krieger School of Arts and Sciences, Johns Hopkins University, Baltimore, Maryland, USA

⁴Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

Acknowledgements We acknowledge the support of Ms Peggy Gross, Ms Monika Kochar and Mr Armaan Rowther in acquiring scientific material and providing editorial assistance.

Contributors AM, HH and YWH conceptualised the study. SA, AM, YWH and HH participated in data extraction and analysis. AM and YWH produced the first draft

of the manuscript, while AMB provided overall guidance and final review of all manuscript drafts.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval This paper is based on detailed literature review; no personal or medical information are included in this study.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement There is no other unpublished data to share.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Global Burden of Disease Study 2013 Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015;386:743–800.
- Mock C, Joshipura M, Arreola-Risa C, et al. An estimate of the number of lives that could be saved through improvements in trauma care globally. *World J Surg* 2012;36:959–63.
- Glance LG, Osler T. Beyond the major trauma outcome study: benchmarking performance using a national contemporary, population-based trauma registry. *J Trauma* 2001;51:725–7.
- Senkowski CK, McKenney MG. Trauma scoring systems: a review. *J Am Coll Surg* 1999;189:491–503.
- Li G, Baker SP. *Injury research*: Springer, 2012.
- Bouillon B, Lefering R, Vorweg M, et al. Trauma score systems: Cologne validation study. *J Trauma* 1997;42:652–8.
- MacKenzie EJ. Injury severity scales: overview and directions for future research. *Am J Emerg Med* 1984;2:537–49.
- Koehler JJ, Baer LJ, Malafa SA, et al. Prehospital Index: a scoring system for field triage of trauma victims. *Ann Emerg Med* 1986;15:178–82.
- Mowafi H, Oranmore-brown R, Cerwensky K, et al. Assessment of the revised kampala trauma score (ktsii) to predict mortality, need for admission, and use of hospital resources at university teaching hospital in lusaka, zambia. *Acad Emerg Med* 2013;20:S322.
- Champion HR. Trauma scoring. *Scand J Surg* 2002;91:12–22.
- Huber-Wagner S, Stegmaier J, Mathonia P, et al. The sequential trauma score - a new instrument for the sequential mortality prediction in major trauma. *Eur J Med Res* 2010;15:185–95.
- Antonelli M, Moreno R, Vincent JL, et al. Application of SOFA score to trauma patients. Sequential organ failure assessment. *Intensive Care Med* 1999;25:389–94.
- Aslar AK, Kuzu MA, Elhan AH, et al. Admission lactate level and the APACHE II score are the most useful predictors of prognosis following torso trauma. *Injury* 2004;35:746–52.
- Imhoff BF, Thompson NJ, Hastings MA, et al. Rapid Emergency Medicine Score (REMS) in the trauma population: a retrospective study. *BMJ Open* 2014;4:004738.
- Maslanka AM. Scoring systems and triage from the field. *Emerg Med Clin North Am* 1993;11:15–27.
- O'Reilly GM, Joshipura M, Cameron PA, et al. Trauma registries in developing countries: a review of the published experience. *Injury* 2013;44:713–21.
- Fryback D, Prokof C, Gustafson D, et al, 1980. The Wisconsin Trauma Index. *Trauma Index Severity Conference* ("The Woodstock Conference"), sponsored by the National Center for Health Services Research Grant No. HS-04149-01, the American Trauma Society, and the University of Wisconsin 1980 Jul
- Knaus WA, Zimmerman JE, Wagner DP, et al. APACHE-acute physiology and chronic health evaluation: a physiologically based classification system. *Crit Care Med* 1981;9:591–7.
- Petrucelli E, States JD, Hames LN. The abbreviated injury scale: Evolution, usage and future adaptability. *Accident Analysis & Prevention* 1981;13:29–35.
- Jat AA, Khan MR, Zafar H, et al. Peer review audit of trauma deaths in a developing country. *Asian J Surg* 2004;27:58–64.
- Lefering R. Development and validation of the revised injury severity classification score for severely injured patients. *Eur J Trauma Emerg Surg* 2009;35:437–47.
- Schluter PJ. Trauma and Injury Severity Score (TRISS): is it time for variable re-categorisations and re-characterisations? *Injury* 2011;42:83–9.
- Boyd CR, Tolson MA, Copes WS. Evaluating trauma care: the TRISS method. Trauma score and the injury severity score. *J Trauma* 1987;27:370–8.
- Khajanchi MU, Kumar V, Gerdin M, et al. Indians fit the Asian trauma model. *World J Surg* 2013;37:705–6.
- Roy N, Gerdin M, Schneider E, et al. Validation of international trauma scoring systems in urban trauma centres in India. *Injury* 2016;47:2459–64.
- Hung YW, He H, Mehmood A, et al. Exploring injury severity measures and in-hospital mortality: A multi-hospital study in Kenya. *Injury* 2017;48:2112–8.
- Weeks SR, Juillard CJ, Monono ME, et al. Is the Kampala trauma score an effective predictor of mortality in low-resource settings? A comparison of multiple trauma severity scores. *World J Surg* 2014;38:1905–11.
- Vidyasagar D. Global notes: the 10/90 gap disparities in global health research. *J Perinatol* 2006;26:55–6.
- Tohira H, Jacobs I, Mountain D, et al. Systematic review of predictive performance of injury severity scoring tools. *Scand J Trauma Resusc Emerg Med* 2012;20:63.
- Gagné M, Moore L, Beaudoin C, et al. Performance of International Classification of Diseases-based injury severity measures used to predict in-hospital mortality: A systematic review and meta-analysis. *J Trauma Acute Care Surg* 2016;80:419–26.
- Keller WK, Dillihunt RC, Fenner HA, et al. Rating the severity of tissue damage. I. The abbreviated scale. *JAMA* 1971;215:277–80.
- Keller WK, Fenner HA, Jolley FL, et al. Rating the severity of tissue damage. II. The comprehensive scale. *JAMA* 1972;220:717–20.
- Baker SP, O'Neill B, Haddon W, et al. The injury severity score: a method for describing patients with multiple injuries and evaluating emergency care. *J Trauma* 1974;14:187–96.
- Krischer JP. Measuring trauma severity: the ESP index. *Health Serv Res* 1978;13:61–5.
- Moore EE, Cogbill TH, Malangoni MA, et al. Organ injury scaling, II: Pancreas, duodenum, small bowel, colon, and rectum. *J Trauma* 1990;30:1427–9.
- Champion HR, Sacco WJ, Lepper RL, et al. An anatomic index of injury severity. *J Trauma* 1980;20:197–202.
- Levy PS, Goldberg J, Rothrock J. The revised estimated survival probability index of trauma severity. *Public Health Rep* 1982;97:452–9.
- Somers RL. The probability of death score: An improvement of the injury severity score. *Accident Analysis & Prevention* 1983;15:247–57.
- Moore EE, Cogbill TH, Malangoni MA, et al. Organ injury scaling, II: Pancreas, duodenum, small bowel, colon, and rectum. *J Trauma* 1990;30:1427–9.
- Moore EE, Shackford SR, Pachter HL, et al. Organ injury scaling: spleen, liver, and kidney. *J Trauma* 1989;29:1664–6.
- Moore EE, Cogbill TH, Jurkovich GJ, et al. Organ injury scaling. III: Chest wall, abdominal vascular, ureter, bladder, and urethra. *J Trauma* 1992;33:337–9.
- Moore EE, Malangoni MA, Cogbill TH, et al. Organ injury scaling. IV: Thoracic vascular, lung, cardiac, and diaphragm. *J Trauma* 1994;36:299–300.
- Champion HR, Copes WS, Sacco WJ, et al. A new characterization of injury severity. *J Trauma* 1990;30:539–46.
- McGonigal MD, Cole J, Schwab CW, et al. A new approach to probability of survival scoring for trauma quality assurance. *J Trauma* 1993;34:863–70.
- Osler T, Rutledge R, Deis J, et al. ICISS: an international classification of disease-9 based injury severity score. *J Trauma* 1996;41:8.
- Osler T, Baker SP, Long W. A modification of the injury severity score that both improves accuracy and simplifies scoring. *J Trauma* 1997;43:922–6.
- Meredith JW, Evans G, Kilgo PD, et al. A comparison of the abilities of nine scoring algorithms in predicting mortality. *J Trauma* 2002;53:621–9.
- Kilgo PD, Osler TM, Meredith W. The worst injury predicts mortality outcome the best: rethinking the role of multiple injuries in trauma outcome scoring. *J Trauma* 2003;55:599–607.
- Günay Y, Yavuz MF, Eşiyok B. Comparison of Turkish Injury Scale (TIS) with the Abbreviated Injury Scale (AIS). *Forensic Sci Int* 2003;132:1–4.

50. Kirkpatrick JR, Youmans RL. Trauma index. An aide in the evaluation of injury victims. *J Trauma* 1971;11:711–4.
51. Teasdale G, Jennett B. Assessment of coma and impaired consciousness. A practical scale. *Lancet* 1974;2:81–4.
52. Bever DL, Veenker CH. An illness-injury severity index for nonphysician emergency medical personnel. *Emt J* 1979;3:45–9.
53. Sacco WJ, Champion HR, Carnazzo AJ, et al. Trauma score. In: AnonymouIn. *Crit Care Med* 1981:672–6.
54. Gormican SP. CRAMS scale: field triage of trauma victims. *Ann Emerg Med* 1982;11:132–5.
55. Rhee KJ, Fisher CJ, Willitis NH. The Rapid Acute Physiology Score. *Am J Emerg Med* 1987;5:278–82.
56. Champion HR, Sacco WJ, Copes WS, et al. A revision of the Trauma Score. *J Trauma* 1989;29:623–9.
57. Kobusingye OC, Lett RR. Hospital-based trauma registries in Uganda. *J Trauma* 2000;48:498–502.
58. Wijdicks EF, Bamlet WR, Maramattom BV, et al. Validation of a new coma scale: The FOUR score. *Ann Neurol* 2005;58:585–93.
59. Yücel N, Lefering R, Maegele M, et al. Trauma Associated Severe Hemorrhage (TASH)-Score: probability of mass transfusion as surrogate for life threatening hemorrhage after multiple trauma. *J Trauma* 2006;60:7.
60. Abib SC, Schettini ST, Figueiredo LF. Prehospital pediatric trauma classification (PHPTC) as a tool for optimizing trauma care resources in the city of São Paulo, Brazil. *Acta Cir Bras* 2006;21:7–11.
61. Rajasekaran S, Naresh Babu J, Dheenadhayalan J, et al. A score for predicting salvage and outcome in Gustilo type-IIIa and type-IIIb open tibial fractures. *J Bone Joint Surg Br* 2006;88:1351–60.
62. Nunez TC, Voskresensky IV, Dossett LA, et al. Early prediction of massive transfusion in trauma: simple as ABC (assessment of blood consumption)? *J Trauma* 2009;66:346–52.
63. Raum MR, Nijsten MW, Vogelzang M, et al. Emergency trauma score: an instrument for early estimation of trauma severity. *Crit Care Med* 2009;37:1972–7.
64. Ordoñez CA, Badiel M, Pino LF, et al. Damage control resuscitation: early decision strategies in abdominal gunshot wounds using an easy "ABCD" mnemonic. *J Trauma Acute Care Surg* 2012;73:1074–8.
65. Jennett B, Bond M. Assessment of outcome after severe brain damage. *Lancet* 1975;1:480–4.
66. Moore EE, Dunn EL, Moore JB, et al. Penetrating abdominal trauma index. *J Trauma* 1981;21:439–45.
67. Le Gall JR, Loirat P, Alperovitch A, et al. A simplified acute physiology score for ICU patients. *Crit Care Med* 1984;12:975–7.
68. Knaus WA, Draper EA, Wagner DP, et al. APACHE II: a severity of disease classification system. *Crit Care Med* 1985;13:818–29.
69. Pollack MM, Ruttimann UE, Getson PR. Pediatric risk of mortality (PRISM) score. *Crit Care Med* 1988;16:1110–6.
70. Johansen K, Daines M, Howey T, et al. Objective criteria accurately predict amputation following lower extremity trauma. *J Trauma* 1990;30:568–73.
71. Knaus WA, Wagner DP, Draper EA, et al. The APACHE III prognostic system. Risk prediction of hospital mortality for critically ill hospitalized adults. *Chest* 1991;100:1619–36.
72. Rady MY, Rivers EP, Martin GB, et al. Continuous central venous oximetry and shock index in the emergency department: use in the evaluation of clinical shock. *Am J Emerg Med* 1992;10:538–41.
73. Rixen D, Raum M, Bouillon B, et al. Base deficit development and its prognostic significance in posttrauma critical illness: an analysis by the trauma registry of the Deutsche Gesellschaft für Unfallchirurgie. *Shock* 2001;15:83–9.
74. Nell V, Yates DW, Kruger J. An extended Glasgow Coma Scale (GCS-E) with enhanced sensitivity to mild brain injury. *Arch Phys Med Rehabil* 2000;81:614–7.
75. Mutooro SM, Mutakooha E, Kyamanywa P. A comparison of Kampala trauma score II with the new injury severity score in Mbarara University Teaching Hospital in Uganda. *East and Central African Journal of Surgery* 2010;15:62–71.
76. Olsson T, Terent A, Lind L. Rapid Emergency Medicine score: a new prognostic tool for in-hospital mortality in nonsurgical emergency department patients. *J Intern Med* 2004;255:579–87.
77. Gomez M, Wong DT, Stewart TE, et al. The FLAMES score accurately predicts mortality risk in burn patients. *J Trauma* 2008;65:636–45.
78. Gance LG, Osler TM, Mukamel DB, et al. TMPM-ICD9: a trauma mortality prediction model based on ICD-9-CM codes. *Ann Surg* 2009;249:1032–9.
79. Sartorius D, Le Manach Y, David JS, et al. Mechanism, glasgow coma scale, age, and arterial pressure (MGAP): a new simple prehospital triage score to predict mortality in trauma patients. *Crit Care Med* 2010;38:831–7.
80. Kondo Y, Abe T, Kohshi K, et al. Revised trauma scoring system to predict in-hospital mortality in the emergency department: Glasgow coma scale, age, and systolic blood pressure score. *Crit Care* 2011;15:R191.
81. Jones JM, Skaga NO, Søvik S, et al. Norwegian survival prediction model in trauma: modelling effects of anatomic injury, acute physiology, age, and co-morbidity. *Acta Anaesthesiol Scand* 2014;58:303–15.
82. Wang MD, Fan WH, Qiu WS, et al. The exponential function transforms the Abbreviated Injury Scale, which both improves accuracy and simplifies scoring. *Eur J Trauma Emerg Surg* 2014;40:287–94.
83. Wang M, Qiu W, Qiu F, et al. Tangent function transformation of the abbreviated injury scale improves accuracy and simplifies scoring. *Arch Med Sci* 2015;11:130–6.