CASE REPORT

Severe Thrombocytopenia in COVID-19: A Conundrum in Dengue-endemic Areas

MB Adarsh¹, Anitha Abraham², P Kavitha³, Meera M Nandakumar⁴, Raman S Vaman⁵

ABSTRACT

Thrombocytopenia in coronavirus disease-2019 (COVID-19) can be attributed to multiple factors. Most often it is disease related. It is usually mild and if severe often associated with severe COVID-19 disease. It can also be due to drugs (Remdesivir, Tocilizumab) or coinfection with other viruses. Here we report two cases of severe thrombocytopenia in COVID-19 due to dengue coinfection. Most often the thrombocytopenia in dengue is self-resolving, and a careful "wait and watch" should suffice unlike COVID-19, where steroids can help if the cytopenia is due to cytokine storm or immune-mediated effects.

Keywords: COVID-19, Dengue, Thrombocytopenia.

Indian Journal of Critical Care Medicine (2021): 10.5005/jp-journals-10071-23778

Introduction

Thrombocytopenia in coronavirus disease-2019 (COVID-19) can be attributed to multiple factors. Most often it is disease related. It can be due to the reduced production of platelets from bone marrow because of cytokine storm or direct viral infiltration of the marrow and due to the increased peripheral destruction from immunemediated effects or excessive peripheral consumption as a result of microthrombi formation.¹ Thrombocytopenia in COVID-19 is usually mild and if severe it is often associated with severe COVID-19 disease. The mean platelet count varied from 164×10^3 /mm³ to 196×10^3 /mm³ among various cohorts of COVID-19 cases. Severe life-threatening thrombocytopenia was not seen in these cohorts. There are many isolated case reports of severe thrombocytopenia in COVID-19 and were mostly attributed to immune-mediated peripheral destruction.^{3–5} It can also be due to drugs (Remdesivir, Tocilizumab) or coinfection with other viruses. Dengue is a mosquitoborne viral illness that is endemic in many countries and its incidence is worrisome in our area during this pandemic.⁶ Here we report two cases of severe thrombocytopenia in COVID-19 due to dengue coinfection. Most often the thrombocytopenia in dengue is selfresolving, and a careful "wait and watch" should suffice unlike COVID-19, where steroids can help if the cytopenia is due to cytokine storm or immune-mediated effects.

Case Description

Two patients presented with dengue coinfection. First was a 34-year male who presented with fever and headache for one day. His oropharyngeal swab for SARS-CoV-2 PCR was positive. His investigations showed a total leucocyte count of 2.9×10^9 /L with 27% lymphocytes and a platelet count of 53×10^9 /L. The second case was a 54-year male with fever and severe body ache for 3 days. His oropharyngeal swab for SARS-CoV-2 PCR was positive. His total leucocyte count was 4.8×10^9 /L with 40% lymphocytes and platelet count was 20×10^9 /L. There were no bleeding manifestations, and both were hemodynamically stable. A rapid ELISA for dengue NS1 antigen was positive in both cases. First case was managed symptomatically with paracetamol and oral fluids. His platelet count normalized in 5 days. The second case

¹Department of General Medicine, Government Medical College, Kasaragod, Kerala, India

²Department of Community Medicine, Government Medical College, Kasaragod, Kerala, India

³Department of Anaesthesia, Government Medical College, Kasaragod, Kerala, India

⁴Department of ENT, Government Medical College, Kasaragod, Kerala, India

⁵Medical Superintendent, Government Medical College, Kasaragod, Kerala, India

Corresponding Author: MB Adarsh, Department of General Medicine, Government Medical College, Kasaragod, Kerala, India, e-mail: adarshmb@gmail.com

How to cite this article: Adarsh MB, Abraham A, Kavitha P, Nandakumar MM, Vaman RS. Severe Thrombocytopenia in COVID-19: A Conundrum in Dengue-endemic Areas. Indian J Crit Care Med 2021;25(4):465–466.

Source of support: Nil
Conflict of interest: None

was given dexamethasone 6 mg for 5 days, and his platelet count normalized in 6 days. Both had no other complications of dengue or COVID-19. The repeat nasopharyngeal swab for rapid antigen test SARS-CoV-2 was negative on the 10th day in both cases and were discharged.

Thrombocytopenia often alerts the physician in COVID-19. Dengue coinfection can be a cause for it. This can cause severe thrombocytopenia and can be life-threatening. Early identification of coinfection can help the physician in anticipating other dengue-related complications like dengue shock syndrome and dengue hemorrhagic fever, which otherwise would have been attributed to severe COVID-19 and the treatment would have been initiated as for severe COVID-19. Few things can point to non-COVID reasons for severe thrombocytopenia in COVID-19 cases. Fever, headache, body ache, and severe fatigue are common to both the viral illnesses. Unlike dengue, skin manifestations are rare in COVID-19.

[©] Jaypee Brothers Medical Publishers. 2021 Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (https://creativecommons.org/licenses/by-nc/4.0/), which permits unrestricted use, distribution, and non-commercial reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

Dengue fever is characterized by maculopapular and diffuse erythematous rash in more than 50% of patients.8 Presence of rash with thrombocytopenia in COVID-19 should alert a physician in this regard. Also, disease-related thrombocytopenia in COVID-19 is often associated with severe disease. If the patient is asymptomatic or mildly symptomatic for COVID-19, other causes like coinfection with dengue as in the index cases should be thought of for severe thrombocytopenia. In both the index, patients were only mildly symptomatic for COVID-19. Despite the dengue coinfection, COVID-19 did not worsen and there was no persistence of antigen positivity. This may be due to the enhanced nonspecific antiviral response elicited by the coinfection. This possible synergistic effect needs to be studied further. But this is not always the case. Few deaths have been reported due to the coinfection. Making the diagnosis of coinfection is also difficult as false-positive serological tests for dengue have been seen in COVID-19 patients. 10 So, we will have to go with antigen-based tests for a proper diagnosis.

In conclusion, dengue coinfection can cause severe thrombocytopenia in COVID-19 and identifying it early can help in the better management of patients.

HIGHLIGHTS

- Dengue coinfection can cause severe thrombocytopenia in COVID-19.
- Dengue–COVID-19 coinfection and severe thrombocytopenia had good outcome at discharge.

ORCID

Meera M Nandakumar © 0000-0003-4733-9135

Raman Swathy Vaman 6 0000-0003-0709-4110

REFERENCES

- Xu P, Zhou Q, Xu J. Mechanism of thrombocytopenia in COVID-19 patients. Ann Hematol 2020;99(6):1205–1258. DOI: 10.1007/s00277-020-04019-0.
- Lippi G, Plebani M, Henry BM. Thrombocytopenia is associated with severe coronavirus disease 2019 (COVID-19) infections: a meta-analysis. Clin Chim Acta 2020;506:145–148. DOI: 10.1016/j. cca.2020.03.022.
- Sadr S, Seyed Alinaghi S, Ghiasvand F, Hassan Nezhad M, Javadian N, Hossienzade R, et al. Isolated severe thrombocytopenia in a patient with COVID-19: a case report. IDCases 2020;21:e00820. DOI: 10.1016/j. idcr.2020.e00820.
- Chen W, Yang B, Li Z, Wang P, Chen Y, Zhou H. Sudden severe thrombocytopenia in a patient in the recovery stage of COVID-19. Lancet Haematol 2020;7(8):e624. DOI: 10.1016/S2352-3026(20)30175-7.
- Malik J, Javaid M, Majedi O, Ishaq U, Zahid T. Paying in blood: a case of thrombocytopenia in Covid-19. Cureus 2020;12(8):e9791. DOI: 10.7759/cureus.9791.
- Varghese B, Kumaran JA. Trend of dengue fever in COVID-19 period: observation from Kerala, India. Int J Commun Med Publ Health 2020;7(8):3307. DOI: 10.18203/2394-6040.ijcmph20203419.
- Wollina U, Karadağ AS, Rowland-Payne C, Chiriac A, Lotti T. Cutaneous signs in COVID -19 patients: a review. Dermatol Ther 2020;33(5):e13549. DOI: 10.1111/dth.13549.
- Thomas E, John M, Kanish B. Mucocutaneous manifestations of dengue fever. Indian J Dermatol 2010;55(1):79. DOI: 10.4103/0019-5154.60359.
- Miah MDA, Husna A. Coinfection, coepidemics of COVID-19, and dengue in dengue-endemic countries: a serious health concern. J Med Virol 2020. Online ahead of print. DOI: 10.1002/jmv.26269.
- Yan G, Lee CK, Lam LTM, Yan B, Chua YX, Lim AYN, et al. Covert COVID-19 and false-positive dengue serology in Singapore. Lancet Infect Dis 2020;20(5):536. DOI: 10.1016/S1473-3099(20)30158-4.

