

RESEARCH ARTICLE

Low Completeness of Bacteraemia Registration in the Danish National Patient Registry

Kim Oren Gradel^{1,2*}, Stig Lønberg Nielsen^{1,3}, Court Pedersen³, Jenny Dahl Knudsen⁴, Christian Østergaard⁴, Magnus Arpi⁵, Thøger Gorm Jensen⁶, Hans Jørn Kolmos⁶, Mette Søgaard⁷, Annmarie Touborg Lassen⁸, Henrik Carl Schønheyder^{9,10}, Danish Collaborative Bacteraemia Network (DACOBAN)[†], Danish Observational Registry of Infectious Syndromes (DORIS)[†]

1 Center for Clinical Epidemiology, South, Odense University Hospital, Odense, Denmark, **2** Research Unit of Clinical Epidemiology, Institute of Clinical Research, University of Southern Denmark, Odense, Denmark, **3** Department of Infectious Diseases, Odense University Hospital, Odense, Denmark, **4** Department of Clinical Microbiology, Copenhagen University Hospital, Hvidovre Hospital, Hvidovre, Denmark, **5** Department of Clinical Microbiology, Copenhagen University Hospital, Herlev Hospital, Herlev, Denmark, **6** Department of Clinical Microbiology, Odense University Hospital, Odense, Denmark, **7** Department of Clinical Epidemiology, Institute of Clinical Medicine, Aarhus University Hospital, Aarhus, Denmark, **8** Department of Emergency Medicine, Odense University Hospital, Odense, Denmark, **9** Department of Clinical Microbiology, Aalborg University Hospital, Aalborg, Denmark, **10** Department of Clinical Medicine, Aalborg University, Aalborg Denmark

^{††} The complete membership of the author group can be found in the Acknowledgments.

* kim.gradel@rsyd.dk



CrossMark
click for updates

OPEN ACCESS

Citation: Gradel KO, Nielsen SL, Pedersen C, Knudsen JD, Østergaard C, Arpi M, et al. (2015) Low Completeness of Bacteraemia Registration in the Danish National Patient Registry. PLoS ONE 10(6): e0131682. doi:10.1371/journal.pone.0131682

Editor: Delmiro Fernandez-Reyes, University College London, UNITED KINGDOM

Received: February 11, 2015

Accepted: June 4, 2015

Published: June 29, 2015

Copyright: © 2015 Gradel et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data from the Danish National Hospital Registry and the Civil Registration System are available from the Danish National Health and Medicines Authority for researchers who meet the criteria for access to confidential data. Researchers who meet these criteria can request the microbiological core data set from the corresponding author who will handle further legal procedures.

Funding: The authors have no support or funding to report.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Bacteraemia is associated with significant morbidity and mortality and timely access to reliable information is essential for health care administrators. Therefore, we investigated the completeness of bacteraemia registration in the Danish National Patient Registry (DNPR) containing hospital discharge diagnoses and surgical procedures for all non-psychiatric patients. As gold standard we identified bacteraemia patients in three defined areas of Denmark (~2.3 million inhabitants) from 2000 through 2011 by use of blood culture data retrieved from electronic microbiology databases. Diagnoses coded according to the International Classification of Diseases, version 10, and surgical procedure codes were retrieved from the DNPR. The codes were categorized into seven groups, ranked a priori according to the likelihood of bacteraemia. Completeness was analysed by contingency tables, for all patients and subgroups. We identified 58,139 bacteraemic episodes in 48,450 patients; 37,740 episodes (64.9%) were covered by one or more discharge diagnoses within the seven diagnosis/surgery groups and 18,786 episodes (32.3%) had a code within the highest priority group. Completeness varied substantially according to speciality (from 17.9% for surgical to 36.4% for medical), place of acquisition (from 26.0% for nosocomial to 36.2% for community), and microorganism (from 19.5% for anaerobic Gram-negative bacteria to 36.8% for haemolytic streptococci). The completeness increased from 25.1% in 2000 to 35.1% in 2011. In conclusion, one third of the bacteraemic episodes did not have a relevant diagnosis in the Danish administrative registry recording all non-psychiatric

contacts. This source of information should be used cautiously to identify patients with bacteraemia.

Introduction

Bacteraemia, with its 30-day mortality of 15–30%, is estimated to be one of the top seven causes of death in the developed countries [1]. Bacteraemia is defined as the presence of viable bacteria or fungi in blood cultures (BCs) [2–4]. In contrast, the term sepsis defines a clinical condition reflecting the host response to infection [5]. However, the two terms are often used interchangeably, which may create some confusion.

Regardless of this, both conditions constitute a severe clinical burden which may favour their easy identification and surveillance through administrative hospital discharge registries. Nevertheless, studies have generally indicated high variation and low completeness in administrative registries for both sepsis [6–8] and bacteraemia [9–14].

The Danish National Patient Registry (DNPR) contains hospital discharge diagnoses and surgical procedures from all non-psychiatric inpatients since 1977 [15]. A prior Danish study found that only 18 of 406 (4.4%) bacteraemic episodes in 1994 were recorded with a relevant bacteraemia or sepsis diagnosis in the DNPR [9]. This may have been due to the newly implemented International Classification of Diseases, version 10 (ICD-10) codes in 1994, and the completeness may have improved since then, possibly also encouraged by the Survival Sepsis Campaign [16]. We therefore investigated the completeness of sepsis/bacteraemia ICD-10 codes in the DNPR, using bacteraemic episodes from 2000 through 2011 derived from positive BC data as gold standard. To assess the impact of completeness on prognostic models we applied multivariate regression analyses according to whether the bacteraemic episodes were recorded in the DNPR, using 30-day mortality as outcome.

Materials and Methods

Setting

The Danish healthcare system is tax financed and provides care free of charge for all residents. The admission of all acutely ill patients to the nearest public hospital in their area of residence prompts a population-based coverage. Our data covered three geographically well-defined areas (North Denmark Region, Capital Region, Funen County) served by four Departments of Clinical Microbiology (DCMs) in hospitals in Aalborg, Herlev, Hvidovre, and Odense (in total 2.3 million inhabitants [17, 18]). BC procedures have been described previously [19–21].

Data linkage

All Danish residents have a unique personal identification number used for all health contacts, which permits unambiguous linkage between health administrative registries [22].

Core dataset

The study periods were 2000–2011 (North Denmark and Capital regions) or 2000–2008 (Funen).

All microbiological results were recorded in an electronic laboratory information system (Aalborg, Herlev, and Hvidovre: ADBakt [Autonik, Sködinge, Sweden]; Odense: the local Patient Administrative System in 2000–2005 and the MADS system [www.madsonline.dk]

thereafter). Key data included dates of draw and receipt of the BC in the DCM, and BC isolates. We retrieved data on all positive BCs and used previously published computer algorithms to exclude likely contaminants and to derive bacteraemic episodes [21, 23]. For each episode we defined the best-estimate baseline date as the date of draw; for bacteraemic episodes with a missing date of draw (9.3%) we used the never-missing date of receipt. We used previously reported computer algorithms to derive incident and non-incident episodes as well as acquisition (community-acquired, healthcare-associated, nosocomial) [21].

Code systems used in the Danish National Hospital Registry

In Denmark, all diagnostic and surgical procedure codes are allocated by physicians when patients are discharged.

Until 1994, diagnoses were coded according to the International Classification of Diseases, version 8 (ICD-8), and thereafter according to the ICD-10, as ICD-9 was never implemented in Denmark [15]. We used the Danish ICD-10 version [24], derived from the WHO classification, vs. 2010 [25] with amendments that more specifically designate bacteraemia (e.g. A49.9A [Bacteraemia, unspecified] found in the Danish, but not in the WHO, version). For each hospitalization, one obligatory principal diagnosis may be supplemented with up to 20 secondary diagnoses.

For surgical procedures, the Nordic Classification of Surgical Procedures [26] (NOMESCO) has been in use since 1996.

Retrieval of diagnoses and surgical procedures related to sepsis or bacteraemia

Two authors (HCS, SLN) independently retrieved codes for diagnoses and surgical procedures that may indicate the presence of bacteraemia, either directly by codes using the term bacteraemia or septicaemia or indirectly by codes indicating focal infections. The authors' codes were combined and consensus was reached with agreement on all included codes, shown in the Appendix.

Linkage to the Danish National Hospital Registry

We linked the core dataset to their DNPR inpatient data and retrieved the date of hospital admission from home which was closest to and equal to or earlier than the best-estimate baseline date. Likewise, we retrieved the date of discharge to home which was closest to and equal to or later than the best-estimate baseline date. For this hospitalization, which covered the bacteraemic episode, we retrieved all relevant diagnosis and surgical procedure codes (see [Appendix](#)).

We then linked the core dataset to the DNPR to retrieve all first-time diagnoses in the Charlson comorbidity index [27] within a 6-year period prior to the best-estimate baseline date. In this index, 19 major disease categories (e.g., malignancy, cardiovascular diseases, and diabetes mellitus) are assigned a score, with higher scores given to prognostically more severe diseases.

Linkage to the Danish Civil Registration System

To obtain mortality data we linked the study population to the Danish Civil Registration System, which comprises daily updated data on the patients' vital status, as well as date of death, disappearance, or emigration, if relevant [28].

Statistical analyses

We categorized the codes for diagnoses and surgical procedures into seven groups and determined the following priority list for the likelihood of representing bacteraemia: 1) infections/bacteraemia; 2) other diagnoses/bacteraemia; 3) other diagnoses/focal infection; 4) surgical procedures/focal infection; 5) infections/systemic infection; 6) other diagnoses/systemic infection; 7) infections/focal infection (see [Appendix](#) for the specific codes and text examples for the three most common codes within each group). If group 1 occurred in the hospitalization comprising the bacteraemic episode, groups 2–7 were annulled. If group 1 did not occur we retrieved group 2 and annulled groups 3–7. If group 2 did not occur we proceeded to group 3, etc. (see [Appendix](#) for examples for two specific bacteraemic episodes).

We used the same seven groups with a different prioritization ($5 > 7 > 3 > 4 > 6 > 1 > 2$) to derive diagnoses/procedures that may indicate the presence of a focal infection, applying the same principles as for bacteraemia (if 5 occurred, 1–4 and 6–7 were annulled, if 5 did not occur we proceeded to 7, etc.). See [Appendix](#) for examples of two specific bacteraemic episodes.

We computed contingency tables for basic patient characteristics in relation to occurrence of all seven groups, group 1 only (most likely bacteraemia diagnosis), and the combination of groups 5, 7, 3, 4, and 6 (most likely diagnosis of or surgery for a focal infection). As basic patient characteristics we selected gender, age group (0–15, 16–64, 65–80, >80 years), Charlson comorbidity index (score 0, 1–2, >2), speciality (defining episodes as either medical, surgical, intensive care unit [ICU], paediatric, or unknown), acquisition of bacteraemia (community, healthcare-associated, nosocomial), main group of microorganisms (*Escherichia coli*, *Enterobacter* spp., *Klebsiella* spp., other Enterobacteriaceae, *Pseudomonas aeruginosa*, anaerobic Gram-negative bacteria, other Gram-negative bacteria, *Staphylococcus aureus*, coagulase-negative staphylococci [CNS], *Streptococcus pneumoniae*, haemolytic streptococci, enterococci, Gram-positive rods, other Gram-positive bacteria, fungi, polymicrobial, undetermined [0.4%]), incident vs. non-incident episodes, 30-day mortality, and, for 2,761 bacteraemic episodes (4.7%), sepsis groups (no sepsis, possibly sepsis (due to missing data), sepsis, severe sepsis/septic shock, and organ dysfunction without sepsis) [18, 20]. We used the chi-square test to assess whether characteristics differed between patients recorded with group 1–7 codes vs. no codes belonging to these groups, for group 1 codes (believed most likely to represent bacteraemia) vs. no codes belonging to this group, and for group 3–7 codes (believed most likely to represent a focal infection) vs. no codes belonging to these groups.

To assess possible time-related aspects, we depicted a histogram with the proportions of groups 1 and 3–7 on the y-axis and calendar year (2000–2011) on the x-axis.

Finally, we used logistic regression analysis to compute odds ratios (ORs) with 95% confidence intervals (CIs) for 30-day mortality, a commonly used outcome in prognostic bacteraemia studies. We adjusted for the above basic patient characteristics except sepsis groups (due to missing data). The analyses covered all bacteraemic episodes as well as subgroup analyses for groups 1–7, group 1, and groups 3–7 during 2000–2008 as data on speciality were incomplete as from 2009.

The program Stata (release 13; StataCorp) was used for all analyses.

Ethical considerations

The study was approved by the Danish Data Protection Agency (record nos. 2007-41-0627, 2013-41-2579). Approval by an ethics committee or consent from participants (including next of kin/caregiver in the case of children) are not required for registry-based research in Denmark. Data were not anonymized prior to analysis.

Table 1. Bacteraemic episodes according to groups of diagnosis/surgical procedure codes that indicate hospitalization with bacteraemia or a focal infection.

Group	No. (%) of bacteraemic episodes	Cumulative no. (%)
1) Infections/bacteraemia	18,786 (32.3)	18,786 (32.3)
2) Other diagnoses/bacteraemia	457 (0.8)	19,243 (33.1)
3) Other diagnoses/focal infection	15,110 (26.0)	34,353 (59.1)
4) Surgical procedures/focal infection	499 (0.9)	34,852 (60.0)
5) Infections/systemic infection	2,027 (3.5)	36,879 (63.4)
6) Other diagnoses/systemic infection	2 (0)	36,881 (63.4)
7) Infections/focal infection	859 (1.5)	37,740 (64.9)
None of the groups 1–7	20,399 (35.1)	58,139 (100)

doi:10.1371/journal.pone.0131682.t001

Results

We identified 58,139 bacteraemic episodes in 48,450 patients of whom 41,633 patients (85.9%) had 1 episode, 5,062 (10.5%) had 2, 1,186 (2.5%) had 3, 561 (1.2%) had 4–10, and eight had 11–16 episodes.

Groups of diagnoses and surgical procedures

Among the 58,139 bacteraemic episodes, 37,740 (64.9%) were related to a hospitalization with one or more of the seven diagnoses/surgery groups we defined as indicative of bacteraemia or a focal infection (Table 1). Among these, 18,786 episodes (32.3%) had a group 1 code (an “infection/bacteraemia” diagnosis, i.e., the highest priority codes representing bacteraemia) with a total of 20,433 of such codes (Table 2). One such code was given to 17,309 episodes (92.1%), two codes to 1,336 episodes (7.1%) and 2–5 codes to the remaining 181 episodes (1.0%). 26,538 episodes (45.7%) had a group 3–7 code indicating a likely focal infection.

For bacteraemia, an increase was detected from 2000 (25.1%) to 2005 (35.2%) after which the completeness varied between 30% and 35% (Fig 1). For infectious foci, an increase was seen from 37.7% in 2000 to 45.8% in 2005 and further from 44.0% in 2007 to 55.1% in 2011.

Characteristics of patients with codes representing bacteraemia or a focal infection

The proportion of bacteraemic episodes that had been assigned a diagnosis/surgery group indicating bacteraemia or a focal infection (64.9%, Table 3) differed within all subgroups ($p < 10^{-4}$) except sepsis groups ($p = 0.06$). Among age groups, the highest completeness was seen for the youngest (0–14 y: 70.7%) and the oldest (>80 y: 68.6%). A higher completeness was seen for females (66.6%) than for males (63.5%), patients without recorded comorbidity (71.1%), paediatric ward patients (71.8%), *E. coli* (73.4%), haemolytic streptococci (76.1%), non-incident bacteraemic episodes (67.2%), and no mortality within 30 days (67.7%). Pertaining to acquisition the completeness declined considerably from community (75.1%), over healthcare-associated (63.1%) to nosocomial (53.4%). For sepsis groups, the lowest completeness (61.9%) was seen for the no sepsis group, whereas higher completeness (70–71%) was encountered for the severer groups (sepsis, severe sepsis/septic shock, organ dysfunction without sepsis).

Characteristics of patients with the most likely bacteraemia codes

The proportion of bacteraemic episodes with a group 1 code (32.3%, Table 3) differed between all subgroups ($p < 10^{-4}$) except for gender ($p = 0.24$). The completeness increased with higher

Table 2. Group 1 (“Infections / bacteraemia”) codes, which most likely represent bacteraemia, given to 18,786 bacteraemic episodes.

Code	Text	Number (%)
DA02.1	<i>Salmonella</i> sepsis	204 (1.0)
DA28.2B	Extraintestinal yersiniosis	1 (0)
DA32.7	Listerial sepsis	43 (0.2)
DA39.2	Acute meningococcaemia	83 (0.4)
DA39.2A	Meningococcal sepsis	20 (0.1)
DA39.3	Chronic meningococcaemia	1 (0)
DA39.4	Meningococcaemia, unspecified	21 (0.1)
DA40.0	Sepsis due to streptococcus, group A	237 (1.2)
DA40.1	Sepsis due to streptococcus, group B	135 (0.7)
DA40.2	Sepsis due to streptococcus, group D	18 (0.1)
DA40.3	Sepsis due to <i>Streptococcus pneumoniae</i>	1,150 (5.6)
DA40.8	Other streptococcal sepsis	271 (1.3)
DA40.9	Streptococcal sepsis, unspecified	555 (2.7)
DA41	Other sepsis	7 (0)
DA41.0	Sepsis due to <i>Staphylococcus aureus</i>	1,611 (7.9)
DA41.1	Sepsis due to other specified staphylococcus	254 (1.2)
DA41.1A	Sepsis due to coagulase-negative staphylococcus	50 (0.2)
DA41.2	Sepsis due to unspecified staphylococcus	639 (3.1)
DA41.3	Sepsis due to <i>Haemophilus influenzae</i>	50 (0.2)
DA41.4	Sepsis due to anaerobes	386 (1.9)
DA41.5	Sepsis due to other Gram-negative organisms	5,741 (28.1)
DA41.8	Other specified sepsis	978 (4.8)
DA41.9	Sepsis, unspecified	7,224 (35.4)
DA42.7	Actinomycotic sepsis	12 (0.1)
DA49.9A	Bacteraemia, unspecified	436 (2.1)
DB37.7	Candidal sepsis	271 (1.3)
DB49.9A	Fungemia, unspecified	35 (0.2)
	Total	20,433 (100)

doi:10.1371/journal.pone.0131682.t002

age groups (from 29.6% for 0–14 y to 35.0% for >80 y), Charlson comorbidity index score (from 30.3% for 0 to 34.7% for >2), and sepsis severity (from 29.3% for no sepsis to 44.3% for severe sepsis/septic shock). It was conspicuously low for surgical ward patients (17.9%) whereas a higher completeness was seen for non-incident bacteraemic episodes (39.3%) and mortality within 30 days (34.7%). Completeness for acquisition declined from community (36.2%), over healthcare-associated (34.9%) to nosocomial (26.0%). For groups of microorganisms (unknown [n = 43] excluded) the completeness varied from 19.5% for anaerobic Gram-negative bacteria to 36.8% for haemolytic streptococci.

Characteristics of patients with the most likely focal infection codes

The proportion of bacteraemic episodes with a group 3–7 code (45.7%, [Table 3](#)) differed between all subgroups (all $p < 10^{-4}$, except for incident vs. non-incident episodes [$p = 0.02$] and sepsis groups [$p = 0.003$]). The completeness increased with age group (from 36.1% for 0–14 y to 48.3% for >80 y) and decreased with higher Charlson comorbidity index score (from 53.8% for 0 to 39.0% for >2). A lower completeness was seen for paediatric ward patients (35.5%) and mortality within 30 days (33.2%). Completeness for acquisition declined from




Fig 1. Annual proportions of bacteraemia episodes captured by ICD-10 or NOMESCO codes designating “Bacteraemia/sepsis” or “Focal infection”, 2000–2011.

doi:10.1371/journal.pone.0131682.g001

community (55.0%), over healthcare-associated (39.9%) to nosocomial (37.4%). For groups of microorganisms (unknown [n = 99] excluded) the completeness varied from 32.3% for CNS to 53.0% for haemolytic streptococci. Among sepsis groups, the trend of higher completeness with increasing severity was less conspicuous than for bacteraemia codes.

30-day mortality in multivariate analyses

The ORs (95% CIs) for 30-day mortality generally varied little whether these were computed for all bacteraemic episodes, episodes with a group 1–7 code, episodes with a group 1 code, or episodes with a group 3–7 code (Table 4) or not. However, surgical episodes with a group 1 code (OR [95% CIs]: 1.07 [0.94–1.20]), episodes with CNS (ranging from 1.24 [1.01–1.52] for episodes with a group 1–7 code to 1.43 [1.11–1.83] for episodes with a group 3–7 code), and in particular paediatric patient episodes with a group 1 code (0.98 [0.59–1.63]) or a group 3–7 code (5.32 [3.48–8.13]) deviated from all the bacteraemic episodes.

Discussion

Even with a comprehensive inclusion of diagnostic ICD-10 codes and NOMESCO procedure codes that could indicate either sepsis/bacteraemia or a focal infection, only 64.9% of bacteraemic episodes had at least one of these codes registered in the relevant hospital contact. With restriction to codes that more likely represented bacteraemia the completeness declined to 32.3%.

Our gold standard was bacteraemic episodes derived from positive BCs recorded in electronic laboratory information systems maintained by DCMs, from which we excluded contamination episodes by generally accepted algorithms [21, 23]. This, as well as the capture of the

Table 3. Patient characteristics in relation to groups of diagnosis/surgical procedure codes that indicate hospitalization with bacteraemia or the presence of a focal infection.

Characteristic	No. episodes	No. (%) with group 1–7	No. (%) with group 1	No. (%) with group 3–7
All	58,139	37,740 (64.9)	18,786 (32.3)	26,538 (45.7)
Gender				
Females	26,848	17,883 (66.6)	8,741 (32.6)	12,793 (47.7)
Males	31,291	19,857 (63.5)	10,045 (32.1)	13,745 (43.9)
Age group, years				
0–14	2,112	1,493 (70.7)	624 (29.6)	763 (36.1)
15–64	19,771	12,383 (62.6)	6,026 (30.5)	8,886 (44.9)
65–80	20,257	12,892 (63.6)	6,543 (32.3)	9,163 (45.2)
>80	15,999	10,972 (68.6)	5,593 (35.0)	7,726 (48.3)
Charlson comorbidity index score				
0	19,684	13,988 (71.1)	5,956 (30.3)	10,587 (53.8)
1–2	22,501	14,014 (62.3)	7,294 (32.4)	9,725 (43.2)
>2	15,954	9,738 (61.0)	5,536 (34.7)	6,226 (39.0)
Speciality⁴				
Medical	29,678	19,291 (65.0)	10,792 (36.4)	12,601 (42.5)
Surgical	10,652	6,110 (57.4)	1,905 (17.9)	5,091 (47.8)
Intensive care unit	3,410	1,976 (58.0)	1,100 (32.3)	1,481 (43.4)
Paediatric	1,627	1,168 (71.8)	513 (31.5)	577 (35.5)
Unknown	115	83 (72.2)	44 (38.3)	52 (45.2)
Acquisition of bacteraemia				
Community	25,383	19,066 (75.1)	9,192 (36.2)	13,971 (55.0)
Healthcare-associated	12,152	7,673 (63.1)	4,244 (34.9)	4,854 (39.9)
Nosocomial	20,604	11,001 (53.4)	5,350 (26.0)	7,713 (37.4)
Group of microorganisms				
<i>Escherichia coli</i>	16,206	11,896 (73.4)	5,895 (36.4)	8,459 (52.2)
<i>Enterobacter</i> spp.	1,232	726 (58.9)	343 (27.8)	499 (40.5)
<i>Klebsiella</i> spp.	3,893	2,468 (63.4)	1,202 (30.9)	1,711 (44.0)
Other Enterobacteriaceae	2,359	1,567 (66.4)	768 (32.6)	1,067 (45.2)
<i>Pseudomonas aeruginosa</i>	1,601	974 (60.8)	561 (35.0)	614 (38.4)
Anaerobic Gram-negative bacteria	1,361	814 (59.8)	265 (19.5)	664 (48.8)
Other Gram-negative bacteria	1,695	930 (54.9)	495 (29.2)	598 (35.3)
<i>Staphylococcus aureus</i>	7,116	4,508 (63.4)	2,605 (36.6)	2,824 (39.7)
Coagulase-negative staphylococci	2,094	960 (45.9)	424 (20.3)	676 (32.3)
<i>Streptococcus pneumoniae</i>	4,912	3,200 (65.2)	1,556 (31.7)	2,489 (50.7)
Haemolytic streptococci	2,096	1,595 (76.1)	772 (36.8)	1,110 (53.0)
Enterococci	2,916	1,865 (64.0)	883 (30.3)	1,397 (47.9)
Other Gram-positive bacteria	2,522	1,391 (55.2)	510 (20.2)	1,057 (41.9)
Gram-positive rods	1,190	619 (52.0)	283 (23.8)	412 (34.6)
Fungi	1,641	990 (60.3)	550 (33.5)	723 (44.1)
Polymicrobial	5,063	3,114 (61.5)	1,631 (32.2)	2,139 (42.3)
Unknown ⁵	242	123 (50.8)	43 (17.8)	99 (40.9)
Incident bacteraemic episode				
Yes	48,437	31,218 (64.5)	14,976 (30.9)	22,215 (45.9)
No	9,702	6,522 (67.2)	3,810 (39.3)	4,323 (44.6)
Sepsis group⁶				
No sepsis	249	154 (61.9)	73 (29.3)	110 (44.2)

(Continued)

Table 3. (Continued)

Characteristic	No. episodes	No. (%) with group 1–7	No. (%) with group 1	No. (%) with group 3–7
Possibly sepsis	421	283 (67.2)	141 (33.5)	205 (48.7)
Sepsis	610	435 (71.3)	208 (34.1)	341 (55.9)
Severe sepsis/septic shock	1,312	921 (70.2)	581 (44.3)	616 (47.0)
Organ dysfunction, no sepsis	169	120 (71.0)	74 (43.8)	84 (49.7)
30-day mortality				
Yes	12,851	7,094 (55.2)	4,464 (34.7)	4,261 (33.2)
No	45,263	30,625 (67.7)	14,314 (31.6)	22,260 (49.2)
Unknown	25	21 (84.0)	8 (32.0)	17 (68.0)

¹ Diagnosis/surgical procedure that indicate hospitalization with bacteraemia

² "Infections / bacteraemia" codes, which most likely represent bacteraemia, cf. Table 2

³ Codes, which represent the presence of a focal infection (see Appendix)

⁴ Only for bacteraemic episodes before 2009 (n = 45,482)

⁵ Mainly due to lack of specification

⁶ Only for incident community-acquired bacteraemic episodes, Funen County, 2000–2008 (n = 2,761)

doi:10.1371/journal.pone.0131682.t003

majority of positive BCs [17], indicates that our study database represents the greater part of detected bacteraemic episodes within well-defined geographic regions.

Bacteraemia is a serious condition [1], which should theoretically encourage its recording in administrative registries. However, many of the diagnoses that most likely capture the aetiological entity bacteraemia actually designate sepsis (Table 2). Sepsis is a clinical entity previously defined as the presence of at least two among four Systemic Inflammatory Response Syndrome (SIRS) criteria as well as infection [5, 29] and currently defined by a broader definition [30]. The correct coding of sepsis is complicated [31–36] and there is no consensus on which code abstraction strategy that will correctly capture septic episodes [37–39]. Two prior studies, a Swedish based on ICD-9 and ICD-10 codes and a US based on ICD-9 codes, compared code abstraction strategies used to retrieve severe sepsis hospitalizations from administrative registries [6, 7]. The number of severe sepsis hospitalizations varied more than three-fold, which plausibly explains the high variation when reporting incidence of sepsis [37–41].

Most administrative data validation studies either retrieved diagnostic codes from administrative registries followed by validation in randomly sampled medical charts [10, 11, 42–46] or they compared administrative registries to assess concordance [11–14, 47–49]. Fewer studies have initially scrutinized data believed to represent the gold standard followed by their completeness in administrative registries [8, 9, 50–53]

To the best of our knowledge, only the Danish ‘predecessor’ study that prompted this study has validated the diagnosis of bacteraemia in administrative registries [9]. That study included 406 bacteraemic episodes from 1994 recorded in a prospectively validated research database of positive BCs and clinical assessments [54]. Only 18 episodes (4.4%) were recorded with a bacteraemia/sepsis diagnosis in the DNPR. The DNPR replacement of ICD-8 by ICD-10 in 1994 [15] may be a reason for this low completeness. Analysis of data for our study population by using the same 30 ICD-10 codes as in the 1994 study [9] yielded a completeness of 32.0% (data not shown), which is virtually identical to the 32.3% reported here (see Appendix, group 1 codes) and thus representing a notable improvement as compared to the 4.4% reported from 1994 [9].

A few studies have validated sepsis in administrative registries, focusing on severe sepsis or septic shock [8, 50, 51]. Comparison to our study is difficult for several reasons: we do not

Table 4. Odds ratios (95% confidence intervals) for 30-day mortality, adjusted for characteristics in first column, 2000–2008.

Characteristic	All episodes (n = 45,472)	With group 1–7 ¹ (n = 28,620)	With group 1 ² (n = 14,350)	With group 3–7 ³ (n = 19,796)
Gender				
Females	1 (ref.)	1 (ref.)	1 (ref.)	1 (ref.)
Males	0.99 (0.95–1.04)	1.01 (0.95–1.08)	1.05 (0.96–1.14)	1.01 (0.93–1.09)
Age, per 1 year increase	1.033 (1.032–1.035)	1.035 (1.032–1.037)	1.031 (1.028–1.034)	1.038 (1.034–1.041)
Charlson comorbidity index score				
0	1 (ref.)	1 (ref.)	1 (ref.)	1 (ref.)
1–2	1.38 (1.30–1.47)	1.36 (1.26–1.47)	1.24 (1.11–1.37)	1.40 (1.27–1.55)
>2	1.95 (1.82–2.08)	1.87 (1.71–2.04)	1.68 (1.50–1.88)	1.94 (1.73–2.16)
Speciality				
Medical	1 (ref.)	1 (ref.)	1 (ref.)	1 (ref.)
Surgical	0.82 (0.78–0.87)	0.82 (0.75–0.89)	1.07 (0.94–1.20)	0.92 (0.83–1.02)
Intensive care unit	3.36 (3.09–3.66)	3.46 (3.09–3.88)	3.32 (2.86–3.86)	3.65 (3.18–4.20)
Paediatric	2.12 (1.64–2.74)	2.86 (2.10–3.90)	0.98 (0.59–1.63)	5.32 (3.48–8.13)
Unknown	0.78 (0.46–1.33)	0.57 (0.27–1.21)	0.52 (0.20–1.34)	0.75 (0.29–1.93)
Acquisition of bacteraemia				
Community	1 (ref.)	1 (ref.)	1 (ref.)	1 (ref.)
Healthcare-associated	1.33 (1.25–1.43)	1.20 (1.09–1.31)	1.12 (1.00–1.26)	1.29 (1.15–1.46)
Nosocomial	1.83 (1.72–1.94)	1.84 (1.70–1.99)	1.71 (1.54–1.89)	1.94 (1.75–2.15)
Group of microorganisms				
<i>Escherichia coli</i>	1 (ref.)	1 (ref.)	1 (ref.)	1 (ref.)
<i>Enterobacter</i> spp.	1.37 (1.16–1.62)	1.36 (1.08–1.73)	1.30 (0.94–1.78)	1.33 (0.97–1.82)
<i>Klebsiella</i> spp.	1.28 (1.16–1.42)	1.33 (1.16–1.53)	1.30 (1.09–1.56)	1.43 (1.19–1.71)
Other Enterobacteriaceae	1.22 (1.07–1.39)	1.36 (1.14–1.62)	1.36 (1.08–1.72)	1.58 (1.27–1.97)
<i>Pseudomonas aeruginosa</i>	1.64 (1.43–1.88)	1.80 (1.49–2.16)	1.88 (1.50–2.36)	1.89 (1.48–2.43)
Anaerobic Gram-negative bacteria	1.99 (1.72–2.32)	1.80 (1.45–2.24)	2.08 (1.51–2.88)	2.02 (1.57–2.61)
Other Gram-negative bacteria	1.35 (1.15–1.58)	1.59 (1.26–2.00)	1.81 (1.35–2.42)	1.62 (1.18–2.22)
<i>Staphylococcus aureus</i>	1.80 (1.66–1.95)	2.05 (1.85–2.28)	1.92 (1.68–2.19)	2.34 (2.03–2.69)
Coagulase-negative staphylococci	0.93 (0.81–1.07)	1.24 (1.01–1.52)	1.39 (1.05–1.85)	1.43 (1.11–1.83)
<i>Streptococcus pneumoniae</i>	1.45 (1.31–1.60)	1.79 (1.57–2.04)	1.91 (1.61–2.26)	1.96 (1.66–2.31)
Haemolytic streptococci	1.41 (1.22–1.63)	1.54 (1.29–1.84)	1.72 (1.37–2.16)	1.45 (1.14–1.85)
Enterococci	1.47 (1.31–1.64)	1.72 (1.49–2.00)	1.83 (1.50–2.22)	1.92 (1.61–2.31)
Other Gram-positive bacteria	1.25 (1.09–1.42)	1.26 (1.04–1.53)	1.37 (1.04–1.80)	1.43 (1.12–1.82)
Gram-positive rods	1.71 (1.45–2.02)	1.94 (1.53–2.46)	2.08 (1.51–2.88)	2.08 (1.53–2.82)
Fungi	2.64 (2.30–3.02)	2.59 (2.16–3.10)	2.60 (2.05–3.30)	2.70 (2.16–3.37)
Polymicrobial	2.30 (2.11–2.50)	2.29 (2.04–2.57)	2.21 (1.91–2.56)	2.36 (2.04–2.74)
Unknown ⁴	0.99 (0.65–1.53)	1.11 (0.58–2.16)	0.82 (0.24–2.81)	1.21 (0.56–2.61)
Incident bacteraemic episode				
Yes	1 (ref.)	1 (ref.)	1 (ref.)	1 (ref.)
No	0.96 (0.90–1.02)	0.96 (0.88–1.04)	0.90 (0.81–1.00)	0.99 (0.89–1.10)

¹ Diagnosis/surgical procedure that may indicate bacteraemia

² "Infections / bacteraemia" codes, which most likely represent bacteraemia, cf. [Table 2](#)

³ Codes that indicate a focal infection (see [Appendix](#))

⁴ Mainly due to lack of speciation

doi:10.1371/journal.pone.0131682.t004

know how much overlap there is between sepsis and bacteraemia, ICD-9 codes (which may differ substantially from ICD-10 codes [32, 55]) were used, capture of sepsis varies up to three-fold depending on the algorithm [6, 7], or the study included emergency department or ICU patients only [8, 51].

Although a low proportion of the bacteraemic episodes was recorded with a relevant diagnosis in the DNPR this may not pose a problem if this capture is non-selective, but this was not the case as proportions varied up to two-fold, e.g., 17.9% for surgical vs. 36.4% for medical patients pertaining to group 1 diagnoses. Likewise, a higher completeness was seen with increasing severity of sepsis, also reported for severe sepsis [8], whereas it declined from community over healthcare-associated to nosocomial acquisition. There were fewer variations between ORs in multivariate 30-day mortality analyses; caveat is still warranted for some subgroups though, e.g., paediatric ward patients. A meta-analysis of 36 severe sepsis trials reported the same declining mortality trend from 1993 through 2009 whether data were from the trials or administrative registries or not, though percent mortality differed considerably [56].

The Danish version of the WHO ICD-10 classification [25] is from 1993 [57] and instructions on coding and registration have not been updated since then. The codes are updated on the official Danish web site [24] and coding is facilitated by private entrepreneurs [58]. Most of the codes that cover bacteraemia include “Sepsis” in their designation, with a few exceptions, such as A49.9A [Bacteraemia, unspecified], used only for 2.1% of the Group 1 codes (Table 2). One reason for this may be that A49.9A is not included in the original code book [57], so only physicians who are aware of the web amendments [24, 58] will probably use this highly relevant code. A future update could alter the designations in the Group 1 codes from “Sepsis” to “Bacteraemia”, as this would be more in accordance with globally accepted definitions [5] and would probably facilitate and increase the recording of bacteraemia in the DNPR.

We used a database which represented clinically important bacteraemic episodes, the study was population-based, and included a high number of episodes that enabled subgroup analyses. However, there were also limitations that warrant further discussion. The main limitation was the inability to report predictive values as we had no information on the use of the 1,079 codes in the DNPR for patients not having bacteraemia. In the Danish ‘predecessor’ study, codes that roughly correspond to our group 1 codes were assessed for all patients in the DNPR, which enabled the reporting of positive predictive values (PPVs), found to be 21.7% [9]. PPVs would probably decline with the inclusion of group 1–7 or group 3–7 codes, in which the prevalence of bacteraemia would not alter the numerator, but the denominator comprising all the said codes would increase. Such low PPVs, and consequently many false positive patients, preclude research on bacteraemia patients based on administrative data. Secondly, we only had clinical data for 4.7% of the bacteraemic episodes. The increasing completeness with higher sepsis severity further indicated the selective recording of patient groups in administrative registries. Thirdly, the multiple ICD-10 codes prompted us to define a limited number of groups based on the likelihood to represent bacteraemia. Although we used a consensus process the classification of codes and the ranking of groups were subjective. Still, the ordinal scale provided a working solution to the conundrum of 5,040 ordered sequences of the 7 groups. Finally, some blood cultures with common skin commensals may represent contamination, and not bacteraemia, which may be reflected in the lower completeness for CNS (Table 3). However, for 9,482 bacteraemic episodes (part of the actual study database) we previously reported 94.6% agreement for bacteraemia vs. contamination when computer algorithms were compared to physicians’ individual clinical assessments [21]. We therefore believe this had minor impact on the results.

In conclusion, our study showed a low completeness of bacteraemic episodes identified in the official Danish administrative registry used for the recording of all hospital contacts.

Further, there were considerable differences in completeness as to whether the acquisition was community, healthcare-associated or nosocomial. Although few studies have shown this for bacteraemia, this is in accordance with sepsis [33, 35] and health-care associated infections [55, 59] for which it has been concluded that their detection should not be based solely on administrative registry data. Hence, bacteraemia studies should preferably be derived from bacteraemia databases based on positive blood cultures [54, 60, 61].

Appendix

Diagnosis/operation groups, their codes, and text examples of the three most common codes within each group, followed by two examples of prioritization

Group 1: Infections/bacteraemia

DA02.1, DA20.7, DA21.7, DA22.7, DA22.9B, DA26.7, DA28.2B, DA32.7, DA39.2, DA39.2A, DA39.3, DA39.4, DA40, DA40.0, DA40.1, DA40.2, DA40.3, DA40.8, DA40.9, DA41, DA41.0, DA41.1, DA41.1A, DA41.2, DA41.3, DA41.4, DA41.5, DA41.8, DA41.9, DA41.9C, DA42.7, DA49.9A, DA54.8G, DB37.7, DB49.9A

The three most common diagnoses from Group 1: See article, [Table 2](#)

Group 2: Other diagnoses/bacteraemia

DO75.3A, DO85.9, DP36, DP36.0, DP36.1, DP36.2, DP36.3, DP36.4, DP36.5, DP36.8, DP36.9

The three most common diagnoses from Group 2 (no./all group codes [%]):

DP36.9 –Bacterial sepsis of newborn, unspecified (2,749/3,816 [72.0])

DP36.2 –Sepsis of newborn due to *Staphylococcus aureus* (904/3,816 [23.7])

DP36.0—Sepsis of newborn due to streptococcus, group B (80/3,816 [2.1])

Group 3: Other diagnoses/focal infection

DD73.3, DE06.0A, DE06.0B, DE10.5C, DE11.5C, DE12.5C, DE13.5C, DE13.5D, DE14.5B, DE14.5C, DE23.6A, DE32.1, DG00, DG00.0, DG00.1, DG00.2, DG00.3, DG00.8, DG00.8A, DG00.8B, DG00.9, DG00.9A, DG01, DG01.9, DG01.9B, DG01.9C, DG01.9D, DG01.9G, DG01.9H, DG01.9I, DG02, DG02.1A, DG02.1C, DG02.8, DG04.2, DG04.2A, DG04.2B, DG05.0, DG05.0E, DG05.0H, DG05.0I, DG05.0O, DG05.0P, DG05.0S, DG05.2, DG05.2L, DG05.2M, DG05.2N, DG06, DG06.0, DG06.0A, DG06.0B, DG06.0C, DG06.0D, DG06.0E, DG06.0F, DG06.1, DG06.1A, DG06.1B, DG06.1C, DG06.2, DG06.2A, DG06.2B, DG06.2C, DG07, DG07.9, DG07.9A, DG07.9B, DG07.9C, DG07.9E, DG08.9A, DG08.9B, DG08.9C, DG08.9D, DG08.9E, DG08.9F, DG08.9G, DG08.9H, DG08.9I, DG08.9J, DG08.9K, DG08.9L, DG08.9M, DG08.9N, DH05.0, DH05.0A, DH05.0B, DH05.0E, DH44.0, DH44.0B, DH70.0, DH70.0A, DH70.0B, DH70.1, DH75.0, DH75.0B, DI30.1, DI30.1A, DI30.1B, DI30.1C, DI30.1D, DI32.0, DI32.0A, DI32.0B, DI32.0C, DI32.1, DI32.1A, DI32.1B, DI32.8A, DI33, DI33.0, DI33.0A, DI33.0B, DI33.0C, DI33.0D, DI33.0E, DI33.0F, DI33.9, DI38, DI39.0, DI39.1, DI39.2, DI39.3, DI39.4, DI39.8, DI39.8A, DI39.8B, DI39.8D, DI39.8E, DI40, DI40.0, DI40.0A, DI41.0, DI41.0B, DI41.0C, DI52.0, DI52.0A, DI68.1, DI68.1A, DJ01.0A, DJ01.0B, DJ01.1A, DJ01.1B, DJ01.2A, DJ01.2B, DJ01.3A, DJ01.3B, DJ01.4A, DJ01.4B, DJ02.9C, DJ03.9D, DJ13, DJ13.9, DJ13.9A, DJ13.9B, DJ14, DJ14.9, DJ14.9A, DJ14.9B, DJ15, DJ15.0, DJ15.1, DJ15.2, DJ15.3, DJ15.4, DJ15.5, DJ15.6, DJ15.6A, DJ15.7, DJ15.8, DJ15.9, DJ16, DJ17.0, DJ17.0A, DJ17.0B, DJ17.0C, DJ17.0D, DJ17.0E, DJ17.0F, DJ17.0G, DJ17.1, DJ18.8, DJ34.0, DJ36.9, DJ39.0, DJ39.0A, DJ39.0B, DJ39.0C, DJ39.1A, DJ85, DJ85.0, DJ85.0A, DJ85.1, DJ85.2, DJ85.3, DJ86, DJ86.0, DJ86.0A, DJ86.9, DJ86.9A, DJ95.0A, DJ98.5D, DK04.1A, DK11.3, DK11.3A, DK11.3B, DK11.3C, DK11.3D, DK12.2, DK12.2A, DK12.2B, DK13.0A, DK14.0A, DK20.9A, DK35, DK35.0, DK35.0A, DK35.1, DK35.1A, DK35.2, DK35.3, DK35.3A, DK35.3B, DK35.8,

DK35.8A, DK35.8B, DK35.8C, DK35.9, DK35.9A, DK35.9B, DK36, DK40.1, DK40.4, DK41.1, DK41.4, DK42.1, DK43.1, DK43.4, DK43.7, DK44.1, DK45.1, DK45.1B, DK45.1C, DK45.1D, DK45.1E, DK45.1F, DK45.1G, DK45.1H, DK45.1I, DK45.1J, DK45.1K, DK45.1L, DK45.1M, DK46.1, DK55.0A, DK55.0B, DK55.0C, DK55.0D, DK55.1B, DK55.1C, DK55.1D, DK57, DK57.0, DK57.0A, DK57.0B, DK57.0C, DK57.1, DK57.2, DK57.2A, DK57.2B, DK57.2C, DK57.3, DK57.4, DK57.4A, DK57.5, DK57.8, DK57.9, DK57.9A, DK59.3C, DK61.0, DK61.0A, DK61.0B, DK61.1, DK61.1A, DK61.2, DK61.3, DK61.4, DK63.0, DK63.8A, DK65, DK65.0, DK65.0A, DK65.0B, DK65.0C, DK65.0D, DK65.0E, DK65.0F, DK65.0G, DK65.0H, DK65.0I, DK65.0J, DK65.0K, DK65.0L, DK65.0M, DK65.0N, DK65.0O, DK65.0P, DK65.8, DK65.8I, DK67.1, DK75.0, DK75.0A, DK75.0B, DK75.0C, DK75.0D, DK80.0, DK80.0A, DK80.0B, DK80.0C, DK80.0D, DK80.1, DK80.1C, DK80.3, DK80.3A, DK80.3B, DK80.3C, DK80.4, DK80.4A, DK80.4B, DK80.4C, DK80.4D, DK80.4E, DK81.0, DK81.0A, DK81.0B, DK81.0C, DK81.0D, DK81.1, DK83.0, DK83.0A, DK83.0B, DK83.0C, DK83.0D, DK83.0E, DK85, DK85.0, DK85.1, DK85.1A, DK85.8A, DK85.8E, DK85.9, DK85.9A, DK85.9B, DK85.9D, DK85.9E, DK86.1A, DL02.0, DL02.1B, DL02.1C, DL02.2G, DL02.2H, DL02.2I, DL02.2J, DL02.2K, DL02.2L, DL02.2M, DL02.2N, DL02.2R, DL02.2S, DL02.2T, DL02.8A, DL02.8B, DL02.8C, DL03.0A, DL03.0F, DL03.1, DL03.1A, DL03.1B, DL03.1C, DL03.1D, DL03.1E, DL03.1F, DL03.1G, DL03.1H, DL03.1I, DL03.3, DL03.3A, DL03.3B, DL03.3C, DL03.3D, DL03.3E, DL03.3F, DL03.8, DL03.8A, DL08.0, DL08.0A, DL08.0a, DL08.8B, DM00, DM00.0, DM00.0A, DM00.0B, DM00.1, DM00.1A, DM00.1B, DM00.2, DM00.2A, DM00.2B, DM00.8, DM00.9, DM01, DM01.0, DM01.1, DM01.3, DM01.3A, DM01.3B, DM01.3C, DM01.6, DM46.2, DM46.3, DM46.3A, DM46.4, DM46.5, DM46.5A, DM49.0, DM49.1, DM49.2, DM49.3, DM49.3A, DM60.0, DM60.0A, DM60.8, DM60.8A, DM60.8A1, DM60.9, DM63.0, DM65.0, DM65.1, DM68.0, DM68.0G, DM68.0H, DM72.5A, DM72.6, DM86, DM86.0, DM86.1, DM86.2, DM86.3, DM86.4, DM86.5, DM86.5A, DM86.6, DM86.8, DM86.8A, DM86.9, DM86.9A, DM90.2, DM90.2C, DN10, DN10.9, DN10.9A, DN10.9B, DN10.9C, DN11, DN11.0, DN11.0A, DN11.1, DN11.1A, DN11.1B, DN11.2, DN11.8, DN12, DN12.9, DN13.6, DN13.6A, DN13.6B, DN13.6C, DN13.6D, DN13.6E, DN15.1, DN15.1A, DN15.1B, DN20.9A, DN30, DN30.0, DN30.1, DN30.8A, DN34.0, DN39.0, DN41.0, DN41.1, DN41.2, DN41.2A, DN41.3, DN41.3A, DN41.8, DN45.0, DN45.0A, DN45.0B, DN45.0C, DN45.0D, DN45.9, DN48.2A, DN48.2B, DN48.2C, DN48.2D, DN48.2E, DN48.2F, DN48.2H, DN48.2I, DN49.8C, DN51.0A, DN51.1E, DN51.1H, DN61.9B, DN61.9E, DN61.9F, DN70.0B, DN70.0C, DN70.0D, DN70.0F, DN70.0G, DN70.1, DN70.1A, DN71.0, DN71.0A, DN71.0B, DN71.0C, DN71.0D, DN71.0E, DN71.0F, DN73.0, DN73.0A, DN73.0B, DN73.0C, DN73.0D, DN73.0E, DN73.1A, DN73.1B, DN73.2A, DN73.2B, DN73.3, DN73.3A, DN73.4A, DN73.5A, DN73.8A, DN73.8B, DN73.8C, DN75.1, DN75.8A, DN76.0A, DN76.0B, DO03.0, DO03.5, DO03.5A, DO03.5B, DO04.0, DO04.5, DO04.5A, DO04.5B, DO07.0, DO07.5, DO08.0, DO08.0A, DO08.0B, DO08.0C, DO08.0D, DO08.0E, DO08.0F, DO08.0G, DO08.0H, DO08.0I, DO08.0J, DO08.0K, DO08.0L, DO08.0M, DO08.0N, DO08.0O, DO08.0P, DO08.0Q, DO08.0R, DO23, DO23.0, DO41.1, DO41.1A, DO41.1B, DO41.1C, DO41.1D, DO75.2, DO75.3, DO85, DO85.9A, DO85.9B, DO86, DO86.0, DO86.0A, DO86.0B, DO86.0C, DO86.0D, DO86.1C, DO86.2, DO86.2B, DO86.3, DO88.3, DO91.1, DO91.1D, DO91.1E, DO91.1F, DO91.1G, DO91.1H, DO91.1I, DO91.1J, DO91.1K, DO91.1L, DO91.2B, DO91.2C, DO91.2D, DO91.2E, DP02.7, DP02.7A, DP02.7B, DP02.7C, DP15.4A, DP23, DP23.2, DP23.3, DP23.4, DP23.5, DP23.6, DP23.6A, DP23.6B, DP23.6C, DP38.9, DP38.9A, DP38.9B, DP38.9C, DP38.9D, DP39.2, DP39.3, DP39.4, DP39.8, DP39.9, DP58.2, DP77, DP77.9, DT79.3, DT80.1A, DT80.1B, DT80.1C, DT80.2, DT80.2A, DT80.2B, DT80.2C, DT80.2D, DT80.2E, DT80.2F, DT81.4, DT81.4A, DT81.4B, DT81.4C, DT81.4D, DT81.4F, DT81.4G, DT81.4H, DT81.4I, DT81.4J, DT81.4P, DT81.4U, DT81.4X, DT82.6, DT82.6A, DT82.7, DT82.7A, DT82.7B,

DT82.7I, DT82.7P, DT83.5A, DT83.5B, DT83.5C, DT83.6, DT83.6A, DT83.6B, DT83.6C, DT84.5, DT84.5A, DT84.6, DT84.6A, DT84.7, DT85.7, DT87.4, DT88.0, DT89

The three most common diagnoses from Group 3 (no./all group codes [%]):

DN30.0—Acute cystitis (1,025/1,936 [52.9])

DN39.0—Urinary tract infection, site not specified (410/1,936 [21.2])

DN10.9—Acute tubulo-interstitial nephritis, unspecified (114/1,936 [5.9])

Group 4: Surgical procedures/focal infection

KAAM, KAAM00, KAAM10, KAAM99, KAWC, KAWC00, KAWC00A, KBWC, KBWC00, KFWC, KFWC00, KFXA10, KGWC, KGWC00, KGWC01, KHWC, KHWC00, KJAJ, KJAJ00, KJJA10A, KJLD13, KJLD13C, KJLD13D, KJWC, KJWC00, KJWC01, KKWC, KKWC00, KKWC01, KLWC, KLWC00, KLWC01, KMWC, KMWC00, KMWC01, KNAS, KNAS1, KNAS10, KNAS11, KNAS12, KNAS13, KNAS14, KNAS15, KNAS16, KNAS19, KNAS2, KNAS20, KNAS21, KNAS22, KNAS23, KNAS24, KNAS25, KNAS26, KNAS29, KNAS4, KNAS40, KNAS41, KNAS42, KNAS43, KNAS44, KNAS45, KNAS46, KNAS49, KNAS5, KNAS50, KNAS51, KNAS52, KNAS53, KNAS54, KNAS55, KNAS56, KNAS59, KNAS6, KNAS60, KNAS61, KNAS62, KNAS63, KNAS64, KNAS65, KNAS66, KNAS69, KNAS9, KNAS90, KNAS91, KNAS92, KNAS93, KNAS94, KNAS95, KNAS96, KNAS99, KNAU89, KNAW69, KNBS, KNBS09, KNBS19, KNBS29, KNBS39, KNBS49, KNBS59, KNBS99, KNBU89, KNBU89A, KNBU89B, KNBW69, KNCS, KNCS09, KNCS19, KNCS29, KNCS39, KNCS49, KNCS59, KNCS99, KNCU89, KNCU89A, KNCU89B, KNCW69, KNDS, KNDS09, KNDS19, KNDS29, KNDS39, KNDS49, KNDS59, KNDS99, KNDU89, KNDU89A, KNDU89B, KNDW69, KNES, KNES19, KNES29, KNES49, KNES59, KNES99, KNEU89, KNEW69, KNFS, KNFS19, KNFS29, KNFS49, KNFS59, KNFS99, KNFU89, KNFU89A, KNFU89B, KNFW69, KNGS, KNGS09, KNGS19, KNGS29, KNGS39, KNGS49, KNGS59, KNGS99, KNGU89, KNGU89A, KNGU89B, KNGW69, KNHS, KNHS09, KNHS19, KNHS29, KNHS39, KNHS49, KNHS59, KNHS99, KNHU89, KNHU89A, KNHU89B, KNHW69, KPJW10, KPWC, KPWC00, KQWB, KQWB00, KQWB10, KQWC, KQWC10, KTJA40, KTJL10, KTKA20

The three most common procedures from Group 4 (no./all group codes [%]):

KTJA40—Percutaneous local drainage of peritoneal cavity (204/538 [37.9])

KJWC00—Reoperation for deep infection in gastroenterological surgery (107/538 [19.9])

KTJL10—Percutaneous drainage of pseudocyst or abscess of pancreas (77/538 [14.3])

Group 5: Infections/systemic infection

DA01, DA01.0, DA01.1, DA01.2, DA01.3, DA01.4, DA20, DA20.9, DA21, DA21.9, DA22, DA22.9, DA23, DA23.0, DA23.1, DA23.2, DA23.3, DA23.8, DA23.9, DA23.9A, DA23.9B, DA23.9C, DA23.9D, DA24, DA24.0, DA24.0A, DA24.1, DA24.2, DA24.3, DA24.4, DA25, DA25.0, DA25.0A, DA25.1, DA25.9, DA26, DA26.9, DA27, DA27.0, DA27.8, DA27.8A, DA27.8B, DA27.8C, DA27.8D, DA27.9, DA28.0, DA32, DA32.8A, DA39, DA39.1, DA41.8, DA41.9, DA41.9A, DA41.9B, DA42.7, DA43.9, DA44.0, DA44.1, DA44.8, DA48.4A, DA49, DA49.1, DB37, DB38.7, DB39.3, DB40.7, DB41.1, DB42.7, DB44.7, DB45, DB45.7, DB46.4, DB95, DB96, DB98.1

The three most common diagnoses from Group 5 (no./all group codes [%]):

DA41.9B –Urosepsis (12/15 [80.0])

DA41.9A –Septic shock (2/15 [13.3])

DA01.0 –Typhus (1/15 [6.7])

Group 6: Other diagnoses/systemic infection

DJ02.0A, DO08.0S, DO08.0T, DO08.0U, DO08.0V, DO08.0X, DO08.0Y, DO08.2J, DO08.2K, DO08.2L, DP36.9, DP37.2, DP37.5

The three most common diagnoses from Group 6 (no./all group codes [%]):

DP37.5—Neonatal candidiasis (3,833/27,976 [13.7])
 DO08.0U –Sepsis after abortion (2,949/27,976 [10.5])
 DO08.0V –Septic shock after molar pregnancy (1,776/27,976 [6.3])

Group 7: Infections/focal infection

DA02, DA02.0, DA02.2, DA02.2A, DA02.2B, DA02.2C, DA02.2D, DA02.2E, DA02.8, DA04.5, DA04.6, DA04.7, DA04.8, DA04.9, DA20.0, DA20.1, DA20.2, DA20.3, DA20.8, DA21.0, DA21.1, DA21.2, DA21.3, DA21.8, DA22.0, DA22.1, DA22.2, DA22.8, DA22.9A, DA22.9C, DA22.9D, DA26.0, DA26.8, DA26.9, DA28.2, DA28.2A, DA32.0, DA32.1, DA32.1A, DA32.1B, DA32.8, DA32.9, DA39.0, DA39.5, DA39.5A, DA39.5B, DA39.8, DA42.8, DA43, DA43.0, DA43.1, DA43.8, DA46, DA46.9, DA48.0, DA54.4, DA54.4A, DA54.4B, DA54.4C, DA54.4D, DA54.8A, DA54.8C, DA54.8D, DA54.8E, DA54.8F, DA69.0C, DA69.1, DA69.1G, DB37.1, DB37.6, DB45.1, DB45.1A, DB45.1B, DI38.9, DI39, DT80.2D1, DT80.2G

The three most common diagnoses from Group 7 (no./all group codes [%]):

DA46.9 –Erysipelas, unspecified (843/2,272 [37.1])
 DI38.9 –Endocarditis, unspecified (287/2,272 [12.6])
 DA39.0—Meningococcal meningitis (190/2,272 [8.4])

Example 1:

A bacteraemic episode with the following 4 codes recorded:

DA41.0—Sepsis due to *Staphylococcus aureus* (Group 1)
 DA41.9A –Septic shock (Group 2)
 DK85.9—Acute pancreatitis, unspecified (Group 6)

KTJL10—Percutaneous drainage of pseudocyst or abscess of pancreas (Group 7)

With a group prioritization of 1 > 2 > 3 > 4 > 5 > 6 > 7, this bacteraemic episode will be recorded as DA41.0—Sepsis due to *Staphylococcus aureus* (Group 1), annulling the three other codes.

With a group prioritization of 5 > 7 > 3 > 4 > 6 > 1 > 2, the same bacteraemic episode will be recorded as KTJL10—Percutaneous drainage of pseudocyst or abscess of pancreas (Group 7), annulling the three other codes.

Example 2:

A bacteraemic episode with the following 3 codes recorded:

DA41.9B –Urosepsis (Group 2)
 DP364—Sepsis of newborn due to *Escherichia coli* (Group 4)
 DG008B - Meningitis from *Escherichia coli* (Group 6)

With a group prioritization of 1 > 2 > 3 > 4 > 5 > 6 > 7, this bacteraemic episode will be recorded as DA41.9B –Urosepsis (Group 2), annulling the two other codes.

With a group prioritization of 5 > 7 > 3 > 4 > 6 > 1 > 2, the same bacteraemic episode will be recorded as DP364—Sepsis of newborn due to *Escherichia coli* (Group 4), annulling the two other codes.

Acknowledgments

Contributing members of DACOBAN: Christian Østergaard (Department of Clinical Microbiology, Copenhagen University Hospital, Hvidovre Hospital, Copenhagen, Denmark), Magnus Arpi (Department of Clinical Microbiology, Copenhagen University Hospital, Herlev Hospital, Copenhagen, Denmark), Kim Oren Gradel (Center for National Clinical Databases—South, Odense University Hospital, Odense, Denmark), Ulrich Stab Jensen (Department of Clinical Microbiology, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark), Sara Thønnings (Department of Clinical Microbiology, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark), Jenny Dahl Knudsen (Department of Clinical

Microbiology, Copenhagen University Hospital, Hvidovre Hospital, Copenhagen, Denmark), Kristoffer Koch (Department of Clinical Microbiology, Aalborg University Hospital, Aalborg, Denmark), Mette Pinholt (Department of Clinical Microbiology, Copenhagen University Hospital, Herlev Hospital, Copenhagen, Denmark), Jesper Smit (Department of Clinical Microbiology, Aalborg University Hospital, Aalborg, Denmark), Henrik Carl Schønheyder (Department of Clinical Microbiology, Aalborg University Hospital, Aalborg, Denmark), Mette Søgaard (Department of Clinical Microbiology, Aalborg University Hospital, Aalborg, Denmark).

Contributing members of DORIS: Anmarie Touborg Lassen (Department of Emergency Medicine, Odense University Hospital, Odense, Denmark), Court Pedersen (Department of Infectious Diseases, Odense University Hospital, Odense, Denmark), Thøger Gorm Jensen (Department of Clinical Microbiology, Odense University Hospital, Odense, Denmark), Hans Jørn Kolmos (Department of Clinical Microbiology, Odense University Hospital, Odense, Denmark), Stig Lønberg Nielsen (Department of Infectious Diseases, Odense University Hospital, Odense, Denmark).

Author Contributions

Conceived and designed the experiments: KOG SLN HCS. Performed the experiments: KOG SLN HCS. Analyzed the data: KOG SLN HCS. Contributed reagents/materials/analysis tools: CP JDK CO MA TGJ HJK ATL. Wrote the paper: KOG SLN CP JDK CO MA TGJ HJK MS ATL HCS.

References

1. Goto M, Al-Hasan MN. Overall burden of bloodstream infection and nosocomial bloodstream infection in North America and Europe. *Clin Microbiol Infect* 2013; 19:501–509
2. Laupland KB, Church DL. Population-based epidemiology and microbiology of community-onset bloodstream infections. *Clin Microbiol Rev* 2014; 27:647–664
3. Schønheyder HC, Paul M. Placing the burden of bacteraemia in perspective. *Clin Microbiol Infect* 2013; 19:489–491
4. Weinstein MP, Reller LB, Murphy JR, Lichtenstein KA. The clinical significance of positive blood cultures: a comprehensive analysis of 500 episodes of bacteremia and fungemia in adults. I. Laboratory and epidemiologic observations. *Rev Infectious Dis* 1983; 5:35–53
5. Bone RC, Balk RA, Cerra FB, Dellinger RP, Fein AM, Knaus WA, et al. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. *Chest* 1992; 101:1644–1655
6. Wilhelms SB, Huss FR, Granath G, Sjöberg F. Assessment of incidence of severe sepsis in Sweden using different ways of abstracting International Classification of Diseases codes: difficulties with methods and interpretation of results. *Crit Care Med* 2010; 38:1442–1449
7. Gaieski DF, Edwards JM, Kallan MJ, Carr BG. Benchmarking the incidence and mortality of severe sepsis in the United States. *Crit Care Med* 2013; 41:1167–1174
8. Whittaker SA, Mikkelsen ME, Gaieski DF, Koshy S, Kean C, Fuchs BD. Severe sepsis cohorts derived from claims-based strategies appear to be biased toward a more severely ill patient population. *Crit Care Med* 2013; 41:945–953
9. Madsen KM, Schønheyder HC, Kristensen B, Nielsen GL, Sørensen HT. Can hospital discharge diagnosis be used for surveillance of bacteremia? A data quality study of a Danish hospital discharge registry. *Infect Control Hosp Epidemiol* 1998; 19:175–180
10. Emori TG, Edwards JR, Culver DH, Sartor C, Stroud LA, Gaunt EE, et al. Accuracy of reporting nosocomial infections in intensive-care-unit patients to the National Nosocomial Infections Surveillance System: a pilot study. *Infect Control Hosp Epidemiol* 1998; 19:308–316
11. Moehring RW, Staheli R, Miller BA, Chen LF, Sexton DJ, Anderson DJ. Central line-associated infections as defined by the Centers for Medicare and Medicaid Services' Hospital-acquired condition versus standard infection control surveillance: why hospital compare seems conflicted. *Infect Control Hosp Epidemiol* 2013; 34:238–244

12. Stevenson KB, Khan Y, Dickman J, Gillenwater T, Kulich P, Myers C, et al. Administrative coding data, compared with CDC/NHSN criteria, are poor indicators of health care-associated infections. *Am J Infect Control* 2008; 36:155–164
13. Stamm AM, Bettacchi CJ. A comparison of 3 metrics to identify health care-associated infections. *Am J Infect Control* 2012; 40:688–691
14. Patrick SW, Davis MM, Sedman AB, Meddings JA, Hieber S, Lee GM, et al. Accuracy of hospital administrative data in reporting central line-associated bloodstream infections in newborns. *Pediatrics* 2013; 131 Suppl 1:S75–80
15. Lynge E, Sandegaard JL, Rebolj M. The Danish National Patient Register. *Scand J Public Health* 2011; 39:30–33
16. Dellinger RP, Levy MM, Rhodes A, Annane D, Gerlach H, Opal SM, et al. Surviving sepsis campaign: international guidelines for management of severe sepsis and septic shock: 2012. *Crit Care Med* 2013; 41:580–637
17. Gradel KO, Schönheyder HC, Arpi M, Knudsen JD, Østergaard C, Søgaard M. The Danish Collaborative Bacteraemia Network (DACOBAN) database. *Clin Epidemiol* 2014; 6:301–308
18. Knudtzen FC, Nielsen SL, Gradel KO, Lassen AT, Kolmos HJ, Jensen TG, et al. Characteristics of patients with community-acquired bacteremia who have low levels of C-reactive protein (≤ 20 mg/L). *J Infect* 2014; 68:149–155
19. Pinholt M, Østergaard C, Arpi M, Bruun NE, Schönheyder HC, Gradel KO, et al. Incidence, clinical characteristics and 30-day mortality of enterococcal bacteraemia in Denmark 2006–2009: a population-based cohort study. *Clin Microbiol Infect* 2014; 20:145–151
20. Gradel KO, Jensen TG, Kolmos HJ, Pedersen C, Vinholt PJ, Lassen AT. Does C-reactive protein independently predict mortality in adult community-acquired bacteremia patients with known sepsis severity? *APMIS* 2013; 121:835–842
21. Gradel KO, Knudsen JD, Arpi M, Østergaard C, Schönheyder HC, Søgaard M. Classification of positive blood cultures: computer algorithms versus physicians' assessment—development of tools for surveillance of bloodstream infection prognosis using population-based laboratory databases. *BMC Med Res Methodol* 2012; 12:139
22. Schmidt M, Pedersen L, Sørensen HT. The Danish Civil Registration System as a tool in epidemiology. *Eur J Epidemiol* 2014; 29:541–549
23. Trick WE, Zagorski BM, Tokars JI, Vernon MO, Welbel SF, Wisniewski MF, et al. Computer algorithms to detect bloodstream infections. *Emerg Infect Dis* 2004; 10:1612–1620
24. SKS-browser, vers 4.02. Available: <http://www.medinfo.dk/sks/brows.php> (Accessed 2014 November 12)
25. WHO, ICD-10 Version: 2010. Available: <http://apps.who.int/classifications/icd10/browse/2010/en#/B49> (Accessed 2014 November 12)
26. NOMESCO Classification of Surgical Procedures (NCSP). Available: http://www.nordclass.se/ncsp_e.htm (Accessed 2014 November 12)
27. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987; 40:373–383
28. Pedersen CB. The Danish Civil Registration System. *Scand J Public Health* 2011; 39:22–25
29. Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D, et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med* 2003; 31:1250–1256
30. Angus DC, van der Poll T. Severe sepsis and septic shock. *N Engl J Med* 2013; 369:840–851
31. Wiedemann LA. Coding sepsis and SIRS. *J AHIMA* 2007; 78:76–78; quiz 79–80
32. Boyle G, Kostick K. Coding UTI to sepsis in ICD-9-CM and ICD-10-CM. *J AHIMA* 2011; 82:52–54
33. Sarrazin MS, Rosenthal GE. Finding pure and simple truths with administrative data. *JAMA* 2012; 307:1433–1435
34. Ljungstrom LR, Steinum O, Brink M, Gardlund B, Martner J, Sjolun J. [Diagnosis and diagnostic coding of severe sepsis and septic shock. ICD-10 should be completed with additional codes]. *Läkartidningen* 2011; 108:276–278
35. Cremer OL. Can administrative data be used to consistently measure the burden of sepsis?. *Crit Care Med* 2014; 42:747–749
36. Klein Klouwenberg PM, Ong DS, Bonten MJ, Cremer OL. Classification of sepsis, severe sepsis and septic shock: the impact of minor variations in data capture and definition of SIRS criteria. *Intensive Care Med* 2012; 38:811–819

37. Angus DC, Linde-Zwirble WT, Lidicker J, Clermont G, Carcillo J, Pinsky MR. Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care. *Crit Care Med* 2001; 29:1303–1310
38. Flaatten H. Epidemiology of sepsis in Norway in 1999. *Crit Care* 2004; 8:R180–184
39. Martin GS, Mannino DM, Eaton S, Moss M. The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med* 2003; 348:1546–1554
40. Wang HE, Shapiro NI, Angus DC, Yealy DM. National estimates of severe sepsis in United States emergency departments. *Crit Care Med* 2007; 35:1928–1936
41. Dombrovskiy VY, Martin AA, Sunderram J, Paz HL. Rapid increase in hospitalization and mortality rates for severe sepsis in the United States: a trend analysis from 1993 to 2003. *Crit Care Med* 2007; 35:1244–1250
42. Schneeweiss S, Robicsek A, Scranton R, Zuckerman D, Solomon DH. Veteran's affairs hospital discharge databases coded serious bacterial infections accurately. *J Clin Epidemiol* 2007; 60:397–409
43. Patkar NM, Curtis JR, Teng GG, Allison JJ, Saag M, Martin C, et al. Administrative codes combined with medical records based criteria accurately identified bacterial infections among rheumatoid arthritis patients. *J Clin Epidemiol* 2009; 62:321–327, 327 e321–327
44. Iwashyna TJ, Odden A, Rohde J, Bonham C, Kuhn L, Malani P, et al. Identifying patients with severe sepsis using administrative claims: patient-level validation of the angus implementation of the international consensus conference definition of severe sepsis. *Med Care* 2014; 52:e39–43
45. Sherman ER, Heydon KH, St John KH, Teszner E, Rettig SL, Alexander SK, et al. Administrative data fail to accurately identify cases of healthcare-associated infection. *Infect Control Hosp Epidemiol* 2006; 27:332–337
46. Grijalva CG, Chung CP, Stein CM, Gideon PS, Dyer SM, Mitchel EF Jr, et al. Computerized definitions showed high positive predictive values for identifying hospitalizations for congestive heart failure and selected infections in Medicaid enrollees with rheumatoid arthritis. *Pharmacoepidemiol Drug Saf* 2008; 17:890–895
47. Ibrahim I, Jacobs IG, Webb SA, Finn J. Accuracy of International classification of diseases, 10th revision codes for identifying severe sepsis in patients admitted from the emergency department. *Crit Care Resusc* 2012; 14:112–118
48. Gedeberg R, Furebring M, Michaelsson K. Diagnosis-dependent misclassification of infections using administrative data variably affected incidence and mortality estimates in ICU patients. *J Clin Epidemiol* 2007; 60:155–162
49. Stone PW, Horan TC, Shih HC, Mooney-Kane C, Larson E. Comparisons of health care-associated infections identification using two mechanisms for public reporting. *Am J Infect Control* 2007; 35:145–149
50. Ollendorf DA, Fendrick AM, Massey K, Williams GR, Oster G. Is sepsis accurately coded on hospital bills? *Value Health* 2002; 5:79–81
51. Poulouse JT, Cartin-Ceba R, Shoja A, Paul A, Trillo-Alvarez C, Kashyap R, et al. Comparison of International Classification of Disease-Ninth Revision (ICD-9) coding with retrospective case review for the diagnosis of septic shock. Conference: C56 Sepsis: From Bench to the Bedside. San Diego Convention Center. *Am J Respir Crit Care Med*, 2009, A4691
52. Henriksen DP, Nielsen SL, Laursen CB, Hallas J, Pedersen C, Lassen AT. How well do discharge diagnoses identify hospitalised patients with community-acquired infections?—a validation study. *PLoS One* 2014; 9:e92891
53. Barber C, Lacaille D, Fortin PR. Systematic review of validation studies of the use of administrative data to identify serious infections. *Arthritis Care Res (Hoboken)* 2013; 65:1343–1357
54. Schønheyder HC, Søgaard M. Existing data sources for clinical epidemiology: The North Denmark Bacteremia Research Database. *Clin Epidemiol* 2010; 2:171–178
55. Jung MA, Banerjee SN. Administrative coding data and health care-associated infections. *Clin Infect Dis* 2009; 49:949–955
56. Stevenson EK, Rubenstein AR, Radin GT, Wiener RS, Walkey AJ. Two decades of mortality trends among patients with severe sepsis: a comparative meta-analysis. *Crit Care Med* 2014; 42:625–631
57. Danish Health and Medicines Authority. Klassifikation af Sygdomme—Systematisk Del. 2nd ed. Copenhagen, Denmark: Munksgaard, 1993
58. Holst AG ICDPedia.dk—diagnosekoder 2.0. Available: <http://icdpedia.dk/> (Accessed 2014 November 12)

59. Goto M, Ohl ME, Schweizer ML, Perencevich EN. Accuracy of administrative code data for the surveillance of healthcare-associated infections: a systematic review and meta-analysis. *Clin Infect Dis* 2014; 58:688–696
60. Voldstedlund M, Haarh M, Mølbak K, MiBa Board of Representatives. The Danish Microbiology Database (MiBa) 2010 to 2013. *Euro Surveill* 2014; 19(1). pii: 20667
61. Skogberg K, Lyytikäinen O, Ollgren J, Nuorti JP, Ruutu P. Population-based burden of bloodstream infections in Finland. *Clin Microbiol Infect* 2012; 18:e170–176