



Mother-to-embryo vitellogenin transport in a viviparous teleost *Xenotoca eiseni*

Atsuo Iida^{a,b,1}, Hiroyuki N. Arai^b, Yumiko Someya^c, Mayu Inokuchi^c, Takeshi A. Onuma^d, Hayato Yokoi^e, Tohru Suzuki^e, Eiichi Hondo^a, and Kaori Sano^f

^aDepartment of Animal Sciences, Graduate School of Bioagricultural Sciences, Nagoya University, Furocho, Chikusa-ku, Nagoya 464-8601, Japan;

^bDepartment of Regeneration Science and Engineering, Institute for Frontier Life and Medical Sciences, Kyoto University, Shogo-in Kawahara-cho 53, Sakyo-ku, Kyoto 606-8507, Japan; ^cDepartment of Life Sciences, Toyo University, Itakura, Gunma 374-0113, Japan; ^dDepartment of Biological Sciences, Graduate School of Science, Osaka University, 1-1 Machikaneyama-cho, Toyonaka, Osaka 560-0043, Japan; ^eLaboratory of Marine Life Science and Genetics, Graduate School of Agricultural Science, Tohoku University, Sendai 980-8577, Japan; and ^fDepartment of Chemistry, Faculty of Science, Josai University, Sakado, Saitama 350-0295, Japan

Edited by John J. Eppig, The Jackson Laboratory, Bar Harbor, ME, and approved September 19, 2019 (received for review July 30, 2019)

Vitellogenin (Vtg), a yolk nutrient protein that is synthesized in the livers of female animals, and subsequently carried into the ovary, contributes to vitellogenesis in oviparous animals. Thus, Vtg levels are elevated during oogenesis. In contrast, Vtg proteins have been genetically lost in viviparous mammals, thus the yolk protein is not involved in their oogenesis and embryonic development. In this study, we identified Vtg protein in the livers of females during the gestation of the viviparous teleost, *Xenotoca eiseni*. Although vitellogenesis is arrested during gestation, biochemical assays revealed that Vtg protein was present in ovarian tissues and lumen fluid. The Vtg protein was also detected in the trophotaeniae of the intraovarian embryo. Immunoelectron microscopy revealed that Vtg protein is absorbed into intracellular vesicles in the epithelial cells of the trophotaeniae. Furthermore, extraneous Vtg protein injected into the abdominal cavity of a pregnant female was subsequently detected in the trophotaeniae of the intraovarian embryo. Our data suggest that the yolk protein is one of the matrotrophic factors supplied from the mother to the intraovarian embryo during gestation in *X. eiseni*.

Goodeidae | reproduction | viviparity | trophotaeniae

Viviparity is a form of reproductive system in which fertilization and embryonic development progress inside the mother's body prior to the birth, as opposed to oviparity where such developmental events are completed outside of the parent. In some viviparous vertebrates, the embryo receives nutrients from the mother in addition to those in the yolk sac. The nutrient supply and embryonic growth in the mother's body are important factors influencing survival in the habitat environment after the birth (1). In eutherian mammals, maternal nutrients including blood plasma pass into the fetus via a placenta and umbilical cords (2). Viviparous species also occur in reptiles, amphibians, and fish (3–5). The embryos of some amphibians and sharks eat each other and develop within the mother's body (6, 7). A viviparous reptile possesses a placenta-like structure similar to those found in eutherians (8). Some viviparous chondrichthyans raise their offspring in the uterus filled with a nutrient-rich liquid during gestation (9, 10). In viviparous teleosts, the mother fish maintains her embryo in the ovarian lumen or oviduct (5). Diverse mechanisms of embryo growth among viviparous vertebrates have been described. However, the nature of the maternal nutrients in viviparous teleosts has not been identified.

The teleost family Goodeidae is currently known to include 42 viviparous species distributed in the lakes and rivers of México (11). Their embryo increases in dry mass in the mother's body during the gestation, suggesting that maternal nutrients are supplied in addition to those in the yolk sac (5). In the mid to late gestation stages, the goodeid embryo possesses a pseudoplacenta, called the trophotaeniae. The trophotaeniae form through hypertrophic development of elongated processes that extend from the hindgut out through the cloaca and are thought to be involved

in nutrient incorporation during gestation. That consists of absorptive epithelium, blood vessels, and mesenchymal connective tissues (12, 13). Previous studies hypothesized that the ovarian fluid components are absorbed into the epithelial cells and then carried into the capillary (12, 14). In some goodeid species, the ovarian fluids include multiple protein components in a similar pattern to that in the blood serum (15, 16). In *Xenotoca eiseni*, the viviparous goodeid species used in this study, the intraovarian embryo develops for approximately 5 wk before birth. The embryo's growth depends on matrotrophic nutrition absorbed from the trophotaeniae for most of its intraovarian development (17–19). The trophotaeniae are regressed by apoptosis during the latest stage of gestation (20); they are no longer required in the postnatal growth phase when oral food intake occurs. The components of the maternal nutrients absorbed from the trophotaeniae have not been identified but are hypothesized to be secreted blood serum proteins.

Vitellogenin (Vtg) is a glycolipophosphoprotein typically present in females, but also in minor amounts in males, that is conserved in nearly all oviparous species including fish, amphibians, reptiles, birds, monotremes, and most invertebrates (21–24). In these vertebrates, *vtg* genes are typically expressed and synthesized in female liver, then the protein product is transported to the ovary via the bloodstream. In the ovary, Vtg protein is cleaved into subdomains and absorbed into the egg

Significance

Viviparity is a type of reproductive system in which the embryo utilizes a maternal nutrient supply until birth. Viviparous species occur in many taxa including bony fish. The mother fish raises her offspring in the ovarian lumen or oviduct during gestation. Embryos of the viviparous bony fish *Xenotoca eiseni* (family Goodeidae) utilize nutrients secreted into the ovarian lumen. However, the source of the maternal nutrients and their mother-to-embryo transport have not been experimentally demonstrated. In this study, we focus on the yolk nutrient protein vitellogenin (Vtg) as a matrotrophic factor. Our results are of fundamental importance to the investigation of viviparous systems in teleosts and in other vertebrates and invertebrates.

Author contributions: A.I. and K.S. designed research; A.I., H.N.A., Y.S., M.I., T.A.O., and E.H. performed research; H.Y., T.S., and K.S. contributed new reagents/analytic tools; A.I. analyzed data; and A.I. wrote the paper.

The authors declare no competing interest.

This article is a PNAS Direct Submission.

This open access article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

¹To whom correspondence may be addressed. Email: tol2.4682@gmail.com.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1913012116/-DCSupplemental.

First published October 8, 2019.

yolk (25, 26). In eutherian mammals, *vlg* genes have been genetically lost through coevolution with casein genes, thus few yolk nutrients are included in their eggs (22, 27). In contrast, some viviparous fish species have maintained *vlg* genes, and a large amount of yolk nutrient is contained in their eggs (28, 29). In these fish, Vtg protein is a potential candidate for one of the maternal nutrients supplied into the intraovarian embryo. However, there is no definitive evidence that maternal Vtg protein is transported into the intraovarian embryo. In this study, we investigated Vtg transport from mother to embryo in the goodeid species, *X. eiseni*, and our results suggest that the yolk protein lost in eutherian mammals is one of the matrotrophic factors supplied to the intraovarian embryo during gestation in the viviparous teleost.

Results

Production and Distribution of Vitellogenin Proteins during Gestation.

At the onset of gestation in goodeid species including *X. eiseni* (Fig. 1A), the main role of the ovary switches from egg production to maintenance of the embryo. During gestation, oogenesis is

arrested at previtellogenesis stages, thus there are no mature eggs in the ovarian lumen (Fig. 1B) and the ovary is dedicated to raising the embryo via nutrient supply. To investigate expression of vitellogenin genes (*vlgA*, *vlgB*, and *vlgC*) in *X. eiseni*, RT-PCR analysis was performed using cDNA purified from the liver and gonad. In vitellogenic females, but not in male fish, *vlgA* and *B* were strongly expressed in the liver. In contrast, there were no detectable signals in testis and ovary. Expression of *vlgC* was minimal in all tissues examined in this study (*SI Appendix, Fig. S1A*). Expression of the *vlg* genes in the liver was not decreased in pregnant females, despite the cessation of vitellogenesis (Fig. 1C). In the ovary, *vlg* genes were not expressed during gestation, as for the vitellogenesis stage (Fig. 1C and *SI Appendix, Fig. S1A*). VtgA and B proteins were detectable by immunohistochemistry in the nonpregnant female liver (*SI Appendix, Fig. S1B*). The strong signals for Vtg proteins in female liver were replicated using another antibody, anti-Vtg#1 (*SI Appendix, Fig. S2A* and *Table S1*). In contrast, specific signals for Vtg protein in male liver were weaker than those in female liver (*SI Appendix, Fig. S1C*). The strong signals for VtgA and B proteins were also observed in the




Fig. 1. Vitellogenin synthesis and supply to ovary during gestation. (A) Whole body photograph of adult *X. eiseni*. The adult male fish exhibits sex-specific characteristics, such as a high body arch and red color in the tail. (Scale bar, 10 mm.) (B) HE-stained transverse sections of vitellogenesis (Upper) and gestation (Lower) stages of an *X. eiseni* female ovary. In the nonpregnant ovary, mature eggs including yolk were observed in the ovarian lumen, and some postvitellogenic oocytes invaginate to the stroma in the ovarian luminal epithelium. In the pregnant ovary, developing embryos filled the ovarian lumen, and there were no mature eggs with yolk. Some immature oocytes can still be observed in the ovarian luminal epithelium. Arrowheads indicate immature oocytes. (Scale bar, 1 mm [whole ovary and embryo] and 200 μ m [immature egg].) (C) RT-PCR for *vlg* genes in female *X. eiseni*. All *vlg* genes were detected in the liver during vitellogenesis. The hepatic expression was also observed in the third week of gestation. No *vlg* expression was observed in any ovary samples. *Nicotinamide adenine dinucleotide (nadh)* was used as an internal control. M, size marker. Vit., vitellogenesis. Ges., gestation. (D) Fluorescence immunohistochemistry for VtgA or B in lobe edges of female liver in the third week of gestation. (Scale bar, 20 μ m.) (E) Fluorescence immunohistochemistry for VtgA or B in the ovarian septum in the third week of gestation. (Scale bar, 20 μ m.) See also *SI Appendix, Figs. S1 and S2*.




Fig. 3. Detection of vitellogenin fragments in ovarian fluid and trophotaeniae. (A) Electrophoresis and CBB staining of ovarian fluid and intraovarian embryo lysate extracted from pregnant female fish in the third week after mating. The ovarian fluid showed 2 major protein bands of 240- and 75-kDa (arrow and arrowhead). The 75-kDa band was also observed in the lysate from whole embryo and trophotaeniae. The 240-kDa protein was not detected in the embryo samples. Ova., ovarian. Vit., vitellogenesis. Ges., gestation. (B) Western blotting (using anti-Vtg#1 antibody) of ovarian fluid and intraovarian embryo lysate extracted from pregnant female fish in the third week after mating. The ovarian fluid indicated 2 major signals against Vtg at 240- and 75-kDa (arrow and arrowhead). In the gestation sample, 6 specific signals (240-, 100-, 75-, 55-, 50-, and 45-kDa) were observed as distinct bands. Magenta boxes indicate the signals against Vtg detected in the both ovarian fluid and trophotaeniae. Blue boxes indicate the signals detected only in ovarian fluid. GAPDH was used to check for contamination of cellular components into the ovarian fluid and nonspecific degradation of extracted proteins during the sample collection. See also *SI Appendix, Fig. S3*.

Fig. S3C). These results indicate that the 100-, 75-, and 50-kDa Vtg fragments are secreted into the ovarian fluid from the ovarian tissues without any changes in their molecular weight, and then absorbed into the embryo via the trophotaeniae.

Vitellogenin Transfer from Mother to Intraovarian Embryo. As an experimental verification the transfer of Vtg from mother to embryo, we performed a tracer assay using a fluorescence-labeled Vtg protein. We prepared a fluorescein isothiocyanate (FITC)-conjugated Vtg protein (FITC-Vtg) that originated from goldfish. The FITC-Vtg injected into the abdominal cavity of female fish was carried and secreted into the ovarian fluid, and then imported into the yolk (*SI Appendix, Fig. S4 A–C*). This indicated that the FITC-Vtg was functional for the mother-to-egg Vtg transfer. To validate a mother-to-embryo Vtg transfer, the tracer was injected into a pregnant female in the third week after mating, and then the intraovarian embryo was extracted and the trophotaeniae were observed using fluorescent microscopy (*Fig. 4A*). The fluorescent dye injected into the abdominal cavity of pregnant female fish exuded to the vascular lumen and was then visualized in the blood vessel network including the caudal fin (*Fig. 4B*). Thus, both the exogenous and endogenous Vtg proteins would be transported via the bloodstream. The

FITC fluorescence was accumulated in the digestive tract of the embryo (*SI Appendix, Fig. S4D*). Confocal microscopy indicated that the FITC fluorescence was also detected in the epithelial layer of the trophotaeniae in the embryo extracted from the FITC-Vtg injected female. In contrast, there were no signals in the PBS-injected control. Furthermore, the fluorescence was merged to immunohistochemistry signals against FITC (*Fig. 4C*). This means that the intraovarian embryo incorporated exogenous Vtg protein into the epithelial cells of the trophotaeniae from the mother's body without loss of the fluorescence and antigenicity of labeled FITC. We also confirmed that a FITC-conjugated dextran (M.W. 250,000) exuded to the vasculature and was absorbed into the trophotaeniae (*SI Appendix, Fig. S5*). These results indicated that maternal proteins and carbohydrates in interstitial fluid and blood serum could be transferred into the intraovarian embryo via the trophotaeniae (*Fig. 5*).

Discussion

In this study, we revealed Vtg synthesis and mother-to-embryo transfer during gestation in the goodeid viviparous teleost *X. eiseni* (*Fig. 5*). In some viviparous surferperches and eelpout, Vtg has been considered not a necessity for gestation, and the need for and roles of Vtg in gestation were mostly elusive in viviparous teleosts. A previous study reported that Vtg-like proteins did not appear in the serum of ovarian fluid during the gestation period of viviparous surferperches (30). Another study indicated that Vtg content in female serum is down-regulated during gestation in a viviparous eelpout (31). The eelpout exhibits a higher Vtg level during vitellogenesis than that during gestation. This suggests that Vtg is required for oogenesis, and that supply into the oocytes could be regulated according to the reproduction cycle. In the family Goodeidae, a previous study indicated that Vtg is not a maternal nutrient supplied to the intraovarian embryo throughout gestation in *Goodea atripinnis* and *Allophorobius robustus* (15). In contrast, other research groups argued that Vtg protein was utilized for intraovarian embryonic growth in *Girardinichthys viviparus* and *Ameiobela splendens*; however, definitive evidence of mother-to-embryo transfer of Vtg protein has not been demonstrated (29). In the present study, we revealed that Vtg gene expression and protein secretion are maintained at a high level during vitellogenesis and gestation in *X. eiseni*. Furthermore, the tracer analysis revealed that FITC-conjugated Vtg protein could be transferred into the trophotaeniae of the intraovarian embryo from the mother. The accumulation of the FITC-fluorescence in the digestive tract might indicate a destination of the Vtg protein or its degradants absorbed from the trophotaeniae. This evidence suggests that Vtg is one of the matrotrophic factors in *X. eiseni*.

We performed the Western blotting analysis to identify the Vtg proteins in the ovarian fluid and the intraovarian embryo by their molecular weight. The molecular weights of the intact VtgA and B proteins in *X. eiseni* were predicted to be ~190 kDa. Following synthesis, Vtg proteins undergo posttranslational modifications like glycosylation and phosphorylation, and subsequently lipidation (32). Thus, the actual weight of native Vtg proteins is difficult to predict. The CBB staining and Western blotting indicated that the 75-kDa Vtg fragment is the major protein secreted into the ovarian fluid, and our data revealed that the 100-, 75-, and 50-kDa Vtg fragments were absorbed into the trophotaeniae. Previous studies indicated that Vtg protein undergoes a specific proteolytic cleavage during uptake into an oocyte (25). Vertebrate Vtg protein is cleaved into the heavy (~120-kDa) and light (~45-kDa) chain lipovitellin, and phosvitin (~34-kDa) (26). We used polyclonal antibodies against whole Vtg sequence purified from blood plasma of 17 β -estradiol treated arctic char or sea bream. Thus, we could not identify a corresponding relationship between the signals and actual amino acid sequences of the cleaved Vtg fragments. Additionally, some




Fig. 4. Tracing for mother-to-embryo vitellogenin transfer in live fish. (A) A scheme for the tracing analysis. Mating: To prepare a pregnant female, a pair of mature *X. eiseni* were mated, and then the male fish was removed after mating. i.p. injection: At 3 wk after mating, fluorescent Vtg solution was injected into the abdominal cavity of the pregnant female. Extraction & observation: At 18 h after injection, the embryo was extracted from the female fish and observed under a fluorescent microscope. (B) Fluorescent microscopy for the injected fluorescent solution in the female fish. The solutions were injected into the abdominal cavity from the gravid spot of pregnant female fish (arrowhead). The blood vessels in the caudal fin were visualized by a green fluorescence for the FITC-Vtg. (Scale bar, 1 mm.) (C) Fluorescent microscopy of the intraovarian embryo. Green fluorescence was detected in the epithelial cell layer of trophotaeniae in the extracted embryo (arrowheads). The green fluorescence was merged to signals for anti-FITC (arrow). E, epithelial layer. M, mesenchyme. (Scale bar, 10 μ m.) See also *SI Appendix, Figs. S4 and S5.*

smaller bands and smear-like signals against the antibodies were observed in the embryo and trophotaeniae. A previous study indicated that Vtg protein in zebrafish embryo was digested and expended during development (33). Therefore, we suggest that the lower signals are Vtg fragments digested to be expended as a maternal nutrient.

Immunoelectron microscopy revealed that trophotaeniae epithelial cells import Vtg fragments via vesicle trafficking. Similar protein uptake by lysosomes or endosomes was previously detected by absorption assay of cationized ferritin (CF) or horseradish peroxidase (HRP) (12, 14). We also demonstrated that high-molecular weight dextran is absorbed into the trophotaeniae. These findings suggest that the epithelial layer of the trophotaeniae functions as an absorptive epithelium that imports maternal proteins and other components as macromolecules.

The trophotaeniae are partially derived from the hindgut and exhibit structural and functional characteristics of intestinal absorptive cells (12, 13). Similar endocytotic absorption of a macromolecule was previously reported in suckling rat absorptive cells (34–36). The suckling stage-specific absorption undergoes changes corresponding with the development of the digestive system (37). A recent study demonstrated dietary protein internalization via receptor-mediated and fluid-phase endocytosis in specialized intestinal cells of zebrafish and mice (38). Therefore, we hypothesize that the macromolecule absorption observed in the postembryonic stage of viviparous fish species is based on a common characteristic between the mammals and the teleostei. The mammals use this mechanism to assimilate macromolecules from the mother's milk, whereas in goodeid fish the trophotaeniae, which are modifications of the juvenile hindgut,




Fig. 5. A model for mother-to-embryo vitellogenin transfer during gestation. In the pregnant female, vitellogenin genes are strongly expressed in the liver, and then the proteins are carried to the ovary via the bloodstream. In the ovarian lumen, full-length and fragmented vitellogenin proteins are secreted into the ovarian fluid, and then absorbed into the embryo via the epithelium of the trophotaeniae. In the epithelium of the trophotaeniae, the fragmented vitellogenin proteins are imported into the epithelial cells and carried as macromolecules via vesicle trafficking.

are utilized for uptake of the intraovarian macromolecules supplied from the mother. To validate the hypothesis for macromolecule absorption, investigation and comparison of the molecular mechanisms of the absorption system in juvenile stages of mammals and viviparous teleosts is required, and also in other species.

Methods

Animal Experiments. This study was approved by the ethics review board for animal experiments of Kyoto University. We killed live animals under anesthesia in minimal numbers according to the institutional guidelines.

Fish Breeding. *X. eiseni* was purchased from Meito Suien Co., Ltd. (Nagoya, Japan). Adult fish were maintained in freshwater at 27 °C under a 14:10-h light: dark photoperiod cycle. Fish were bred in a mass-mating design, and ~50 fish were maintained for this study. The juveniles were fed live brine shrimp larvae and Hikari Rabo 450 fish food (Kyorin Co., Ltd., Himeji, Japan), and the adults were fed Hikari Crest Micro Pellets (Kyorin). To accurately track the pregnancy period, the laboratory-born fish were crossed in a pair-mating design, and the mating behavior was recorded.

Sample Collection. Fish samples were anesthetized using tricaine on ice, prior to surgical extraction of tissues or embryos. The obtained samples were stored on ice until the subsequent manipulations. In this study, we dissected ~20 adult fish and 30 pregnant females, and extracted 15–30 embryos in each operation.

Antibodies. Polyclonal antibodies against VtgA and B were generated in this study. The antigen sequences are aa 901–1,018 (VtgA, GenBank: AC130217.1) and aa 902–993 (VtgB, GenBank: AC130218.1). Inclusion bodies of the antigen peptides were harvested from the transformant *Escherichia coli* BL21 Star (DE3) (Thermo Fisher Scientific, Waltham, MA) (39). For determination of N-terminal sequences of the antigen protein, the inclusion bodies loaded onto Ni-NTA Superflow (Qiagen, Valencia, CA) were subjected to a protein sequencer (Procise 491HT; Applied Biosystems, Foster City, CA). Milligrams (0.2 mg) of each inclusion body diluted in urea/PBS were injected into Jcl:ICR mice (CLEA Japan, Inc., Meguro, Japan) 4 times every 2 wk. Serum samples were harvested from the antigen-injected mice. The sera, diluted to 50% in glycerol with 0.1% sodium azide, were used for immunohistochemistry. Information for the commercial antibodies used is listed in [SI Appendix, Table S1](#).

RT-PCR. Total RNA was extracted from tissues or whole embryo using the RNeasy Mini kit (Qiagen) and reverse-transcribed using SuperScript III reverse transcriptase (Thermo Fisher Scientific). PCR was carried out using KOD-FX (Toyobo, Osaka, Japan) under the following conditions: 100 s at 94 °C, followed by 28 cycles of 20 s at 94 °C, 20 s at 60 °C, and 20 s at 72 °C; and 40 s at 72 °C. Primer sequences are listed in [SI Appendix, Table S1](#).

Fluorescent Immunohistochemistry. Tissue samples were fixed in 4.0% paraformaldehyde/PBS (PFA/PBS) at 4 °C overnight. When using the anti-Vtg#1, fixed samples were incubated in 10 mM Tris-EDTA (pH 9.5) at 95 °C for 20 min as an antigen activation. The activation is not required for the other antibodies. Samples were permeabilized using 0.5% TritonX-100/PBS at room temperature for 30 min, and then treated with Blocking-One solution (Nacalai Tesque, Kyoto, Japan) at room temperature for 1 h. Primary antibodies were used at 1:100 (anti-FITC) or 1:500 (the others) dilution with

Blocking-One solution. Samples were reacted with primary antibodies at 4 °C overnight. Secondary antibodies ([SI Appendix, Table S1](#)) were used at 1:500 dilution in 0.1% Tween-20/PBS with DAPI (Sigma-Aldrich, St. Louis, MO). Samples were treated in the secondary antibody solution at 4 °C overnight. Microscopic observation was performed using Leica TCS SP8 microscopes (Leica Microsystems, Wetzlar, Germany).

Immunoelectron Microscopy. Embryo samples were fixed in 4.0% PFA/PBS. Fixed samples were washed in PBS and then incubated in 10 mM Tris-EDTA (pH 9.5) at 95 °C for 20 min as an antigen activation. Activated samples were reacted with primary antibody (anti-Vtg#1) at 4 °C overnight, and then reacted with biotinylated anti-rabbit IgG (Vector, Burlingame, CA) at room temperature for 2 h. Samples were performed with the avidin–biotin–peroxidase complex kit (Vector), and visualized with 0.05% 3,3'-diaminobenzidine (DAB; Dojindo Laboratories, Kumamoto, Japan) and 0.01% hydrogen peroxide in 50 mM Tris buffer (pH 7.2) at room temperature for 10 min. The stained samples were fixed in 1% osmium tetroxide in 0.1 M sodium phosphate buffer (pH 7.4) at room temperature for 1 h. The fixed samples were dehydrated in ethanol, transferred to propylene oxide, and embedded in Spurr's resin (Polysciences, Warrington, PA). Ultrathin sections were cut with a diamond knife every 70 nm and mounted on grids. The sections were stained with uranyl acetate and lead citrate, and examined with a transmission electron microscope (JEM2100, JEOL, Tokyo, Japan).

Western Blotting. Samples were washed with cold PBS and lysed in RIPA Buffer (10 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1 mM EDTA, 0.1% SDS, 1% Nonidet P-40, 0.1% sodium deoxycholate) containing 1% protease inhibitor mixture (Nacalai Tesque). The lysates were diluted by 50% in 2× SDS buffer (4% SDS, 20% glycerol, 0.002% bromophenol blue, 125 mM Tris-HCl, pH 6.8) containing 10% 2-mercaptoethanol (2-ME), and denatured at 95 °C for 5 min. The sample solutions were loaded and separated on 10% polyacrylamide gels, after which the proteins were transferred to polyvinylidene difluoride membranes (Immobilon-P; Millipore, Billerica, MA). After blocking with Blocking-One solution (Nacalai Tesque), the membranes were incubated at 4 °C overnight with primary antibodies (Vtg#1, 1:2,000, Vtg#2, 1:2,000, anti-GAPDH, 1:2,000). After washing, membranes were reacted with secondary antibody (anti-rabbit IgG, 1:5,000, Cell Signaling Technology, Danvers). The signals were visualized using ECL Prime Western Blotting Detection Reagent (GE Healthcare, Chicago, IL). Image data were acquired with an ImageQuant LAS 4000 (GE Healthcare).

Tracing of Mother-to-Embryo Transport. A FITC-conjugated goldfish Vtg protein was prepared according to the previous study (40). The FITC-Vtg, FITC-dextran (MW 250,000, Sigma-Aldrich) diluted in PBS (final conc. 1.0 mg/mL), or control solvent (PBS) was injected into the abdominal cavity of adult females under anesthesia. The injected females were incubated in a separate tank. For angiography of the injected dye, the female fish were observed under anesthesia using a Leica M205C microscope at 30 min postinjection. To investigate absorption into the ovary or intraovarian embryo, the sample was surgically harvested from the females under anesthesia at 18-, 42-, or 90-h postinjection. The extracted samples were observed using a Leica M205C, MZ16FA, or TCS-SP8 microscope.

ACKNOWLEDGMENTS. Breeding and experiments were performed in the laboratory of Dr. Atsuko Sehara-Fujisawa. We thank Yoshiko Aihara for helpful discussion. Ryo Kurokawa helped with the sample collection. This work was supported by crowdfunding via The Academist, Inc. The crowdfunding investors for this project were Nobuo Ishii, Kyoko Kawamura, Mayu Miyamoto,

Eiri Ono, Shoko Saito, Kohei Shibata, Tomosato Takabe, Yuma Nihata, Tokuyoshi Wakamatsu, Yoshinori Wakamatsu, Shoichi Tamura, Keiko Grace

Kobori, Tsuneaki Hasekura, and the Patchwork Club of Fujimida Elementary School (Nagoya, Japan).

- D. G. Blackburn, "Viviparity and oviparity: Evolution and reproductive strategies" in *Encyclopedia of Reproduction*, T. E. Knobil, J. D. Neill, Eds. (Academic Press, 1999), vol. 4, pp. 994–1003.
- A. W. Bell, R. A. Ehrhardt, Regulation of placental nutrient transport and implications for fetal growth. *Nutr. Res. Rev.* **15**, 211–230 (2002).
- J. U. Van Dyke, M. C. Brandley, M. B. Thompson, The evolution of viviparity: Molecular and genomic data from squamate reptiles advance understanding of live birth in amniotes. *Reproduction* **147**, R15–R26 (2013).
- M. H. Wake, Fetal adaptations for viviparity in amphibians. *J. Morphol.* **276**, 941–960 (2015).
- J. P. Wourms, Viviparity: The maternal-fetal relationship in fishes. *Am. Zool.* **21**, 473–515 (1981).
- D. Buckley, M. Alcobendas, M. García-París, M. H. Wake, Heterochrony, cannibalism, and the evolution of viviparity in *Salamandra salamandra*. *Evol. Dev.* **9**, 105–115 (2007).
- J. P. Wourms, Reproduction and development in chondrichthyan fishes. *Am. Zool.* **17**, 379–410 (1977).
- D. G. Blackburn, A. F. Flemming, Invasive implantation and intimate placental associations in a placentotrophic African lizard, *Trachylepis ivensi* (scincidae). *J. Morphol.* **273**, 137–159 (2012).
- J. I. Castro, K. Sato, A. B. Bodine, A novel mode of embryonic nutrition in the tiger shark, *Galeocerdo cuvier*. *Mar. Biol. Res.* **12**, 200–205 (2016).
- K. Sato et al., How great white sharks nourish their embryos to a large size: Evidence of lipid histology in lamnid shark reproduction. *Biol. Open* **5**, 1211–1215 (2016).
- K. L. Foster, K. R. Piller, Disentangling the drivers of diversification in an imperiled group of freshwater fishes (Cyprinodontiformes: Goodeidae). *BMC Evol. Biol.* **18**, 116 (2018).
- J. Lombardi, J. P. Wourms, The trophotaenial placenta of a viviparous goodeid fish. III: Protein uptake by trophotaeniae, the embryonic component. *J. Exp. Zool.* **236**, 165–179 (1985).
- J. P. Wourms, J. Lombardi, Reflections on the evolution of piscine viviparity. *Am. Zool.* **32**, 276–293 (1992).
- L. Grosse-Wichtrup, H. Greven, Uptake of ferritin by the trophotaeniae of the goodeid fish, *Ameia splendens* Miller and Fitzsimons 1971 (Teleostei, Cyprinodontiformes). *Cytobios* **42**, 33–40 (1985).
- F. Hollenberg, J. P. Wourms, Embryonic growth and maternal nutrient sources in goodeid fishes (Teleostei: Cyprinodontiformes). *J. Exp. Zool.* **271**, 379–394 (1995).
- J. F. Schindler, Structure and function of placental exchange surfaces in goodeid fishes (Teleostei: Atheriniformes). *J. Morphol.* **276**, 991–1003 (2015).
- G. Mendoza, The ovary and anal processes of "*Characodon*" *eiseni*, a viviparous cyprinodont teleost from Mexico. *Biol. Bull.* **129**, 303–315 (1965).
- G. Mendoza, The fine structure of an absorptive epithelium in a viviparous teleost. *J. Morphol.* **136**, 109–115 (1972).
- M. C. Uribe, H. J. Grier, S. A. Avila-Zúñiga, A. García-Alarcón, Change of lecithotrophic to matrotrophic nutrition during gestation in the viviparous teleost *Xenotoca eiseni* (Goodeidae). *J. Morphol.* **279**, 1336–1345 (2018).
- A. Iida, T. Nishimaki, A. Seihara-Fujisawa, Prenatal regression of the trophotaenial placenta in a viviparous fish, *Xenotoca eiseni*. *Sci. Rep.* **5**, 7855 (2015).
- A. Canapa et al., Vitellogenin gene expression in males of the Antarctic fish *Trematomus bernacchii* from Terra Nova Bay (Ross Sea): A role for environmental cadmium? *Chemosphere* **66**, 1270–1277 (2007).
- R. Robinson, For mammals, loss of yolk and gain of milk went hand in hand. *PLoS Biol.* **6**, e77 (2008).
- F. Engelmann, Insect vitellogenin: Identification, biosynthesis, and role in vitellogenesis. *Adv. Insect. Phys.* **14**, 49–108 (1979).
- G. Jubeaux et al., Vitellogenin-like proteins among invertebrate species diversity: Potential of proteomic mass spectrometry for biomarker development. *Environ. Sci. Technol.* **46**, 6315–6323 (2012).
- R. A. Wallace, "Vitellogenesis and oocyte growth in nonmammalian vertebrates" in *Oogenesis*, L. W. Browder, Ed. (Springer, 1985), pp. 127–177.
- B. M. Byrne, M. Gruber, G. Ab, The evolution of egg yolk proteins. *Prog. Biophys. Mol. Biol.* **53**, 33–69 (1989).
- D. Brawand, W. Wahli, H. Kaessmann, Loss of egg yolk genes in mammals and the origin of lactation and placentation. *PLoS Biol.* **6**, e63 (2008).
- S. Sawaguchi et al., Multiple vitellogenins (Vgs) in mosquitofish (*Gambusia affinis*): Identification and characterization of three functional Vg genes and their circulating and yolk protein products. *Biol. Reprod.* **72**, 1045–1060 (2005).
- A. Vega-López et al., The role of vitellogenin during gestation of *Girardinichthys viviparus* and *Ameia splendens*; two goodeid fish with matrotrophic viviparity. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **147**, 731–742 (2007).
- V. Devlaming et al., Aspects of embryo nutrition and excretion among viviparous embiotocid teleosts: Potential endocrine involvements. *Comp. Biochem. Physiol. Part A Physiol.* **76**, 189–198 (1983).
- Y. Koya, T. Matsubara, T. Ikeuchi, S. Adachi, K. Yamauchi, Annual changes in serum vitellogenin concentrations in viviparous eelpout, *Zoarces elongatus*. *Comp. Biochem. Physiol. A Physiol.* **118**, 1217–1223 (1997).
- R. N. Finn, Vertebrate yolk complexes and the functional implications of phosphovitins and other subdomains in vitellogenins. *Biol. Reprod.* **76**, 926–935 (2007).
- U. Gündel, D. Benndorf, M. von Bergen, R. Altenburger, E. Küster, Vitellogenin cleavage products as indicators for toxic stress in zebra fish embryos: A proteomic approach. *Proteomics* **7**, 4541–4554 (2007).
- M. Fujita, F. Reinhart, M. Neutra, Convergence of apical and basolateral endocytic pathways at apical late endosomes in absorptive cells of suckling rat ileum in vivo. *J. Cell Sci.* **97**, 385–394 (1990).
- R. Baba, R. Tanaka, M. Fujita, M. Miyoshi, Cellular differentiation of absorptive cells in the neonatal rat colon: An electron microscopic study. *Med. Electron Microsc.* **32**, 105–113 (1999).
- R. Baba, M. Fujita, C. E. Tein, M. Miyoshi, Endocytosis by absorptive cells in the middle segment of the suckling rat small intestine. *Anat. Sci. Int.* **77**, 117–123 (2002).
- R. Baba, M. Yamami, Y. Sakuma, M. Fujita, S. Fujimoto, Relationship between glucose transporter and changes in the absorptive system in small intestinal absorptive cells during the weaning process. *Med. Mol. Morphol.* **38**, 47–53 (2005).
- J. Park et al., Lysosome-rich enterocytes mediate protein absorption in the vertebrate gut. *Dev. Cell*, S1534–5807(19)30661–6 (2019).
- K. Sano et al., Purification and characterization of zebrafish hatching enzyme—An evolutionary aspect of the mechanism of egg envelope digestion. *FEBS J.* **275**, 5934–5946 (2008).
- Y. Matsuda, Y. Ito, H. Hashimoto, H. Yokoi, T. Suzuki, Detection of vitellogenin incorporation into zebrafish oocytes by FITC fluorescence. *Reprod. Biol. Endocrinol.* **9**, 45 (2011).