

# BMJ Open Impact of aeroplane noise on mental and physical health: a quasi-experimental analysis

Scarlett Sijia Wang <sup>1</sup>, Sherry Glied,<sup>1</sup> Sharifa Williams,<sup>2</sup> Brian Will,<sup>3</sup> Peter Alexander Muennig<sup>4</sup>

**To cite:** Wang SS, Glied S, Williams S, *et al.* Impact of aeroplane noise on mental and physical health: a quasi-experimental analysis. *BMJ Open* 2022;**12**:e057209. doi:10.1136/bmjopen-2021-057209

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-057209>).

Brian Will deceased

Received 13 September 2021  
Accepted 21 February 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Robert F. Wagner Graduate School of Public Service, New York University, New York, New York, USA

<sup>2</sup>Center for Research on Cultural and Structural Equity in Behavioral Health, Nathan S Kline Institute for Psychiatric Research, Orangeburg, New York, USA

<sup>3</sup>Queens Quiet Skies, New York, New York, USA

<sup>4</sup>Health Policy and Management, Columbia University Mailman School of Public Health, New York, New York, USA

## Correspondence to

Scarlett Sijia Wang;  
[scarlett@nyu.edu](mailto:scarlett@nyu.edu)

## ABSTRACT

**Objectives** Historically, departures at New York City's LaGuardia airport flew over a large sports complex within a park. During the US Open tennis games, flights were diverted to fly over a heavily populated foreign-born neighbourhood for roughly 2 weeks out of the year so that the tennis match was not disturbed (the 'TNNIS' departure). In 2012, the use of the TNNIS departure became year-round to better optimise flight patterns around the metropolitan area.

**Methods** We exploited exogenously induced spatial and temporal variation in flight patterns to examine difference-in-difference effects of this new exposure to aircraft noise on the health of individual residents in the community relative to individuals residing within a demographically similar community that was not impacted. We used individual-level Medicaid records, focusing on conditions associated with noise: sleep disturbance, psychological stress, mental illness, substance use, and cardiovascular disease.

**Results** We found that increased exposure to aeroplane noise was associated with a significant increase in insomnia across all age groups, but particularly in children ages 5–17 (OR=1.64, 95% CI=1.12 to 2.39). Cardiovascular disease increased significantly both among 18–44-year-old (OR=1.45, 95% CI=1.41 to 1.49) and 45–64-year-old Medicaid recipients (OR=1.15, 95% CI=1.07 to 1.25). Substance use and mental health-related emergency department visits also increased. For ages 5–17, rate ratio (RR) was 4.11 (95% CI=3.28 to 5.16); for ages 18–44, RR was 2.46 (95% CI=2.20 to 2.76); and for ages 45–64, RR was 1.48 (95% CI=1.31 to 1.67).

**Conclusion** We find that increased exposure to aeroplane noise was associated with an increase in diagnosis of cardiovascular disease, substance use/mental health emergencies and insomnia among local residents.

## INTRODUCTION

LaGuardia's airspace originally utilised departures over areas that were less populated, such as waterways, parks or areas with warehouses or manufacturing.<sup>1</sup> However, as air traffic increased over time, the airspace used for traditional routes of arrivals and departures became crowded and conflicted with that of a nearby airport, John F. Kennedy.<sup>2</sup> As with LaGuardia, other airports sometimes

## Strengths and limitations of this study

- We used a quasiexperimental design to study before and after impacts of a flight pattern change in two matched zip code clusters within New York City (a difference-in-difference design).
- We used a large health insurance claims database that allowed us to capture diagnoses for most residents in both impacted and unimpacted zip code clusters.
- Despite the difference-in-difference design, it is possible that participants self-segregated after the increase in aircraft noise or that other unmeasured factors influenced the observed outcomes.
- We were unable to compute a dose–response relationship due to the use of aggregated noise data.
- We find that a sudden and dramatic change in aircraft noise was associated with increased diagnoses of insomnia, cardiovascular disease, substance abuse and mental illness.

manage increases in traffic by optimising flight patterns with less regard for the populations on the ground.<sup>2</sup> Almost invariably, these new flight patterns require routing aircraft over populated areas that were not previously exposed to aircraft noise.

Noise, and aircraft noise in particular, is associated with a number of health problems, particularly sleep disturbances, mental illness and substance use.<sup>3–8</sup> The sleep disturbances and psychological stress associated with aircraft noise are, in turn, thought to produce a cascade of biological effects that result in premature ageing via endocrinologic changes.<sup>9–14</sup>

Noise is thought to produce stress by activating the central nervous system and by interfering with sleep.<sup>3 6 8 15 16</sup> This stress produces predictable changes in biochemical pathways in human and animal studies that accelerate the rate of ageing.<sup>14 17 18</sup> This accelerated ageing process has been linked to the premature onset of age-related diseases, including cardiovascular disease.<sup>9 19 20</sup>

While the pathways linking poor sleep and psychological stress to premature ageing and chronic disease are understood, few studies have experimentally examined interventions that alter noise exposure in human populations.<sup>21</sup> Most of our knowledge about the health impact of aircraft noise in humans is based on associational studies.<sup>7</sup> These studies suffer from a number of limitations. On one hand, people who live near airports may self-select, such that those who are less sensitive to noise can take advantage of lower home prices on purchases or rentals for homes.<sup>13 19 20</sup> On the other hand, those who live near airports tend to have lower than average income, a major risk factor for premature disease and death.<sup>19 22–24</sup> There is limited evidence based on the impact of aircraft noise on premature ageing and health based on experimental or quasiexperimental analysis.<sup>12 13 23 25</sup>

Flight pattern changes afford a unique opportunity for studying the health impact of aircraft noise in humans. In the past decade, flight patterns have shifted, and these shifts have increasingly been accompanied by resident complaints.<sup>26</sup> As they do so, it becomes possible to identify areas that are impacted by new aircraft noise. In general, the point of maximum noise from an aircraft happens immediately after take-off as the aircraft is on full power. This is the for the experimental group in our study.

We conducted a longitudinal case/control study of one well-documented flight pattern change in New York City. LaGuardia Airport (LGA) is one of three major airports in the greater New York City area. One of its departure patterns utilises Flushing Meadows Park, a route known as the ‘Whitestone Climb.’<sup>24</sup> Because it is over greenspace, the Whitestone Climb has little impact on humans living in nearby dwellings. However, this park is also the location of the US Open Tennis match. During games, a seldom-used departure route called the ‘Flushing Climb’ (now called TNNIS) was used so that the tennis games were not disturbed by aircraft noise. The TNNIS climb routes aircraft over densely populated Flushing, Queens, greatly increasing the exposure of residents to noise on the ground.<sup>24</sup>

A Freedom of Information Act (FOIA) request by a local group opposed to aeroplane noise in Queens demonstrated that flight patterns using the TNNIS climb have increased to year-round use since 2012. Before that time, the use of the TNNIS climb was rare outside of the US Open.<sup>27</sup>

Previous work found that the year-round use of the TNNIS climb was costly, both in terms of money and lives.<sup>24</sup> However, this economic analysis was primarily based on associational data. Using data on flight patterns over Flushing obtained using the FOIA as well as Medicaid utilisation data, we conducted a quasiexperimental analysis of the health impacts of the aeroplane noise associated with this new route. In the USA, Medicaid is a safety-net health insurance programme for the low-income population. In New York State, over five million low-income individuals enrolled in the Medicaid program in 2012.

## METHODS

### Data

The data used in this study are New York City Medicaid claims prepared by the New York University Health Evaluation and Analytics Lab. The data include Medicaid member demographic information, address history, eligibility, medical services, and diagnostic information. The database consists of Medicaid fee for service claims and managed care encounters; both are comparable in quality.<sup>28</sup>

### A priori specifications and hypotheses


We hypothesised that exposure to aeroplane noise would increase healthcare utilisation, insomnia, mood disorder, anxiety, depression and cardiovascular disease depending on the age group.<sup>3 6 8 9 12 13</sup> Specifically, exposure to aeroplane noise would produce sleep disorders across all age groups,<sup>29</sup> would lead to emotional or behavioural disturbances including substance abuse, mood disorder, depression and developmental disorders among young adults aged 18–45 years who tend to be more at risk of these stress-associated disorders,<sup>30</sup> and would produce or exacerbate cardiovascular disease among older adults aged 45 and over when heart disease begins to increase in prevalence.<sup>31</sup> Noise studies suggest wide-ranging psychoneuroendocrinological effects (allostatic load) producing hypertension, hyperglycaemic and hypercholesterolaemia.<sup>3 6 8 9 12 13 30 31</sup> These biological changes are linked to cardiovascular disease, a correlate of exposure to aeroplane noise as well as other forms of nighttime noise.<sup>7 10 11 32</sup>

### Study design

We used individual-level data at the member-cohort level for the analysis. We selected samples of Medicaid members residing in each of the two neighbourhoods at two points in time. The pre-cohort was defined as Medicaid recipients living in the study neighbourhoods between 2009–2011 (pre-cohort) and 2013–2016 (post-cohort). About 64% of the Flushing pre-cohort and 63% of the Sunset Park pre-cohort were also in the post-cohort. We used the difference-in-difference models to analyse the results.

### Exposure

To determine exposure, we used data extracted under a FOIA request for flight patterns over Flushing, Queens, New York and from Part 150 Study Technical Advisory Committee Meeting No. 8 documents.<sup>33</sup> These documents were derived from a 2014 study conducted and funded by the Port Authority of NY & NJ (Port Authority) in partnership with the Federal Aviation Administration (FAA). In these documents, the Port Authority presents estimated noise exposure in geographies surrounding LGA. Exposure is quantified using the Integrated Noise Model in DNL (day-night average sound level) units. We also visually inspected changes in sound related to aircraft flight over sound monitors on the ground in Flushing using Flight Aware, a publicly available flight-tracking website and visited the area.<sup>34</sup> This was done to ensure that the estimates from the Port Authority had face validity.



**Figure 1** Intensity of noise exposure over a 24-hour period by census tract. The DNL 55 corridor (outer line) is demarcated separately from the DNL 60 corridor (inner line). DNL, day night level.

These geographic regions or corridors were stratified according to intensity of noise exposure over a 24-hour period. We divided census tracts into >60DNL, 55–60 DNL and <55DNL (figure 1). We aggregated census tracts and for regions of Flushing, Queens with noise exposure levels of 55 DNL or greater after 2012.<sup>19</sup> These tracts after 2012 are therefore identified as the treatment condition in this quasi-experimental analysis.

Flushing, Queens is a vibrant, predominantly immigrant neighbourhood.<sup>24</sup> It is increasingly populated by Asians immigrants, particularly those of Chinese descent. The English proficiency in 2018 was 49%, and the population was 54% Asian and 26% white. While the neighbourhood ranked as one of the poorest, the rates of education are higher than average and the rates of crime, obesity and hypertension are much lower than New York City as a whole.<sup>24</sup>

Sunset Park in Brooklyn, New York was identified as an appropriate control neighbourhood as the neighbourhood (1) did not experience a change in exposure to aircraft noise after 2012 when the TNNIS climb began frequent use, and (2) is similar to Flushing with respect to the distribution of sociodemographic and economic characteristics.<sup>35 36</sup> Like Flushing, Sunset Park is increasingly populated by those of Chinese descent with 32% of the population identifying as Asian and 23% identifying as white. About 48% of the residents were born outside of the USA and the English proficiency in 2018 was 51%.<sup>25</sup> Sunset Park also has high poverty rates with relatively low rates of crime, obesity and hypertension, and high levels of education relative to New York City as a whole.<sup>24</sup> Census tracts in Sunset Park were matched to those identified in Flushing based on race, foreign-born status, and age distribution.

### Key outcomes

We used International Classification for Disease revision (ICD-9 and ICD-10) codes as well as Clinical Classifications Software (CCS) codes for diagnostic groups to identify the following conditions of interest: insomnia (ICD-9=3270, 78052 or ICD-10=G470), cardiovascular disease (CCS=109–113), alcohol use disorder (CCS=660), substance use disorder (CCS=661), anxiety (CCS=651), depression (ICD-9=311 or ICD-10=F33), mood disorder (CCS=657) and disorders usually diagnosed in infancy, childhood adolescence (CCS=655), which includes autism, childhood emotional disorder and separation anxiety.

We linked census tracts of Flushing and Sunset Park to geocoded Medicaid addresses. If a recipient had a Medicaid-registered address within a given census tract, they were assigned to that census tract. Participants were excluded if they had invalid addresses, dual Medicare status, did not have a valid date of birth or were not officially enrolled in Medicaid during the study period (table 1). Participant samples were then defined as Medicaid recipients in the period 2009–2011 (pre-implementation period) and 2013–2015 (the TNNIS use period) and who resided within census tracts in Flushing and Sunset Park.

For these identified records, indicator variables were created to identify type of medical claims inpatient, outpatient and emergency department (ED) visits as well as prescription drugs, both overall and for visits related to substance use and mental health disorders (650–663, 670). We additionally obtained information on the age of the subscriber associated with each record. Because we did not have access to Medicare records, and did not include dual eligible participants due to the high likelihood of



**Table 1** Demographic characteristics and outcome prevalence and rates for Flushing, New York and Sunset Park, New York before (earlier than 2012) and after (after 2013) aeroplane noise increased in Flushing, New York

| Baseline characteristics                    | Post period: 2013–2015 |             |  |           |             |          |             |          |             |          |             |          |             |          |             |           |             |  |
|---|------------------------|-------------|--|-----------|-------------|----------|-------------|----------|-------------|----------|-------------|----------|-------------|----------|-------------|-----------|-------------|--|
|   | Pre period: 2009–2011  |             |  | Age 18–44 |             |          | Age 45–64   |          |             | Age 5–17 |             |          | Age 18–44   |          |             | Age 45–64 |             |  |
|   | Flushing               | Sunset Park |  | Flushing  | Sunset Park | Flushing | Sunset Park | Flushing | Sunset Park | Flushing | Sunset Park | Flushing | Sunset Park | Flushing | Sunset Park | Flushing  | Sunset Park |  |
| <b>Demographics</b>                         |                        |             |  |           |             |          |             |          |             |          |             |          |             |          |             |           |             |  |
| Total N                                     | 20120                  | 21597       |  | 57089     | 52016       | 36472    | 18681       | 24552    | 26009       | 76278    | 60774       | 50806    | 24421       |          |             |           |             |  |
| Age (mean)                                  | 11.69                  | 11.12       |  | 31.07     | 29.86       | 53.44    | 53.29       | 11.43    | 10.71       | 30.95    | 30.35       | 53.99    | 54.22       |          |             |           |             |  |
| Age (SD)                                    | 3.94                   | 3.92        |  | 8.19      | 7.54        | 5.59     | 5.5         | 3.94     | 3.87        | 7.84     | 7.25        | 5.44     | 5.57        |          |             |           |             |  |
| Female (%)                                  | 48                     | 48          |  | 58        | 57          | 54       | 51          | 48       | 48          | 56       | 54          | 54       | 52          |          |             |           |             |  |
| Asian (%)                                   | 50                     | 46          |  | 60        | 62          | 63       | 60          | 52       | 47          | 59       | 59          | 63       | 62          |          |             |           |             |  |
| Black (%)                                   | 6                      | 2           |  | 5         | 1           | 4        | 2           | 4        | 1           | 3        | 1           | 3        | 1           |          |             |           |             |  |
| Hispanic (%)                                | 17                     | 15          |  | 11        | 9           | 11       | 11          | 14       | 14          | 7        | 7           | 7        | 8           |          |             |           |             |  |
| White (%)                                   | 11                     | 24          |  | 10        | 15          | 10       | 14          | 10       | 24          | 8        | 14          | 8        | 12          |          |             |           |             |  |
| Other (%)                                   | 5                      | 3           |  | 5         | 4           | 6        | 7           | 5        | 3           | 4        | 3           | 5        | 6           |          |             |           |             |  |
| Unknown (%)                                 | 12                     | 11          |  | 10        | 8           | 6        | 5           | 14       | 12          | 20       | 16          | 13       | 11          |          |             |           |             |  |
| Average months on Medicaid per year         | 9                      | 10          |  | 8         | 8           | 9        | 10          | 9        | 10          | 8        | 8           | 9        | 9           |          |             |           |             |  |
| Total Medicaid spending per person per year | US\$1911               | US\$1904    |  | US\$3818  | US\$3954    | US\$6754 | US\$6076    | US\$1783 | US\$1972    | US\$3398 | US\$3914    | US\$6520 | US\$6115    |          |             |           |             |  |
| <b>Prevalence per 100000</b>                |                        |             |  |           |             |          |             |          |             |          |             |          |             |          |             |           |             |  |
| Insomnia                                    | 398                    | 477         |  | 4208      | 6096        | 8036     | 9143        | 623      | 450         | 4755     | 5873        | 11034    | 10843       |          |             |           |             |  |
| Cardiovascular disease*                     | NA*                    | NA*         |  | 1955      | 1576        | 9934     | 9073        | NA*      | NA*         | 3575     | 2040        | 13260    | 10786       |          |             |           |             |  |
| Alcohol use disorder                        | NA*                    | NA*         |  | 2114      | 1173        | 2470     | 2184        | NA*      | NA*         | 2264     | 1358        | 2870     | 2199        |          |             |           |             |  |
| Substance use disorder                      | NA*                    | NA*         |  | 2265      | 1517        | 1799     | 2098        | NA*      | NA*         | 3799     | 2926        | 4250     | 4058        |          |             |           |             |  |
| Anxiety                                     | NA*                    | NA*         |  | 5124      | 4639        | 6279     | 6279        | NA*      | NA*         | 5726     | 5265        | 7537     | 7416        |          |             |           |             |  |
| Depression                                  | NA*                    | NA*         |  | 3782      | 2874        | 6007     | 5867        | NA*      | NA*         | 3191     | 2272        | 5637     | 4656        |          |             |           |             |  |
| Mood disorder                               | NA*                    | NA*         |  | 6371      | 4900        | 9399     | 8891        | NA*      | NA*         | 5607     | 4410        | 8375     | 7297        |          |             |           |             |  |
| Disorders diagnosed young                   | 1983                   | 1394        |  | 289       | 212         | 170      | 112         | 2480     | 2219        | 307      | 244         | 163      | 188         |          |             |           |             |  |

Continued

Table 1 Continued

| Baseline characteristics              | Pre period: 2009–2011 |             |           |             |           |             | Post period: 2013–2015 |             |           |             |           |             |
|---------------------------------------|-----------------------|-------------|-----------|-------------|-----------|-------------|------------------------|-------------|-----------|-------------|-----------|-------------|
|                                       | Age 5–17              |             | Age 18–44 |             | Age 45–64 |             | Age 5–17               |             | Age 18–44 |             | Age 45–64 |             |
|                                       | Flushing              | Sunset Park | Flushing  | Sunset Park | Flushing  | Sunset Park | Flushing               | Sunset Park | Flushing  | Sunset Park | Flushing  | Sunset Park |
| Visits per 1000 per year              |                       |             |           |             |           |             |                        |             |           |             |           |             |
| Emergency department                  | 328                   | 216         | 335       | 257         | 288       | 237         | 375                    | 188         | 360       | 217         | 332       | 216         |
| Emergency department (SM)             | 13                    | 20          | 26        | 21          | 31        | 39          | 20                     | 7           | 32        | 12          | 45        | 36          |
| Inpatient stays                       | 70                    | 53          | 267       | 319         | 299       | 245         | 60                     | 49          | 231       | 300         | 234       | 190         |
| Inpatient stays (SM)                  | 14                    | 7           | 43        | 24          | 45        | 32          | 11                     | 5           | 37        | 21          | 37        | 21          |
| Outpatient visits per person per year |                       |             |           |             |           |             |                        |             |           |             |           |             |
| Total outpatient                      | 3.4                   | 4.1         | 4.1       | 4.6         | 6.7       | 7.3         | 3.9                    | 5.2         | 4.5       | 5.5         | 7.6       | 8.2         |
| Outpatient (SM)                       | 0.2                   | 0.2         | 0.3       | 0.2         | 0.4       | 0.3         | 0.3                    | 0.3         | 0.3       | 0.3         | 0.5       | 0.4         |

\*We adopted a longer follow-up period 2008–2011 (preimplementation) and 2013–2016 (TNNIS implementation) to allow for lag time in disease manifestation.

pre-existing medical conditions and smaller sample size, our sample does not include adults aged 65 or older. Age in years was defined as the calculated age on 1 January 2009 and 1 January 2013, and stratified into three age cohorts, 5–17, 18–44 and 45–64 years.

### Statistical analyses

Our focus is on the rate of diagnoses for the hypothesised conditions. We first assess whether there were significant changes in utilisation overall between the baseline and TNNIS use periods and whether the observed changes differed by neighbourhood (ie, exposure) after considering other changes over time between these neighbourhoods. We use Poisson regression (see equation 1) to model the number of overall and substance use and mental health-related inpatient, ED and outpatient visits for those months in which participants were enrolled in Medicaid.

For our primary analyses, we use logistic regression (see equation 2) to examine the odds of receiving a diagnosis for the hypothesised conditions. Before implementing these regression analyses, we examined trends in socio-demographic characteristics as well as trends in Medicaid enrolment to ensure that no divergent patterns were noted around 2012. Because racial composition varied somewhat between the two neighbourhoods (table 1), we controlled for race in our analyses to ensure that compositional changes by race did not influence the analysis. We also stratified by age so that we could better test our a priori hypotheses by condition. For chronic diseases, that is, cardiovascular disease, we adopted a longer follow-up period 2008–2011 (preimplementation) and 2013–2016 (TNNIS implementation) to allow for lag time in disease manifestation.

$$\log(E(Y|x)) = \alpha + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_1 x_2 + \beta_4 x_3 \quad (1)$$

where Y=number of Medicaid claims for condition of interest and offset=number of Medicaid enrolment month.

$$\log\left(\frac{p}{1-p}\right) = \alpha + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_1 x_2 + \beta_4 x_3, \quad (2)$$

where p= Pr left (Y=1 right) is the probability of having Medicaid claim for condition of interest

Here,  $x_1$  was the indicator for neighbourhood exposure condition (Sunset Park=0 vs Flushing=1);  $x_2$  indicated implementation period (preimplementation=0 vs TNNIS implementation=1); and  $x_3$  race/ethnicity group membership (Asian=1, black=2, Hispanic=3, white=4 (reference), other=5, nknown=6).

### Patient and public involvement

The research question was inspired by the work of a non-profit community organisation called Queens Quiet Skies that works to mitigate aeroplane noise. One of the coauthors of the paper was a member of this organisation and obtained the FOIA requests for FAA documents. These documents were used to identify the treatment

**Table 2** Model results and 95% CI

|                             | Rate ratios from the difference-in-difference Poisson model  |                     |                     |
|-----------------------------|--|---------------------|---------------------|
|                             | Age 5–17   | Age 18–44           | Age 45–64           |
| Inpatient visits            | 0.92 (0.83 to 1.03)  | 1.05 (1.02 to 1.08) | 0.93 (0.88 to 0.97) |
| Emergency department visits | 1.31 (1.24 to 1.37)  | 1.45 (1.41 to 1.49) | 1.16 (1.11 to 1.21) |
| Outpatient visits           | 0.86 (0.85 to 0.87)  | 1.04 (1.04 to 1.05) | 0.92 (0.92 to 0.93) |
| Pharmacy claims             | 0.94 (0.94 to 0.95)  | 1.06 (1.06 to 1.06) | 0.93 (0.92 to 0.93) |
|                             | Rate ratios from the difference-in-difference Poisson model<br>substance use and mental health related |                     |                     |
|                             | Age 5–17   | Age 18–44           | Age 45–64           |
| Inpatient visits            | NA*  | 1.11 (1.01 to 1.22) | 1.19 (1.04 to 1.36) |
| Emergency department visits | 4.11 (3.28 to 5.16)  | 2.46 (2.20 to 2.76) | 1.48 (1.31 to 1.67) |
| Outpatient visits           | 1.12 (1.09 to 1.16)  | 0.93 (0.92 to 0.95) | 0.87 (0.85 to 0.89) |
|                             | ORs from the difference-in-difference logistic model   |                     |                     |
|                             | Age 5–17   | Age 18–44           | Age 45–64           |
| Insomnia                    | 1.64 (1.12 to 2.39)  | 1.17 (1.09 to 1.26) | 1.18 (1.09 to 1.28) |
| Cardiovascular disease†     | NA*  | 1.45 (1.30 to 1.62) | 1.15 (1.07 to 1.25) |
| Alcohol use disorder        | NA*  | 0.97 (0.86 to 1.11) | 1.16 (0.99 to 1.35) |
| Substance use disorder      | NA*  | 0.92 (0.83 to 1.03) | 1.24 (1.07 to 1.44) |
| Depression                  | NA*  | 1.12 (1.02 to 1.24) | 1.20 (1.08 to 1.33) |
| Anxiety                     | NA*  | 1.02 (0.95 to 1.10) | 1.01 (0.92 to 1.11) |
| Mood disorder               | NA*  | 1.03 (0.95 to 1.10) | 1.10 (1.00 to 1.20) |
| Disorders diagnosed young   | 0.80 (0.66 to 0.97)  | 0.99 (0.72 to 1.37) | 0.56 (0.31 to 1.04) |

\*These diseases and conditions are rare or difficult to diagnose in children.

†We adopted a longer follow-up period 2008–2011(pre-implementation) and 2013–2016 (TNNIS implementation) to allow for lag time in disease manifestation.

census tracts and measuring the level of aeroplane noise exposure.

## RESULTS

Participants were generally similar across both groups over the two points in time (table 1), but healthcare utilisation varied over time by age group and treatment status.

The increased use of the TNNIS climb occurred in 2012.<sup>27</sup> Prior to that date the climb was only used for the US Open or unexpected weather/runway repairs.<sup>27</sup> We were only able to obtain data on TNNIS departures after the New York Port Authority's Fiscal Year 2013 because the Port Authority indicated that the earlier data had been lost. There were roughly 1278 TNNIS departures/year on average during US Open events in the 2013–2019 period, providing a point of reference. Since 2013, the total annual number of TNNIS climbs ranged between 9349 and 29 676, with an average of 18653/year. The DNL figures (figure 1) reflect the average noise exposure by census tract across the 2013–2019 period, and may not reflect the actual change in aircraft noise in Flushing, New York in the pre-2012 and post-2012 periods.

## Overall medical utilisation

Table 2 provides results from regression models assessing period-related changes in medical utilisation and diagnoses. The effects of the change in flight patterns on overall utilisation were inconsistent across types of utilisation and age. Overall, outpatient visits increased slightly in Flushing relative to Sunset Park for those aged 18–44 (rate ratio (RR)=1.04, 95% CI=1.04 to 1.05). Prescription drug claims also increased by a similar amount for this group (RR=1.06, 95% CI=1.06 to 1.06). However, outpatient visits and prescription drug use for children in Flushing aged 5–17 (outpatient RR=0.86, 95% CI=0.85 to 0.87; prescription drug claims RR=0.94, 95% CI=0.94 to 0.95) as well as for older adults 45–64 declined (outpatient RR=0.92, 95% CI=0.92 to 0.93; prescription drug claims RR=0.93, 95% CI=0.92 to 0.93).

While the general pattern for outpatient visits indicates decreased medical utilisation in Flushing compared with Sunset Park over time, ED visits in Flushing increased in the post TNNIS period among all age groups. For ages 5–17, the RR was 1.31 (95% CI=1.24 to 1.37); for ages 18–44, the RR was 1.45 (95% CI=1.41 to 1.49); and for ages 45–64, the RR was 1.16 (95% CI=1.11 to 1.21). Substance use and mental health-related ED visits also increased in Flushing

in the post period relative to Sunset Park, with rate ratios ranging between 2.5 and 4.1. For ages 5–17, RR was 4.11 (95% CI=3.28 to 5.16); for ages 18–44, RR was 2.46 (95% CI=2.20 to 2.76); and for ages 45–64, RR was 1.48 (95% CI=1.31 to 1.67).

Relative to Sunset Park, inpatient visits in Flushing also show statistically significant increases for overall visits for ages 18–44 (RR=1.05, 95% CI=1.02 to 1.08). However, statistically significant decreases were observed for ages 45–64 (RR=0.93, 95% CI=0.88 to 0.97).


### Changes by diagnosis

Relative to Sunset Park, implementation of the TNNIS climb was associated with increases in insomnia diagnoses, particularly for children. For example, the crude prevalence of insomnia for children increased by 57% from 398 per 100 000 in Flushing, compared with a 6% decrease from 477 per 100 000 in Sunset Park. For children in this age group, the OR for insomnia was 1.64 (95% CI=1.12 to 2.39). For older ages, the effect sizes were somewhat less striking (ie, for the 18–44 age group, the OR was 1.17 (95% CI=1.09 to 1.26), and for ages 45–64 the OR=1.1 (95% CI=1.09 to 1.28)).

Cardiovascular disease diagnoses increased significantly in Flushing relative to Sunset Park in the post-2012 period.

For 18–44-year-olds, the crude prevalence of cardiovascular disease increased in both neighbourhoods due to ageing of the samples, by 83% from 1955 per 100 000 in Flushing and by 29% from 1576 per 100 000 in Sunset Park. The OR for cardiovascular disease diagnoses in Flushing relative to Sunset Park in this age group was 1.45 (95% CI=1.30 to 1.62). For 45–64-year-olds, the crude prevalence increased by 33% from 9934 per 100 000 in Flushing and 19% from 9073 per 100 000 in Sunset Park. For this age group, the OR was 1.15 (95% CI=1.07 to 1.25). Substance use disorder only increased significantly for the 45–64 age group in Flushing relative to Sunset Park (OR=1.24, 95% CI=1.07 to 1.44).

Figure 2 shows the monthly prevalence of insomnia and cardiovascular disease diagnoses for the 45–64 age group. Age is measured at the beginning of each period, 1 January 2009 for the pre period and 1 January 2013 for the post period. The numerator is the number of unique individuals with one or more diagnosis from inpatient, emergency room or outpatient claims and the denominator is the number of Medicaid-enrolled patients. The trends of both conditions increased throughout the study periods, because people are getting older, but Flushing showed increases that were larger in magnitude in the post period relative to Sunset Park.



**Figure 2** Prevalence of insomnia and cardiovascular disease diagnoses per 100 000 among 45–64 age group. \*Age is measured at the beginning of each period, 1 January 2009 for the pre period and 1 January 2013 for the post period.

Results for other conditions were more mixed. Clinical depression diagnoses increased for the two older age groups (ages 18–44 OR=1.12, 95% CI=1.02 to 1.24; ages 45–64 OR=1.20, 95% CI=1.08 to 1.33). Broader mood disorder diagnoses, however, only showed statistically significant increases for the 45–65 age group (OR=1.10, 95% CI=1.00 to 1.20). For 5–17-year-olds, developmental disorder diagnoses significantly decreased (OR=0.80, 95% CI=0.66 to 0.97) in Flushing relative to Sunset Park after the implementation of TNNIS.

## DISCUSSION

We find that increases in aeroplane noise at DNL levels greater than 55 were associated with increases in insomnia, depression, substance abuse and cardiovascular disease across most age groups. These diagnoses are generally consistent with our a priori hypotheses regarding the relationship between exposure to aeroplane noise and health.<sup>3 6 8 9 12 13 30 31</sup> Specifically, aeroplane noise may produce disruptions in sleep and psychological stress, thereby producing neuroendocrine disruptions that lead to mental health disorders and cardiovascular disease.

The biological pathways through which aeroplane noise impacts health have been elucidated.<sup>9–14</sup> Numerous associational studies suggest that aeroplane noise produces real-world health impacts, and experimental animal models show a wide range of health impacts associated with noise-induced stress as well.<sup>3–9 11–13 15–18 32</sup> Our study adds quasiexperimental evidence in humans to this substantial body of research showing that increasing aeroplane noise will have detrimental health impacts on communities surrounding airports. The magnitude of our findings is not strictly comparable to those in associational studies because lagged health effects (eg, the time required for psychological stress to manifest as cardiovascular disease) tend to mute the measured impacts.

Nevertheless, the magnitude of the impact of aeroplane noise on the health outcomes we observe are generally in line with previous work. For instance, an earlier analysis of associational studies of the health impact of aircraft noise in Flushing, New York estimated that aircraft noise would produce a weighted increase in cardiovascular disease of 14% (RR=1.14, 95% CI=1.08 to 1.22) and a weighted increase in anxiety of 79% (RR=1.79, 95% CI=1.0 to 3.1).<sup>11 24</sup> We observe an OR for cardiovascular disease among 18–64-year-olds in the range of 1.12–1.40. While the studies examine incident cardiovascular disease and we measure both incident and prevalent cardiovascular disease, it is reasonable to assume that the OR we estimate does not greatly overestimate the adjusted RR computed using associational studies.<sup>37</sup>

In the international literature, the self-reported annoyance, health, health-related quality of life and cardiovascular disease rates for those who live close to airports is significantly lower than for matched individuals living in quieter areas.<sup>38–40</sup> In this literature, these latter findings are particularly true for noise-sensitive individuals.<sup>38 39</sup> This

suggests that self-selection by noise may mute previously observed effects in ecological studies, which control for socioeconomic status but not always noise sensitivity. One strength of our study is that the change in aircraft noise was exogenous and moving out of a neighbourhood requires time and effort.

Our study was subject to a number of limitations. First, the health effects in a predominantly Chinese-American population may not be generalisable to other populations. Chinese-Americans in New York City are unusually healthy.<sup>41</sup> Medicaid data also present unique challenges. Participants can enter and exit the programme, for example. If there are more participants exiting the programme in one area relative to another, the observed outcomes will also change. We addressed this problem by adjusting for the months a participant was enrolled in Medicaid within a calendar year.

Next, we use DNL as a measure. Frequency of noise exposure may be superior at predicting health outcomes, but frequency data were not available. Finally, it is possible that the change in neighbourhood composition over time differed before and after the implementation of year-round TNNIS departures in Flushing relative to Sunset Park. However, we did not observe any trends in the available data that suggested this was the case, and there were no major events in 2012 that clearly serve as an alternative causal factor for either the primary or unexpected findings. Moreover, our findings apply only to the zip codes directly under the DNL zones defined by our analysis.

Cost-effectiveness analyses (based partly on earlier associational data) show that the benefits of noise-mitigation strategies (reduced illness and discomfort) tend to outweigh the costs.<sup>24 42</sup> Given that these earlier studies did not include the full range of health outcomes that we measure here, it is likely that these studies understate the already substantial benefits of aircraft noise mitigation strategies.

Much more comprehensive quasiexperimental and economic analyses are required to determine the extent to which policymakers may wish to act. The costliest options—building airports far from populated areas and providing high-speed transit and freeways—can increase the cost of mitigation by billions of dollars.

**Acknowledgements** The authors thank NYU Health Evaluation and Analytics Lab and the New York State Department of Health for making the Medicaid claims data available and gratefully acknowledge the funding for this research from the Robert Wood Johnson Foundation's Policies for Action program. Disclaimer: The views and opinions expressed in this article are those of the authors and do not necessarily reflect the official policy or position of the New York State Department of Health. Example of analysis performed within this article are only examples. They should not be utilized in real-world analytic products.

**Contributors** SSW, SG, SW and PAM approved the final draft and agreed to be accountable for all aspects of the work. SSW contributed to study design, data linkage, analysis, interpretation of the data, drafting the methods and results sections. SG contributed to the acquisition of data, study design, analysis and interpretation of data. SZW contributed to data analysis and interpretation of data. PAM contributed to study conception, study design, analysis, interpretation of data and drafting the introduction and discussion sections. Though BW passed away prior to the submission of the manuscript, he had significant contributions in the study conception, the acquisition of data and sample identification. PAM is the guarantor of the study.

**Funding** Robert Wood Johnson Foundation (75822).



**Disclaimer** The views and opinions expressed in this article are those of the authors and do not necessarily reflect the official policy or position of the New York State Department of Health. Example of analysis performed within this article are only examples. They should not be utilised in real-world analytic products.

**Map disclaimer** The inclusion of any map (including the depiction of any boundaries therein), or of any geographic or locational reference, does not imply the expression of any opinion whatsoever on the part of BMJ concerning the legal status of any country, territory, jurisdiction or area or of its authorities. Any such expression remains solely that of the relevant source and is not endorsed by BMJ. Maps are provided without any warranty of any kind, either express or implied.

**Competing interests** At the time of the study, Brian Will worked at a non-profit organisation called Queens Quiet Skies who are a grass-roots group aiming to address airport noise.

**Patient and public involvement** Patients and/or the public were involved in the design, or conduct, or reporting or dissemination plans of this research. Refer to the Methods section for further details.

**Patient consent for publication** Not applicable.

**Ethics approval** This study is approved by the Institutional Review Board at New York University Washington Square under IRB-FY2016-1101.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** No data are available. We used individual-level claims data that contain protected Patient Health Information (PHI). Therefore, the data cannot be made available publicly as required by the Health Insurance Portability and Accountability Act (HIPAA).

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iD

Scarlett Sijia Wang <http://orcid.org/0000-0003-2687-0365>

## REFERENCES

- Nolan M. *Fundamentals of air traffic control: Cengage learning*, 2010.
- Prevot T, Homola J, Mercer J, et al. Initial evaluation of NextGen air/ground operations with ground-based automated separation assurance. *ATM2009, Napa, California* 2009.
- Zaharna M, Guilleminault C. Sleep, noise and health: review. *Noise Health* 2010;12:64.
- Clark C, Stansfeld SA. The effect of transportation noise on health and cognitive development: a review of recent evidence. *Int J Comp Psychol* 2007;20.
- Stansfeld SA, Haines MM, Burr M, et al. A review of environmental noise and mental health. *Noise Health* 2000;2:1.
- Passchier-Vermeer W, Passchier WF. Noise exposure and public health. *Environ Health Perspect* 2000;108:123.
- Morrell S, Taylor R, Lyle D. A review of health effects of aircraft noise. *Aust N Z J Public Health* 1997;21:221–36.
- Beutel ME, Jünger C, Klein EM, et al. Noise annoyance is associated with depression and anxiety in the general population- the contribution of aircraft noise. *PLoS One* 2016;11:e0155357.
- Münzel T, Schmidt FP, Steven S, et al. Environmental noise and the cardiovascular system. *J Am Coll Cardiol* 2018;71:688–97.
- Schmidt FP, Basner M, Kröger G, et al. Effect of nighttime aircraft noise exposure on endothelial function and stress hormone release in healthy adults. *Eur Heart J* 2013;34:3508–14.
- Babisch W. Cardiovascular effects of noise. *Noise Health* 2011;13:201.
- Maschke C, Harder J, Ising H. Stress hormone changes in persons exposed to simulated night noise. *Noise Health* 2002;5:35.
- Melamed S, Bruhis S. The effects of chronic industrial noise exposure on urinary cortisol, fatigue and irritability: a controlled field experiment. *J Occup Environ Med* 1996;38:252–6.
- McEwen BS. Protective and damaging effects of stress mediators. *N Engl J Med* 1998;338:171–9.
- Basner M, Babisch W, Davis A, et al. Auditory and non-auditory effects of noise on health. *Lancet* 2014;383:1325–32.
- Perron S, Tétreault L-F, King N, et al. Review of the effect of aircraft noise on sleep disturbance in adults. *Noise Health* 2012;14:58.
- Xue L, Zhang D, Yibulayin X. Effects of high frequency noise on female rat's multi-organ histology. *Noise Health* 2014;16:213.
- Ising H, Babisch W, Kruppa B. Noise-induced endocrine effects and cardiovascular risk. *Noise Health* 1999;1:37.
- Cohen JP, Coughlin CC. Changing noise levels and housing prices near the Atlanta Airport. *Growth Change* 2009;40:287–313.
- Cohen JP, Coughlin CC. Spatial hedonic models of Airport noise, proximity, and housing PRICES\*. *J Reg Sci* 2008;48:859–78.
- Kawachi I, Adler NE, Dow WH. Money, schooling, and health: mechanisms and causal evidence. *Ann N Y Acad Sci* 2010;1186:56–68.
- Muennig P, Franks P, Jia H, et al. The income-associated burden of disease in the United States. *Soc Sci Med* 2005;61:2018–26.
- Boes S, Nüesch S, Stillman S. Aircraft noise, health, and residential sorting: evidence from two quasi-experiments. *Health Econ* 2013;22:1037–51.
- Zafari Z, Jiao B, Will B, et al. The trade-off between optimizing flight patterns and human health: a case study of aircraft noise in queens, NY, USA. *Int J Environ Res Public Health* 2018;15:15081753. doi:10.3390/ijerph15081753
- Nüesch S. Aircraft noise, health, and residential sorting: evidence from two quasi-experiments. *Standardization news: SN* 2012; (6744).
- Eagan ME, Hanrahan R, Miller R. Implementing performance based navigation procedures at US Airport: improving community noise exposure. INTER-NOISE and NOISE-CON Congress and conference proceedings; Institute of noise control engineering 2013:1577–86.
- Buckley C. A rumble in the sky and grumbles below, 2020. Available: <https://www.nytimes.com/2013/08/26/nyregion/planes-roar-from-above-creates-a-rumble-on-the-ground.html> [Accessed 17 Nov 2020].
- Mathematica. *Medicaid managed care and integrated delivery systems: technical assistance to states and strengthening federal oversight*. Washington, DC: Mathematica Policy Research, 2013.
- Bhaskar S, Hemavathy D, Prasad S. Prevalence of chronic insomnia in adult patients and its correlation with medical comorbidities. *J Family Med Prim Care* 2016;5:780–4.
- Bernstein GA, Borchardt CM. Anxiety disorders of childhood and adolescence: a critical review. *J Am Acad Child Adolesc Psychiatry* 1991;30:519–32.
- Mensah GA, Brown DW. An overview of cardiovascular disease burden in the United States. *Health Aff* 2007;26:38–48.
- Halonen JI, Hansell AL, Gulliver J, et al. Road traffic noise is associated with increased cardiovascular morbidity and mortality and all-cause mortality in London. *Eur Heart J* 2015;36:2653–61.
- Technical Advisory Committee (TAC) meeting 8. Available: <http://www.panyjpart150.com/AdminPages/GetProjectFile.asp?a=LGA&f=LGA%20TAC%20Meeting%20No.%208%20Summary,%20August%2016,%202016.pdf> [Accessed 17 Jun 2020].
- Flight Aware. Live flight tracking. Available: <https://flightaware.com> [Accessed 20 Jun 2020].
- NYC Health. Community health profile: Sunset Park. Available: <https://www1.nyc.gov/assets/doh/downloads/pdf/data/2018chp-bk7.pdf> [Accessed 2 Mar 2020].
- NYC Health. Community health profile: flushing and Whitestone. Available: <https://www1.nyc.gov/assets/doh/downloads/pdf/data/2018chp-qn7.pdf> [Accessed 2 Mar 2020].
- Knol MJ, Le Cessie S, Algra A, et al. Overestimation of risk ratios by odds ratios in trials and cohort studies: alternatives to logistic regression. *CMAJ* 2012;184:895–9.
- Welch D, Shepherd D, McBride D. Health-related quality of life is impacted by proximity to an Airport in noise sensitive people. INTER-NOISE and NOISE-CON Congress and conference proceedings; 2016: Institute of Noise Control Engineering 2016:5003–10.
- Shepherd D, McBride D, Dirks KN. Annoyance and health-related quality of life: a cross-sectional study involving two noise sources. *J Environ Prot* 2014.
- Bronzaft AL, Dee Ahern K, McGinn R. Aircraft noise: a potential health hazard. *Environ Behav* 1998;30:101–13.
- Muennig P, Wang Y, Jakubowski A. The health of immigrants to New York City from mainland China: evidence from the new York health examination and nutrition survey. *J Immigr Refug Stud* 2012;10:131–7.
- Jiao B, Zafari Z, Will B, et al. The cost-effectiveness of lowering permissible noise levels around U.S. Airports. *Int J Environ Res Public Health* 2017;14:1497.