

RESEARCH

Open Access

Transport genes and chemotaxis in *Laribacter hongkongensis*: a genome-wide analysis

Susanna KP Lau^{1,2,3,4*†}, Rachel YY Fan^{4†}, Gilman KM Wong^{4†}, Jade LL Teng⁴, Kong-Hung Sze⁵, Herman Tse^{1,2,3,4}, Kwok-Yung Yuen^{1,2,3,4} and Patrick CY Woo^{1,2,3,4*}

Abstract

Background: *Laribacter hongkongensis* is a Gram-negative, sea gull-shaped rod associated with community-acquired gastroenteritis. The bacterium has been found in diverse freshwater environments including fish, frogs and drinking water reservoirs. Using the complete genome sequence data of *L. hongkongensis*, we performed a comprehensive analysis of putative transport-related genes and genes related to chemotaxis, motility and quorum sensing, which may help the bacterium adapt to the changing environments and combat harmful substances.

Results: A genome-wide analysis using Transport Classification Database TCDB, similarity and keyword searches revealed the presence of a large diversity of transporters ($n = 457$) and genes related to chemotaxis ($n = 52$) and flagellar biosynthesis ($n = 40$) in the *L. hongkongensis* genome. The transporters included those from all seven major transporter categories, which may allow the uptake of essential nutrients or ions, and extrusion of metabolic end products and hazardous substances. *L. hongkongensis* is unique among closely related members of *Neisseriaceae* family in possessing higher number of proteins related to transport of ammonium, urea and dicarboxylate, which may reflect the importance of nitrogen and dicarboxylate metabolism in this saccharolytic bacterium. Structural modeling of two C⁴-dicarboxylate transporters showed that they possessed similar structures to the determined structures of other DctP-TRAP transporters, with one having an unusual disulfide bond. Diverse mechanisms for iron transport, including hemin transporters for iron acquisition from host proteins, were also identified. In addition to the chemotaxis and flagella-related genes, the *L. hongkongensis* genome also contained two copies of *qseB/qseC* homologues of the AI-3 quorum sensing system.

Conclusions: The large number of diverse transporters and genes involved in chemotaxis, motility and quorum sensing suggested that the bacterium may utilize a complex system to adapt to different environments. Structural modeling will provide useful insights on the transporters in *L. hongkongensis*.

Background

Laribacter hongkongensis is a Gram-negative, sea gull-shaped, rod that belongs to the *Neisseriaceae* family of β -proteobacteria [1,2]. The bacterium was first isolated from the blood and empyema pus of a man with alcoholic cirrhosis and bacteremic empyema thoracis in Hong Kong [1]. Using the selective medium, cefoperazone MacConkey agar, the bacterium was subsequently isolated from the stool of patients with gastroenteritis [3,4]. In a multicenter case-control study, *L. hongkongensis* was shown to be associated with community-acquired

gastroenteritis, with recent travel and eating fish being risk factors [5]. Apart from the human gut, *L. hongkongensis* has also been isolated from gut of freshwater animals including fish and Chinese tiger frogs as well as water from drinking water reservoirs [2,5-9]. In order to adapt to the changing environments and intestines of different animal hosts including human, fish and amphibians, *L. hongkongensis* must possess mechanisms to combat harmful substances in the environment and immune defense of animal hosts.

Transport-related proteins of bacteria are important in allowing the uptake of essential nutrients or ions, and extrusion of metabolic end products and hazardous substances. Bacteria employ different mechanisms for transport of different chemicals and these mechanisms have

* Correspondence: skplau@hkucc.hku.hk; pcywoo@hkucc.hku.hk

† Contributed equally

¹State Key Laboratory of Emerging Infectious Diseases, Hong Kong
Full list of author information is available at the end of the article

been classified into seven major categories according to the Transport Protein Database (TCDB): channels and pores (class 1), electrochemical potential-driven transporters (class 2), primary active transporters (class 3), group translocators (class 4), transmembrane electron carriers (class 5), accessory factors involved in transport (class 8), and incompletely characterized transport systems (class 9).

Bacteria also possess sophisticated signaling systems to sense and adapt to various substances in the environment. Depending on whether the environmental substances are attractants or repellents, the bacterium may migrate towards or away from the substances, which include certain amino acids, sugars, and metal ions [10-12]. This sense-and-swim ability is important for bacteria to be able to find the suitable environment for optimal growth. Chemotaxis involves two separate systems, the chemoreceptors located in the bacterial cell membrane which are important for sensing the binding compounds, and the transduction proteins which are involved in the downstream signal transduction in response to the stimuli. The chemoreceptors are also called methyl-accepting chemotaxis proteins (MCPs), which are reversibly methylated and function as homodimers [11,13].

The availability of the complete genome sequence of *L. hongkongensis* has allowed an opportunity to study its biology and important factors for adaptation to the changing environment [14]. We have previously found that transport-related proteins, including all seven major categories of transporters, account for about 14.1% of all coding sequences in the *L. hongkongensis* genome, suggesting that this group of proteins may be important for survival of the bacterium in the various environments and hosts [14]. Genes related to motility and chemotaxis were also identified [14]. Except for the first strain isolated from blood culture and empyema pus of a patient which was likely a non-motile variant, all strains from patients with gastroenteritis, animals or environmental water samples are motile with polar flagellae [1,4-7,10], suggesting that chemotaxis and motility may be an important mechanism for environmental adaptation in most isolates of *L. hongkongensis*. In this study, a comprehensive analysis of putative transport-related genes and genes related to chemotaxis, motility and quorum sensing in the *L. hongkongensis* genome is performed.

Results and discussion

Transport genes in *L. hongkongensis* genome

A huge diversity of transporters, including those from all seven major categories, were identified in the *L. hongkongensis* genome, as described in our previous complete genome report [14]. This may reflect its ability to adapt to various environments, including freshwater animals, water and human intestines. These transporters included: (1) 48

channels and pores, (2) 134 electrochemical potential-driven transporters, (3) 194 primary active transporters, (4) 9 group translocators, (5) 16 transmembrane electron carriers, (6) 7 accessory factors involved in transport and (7) 49 transporters of incompletely characterized transport systems (Table 1).

Channels and pores

The outer membranes of lipid bilayer envelopes of Gram-negative bacteria contain large numbers of water-filled transmembrane protein channels known as porins [15]. They serve as a molecular filter allowing for permeation of hydrophilic molecules up to a certain size or specific solutes into the periplasmic space. Some bacterial porins also serve as receptor for phage and bacteriocin binding [16]. X-ray crystallography studies and atomic structures have revealed that porin molecules exist as trimers, with the transmembrane core composed of mostly β -sheets and some α -helices [15]. The *L. hongkongensis* genome contained 48 coding sequences (CDSs) belonging to channels and pores, of which 17 were α -type channels, 29 were β -barrel porins and 2 were holins (Table 1).

Among the 17 α -type channels, five were mechanosensitive channels, including one large conductance mechanosensitive channel (LHK_02562) and four small conductance mechanosensitive channels (LHK_01830, LHK_01942, LHK_02394 and LHK_02965), which are responsible for mediating resistance to mechanophysical changes [17]. Interestingly, three CDSs encoding proteins of the ammonium transporter family were identified in the *L. hongkongensis* genome, as compared to only one copy such genes in *Chromobacterium violaceum*, the most closely related bacterial species of the *Neisseriaceae* family with complete genome sequence available (Table 2). Moreover, a homologue of urea transporter responsible for urea uptake (LHK_01044) was also present in *L. hongkongensis* (Table 2), while this protein was absent in *C. violaceum* and the pathogenic *Neisseria* spp., *Neisseria gonorrhoeae* and *Neisseria meningitidis*. This may reflect the importance of nitrogen metabolism of the bacterium, as *L. hongkongensis* is assacharolytic and has been shown to use different pathways for arginine synthesis regulated at different temperatures [14]. In fact, the habitats of the closely related bacterial species are quite different from that of *L. hongkongensis*, where the latter can survive in human intestine in addition to diverse freshwater environment. This may also explain its unique ability in maximizing nitrogen metabolism. Among the β -barrel porins, the OmpA-OmpF-type porins are most well known in bacteria to allow passive diffusion of hydrophilic substrates across the outer membrane. Three CDSs coding for putative OmpA-OmpF-type porins were identified in the *L. hongkongensis* genome. Interestingly, two homologues of another β -barrel porin, fatty acid transporter gene

Table 1 Transporters in *L. hongkongensis* and *C. violaceum*

| Category | <i>L. hongkongensis</i> | | | <i>C. violaceum</i> | | |
|--|-------------------------|-----------------|---------------------|---------------------|-----------------|---------------------|
| | No. of CDSs | % of total CDSs | % of transport CDSs | No. of CDSs | % of total CDSs | % of transport CDSs |
| Channel and Pores | 48 | 1.5 | 10.5 | 63 | 1.4 | 11.3 |
| α -type channels | 17 | | | 26 | | |
| β -barrel porins | 29 | | | 43 | | |
| Pore-forming toxins (proteins and peptides) | 0 | | | 3 | | |
| Holins | 2 | | | 2 | | |
| Electrochemical Potential-driven Transporters | 134 | 4.1 | 29.3 | 161 | 3.7 | 28.8 |
| Porters (uniporters, symporters and antiporters) | 132 | | | 159 | | |
| Ion-gradient-driven energizers | 2 | | | 2 | | |
| Primary Active Transporters | 194 | 6.0 | 42.5 | 252 | 5.7 | 45.0 |
| P-P-bond-hydrolysis-driven transporters | 150 | | | 206 | | |
| Decarboxylation-driven transporters | 5 | | | 7 | | |
| Oxidoreduction-driven transporters | 39 | | | 39 | | |
| Group Translocators | 9 | 0.3 | 2.0 | 18 | 0.4 | 3.2 |
| Phosphotransfer-driven group translocators | 2 | | | 8 | | |
| Acyl CoA ligase-coupled transporters | 7 | | | 10 | | |
| Transmembrane Electron Carriers | 16 | 0.5 | 3.5 | 13 | 0.3 | 2.3 |
| Transmembrane 2-electron transfer carriers | 14 | | | 12 | | |
| Transmembrane 1-electron transfer carriers | 2 | | | 1 | | |
| Accessory Factors Involved in Transport | 7 | 0.2 | 1.5 | 20 | 0.5 | 3.6 |
| Auxiliary transport proteins | 7 | | | 20 | | |
| Incompletely Characterized Transport Systems | 49 | 1.5 | 10.7 | 33 | 0.7 | 5.9 |
| Recognized transporters of unknown biochemical mechanism | 15 | | | 14 | | |
| Putative transport proteins | 34 | | | 19 | | |

Table 2 α -type channels in *L. hongkongensis* and their closest homologues

| CDS | Protein | Closest match organism | Best E-value | Amino acid identity (%) |
|-----------|--|--|--------------|-------------------------|
| LHK_02933 | Ammonium transporter | <i>L. nitroferrum</i> | 2.00E-146 | 73.18 |
| LHK_03249 | Ammonium transporter | <i>Shewanella halifaxensis</i> | 2.00E-118 | 62.32 |
| LHK_03154 | Ammonium transporter family protein | <i>L. nitroferrum</i> | 1.00E-163 | 78.99 |
| LHK_02207 | Flagellar motor protein MotA | <i>L. nitroferrum</i> | 1.00E-122 | 74.48 |
| LHK_00970 | Ion transporter | <i>C. violaceum</i> | 5.00E-78 | 58.96 |
| LHK_02562 | Large-conductance mechanosensitive channel | <i>Pelodictyon luteolum</i> | 2.00E-43 | 56.95 |
| LHK_01830 | Transmembrane protein | <i>C. violaceum</i> | 2.00E-109 | 57.52 |
| LHK_01942 | Mechanosensitive ion channel protein | <i>Janthinobacterium</i> sp. Marseille | 5.00E-79 | 41.26 |
| LHK_02394 | MscS Mechanosensitive ion channel | <i>L. nitroferrum</i> | 7.00E-55 | 48.95 |
| LHK_02965 | Transporter, small conductance mechanosensitive ion channel family | <i>E. coli</i> O157:H7 | 5.00E-73 | 61.04 |
| LHK_02739 | Molecular chaperone DnaK | <i>C. violaceum</i> | 0 | 85.98 |
| LHK_02206 | OmpA/MotB domain protein | <i>L. nitroferrum</i> | 4.00E-97 | 75.46 |
| LHK_01044 | Urea transporter | <i>Methylobacterium extorquens</i> PA1 | 1.00E-65 | 50.46 |
| LHK_00053 | TolQ-related transport transmembrane protein | <i>C. violaceum</i> | 1.00E-86 | 74.66 |
| LHK_03174 | TolR protein | <i>C. violaceum</i> | 5.00E-30 | 51.88 |
| LHK_00499 | Probable exbB-like biopolymer transport | <i>C. violaceum</i> | 4.00E-55 | 59.31 |
| LHK_00498 | Biopolymer transport <i>exbD</i> transmembrane protein | <i>Burkholderia pseudomallei</i> 112 | 7.00E-36 | 55.88 |

(*fadL*), were also found, which may be important for uptake of long-chain fatty acids in freshwater environments poor in lipids or fatty acids.

Electrochemical potential-driven transporters

The *L. hongkongensis* genome possessed a large number of CDSs ($n = 134$) encoding for putative electrochemical potential-driven transporters, among which the majority (132 CDSs) were porters including uniporters, symporters and antiporters, while the remaining two CDSs were ion-gradient-driven energizers (Table 1). Of the 132 porters, 19 (14.3%) belonged to the major facilitator superfamily (MFS). MFS proteins are important transporters in bacteria, which allow transport of molecules by an electrochemical ion gradient and typically contain a single subunit with 12 membrane-spanning helices [18]. The MFS proteins of *L. hongkongensis* were predicted to mediate transport of diverse substrates including ions, drugs and metabolites. Another major family of porters were the resistance-nodulation-cell division (RND) superfamily (28 CDSs), which are responsible for transporting a wide variety of substrates including antibiotics, dyes, detergents, fatty acids, bile salts, organic solvents, heavy metals, auto-inducers and lipooligosaccharides in Gram-negative bacteria [19,20]. Other porters belonged to diverse families of proteins which facilitate the transport of diverse substances including ions, amino acids, drugs, heavy metal such as nickel and cobalt, nucleobase, C_4 -dicarboxylates and other metabolites. The presence of various porters may be involved in acquisition of essential substances for metabolism and bacterial resistance to environmental toxic substances including heavy metals. Interestingly, a total of 11 porters for dicarboxylate transport were found in *L. hongkongensis* genome, as compared to only 6 in *C. violaceum* and 1 each in *N. meningitidis* and *N. gonorrhoeae* genomes (Table 3). C_4 -dicarboxylates are intermediates in TCA cycle that can be utilized by bacteria as nonfermentable carbon and/or energy sources under aerobic or anaerobic conditions [21]. Some C_4 -dicarboxylates, such as succinate, oxalate and malate, can also be found in nature [22]. The presence of high number of C_4 -dicarboxylates transporters may reflect the ability of using C_4 -dicarboxylates as carbon sources in *L. hongkongensis*, as the bacterium is assacharolytic, lacking a complete glycolytic pathway, and is in line with our experiments

showing that L-malate can be used as its sole carbon source [14].

Six of the 11 porters for dicarboxylate transport found in *L. hongkongensis* genome were believed to form two DctP-type tripartite ATP-independent periplasmic (TRAP) transporters which belong a heterogeneous group of substrate-binding protein (SBP)-dependent secondary transporters of a diverse range of substrates found in bacteria and archaea [23-25]. The genes encoding the 3 subunits were arranged in an operon, with two membrane proteins DctQ and DctM associating with DctP to form a C_4 -dicarboxylate TRAP transporter [26]. Several TRAP transporters have been characterized in detail, with the structures of at least seven DctP-type SBP subunits determined [25]. These studies revealed significant structural and architectural similarities among the different SBPs, while highlighting the differences that permitted these proteins to bind their respective substrates with high affinity and specificity. Besides substrate recognition, it was also found that the SBP performs other essential functions [27], and likely interacts with the integral membrane components in a hitherto undiscovered manner. One operon (LHK_00983-00984-00985), encoding C_4 -dicarboxylate transporter, was found downstream of several genes related to allantoin regulation and utilization; while the other operon (LHK_01394-01393-01392) was located upstream of the *maeB* gene encoding NADP-dependent malate dehydrogenase. The SBP encoded by LHK_00983 (DctP_00983) was a 331 aa protein containing a 22 aa N-terminal signal peptide, with a predicted molecular weight of 33.9 kDa. It possessed 48% amino acid identity to the closest homolog in *Roseovarius* sp. TM1035 (NCBI accession no.: ZP_01881277). The SBP encoded by LHK_01394 (DctP_01394) was a 335 aa protein containing a 24 aa N-terminal signal peptide, with a predicted molecular weight of 34.3 kDa. It possessed 74% amino acid identity to the closest homolog in *C. violaceum* ATCC12472. The homology model and structural alignment of the homology model showed that the overall structure of DctP_00983 and DctP_01394 was very similar to the determined structures of other DctP-type SBPs (Figure 1 and 2, and see Supplementary material). Similar to other DctP homologs, they were divided into two domains with conserved arrangements of α -helices and β -sheets, which are

Table 3 Porters for dicarboxylates in *L. hongkongensis* and related bacteria

| Family | <i>L. hongkongensis</i> | <i>C. violaceum</i> | <i>N. meningitidis</i> | <i>N. gonorrhoeae</i> |
|--|-------------------------|---------------------|------------------------|-----------------------|
| C_4 -Dicarboxylate Uptake (Dcu) Family | 0 | 2 | 0 | 0 |
| Dicarboxylate/Amino Acid:Cation (Na or H) Symporter (DAACS) Family | 3 | 1 | 0 | 0 |
| Tripartite ATP-independent Periplasmic Transporter (TRAP-T) Family | 6 | 3 | 0 | 0 |
| Divalent Anion:Na+ Symporter (DASS) Family | 1 | 0 | 1 | 1 |
| C_4 -dicarboxylate Uptake C (DcuC) Family | 1 | 0 | 0 | 0 |
| Total | 11 | 6 | 1 | 1 |




Figure 1 Homology model of DctP_00983 (panel A) and DctP_01394 (panel B), putative DctP TRAP transporters for C₄-dicarboxylate in *L. hongkongensis*. For DctP_00983, the C-score of the model was 1.49, which approximately corresponded to an expected TM-score of 0.92 ± 0.06 and an expected root-mean-square deviation (RMSD) of 3.2 ± 2.3 Å from the native structure. The Ramachandran plot showed that 99.6% of aa are in the favored and allowed regions. Calculated G-factors for dihedral angles and main-chain covalent forces are 0.11 and -0.17 respectively, with an overall average of 0.01. The Z-score of the model is -7.84, which is comparable to other experimentally determined protein chains of a similar size in the PDB. Local model quality analysis by plot of residue scores in ProSA-web did not reveal any problematic regions in the structure. The quality analysis results suggested that the homology model is mostly reliable with good structural qualities. For DctP_01394, the C-score of the model was 1.36, which approximately corresponded to an expected TM-score of 0.90 ± 0.06 and an expected RMSD of 3.5 ± 2.4 Å from the native structure. The Ramachandran plot showed that 99.0% of aa are in the favored and allowed regions. Calculated G-factors for dihedral angles and main-chain covalent forces are 0.09 and -0.17 respectively, with an overall average of 0.00. The Z-score of the model is -8.15, which is comparable to other experimentally determined protein chains of a similar size in the PDB. Local model quality analysis by plot of residue scores in ProSA-web did not reveal any problematic regions in the structure. The quality analysis results suggested that the homology model of DctP_01394 is also reliable with good structural qualities.

connected by a characteristic hinge made up of two β -strands and an α -helix. A highly conserved arginine residue in domain II is present in both proteins (Arg145 of DctP_00983 and Arg147 of DctP_01394), which

corresponds to Arg147 in SiaP of *H. influenzae* essential to SBP function by forming a salt bridge with the carboxylate group of the ligand [28]. Interestingly, a disulfide bond was predicted between the cysteine residues at positions




Figure 2 Structural alignment of the homology model of DctP_00983 and DctP_01394, showing similar structures to other DctP-type SBPs (panel A) and a disulfide bond predicted between the cysteine residues at positions 129 and 182 of DctP_00983 (panel B). RMSD between DctP_00983 and the related structures ranged from 0.761 to 1.290 Å. RMSD between DctP_01394 and the related structures ranged from 0.891 to 1.377 Å.

129 and 182 for DctP_00983 (Figure 2) by homology modeling and sequence analysis. This structural feature was also found in the closest homolog in *Roseovarius* sp. TM1035, but absent from other related DctP-type SBP homologs including DctP_01394.

Primary active transporters

Primary active transporters mediate energy-driven transport of substances in and out of bacterial cells by using ATP hydrolysis, photon absorption, electron flow, substrate decarboxylation, or methyl transfer [29]. Primary active transporters were the most abundant class of transporters (194 CDSs), constituting 6% of CDSs in the *L. hongkongensis* genome, among which 150 belonged to P-P-bond-hydrolysis-driven transporters (Table 1). Of the 150 P-P-bond-hydrolysis-driven transporters, 109 were ATP-binding cassette (ABC) transporters which are one of the largest groups of membrane proteins using energy from ATP hydrolysis for transport. In bacteria, they reside in the inner membrane and are involved in both uptake and export of a wide range of substances. All ABC transporters share a common basic structure which consists of four domains: two transmembrane domains, typically with six transmembrane spans per domain, and two cytoplasmic nucleotide-binding domains which catalyze nucleotide hydrolysis [30]. In bacteria, these domains are encoded as separate polypeptides. Determined by the structure of the transmembrane domain, ABC transporters are typically specific for the substrates that they are responsible for, although some may transport for multiple related substances. As a result, the numbers of ABC transporters in different bacterial species vary widely, depending on its need for adaptation to varying environmental conditions [31]. The ABC transporters in the *L. hongkongensis* are likely involved in the active transport of diverse substances, including carbohydrate, amino acids or peptides, ions, vitamins, lipids, drugs and heavy metals including molybdenum, iron, zinc, cobalt, magnesium, copper, cadmium, mercury, lead, arsenite and nickel. These systems were often arranged in gene clusters comprising the ATP-binding protein and two auxiliary proteins, a permease and a substrate-binding protein. Compared to the 70 ABC transporters found in *E. coli* [31], the *L. hongkongensis* genome contained a large number of such proteins, reflecting its ability to adapt to different hosts and environment.

Apart from P-P-bond-hydrolysis-driven transporters, other primary active transporters identified in the *L. hongkongensis* genome included oxidoreduction-driven transporters (39 CDSs) and decarboxylation-driven transporters (5 CDSs), which use chemical energy to perform transport of charged or uncharged molecules across the membrane against the concentration gradient [32].

Group translocators

Of the nine group translocators, two were phosphotransfer-driven group translocators and seven were acyl CoA ligase-coupled transporters belonging to the fatty acid transporter (FAT) family. The phosphotransferase group translocators are components of the bacterial phosphotransferase system (PTS), which catalyzes translocation of sugars and hexitols with concomitant phosphorylation, and regulates the metabolism in response to the availability of carbohydrates. PTSs consist of two cytoplasmic proteins, enzyme I (EI) and HPr, and a variable number of sugar-specific transport complexes (Enzymes II^{sugar}) belonging to the group translocators. While the *Escherichia coli* genome encoded 38 different PTS proteins, the *L. hongkongensis* genome encoded only one gene for EI and HPr each and two genes for transporters, one containing protein-N p-phosphohistidine-sugar phosphotransferase IIA domain and the other containing nitrogen-regulatory fructose-specific IIA domain [33]. This is likely related to the relative unimportance of sugar metabolism in *L. hongkongensis*.

Transmembrane electron carriers

There were 16 transmembrane electron carriers in the *L. hongkongensis* genome, including 14 transmembrane 2- and two transmembrane 1-electron transfer carriers. Among the 14 transmembrane 2-electron transfer carriers, 12 belonged to the prokaryotic molybdopterin-containing oxidoreductase (PMO) family, and the other 2 belonged to the disulfide bond oxidoreductase D (DsbD) and B (DsbB) family respectively.

Accessory factors involved in transport

There were seven accessory factors belonging to auxiliary transport proteins in the *L. hongkongensis* genome, 3 belonging to the membrane fusion protein (MFP) family, 2 to the phosphotransferase system enzyme I (EI) family, 1 to the phosphotransferase system HPr (HPr) family and 1 to the stomatin/podocin/band 7/nephrin.2/SPFH (stomatin) family.

Incompletely characterized transport systems

Of the 49 CDSs belonging to incompletely characterized transport system, 15 were recognized transporters of unknown biochemical mechanism, with 6 belonging to the putative type VI symbiosis/virulence secretory pathway (VISP) family, 2 to the HlyC/CorC (HCC) family, 2 to the capsular polysaccharide exporter (CPS-E) family, 1 to the tellurium ion resistance (TerC) family and the remaining 4 being metal ion transporters. The other 34 CDSs were putative transport proteins, including 2 CDSs of the camphor resistance (CrcB) family and 1 probable hemolysin III.

Iron Transport in *L. hongkongensis*

Iron is an essential metal for most microorganisms used in many key molecules involved in metabolism. In bacteria,

iron metabolism has been shown to be important in adaptation to the environment especially within the host and as a result related to virulence. Diverse mechanisms for iron transport were identified in the *L. hongkongensis* genome, suggesting that the bacterium is able to adapt to iron limitation present in human body which represents one of the non-specific immune response called induced hypoferrremia [34,35]

Siderophores and iron uptake

Siderophores are low molecular mass compounds with high affinity for ferric iron. In contrast to *C. violaceum* which produced siderophores for iron acquisition, proteins related to siderophore production were not found in *L. hongkongensis* genome. However, a homolog of TonB-dependent siderophore receptor (LHK_00497) was present, as described in our previous report [14]. Although *Listeria monocytogenes* also did not produce siderophores for iron acquisition, it was able to obtain iron by using either exogenous siderophores produced by various microorganisms or natural catechol compounds widespread in the environment [36,37]. It remains to be determined if *L. hongkongensis* can utilize exogenous siderophores or other natural iron-binding compounds for iron acquisition.

Hemin transport

Despite the inability to produce siderophores, a set of genes related to the transport of hemin were identified in *L. hongkongensis* genome (8 CDSs compared to 6 CDSs in *C. violaceum*). The 8 CDSs included TonB-dependent receptor (LHK_01193), hemin degrading factor (LHK_01192), ABC transporter permease (LHK_01189), ferric citrate transport system ATP-binding protein (LHK_01188), hemin-binding periplasmic protein (LHK_01190), hemin importer ATP-binding subunit (LHK_01427), hemin ABC transporter permease protein (LHK_01428) and Fur family ferric uptake regulator (LHK_01431). The conserved domains for hemin receptor, FRAP and NPFL, were also identified in the TonB-dependent receptor [38]. This suggests that *L. hongkongensis* is able to utilize iron source from host proteins, which may be important for survival in its hosts. Three other CDSs, homologous to *fbpA* (LHK_02634), *fbpB* (LHK_02635) and ATP-binding protein (LHK_02636), ABC transporters for transferrin and lactoferrin, were also present, although the outer membrane receptor is not found.

ABC transporters of the metal type

A cluster of three genes encoding an ABC transporter of the metal type (homologous to that identified in *C. violaceum*) was identified in the *L. hongkongensis* genome. They encoded a periplasmic Mn²⁺/Zn²⁺-binding (lipo)protein (surface adhesion A) (*znuA*), a Mn²⁺/Zn²⁺ permease component (*znuB*) and the ATPase component (*znuC*). In addition, a gene encoding a putative cadmium-translocating ATPase component (cadmium-translocating P-type

ATPase) (*CadA*) (LHK_00449) was also present. A similar gene was also found in *C. violaceum* (CV1154), which was thought to be a surface adhesion A component for Mn²⁺/Zn²⁺ binding. The Fur family ferric uptake regulator (*zur*) (LHK_01344) was also present.

Other transporters

In addition to the above transporters, two CDSs encoding ferrous iron transport proteins, *feoA* (LHK_03044) and *feoB* (LHK_03045), were identified in *L. hongkongensis* genome, which are believed to provide iron supply under anaerobic or low pH conditions in bacteria [39]. Three other CDSs homologous to iron uptake ABC transporter periplasmic solute-binding protein (LHK_01590), ABC transporter permease (LHK_01593) and ABC transporter ATP-binding protein (LHK_01591) were also found.

Iron storage

Mechanism required for storage of iron after its acquisition from the environment was present in *L. hongkongensis*, which mainly depends on two proteins: bacterioferritin (BFR) (LHK_01239, homologous to CV3399 in *C. violaceum*) and frataxin-like homolog (LHK_00023, homologous to Daro_0208 in *Dechloromonas aromatica*). The BFR is an iron-storage protein with close similarity to the ferritins found in both eukaryotes and prokaryotes [40]. The frataxin-like homolog has been implicated in iron storage in other bacteria. The frataxin-like domain is related to frataxin, the protein mutated in Friedreich's ataxia which is therefore proposed to result from decreased mitochondrial iron storage [41,42].

Regulation of iron transport

Fur protein is a global repressor protein by forming Fur-Fe²⁺ complexes that bind to iron-dependent promoter during iron-rich conditions. It regulates ferrichrome (*fhuABCDG*), ferric citrate (*fecABCDE*) and ferrous iron (*feoABC*) uptake systems. The Fur protein in *L. hongkongensis* was encoded in CDS LHK_01431 (homologous to FuraDRAFT_2340 in *Lutella nitroferum*).

Chemotaxis in *L. hongkongensis*

Methyl-accepting chemotaxis and chemosensory transducer proteins

A total of 52 open reading frames (CDSs) were related to chemotaxis, of which 29 encoded MCPs and 22 were chemosensory transducer proteins. Most genes encoding MCPs were scattered throughout the *L. hongkongensis* genome, while the genes encoding transducer proteins were mostly arranged in three gene clusters as described in our previous report (Table 4) [14].

All the predicted MCPs in *L. hongkongensis* possessed a transmembrane domain, which is compatible with their anticipated location in the bacterial cell membrane and function as receptors. Conserved domain structures were also identified in some of the MCPs. The plasmid

Table 4 CDSs related to chemotaxis in *L. hongkongensis* genome

| CDS | Gene | Product | Organism with the closest matching sequences | E-value | Identities | Cluster ^a |
|-----------|--------------|--|--|---------|---------------|----------------------|
| LHK_00115 | | histidine kinase, HAMP region: chemotaxis sensory transducer | <i>D. aromatica</i> | 1e-96 | 242/680 (35%) | |
| LHK_00482 | | methyl-accepting chemotaxis sensory transducer | <i>L. nitroferum</i> | 4e-55 | 164/543(30%) | |
| LHK_00516 | | methyl-accepting chemotaxis sensory transducer | <i>L. nitroferum</i> | 8e-129 | 265/513 (51%) | |
| LHK_00553 | | diguanylate phosphodiesterase | <i>C. violaceum</i> | 6e-111 | 211/406 (51%) | CA |
| LHK_00554 | <i>cheA1</i> | CheA signal transduction histidine kinase | <i>L. nitroferum</i> | 0 | 443/613 (72%) | CA |
| LHK_00555 | <i>cheZ1</i> | chemotaxis phosphatase, CheZ | <i>L. nitroferum</i> | 2e-69 | 139/244(59%) | CA |
| LHK_00556 | <i>cheY1</i> | chemotaxis regulator protein CheY | <i>C. violaceum</i> | 4e-61 | 109/130 (83%) | CA |
| LHK_00557 | <i>cheV1</i> | chemotaxis protein CheV | <i>C. violaceum</i> | 1e-138 | 240/314 (76%) | CA |
| LHK_00558 | <i>cheV2</i> | chemotaxis protein CheV | <i>C. violaceum</i> | 5e-147 | 251/313 (80%) | CA |
| LHK_00559 | | two-component sensor histidine kinase | <i>L. nitroferum</i> | 2e-59 | 169/381 (44%) | CA |
| LHK_00560 | | chemotaxis sensory transducer | <i>D. aromatica</i> | 6e-24 | 100/320 (31%) | CA |
| LHK_00561 | <i>cheY2</i> | chemotaxis protein cheY | <i>D. aromatica</i> | 8e-46 | 85/121 (70%) | CA |
| LHK_00562 | <i>cheA2</i> | chemotaxis protein CheA | <i>C. violaceum</i> | 2e-161 | 358/746 (47%) | CA |
| LHK_00563 | <i>cheW</i> | CheW protein | <i>Burkholderia phytofirmans</i> | 1e-40 | 95/153 (62%) | CA |
| LHK_00564 | | methyl-accepting chemotaxis protein | <i>C. violaceum</i> | 4e-143 | 315/475 (66%) | CA |
| LHK_00565 | <i>cheR</i> | CheR chemotaxis protein methyltransferase | <i>Janthinobacterium</i> sp. Marseille | 5e-68 | 125/273 (45%) | CA |
| LHK_00566 | <i>cheB1</i> | chemotaxis-specific methylesterase | <i>Nitrosomonas europaea</i> | 2e-99 | 186/355 (52%) | CA |
| LHK_00567 | <i>cheD</i> | chemoreceptor glutamine deamidase CheD | <i>D. aromatica</i> | 5e-59 | 108/189 (57%) | CA |
| LHK_00603 | | methyl-accepting chemotaxis protein | <i>C. violaceum</i> | 7e-103 | 242/624 (38%) | |
| LHK_00617 | | methyl-accepting chemotaxis protein IV | <i>C. violaceum</i> | 2e-100 | 223/481 (46%) | |
| LHK_00700 | | methyl-accepting chemotaxis sensory transducer | <i>Allochroamatium vinosum</i> | 0 | 384/715 (53%) | |
| LHK_00726 | <i>aer1</i> | methyl-accepting chemotaxis sensory transducer with Pas/Pac sensor | <i>L. nitroferum</i> | 7e-114 | 232/528 (43%) | |
| LHK_00935 | <i>cheR</i> | MCP methyltransferase, CheR-type | <i>L. nitroferum</i> | 2e-92 | 170/282 (60%) | |
| LHK_01020 | | putative aromatic hydrocarbon chemotaxis transducer | <i>Azoarcus</i> sp. | 4e-62 | 140/338 (41%) | |
| LHK_01116 | | methyl-accepting chemotaxis protein | <i>Denitrovibrio acetiphilus</i> | 1e-59 | 152/461 (32%) | |
| LHK_01212 | | methyl-accepting chemotaxis sensory transducer | <i>L. nitroferum</i> | 1e-135 | 261/476 (54%) | |
| LHK_01359 | <i>cheY3</i> | chemotaxis regulator protein CheY | <i>C. violaceum</i> | 1e-56 | 102/127 (80%) | CB |
| LHK_01360 | <i>cheV3</i> | chemotaxis protein CheV | <i>C. violaceum</i> | 1e-134 | 231/309 (74%) | CB |
| LHK_01361 | | methyl-accepting chemotaxis sensory transducer | <i>L. nitroferum</i> | 6e-47 | 157/506 (31%) | CB |
| LHK_01372 | | chemotaxis sensory transducer | <i>D. aromatica</i> | 4e-49 | 166/534 (31%) | |
| LHK_01470 | | putative aromatic hydrocarbon chemotaxis transducer | <i>Azoarcus</i> sp. | 2e-93 | 222/539 (41%) | |
| LHK_01602 | | methyl-accepting chemotaxis sensory transducer | <i>L. nitroferum</i> | 0 | 339/601 (56%) | |
| LHK_01618 | | methyl-accepting chemotaxis sensory transducer | <i>L. nitroferum</i> | 2e-87 | 209/525 (39%) | |
| LHK_01706 | | methyl-accepting chemotaxis protein IV | <i>C. violaceum</i> | 1e-121 | 247/481 (51%) | |
| LHK_01721 | | methyl-accepting chemotaxis protein | <i>C. violaceum</i> | 4e-113 | 240/627 (38%) | |
| LHK_02037 | | methyl-accepting chemotaxis sensory transducer | <i>Leptospirillum ferrodiazotrophum</i> | 5e-63 | 137/327 (41%) | |
| LHK_02158 | <i>aer2</i> | methyl-accepting chemotaxis sensory transducer with Pas/Pac sensor | <i>Ralstonia pickettii</i> | 6e-39 | 98/276 (35%) | |
| LHK_02165 | | methyl-accepting chemotaxis protein | <i>C. violaceum</i> | 8e-146 | 275/631 (43%) | |
| LHK_02364 | <i>cheB2</i> | response regulator receiver modulated CheB methylesterase | <i>Geobacter bemidjiensis</i> | 1e-63 | 122/206 (59%) | |
| LHK_02427 | | methyl-accepting chemotaxis protein | <i>C. violaceum</i> | 6e-110 | 227/629 (36%) | CC |
| LHK_02428 | | Hypothetical protein | No | | | CC |
| LHK_02429 | <i>cheV4</i> | response regulator receiver modulated CheW protein | <i>L. nitroferum</i> | 2e-145 | 248/313 (79%) | CC |
| LHK_02430 | <i>cheV5</i> | chemotaxis protein CheV | <i>C. violaceum</i> | 3e-137 | 237/314 (75%) | CC |
| LHK_02431 | <i>cheY4</i> | chemotaxis regulator protein CheY | <i>C. violaceum</i> | 2e-58 | 105/127 (82%) | CC |
| LHK_02432 | <i>cheZ2</i> | chemotaxis phosphatase, CheZ | <i>L. nitroferum</i> | 2e-63 | 129/248 (52%) | CC |

Table 4 CDSs related to chemotaxis in *L. hongkongensis* genome (Continued)

| | | | | | | |
|-----------|--------------|--|---|--------|---------------|----|
| LHK_02433 | <i>cheA3</i> | CheA signal transduction histidine kinase | <i>L. nitroferum</i> | 0 | 420/611 (68%) | CC |
| LHK_02455 | | methyl-accepting chemotaxis sensory transducer | <i>Candidatus Accumulibacter phosphates</i> | 1e-74 | 154/326 (47%) | |
| LHK_02575 | | putative Methyl-accepting or sensory transducer chemotaxis protein | <i>Alteromonadales bacterium</i> | 1e-83 | 172/407 (42%) | |
| LHK_02814 | <i>aer3</i> | chemotaxis sensory transducer | <i>Rhodopseudomonas palustris</i> | 8e-42 | 138/425 (32%) | |
| LHK_02834 | | methyl-accepting chemotaxis protein | <i>Pseudomonas syringae</i> | 1e-45 | 148/437 (33%) | |
| LHK_03026 | | methyl-accepting chemotaxis protein | <i>C. violaceum</i> | 6e-145 | 275/627 (43%) | |
| LHK_03119 | | methyl-accepting chemotaxis sensory transducer | <i>L. nitroferum</i> | 2e-133 | 273/514 (53%) | |
| LHK_03163 | | methyl-accepting chemotaxis sensory transducer | <i>Candidatus Accumulibacter phosphatis</i> | 9e-50 | 167/494 (33%) | |

^aThe Che proteins were encoded in three gene clusters, named CA, CB and CC (chemotaxis A, B and C clusters)

achromobacter secretion (PAS) domain was found in four MCPs (LHK_00564, LHK_00726, LHK_02158 and LHK_02814). PAS domains are energy-sensing modules that are found in proteins from archaea to humans [43]. The histidine kinase adenyl cyclase MCP and phosphatase (HAMP) domain was present in 22 of the 29 MCPs. The HAMP domain interacts with the PAS domain for signal transduction in aerotaxis (oxygen-sensing) receptor in *Escherichia coli* [43], and possesses roles of regulating the phosphorylation or methylation of homodimeric receptors by transmitting the conformational changes in periplasmic ligand-binding domains to cytoplasmic signaling kinase and methyl-acceptor domains [44].

These chemosensory transducer proteins work as two-component regulatory systems which typically consist of a sensory histidine kinase and a response regulator. The histidine kinase is usually a transmembrane receptor and the response regulator a cytoplasmic protein [45]. Following autophosphorylation at a conserved histidine residue in response to changes in chemoreceptor occupancy, the histidine kinase serves as a phospho-donor for the response regulator. Once phosphorylated, the response regulator mediates changes in gene expression or cell motility. CheA is a typical sensory histidine kinase while CheY is a downstream regulator protein [46]. Upon phosphorylation, CheY binds to the FlIM component at the base of the flagellar motor switch to induce clockwise rotation [47]. In contrast to the single copies of CheA and CheY in *E. coli*, the presence of 22 chemosensory transducer proteins, many with multiple copies including three CheA, one CheB, one CheD, two CheR, five CheV, one CheW, four CheY, and two CheZ, suggested that *L. hongkongensis* may utilize a complex transducer system to mediate chemotaxis response and adapt to environmental changes (Table 4). These Che proteins were encoded in three gene clusters, named CA, CB and CC. The first and largest cluster, CA, encoded two CheA, one


CheR, two CheY, two CheV, one CheZ, and the single CheD and CheW. The second and smallest cluster, CB, encoded one CheV and CheY. The third cluster, CC, encoded one CheA, one CheY, two CheV and one CheZ. Phylogenetic analysis of CheAs, CheVs and CheYs of *L. hongkongensis* suggested that the multiple copies are the result of both horizontal transfer events and gene duplication, as some of the copies were more closely related to the corresponding proteins in other bacteria while others were more closely related among the homologues of *L. hongkongensis* (Figure 3).

The CheA proteins of *L. hongkongensis* were most closely related to homologues in the closely related *Chromobacterium violaceum* and *Lutiella nitroferum* with 47% to 72% amino acid identities. CheA has five domains, P1 to P5 [46]. All the three CheA proteins in *L. hongkongensis* contained these conserved domains. In the P1 domain, the invariant histidine residue, which undergoes phosphorylation by the P4 domain, was also present. In the kinase domain P4, the four conserved regions designated the N, G1, F and G2 boxes were also present in the three CheAs (Figure 4).


The CheY proteins of *L. hongkongensis* were highly similar to the homologues in *C. violaceum* and *Dechloromonas aromatica*, with 70% to 83% amino acid identities. Multiple alignment of the four CheY with that of *E. coli* showed the presence of all five amino acid residues conserved among response regulators [46,48]: aspartate at positions 12, 13 and 57; threonine at position 87, and lysine at position 109, with the aspartate at position 57 representing the phosphorylation site (Figure 5). Residues that interact with P2 domain of CheA were identified.

Other Che proteins are believed to be involved in the regulation of bacterial chemotaxis, although the exact function of some are not fully understood. Among them, CheB is known to work in conjunction with CheR in the reversible methylation of the MCPs. CheR is a

CheA



CheV



CheY




Figure 3 Phylogenetic tree showing the relationships of the CheAs, CheVs and CheYs from *L. hongkongensis* to those from other bacteria. The unrooted trees are constructed by using the neighbor-joining method using Kimura's two-parameter correction, with bootstrap values calculated from 1000 trees. The scale bar indicates the estimated number of substitutions per 20 bases. Bacterial names and accession numbers are given as cited in the GenBank database.

| | | | | | |
|-----------|------------|-------------|-------------|------------|---------------------------|
| | P1 domain | | | | |
| Eco_cheA | MHRSVFPQC | HQPNRGSVS | MDISDFYQTF | FDEADELLAD | MEQHLLVLQP -EAPDAEQLN 59 |
| LHK_00554 | ----- | -----MSDF | AGMELLQDF | LTESELLED | VDNKLVELEK -YEDKGLLN 43 |
| LHK_00562 | ----- | -----MS | IDLSQFHAAF | FDEAAEHLET | IERLLESQF GQTTDGEALN 42 |
| LHK_02433 | ----- | -----MSDF | NGMELLQDF | LTESELLED | VDNKLVELEK -YEDKGLLN 43 |
| | P1 domain | | | | |
| Eco_cheA | AIFRAAHSIK | GGAGTFGFSV | LQETTHLMEN | LLDEARRGEM | QLNTDIINLF LETKDIMQE 119 |
| LHK_00554 | DIFRGRFTIK | GGAGFLNATP | LVTLCRHTEN | LFDKLRNGEL | KLNSHVMDVI LDATGVVRDM 103 |
| LHK_00562 | AIFRAAHSIK | GSAGTFGFAD | IAGTFHGLEN | LLDRIRRGEL | PLTAGRISVC LKSRDVIADQ 102 |
| LHK_02433 | DIFRGRFTIK | GGAGFLNATP | LVTLCRHTEN | LFDKLRNGEL | QLNSHVMDVI LDATGVVRDM 103 |
| | P1 domain | | | | |
| Eco_cheA | LDAYKQ-SQE | PDAASFNYIC | QALRQLALEA | KGETPSAVTR | LS-----VVA KSEPQDEQSR 173 |
| LHK_00554 | FGDLAQSRMP | APAP----- | ----- | ----- | ----- 117 |
| LHK_00562 | LAHRDGATA | VDPRLAEIE | AALLAAHDEG | T-DMPVPTVS | SRPDGGHYLL CWRDQLGHPE 161 |
| LHK_02433 | FGDLAQSRMP | APAP----- | ----- | ----- | ----- 117 |
| | P2 domain | | | | |
| Eco_cheA | SQSPRRILS | RLKAGEVDLL | EEELGHLTTL | TDVVKGADSL | SAILP-----D DIAEDDIT 227 |
| LHK_00554 | ----- | ---QHILDNL | DAVLSGQTAP | VAAAPAAAAS | QPGEPDWQL YQAVVVPADA 164 |
| LHK_00562 | GLQDRLAALG | TVQP-CGEQG | EELVFQLQSG | LSATDIAESL | AFWLPFGQFR LEHLESQGE 220 |
| LHK_02433 | ----- | ---QHILDNL | DAMLAGQDMV | RTVPLPAVP | DP-----ASVSQPASA 155 |
| | P2 domain | | | | |
| Eco_cheA | AVLCFVIEAD | QITFETVD-- | ----- | VSPKISTPPM | LKLAAEQAPT GRVEREKTTR 275 |
| LHK_00554 | AAQLTKSAP | STAVAAAP-- | ----- | ----APVA | ATPAASSPPP AQSRDRSPAS 207 |
| LHK_00562 | AFGVVDTQR | VVATSATEPA | APVDEGWGLF | APPAATADV | SPVPAEPAAT GSTPALQAT 280 |
| LHK_02433 | PP-----AAS | TVAVQAGF-- | ----- | ----ATTAA | AQPPE-----RERNISG 185 |
| | P3 domain | | | | |
| Eco_cheA | SSESTSIRVA | VEKVDQLNL | VGELVITQSM | LAQR-----S | SELDPVNHGD LITSMGQLQR 330 |
| LHK_00554 | SFQEATIRID | TQRLDQVLNL | SGEIGLTKNR | LTIRTEIMQ | GNLGANTLRS LDEAVSQLDL 267 |
| LHK_00562 | FQETSIRVN | VEKVDQLLNR | IGELVITQSM | LAQQ-----V | ERLGLASEE LQRGMAQLER 335 |
| LHK_02433 | NQQTTLRVD | AQRLDQVLNL | SGEIGLTKNR | LTIRTEIIQ | GNLGADVLHS LDEAVSQLDL 245 |
| | P3 domain | | P4 domain | | |
| Eco_cheA | NARDLQESVM | SIRMPMEYV | FSRYPRLRVD | LAGKLGKQVE | LTIVGSSTEL DKSLIERIID 390 |
| LHK_00554 | LVGDLQNAV | KTRMQPIGRL | FQKYPRLRAR | LARQLGKEVE | LVLSGEETEL DKTMIEDLND 327 |
| LHK_00562 | TTRELQEAVM | SVRMLPVASV | FGRFPRLVRE | LQKLGKRAVE | LQVIGEQTEI DKSFVEKLT 395 |
| LHK_02433 | LVGDLQNAV | KTRMQPIGRL | FQKYPRLRAR | LARQLGKVDV | LVLSGEETEL DKTMIEDLND 305 |
| | P4 domain | | | | |
| Eco_cheA | FLTHLVRNSL | DHGETELPEKR | LAAGKNSVGN | LILSAEHQGG | NICIEVTDG AGLNREIRILA 450 |
| LHK_00554 | FLVHLVRNAV | DHGETETPEER | LAAGKSAQSI | VELSAQQVGD | HIVIEVADG RGMNAGMLRK 387 |
| LHK_00562 | FLTHLVRNSL | DHGETESAAGR | AGAGKPPVGR | LTLRAFHQGG | HIVIEVSDG AGLQRDRILA 455 |
| LHK_02433 | FLVHLVRNAV | DHGETETPDER | QAAGKPVQSA | VKLSAQQVGD | HIVIEVADG RGMNAGMLRK 365 |
| | N | | | G1 | |
| | P4 domain | | | | |
| Eco_cheA | KAASQ---GL | TVSENMSDDE | VAMLI FAPGF | STAEQVTDVS | GRGVGMDVVK RNIQEMGGHV 507 |
| LHK_00554 | KALEKGLIDL | EQANSLDDKQ | ALHILFLPGF | STKDQISSVS | GRGVGMDVVR TNIQKLNRI 447 |
| LHK_00562 | KAREQ---GL | NVSDTMNDAE | VWQLIFEPGF | STAQAVTDVS | GRGVGMDVVR RNIEAMGSI 512 |
| LHK_02433 | KALEKGLIDL | EQANSMDDKQ | ALHILFLPGF | TTKSEISSVS | GRGVGMDVVR TNIQKLNRI 425 |
| | | | F | | G2 |
| | P4 domain | | P5 domain | | |
| Eco_cheA | EIQSKQGTGT | TIRILLPLTL | AILDGMSVRV | ADEVFILPLN | AVMESLQPRE ADLHPLAGGE 567 |
| LHK_00554 | DISSAPNEG | KISISLPLTL | AILPVLVVKV | CNQPFAVPLA | MVREIIPFS DSIQEVSGRP 507 |
| LHK_00562 | RIESLAGVGT | TISLHLPLTL | AILDGMSIAI | GNEIYILPLS | QVVESLQPR ADEVFTLAGQ 572 |
| LHK_02433 | DIRSAPNEG | RISISLPLTL | AILPVLVARV | CDQSFAMPLA | MVREIIPAT DAIQEVSGRP 485 |
| | P5 domain | | | | |
| Eco_cheA | RVLEVRGEYL | PIVELWKVFN | VAGAKTEATQ | GIVVILQSGG | RRYALLVDQL IGQHVVVKN 627 |
| LHK_00554 | -TIVRDEIL | PVRRLELLG | WKATQEPCEG | ---VLMQSAE | KTFILAI DSIQEVSGRP 563 |
| LHK_00562 | RLLRVRGECL | PLLALWEIFD | IQPRVREAEH | GLVITIVAS | QRVALLVDDL VSQQQVVKV 632 |
| LHK_02433 | -TIVRDEVL | PVRRLELLG | WKS DREPFGF | ---ILMQAE | KTFILAVDTF VSGREVVVKS 541 |
| | P5 domain | | | | |
| Eco_cheA | LESNYRKVPG | ISAATILGDG | SVALIVDVSA | LQAINREQRM | ANTAA----- 672 |
| LHK_00554 | LQN--IRPKG | VAGATLSGDG | AVVLVLDMD | LLSNDRGDHK | LARAGNMPL EAAAV 616 |
| LHK_00562 | LETNYRVRG | AAAATILGDG | QVAFILDGAE | VVRMGQGG-- | ----- 670 |
| LHK_02433 | LQN--VSLKG | VAGATLSGDG | AVVLVLDMS | LLNTDK--AH | AAVHGAL TPL PA--- 589 |

Figure 4 Amino acid sequence alignments of *L. hongkongensis* and *E. coli* CheAs. The conserved P1 to P5 domains are marked above the sequences. The histidine residue at potential phosphorylation site is shaded. The four conserved regions designated the N, G1, F and G2 boxes within P4 domain are marked in open boxes.

constitutively active methyltransferase which methylates the conserved glutamine residues of MCPs, while the methyltransferase CheB is responsible for demethylation [49,50]. Similar to CheY, the CheB of *L. hongkongensis*

also contained the five conserved amino residues of response regulators. In addition, three conserved residues of the catalytic site, serine at position 164, histidine at position 190 and aspartate at position 286, and the

| | | | | | | | |
|------------------|------------|------------|------------|------------|------------|------------|-----|
| Eco_cheY | ---MADKELK | FLVVDDFSTM | RRIVRNLLKE | LGFNVEEAE | DGLDALNKLQ | AGGYGFVISD | 57 |
| LHK_00556 | MIEAADKNLR | FLVVDDFSTM | RRILRNLLKE | LGFTNVDEAE | DGQVALHKLR | SQPYEFVSD | 60 |
| LHK_00561 | -----MAK | ILAVDDSPSI | RQMTFTLKN | AGYD-IISAP | DGLAGLKEAN | SHRVDLVLT | 53 |
| LHK_01359 | MLDEVNPNLR | FLVVDDFSTM | RRILRNLFKE | LHFTNIDEAE | DGQVALHKLR | SQPYEFVSD | 60 |
| LHK_02431 | MLDEVNPNLR | FLVVDDFSTM | RRILRNLLKE | LHFTNIDEAE | DGQVALHKLR | SQPYEFVSD | 60 |
| ▲ ▲ ▲ ▲ ▲ ▲ ▲ ▲ | | | | | | | |
| Eco_cheY | WNMPNMDGLE | LLKTIRADGA | MSALPVLMT | AEAKKENIIA | AAQAGASGYV | VKPFTAATLE | 117 |
| LHK_00556 | WNMPNMTGIE | LLRAVRADAQ | LRHLPFLMIT | AEAKRENIIE | AAQAGASGYI | VKPFTAATLE | 120 |
| LHK_00561 | QNMPGMDGLT | LIRELRQLPA | YRATPILMLT | TEAGDDMKAQ | GRAAGASGWM | VKPFDPQKLV | 113 |
| LHK_01359 | WNMPNMTGIE | LLRAVRADAQ | LRHLPFLMIT | AEAKRENIIE | AAQAGASGYI | VKPFTVVTLE | 120 |
| LHK_02431 | WNMPNMTGIE | LLRAVRADAQ | LRHLPFLMIT | AEAKRENIIE | AAQAGASGYI | VKPFTAATLE | 120 |
| Eco_cheY | EKLNKIFEKL | GM--- | 129 | | | | |
| LHK_00556 | EKLAKIFQST | SRQAG | 135 | | | | |
| LHK_00561 | DVVRKLLG-- | ----- | 121 | | | | |
| LHK_01359 | EKLAKIFQST | SRQAG | 135 | | | | |
| LHK_02431 | EKLAKIFQST | SRQAG | 135 | | | | |

Figure 5 Amino acid sequence alignments of *L. hongkongensis* and *E. coli* CheYs. The conserved aspartate, threonine and lysine residues are shaded. The aspartate residue at potential phosphorylation site is marked by black square, and residues of *E. coli* CheY that interact with the P2 domain of *E. coli* Che A are marked by black triangles above the residues.

GXGXXG nucleotide-binding-fold sequences conserved among CheB proteins were also present (Figure 6) [51].

Similar multiple copies of chemosensory transducer proteins have also been reported in *C. violaceum* and *Rhodobacter sphaeroides* [46,48]. Interestingly, the organization of the first cluster in *L. hongkongensis*, CA, was similar to one of the three clusters, cluster 3, in *C.*

violaceum, although some of the genes were in opposite coding direction. In *R. sphaeroides*, it has been shown that some of the multiple copies of Che proteins are essential (e.g. CheA2) while others are not (e.g. CheA1) although the multiple chemosensory protein homologues are not redundant [46,52]. Further studies are required to investigate the differential function of the

| | | | | | | | | |
|------------------|-------------|------------|-------------|-------------|-------------|-------------|------------|----|
| Eco_cheB | MSK-IRVLSV | DD | SALMRQIM | TEINSHSDM | EMVATAPDPL | VARDLIKKFN | PDVLTLDVEM | 59 |
| LHK_00566 | MSQPVRVLV | DD | SALMRNLL | AELINACDGM | CCVQQAADPL | QARESIRLLA | PDVVTLDVEM | 60 |
| Eco_cheB | PRMDGLDFLE | KLMRLRMPV | VMVSSLTGKG | SEVTLRALEL | GAIQDFVTKPQ | LGIREGMLAY | 119 | |
| LHK_00566 | PHMDGLEFLR | RLMRLRPTPV | VMVSSLTARG | SEVAIEALAL | GAVEVVEKPG | AGLGQAIPIRF | 120 | |
| Eco_cheB | SEMIAEKVRT | AAKASLAAHK | PLSAPTTLKA | GPLLSSEKLI | AIGASTGGTE | AIRHVLQPLP | 179 | |
| LHK_00566 | AMQLTGAIQK | AAQANLSRIC | NLNKSPAMPS | LRPPARDALV | LIGASTGGTE | ALVTLLSALP | 180 | |
| Eco_cheB | LSSPALLITQ | HMPPGFTRSF | ADRLNKLCQI | GVKEAEDGER | VLPGHAYIAP | GDRHMELARS | 239 | |
| LHK_00566 | AQMPPVLIVQ | HMPAGFTASF | ARRLDQACVL | QVREAQGG EK | LAPGQVWVAP | GHAHLQLSAQ | 240 | |
| Eco_cheB | GANYQIKIHD | GPAVNRHRPS | VDVLFHVSVAK | QAGRNAVGV I | LTGMGNDGAA | GMLAMRQAGA | 299 | |
| LHK_00566 | AGDWR TQLVD | SDPVNRHRPS | VDVLFH SALK | VAGRHTVAVL | LTGMGRDGAQ | GLLALRKAGA | 300 | |
| | | | | | GXGXXG | | | |
| Eco_cheB | WTLAQNEASC | VVFGMPREAI | NMGGVCEVVD | LSQVSQOMLA | KISAGQAIRI | ----- | 349 | |
| LHK_00566 | YTYAQDKASS | VVFGMPREAI | EIGAACEVAS | LGDMAHQMVT | RMAGGAGGRA | EEGKGVAAAPH | 360 | |

Figure 6 Amino acid sequence alignment of *L. hongkongensis* and *E. coli* CheBs. The 5 conserved aspartate, threonine and lysine residues also found in CheY are shaded. The three conserved residues of the catalytic site Ser164, His190 and Asp286 in *E. coli* CheB are marked by triangles above the residues and the GXGXXG nucleotide-binding-fold consensus sequences of other CheB marked in open box.

multiple copies of chemosensory transducer proteins in *L. hongkongensis*.

Flagellar proteins in *L. hongkongensis*

A total of 40 CDSs, arranged in six gene clusters, were likely involved in the biosynthesis of flagella in *L. hongkongensis* (Table 5). These six clusters, FA, FB, FC, FD,

FE and FF, encoded 11, 3, 5, 2, 16 and 3 genes respectively. The organization and gene contents of the first five clusters were highly similar to five of the seven clusters of flagellar genes (clusters 1, 2, 4, 5 and 7) previously found in *C. violaceum* [48], which is also a motile bacterium found in multiple ecosystems, including water and

Table 5 CDSs involved in flagella biosynthesis in *L. hongkongensis* genome

| CDS | Gene | Product | Organism with the closest matching sequences | E-value | Identities | Cluster ^a |
|-----------|-------------|--|--|---------|---------------|----------------------|
| LHK_00436 | <i>flgL</i> | flagellar hook-associated protein 3 | <i>L. nitroferum</i> | 1e-59 | 127/312 (40%) | FA |
| LHK_00437 | <i>flgK</i> | flagellar hook-associated protein FlgK | <i>L. nitroferum</i> | 1e-109 | 258/634 (40%) | FA |
| LHK_00438 | <i>flgJ</i> | flagellar rod assembly protein/muramidase FlgJ | <i>L. nitroferum</i> | 3e-68 | 144/296 (48%) | FA |
| LHK_00439 | <i>flgI</i> | flagellar basal body P-ring protein | <i>C. violaceum</i> | 1e-95 | 197/294 (67%) | FA |
| LHK_00440 | <i>flgH</i> | flagellar L-ring protein | <i>L. nitroferum</i> | 4e-60 | 122/231 (52%) | FA |
| LHK_00441 | <i>flgG</i> | flagellar basal-body rod protein FlgG | <i>Ralstonia pickettii</i> | 2e-92 | 162/260 (62%) | FA |
| LHK_00442 | <i>flgF</i> | flagellar basal-body rod protein FlgF | <i>L. nitroferum</i> | 1e-75 | 143/246 (58%) | FA |
| LHK_00443 | <i>flgE</i> | flagellar basal body FlaE domain-containing protein | <i>Pseudomonas putida</i> | 4e-76 | 212/598 (35%) | FA |
| LHK_00444 | <i>flgD</i> | flagellar hook capping protein | <i>L. nitroferum</i> | 8e-38 | 88/240 (36%) | FA |
| LHK_00445 | <i>flgC</i> | flagellar basal-body rod protein flgC | <i>C. violaceum</i> | 2e-49 | 92/136 (67%) | FA |
| LHK_00446 | <i>flgB</i> | flagellar basal-body rod protein FlgB | <i>L. nitroferum</i> | 2e-41 | 89/136 (65%) | FA |
| LHK_00584 | <i>flgN</i> | FlgN family protein | <i>C. violaceum</i> | 2e-15 | 48/131 (36%) | FB |
| LHK_00585 | <i>flgM</i> | anti-sigma-28 factor, FlgM | <i>L. nitroferum</i> | 4e-09 | 36/59 (61%) | FB |
| LHK_00586 | <i>flgA</i> | flagella basal body P-ring formation protein FlgA | <i>L. nitroferum</i> | 2e-36 | 85/206 (41%) | FB |
| LHK_00781 | <i>fliA</i> | RNA polymerase sigma factor for flagellar operon | <i>C. violaceum</i> | 5e-89 | 165/242 (68%) | FC |
| LHK_00782 | <i>fleN</i> | flagellar synthesis regulator FleN | <i>L. nitroferum</i> | 3e-49 | 121/268 (45%) | FC |
| LHK_00783 | <i>fihF</i> | flagellar biosynthesis regulator FihF | <i>C. violaceum</i> | 1e-119 | 250/504 (49%) | FC |
| LHK_00784 | <i>fihA</i> | flagellar biosynthesis protein FihA | <i>L. nitroferum</i> | 0 | 519/682 (76%) | FC |
| LHK_00785 | <i>fihB</i> | flagellar biosynthetic protein FihB | <i>L. nitroferum</i> | 2e-136 | 226/378 (59%) | FC |
| LHK_02206 | <i>motB</i> | OmpA/MotB domain protein | <i>L. nitroferum</i> | 6e-111 | 206/273 (75%) | FD |
| LHK_02207 | <i>motA</i> | flagellar motor protein MotA | <i>L. nitroferum</i> | 9e-123 | 213/286 (74%) | FD |
| LHK_02348 | <i>fliR</i> | flagellar biosynthetic protein FliR | <i>L. nitroferum</i> | 1e-60 | 142/258 (55%) | FE |
| LHK_02349 | <i>fliQ</i> | flagellar biosynthetic protein FliQ | <i>L. nitroferum</i> | 6e-24 | 65/89 (73%) | FE |
| LHK_02350 | | GCN5-related N-acetyltransferase | <i>Methylocella silvestris</i> | 5e-09 | 47/150 (31%) | FE |
| LHK_02351 | <i>fliP</i> | flagellar biosynthesis protein FliP | <i>C. violaceum</i> | 7e-95 | 178/252 (70%) | FE |
| LHK_02352 | <i>fliO</i> | flagellar protein FliO | <i>C. violaceum</i> | 2e-16 | 52/100 (52%) | FE |
| LHK_02353 | <i>fliN</i> | flagellar motor switch protein FliN | <i>L. nitroferum</i> | 2e-54 | 111/140 (79%) | FE |
| LHK_02354 | <i>fliM</i> | flagellar motor switch protein FliM | <i>L. nitroferum</i> | 3e-160 | 272/327 (83%) | FE |
| LHK_02355 | <i>fliL</i> | flagellar fliL transmembrane protein | <i>C. violaceum</i> | 2e-28 | 64/136 (47%) | FE |
| LHK_02356 | <i>fliK</i> | flagellar hook-length control protein | <i>Nitrosomonas europaea</i> | 5e-18 | 41/108 (37%) | FE |
| LHK_02357 | <i>fliJ</i> | flagellar export protein FliJ | <i>L. nitroferum</i> | 3e-20 | 64/142 (45%) | FE |
| LHK_02358 | <i>fliI</i> | flagellar protein export ATPase FliI | <i>L. nitroferum</i> | 0 | 331/453 (73%) | FE |
| LHK_02359 | <i>fliH</i> | flagellar assembly protein FliH | <i>L. nitroferum</i> | 8e-32 | 109/275 (39%) | FE |
| LHK_02360 | <i>fliG</i> | flagellar motor switch protein FliG | <i>L. nitroferum</i> | 2e-148 | 261/332 (78%) | FE |
| LHK_02361 | <i>fliF</i> | flagellar M-ring protein FliF | <i>L. nitroferum</i> | 0 | 339/585 (57%) | FE |
| LHK_02362 | <i>fliE</i> | flagellar hook-basal body complex subunit FliE | <i>L. nitroferum</i> | 5e-27 | 69/110 (62%) | FE |
| LHK_02363 | | two component, sigma54 specific, transcriptional regulator, Fis family | <i>L. nitroferum</i> | 2e-143 | 279/450 (62%) | FE |
| LHK_02703 | <i>fliD</i> | flagellar hook-associated 2 domain protein | <i>L. nitroferum</i> | 5e-45 | 136/445 (30%) | FF |
| LHK_02704 | <i>flaG</i> | FlaG flagellar protein | <i>Janthinobacterium</i> sp. Marseille | 2e-11 | 38/105 (36%) | FF |
| LHK_02705 | <i>fliC</i> | flagellin domain-containing protein | <i>Acidovorax</i> sp. | 2e-73 | 159/288 (55%) | FF |

^aThe flagellar proteins were arranged in six gene clusters, FA, FB, FC, FD, FE and FF (flagellar A, B, C, D, E and F clusters)

soil. On the other hand, the pathogenic *Neisseria* species, *Neisseria gonorrhoeae* and *Neisseria meningitidis*, which also belong to the same *Neisseriaceae* family, are non-motile with humans being the only host and reservoir, and do not possess flagellar genes.

A bacterial flagellum is typically composed of three parts, the filament formed by flagellin subunits, basal body attached to the bacterial cell membrane, and the hook which links between the filament and basal body [53]. All the major proteins that form these flagellar components were present in the *L. hongkongensis* genome. They included FliC and FliD which form the major part of the filament; FlgE, FlgK and FlgL which form the hook and hook-filament junction; and Flg B, FlgC, FlgH, FlgI, FlhA, FlhB, FliF, FliG, FliH, FliI, FliM, FliN, FliO, FliP, FliQ, FliR, MotA and MotB which form the basal body and flagellar-motor complex. Putative regulators of these flagellar proteins were also identified. FlgD and FliK are regulators of the hook component FlgE. FlgA, FlgN (both being chaperon proteins) and FliJ are involved in export of flagellar components. The anti-sigma factor gene FlgM and σ^{28} FliA that regulates late gene products were also present. However, similar to *C. violaceum*, the *L. hongkongensis* genome lacked the FlhDC operon genes, suggesting that the regulation of flagellar protein expression is controlled by FlgM/FliA in this group of bacteria.

Quorum sensing in *L. hongkongensis*

In addition to chemotaxis through which bacteria can rapidly adapt to environmental changes, quorum sensing is another way to assess the environment and to recognize the host. Quorum sensing is a signaling system through which bacteria can communicate among themselves by the production of and response to chemical signals called autoinducers [54]. In response to the changing concentrations of these autoinducers, downstream gene expression can be regulated. This cell-to-cell communication system, first identified in *Vibrio harveyi* in the regulation of bioluminescence, is now known to exist in diverse bacteria, especially those that reside in the gastrointestinal tract where recognition of the host may be important for survival and virulence gene expression [54,55]. Among the three major quorum-sensing mechanisms, including the LuxR-I, LuxS/AI-2, and AI-3/epinephrine/norepinephrine systems, known to be utilized by enteric bacteria, only the latter was found in the *L. hongkongensis* genome, suggesting that this system played a major role in quorum-sensing in the bacterium [14].

The AI-3/epinephrine/norepinephrine system is involved in inter-kingdom cross-signaling and regulation of virulence gene transcription and motility [54]. This mechanism is best characterized in enterohemorrhagic *E. coli* (EHEC) which causes fatal hemorrhagic colitis and hemolytic uremic syndrome. It has been shown that the locus of enterocyte effacement (LEE), an important

virulence factor in EHEC, and the flagellar genes of EHEC are regulated by the AI-3 system which involves AI-3 produced by the commensal gastrointestinal microflora and/or epinephrine/norepinephrine produced by the host [56,57]. The AI-3 system has also been implicated in biofilm formation in enteropathogenic *E. coli* (EPEC) [58]. Clarke et al. have recently identified the protein, QseC that binds to AI-3 and epinephrine/norepinephrine, suggesting its involvement in the AI-3 system [59]. QseC belongs to a two-component system, QseB/C, in which QseC is the sensor kinase and QseB the response regulator. QseB/C has also been shown to be involved in activation of the flagella regulon and virulence in a rabbit model for EHEC [59,60]. The *L. hongkongensis* genome contained two sets of genes, LHK_00329/LHK_00328 and LHK_1812/LHK_1813, homologous to *qseB/qseC* [14], most closely related to homologues in *C. violaceum* and *Azoarcus* sp. strain BH72 respectively. The two *qseB* genes in *L. hongkongensis* possessed the response regulator receiver domain (PF00072) and the C-terminal domain of transcriptional regulatory protein (PF00486) previously found in the QseB of *E. coli*. The two *qseC* genes in *L. hongkongensis* also contained the His Kinase A (phosphoacceptor) domain (PF00512) and the histidine kinase-, DNA gyrase B-, and HSP90-like ATPase domain (PF02518) previously identified in the QseC of *E. coli*. The presence of two copies of *qseB/qseC* suggested that the AI-3 system may be an important mechanism for adaptation to the changing environment and animal hosts for *L. hongkongensis*.

Conclusions

A large number of diverse transporters ($n = 457$), including those from all seven major transporter categories, were identified in the *L. hongkongensis* genome. A diversity of genes involved in chemotaxis, motility and quorum sensing were also found. This suggested that the ability to transport various substances plays an important role in the physiology or survival of *L. hongkongensis*, which may also utilize a complex system to mediate chemotaxis response and adapt to and survive in the rapidly changing environments. In particular, the bacterium is unique among closely related members of *Neisseriaceae* family in possessing higher number of proteins related to transport of ammonium, urea and dicarboxylate, which may reflect the importance of nitrogen and dicarboxylate metabolism in *L. hongkongensis* which is assacharolytic. Structural modeling of two C_4 -dicarboxylate transporters showed that they possessed similar structures to the determined structures of other DctP-TRAP transporters, but one with a rarely seen disulfide bond. A large number of ABC transporters were also identified. These suggest that the bacterium may be able to transport a wide variety of substrates including antibiotics, dyes, detergents,

fatty acids, bile salts, organic solvents, ions, amino acids, drugs, heavy metals such as nickel and cobalt, nucleobase, C₄-dicarboxylates and other metabolites. Diverse mechanisms for iron transport, including hemin transporters for iron acquisition from host proteins, were identified, suggesting that the bacterium may adapt to iron limitation present in human host. Using blastp of all transporters against rcsb pdb, many of these genes were also found to have homologous proteins of high sequence identities with known structures (data not shown). The large number of chemosensory transducer proteins, many having multiple copies arisen from both horizontal transfer events and gene duplications, may constitute a complex transducer system for mediating chemotaxis response and adapt to environmental changes. The presence of two copies of *qseB/qseC* homologs suggests that *L. hongkongensis* may use the AI-3 system for cross-kingdom quorum-sensing and regulation of potential virulence factors. Further studies are required to better characterize the precise target substance for transport proteins of interest, and the targets regulated by *qseB/qseC* in *L. hongkongensis*, which may shed light on its potential mechanisms for pathogenicity. Structural modeling can be a useful tool to provide useful structural insights about these genes in *L. hongkongensis*.

Methods

Transport genes were identified and classified according to Transport Classification Database TCDB <http://www.tcdb.org/> and manual annotation. These CDSs were from COG C (Energy production and conversion), COG D (Cell cycle control, cell division, chromosome partitioning), COG E (Amino acid transport and metabolism), COG F (Nucleotide transport and metabolism), COG G (Carbohydrate transport and metabolism), COG H (Coenzyme transport and metabolism), COG I (Lipid transport and metabolism), COG J (Translation, ribosomal structure and biogenesis), COG K (Transcription), COG L (Replication, recombination and repair), COG M (Cell wall/membrane/envelope biogenesis), COG N (Cell motility), COG O (post-translational modification, protein turnover, chaperones), COG P (Inorganic ion transport and metabolism), COG Q (Secondary metabolites biosynthesis, transport and catabolism), COG R (General function prediction only), COG S (Function unknown), COG T (Signal transduction mechanisms), COG U (Intracellular trafficking, secretion and vesicular transport) and COG V (Defense mechanisms). CDSs that were classified to COG N (cell motility) and COG T (signal transduction mechanisms), and COG M (cell wall/membrane/envelope biogenesis) were manually annotated for identification of genes related to chemotaxis, motility and quorum sensing. CDSs from other COGs were searched for additional genes using

keywords: chemotaxis, che, MCP, flagellar etc. All putative genes were studied by manual curation based on the BLASTx result or multiple alignments. Phylogenetic relationships were determined using Clustal x version 1.81. Protein family analysis was performed using PFAM [61]. Results were also compared to those of *N. gonorrhoeae*, *N. meningitidis*, *C. violaceum*, which were the other bacterial species in the *Neisseriaceae* family with complete genome sequences available, where appropriate [29,62-70]. Genes encoding TRAP transporters were located and annotated as described above. Sequence analysis for the presence of signal peptide and transmembrane domains were performed using SignalP v3.0 and TMHMM v2.0 servers respectively [71,72]. Identification of homologs in other bacteria was performed by using BLASTP sequence similarity search against the nr database in NCBI GenBank. The predicted sequences of mature SBPs were submitted to the I-TASSER server for homology modeling using default parameters and available structures of several DctP-type SBP homologs (PDB code: 3B50, 2XA5, 3GYG, 3FXB, 2HPG, and 2CEY) as templates [73]. If multiple homology models were returned, then the best model was selected for further analysis based on the C-score. Quality assessment of the homology model was performed using PROCHECK [74] and ProSA-web [75]. Presence and connectivity of disulfide bonds in the protein were predicted using the DiANNA v1.1 server [76]. Structural alignment of the homology models of SBPs in *L. hongkongensis* and related structures in Protein Data Bank (<http://www.pdb.org>) was performed using the MatchMaker tool of UCSF Chimera with selected structures (PDB code: 2HZK, 2CEY, 2VPN, 2PFZ, 2PFY, and 2ZZV) [77]. Molecular images were generated using UCSF Chimera.

List of abbreviations

ABC: ATP-binding cassette; ATP: Adenosine-5'-triphosphate; BFR: Bacterioferritin; CDS(s): Coding sequence(s); COG: Clusters of orthologous group; CPS-E: Capsular polysaccharide export; CrcB: Camphor resistance; DAACS: Dicarboxylate/amino acid:cation (Na or H) Symporter; DASS: Divalent Anion:Na⁺ Symporter; Dcu: C₄-dicarboxylate uptake; DNA: Deoxyribonucleic acid; DsbB: Disulfide bond oxidoreductase B; DsbD: Disulfide bond oxidoreductase D; EHEC: Enterohemorrhagic *E. coli*; EPEC: Enteropathogenic *E. coli*; EI: Enzyme I; FAT: Fatty acid transporter; G: Guanine; HAMP: Histidine kinase adenyllyl cyclase MCP and phosphatase; HCC: HlyC/CorC; LEE: Locus of enterocyte effacement; MCP(s): Methyl-accepting chemotaxis protein(s); MFP: Membrane fusion protein; MFS: Major facilitator superfamily; PAS: Plasmid achromobacter secretion; PMO: Prokaryotic molybdopterin-cont; P-P-bond: Diphosphate bond; PTS: Phosphotransferase system; RND: Resistance-nodulation-cell-division; TCDB: Transport protein database; TerC: Tellurium ion resistance; TRAP-T: Tripartite ATP-independent periplasmic transporter; VSP: Putative type VI symbiosis/virulence secretory pathway.

Acknowledgements

This work is partly supported by the Research Grant Council Grant, Committee for Research and Conference Grant and University Development Fund, The University of Hong Kong; the HKSAR Research Fund for the Control of Infectious Diseases of the Health, Welfare and Food Bureau. We are grateful to support from the Genome Research Centre, The University of

Hong Kong, and the generous support of Mrs. Carol Yu, Professor Richard Yu, Mr. Hui Hoy and Mr. Hui Ming in the genomic sequencing platform.

Author details

¹State Key Laboratory of Emerging Infectious Diseases, Hong Kong.
²Research Centre of Infection and Immunology, The University of Hong Kong, Hong Kong. ³Carol Yu Centre for Infection, The University of Hong Kong, Hong Kong. ⁴Department of Microbiology, The University of Hong Kong, Hong Kong. ⁵Department of Chemistry, The University of Hong Kong, Hong Kong.

Authors' contributions

PCYW, KYK and SKPL designed and supervised the study. RYYF, GKMW and JLLT annotated the genome. HT and KHS performed bioinformatics analysis. SKPL, RYYF and GKMW drafted the manuscript. All authors read, corrected and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 22 February 2011 Accepted: 17 August 2011

Published: 17 August 2011

References

1. Yuen KY, Woo PCY, Teng JLL, Leung KW, Wong MKM, Lau SKP: *Laribacter hongkongensis* gen. nov., sp. nov., a novel Gram-negative bacterium isolated from a cirrhotic patient with bacteremia and empyema. *J Clin Microbiol* 2001, **39**:4227-4232.
2. Woo PCY, Lau SKP, Teng JLL, Yuen KY: Current status and future directions of *Laribacter hongkongensis*, a novel bacterium associated with gastroenteritis and traveller's diarrhoea. *Curr Opin Infect Dis* 2005, **18**:413-419.
3. Lau SKP, Woo PCY, Hui WT, Li MWS, Teng JLL, Que TL, Yung RWH, Luk WK, Lai RWM, Yuen KY: Use of cefoperazone MacConkey agar for selective isolation of *Laribacter hongkongensis*. *J Clin Microbiol* 2003, **41**:4839-4841.
4. Woo PCY, Kuhnert P, Burnens AP, Teng JLL, Lau SKP, Que TL, Yau HH, Yuen KY: *Laribacter hongkongensis*: a potential cause of infectious diarrhea. *Diagn Microbiol Infect Dis* 2003, **47**:551-556.
5. Woo PCY, Lau SKP, Teng JLL, Que TL, Yung RWH, Luk WK, Lai RWM, Hui WT, Wong SSY, Yau HH, Yuen KY: Association of *Laribacter hongkongensis* in community-acquired gastroenteritis with travel and eating fish: a multicentre case-control study. *Lancet* 2004, **363**:1941-1947.
6. Teng JL, Woo PC, Ma SS, Sit TH, Ng LT, Hui WT, Lau SK, Yuen KY: Ecoepidemiology of *Laribacter hongkongensis*, a novel bacterium associated with gastroenteritis. *J Clin Microbiol* 2005, **43**:919-922.
7. Lau SK, Woo PC, Fan RY, Ma SS, Hui WT, Au SY, Chan LL, Chan JY, Lau AT, Leung KY, Pun TC, She HH, Wong CY, Wong LL, Yuen KY: Isolation of *Laribacter hongkongensis*, a novel bacterium associated with gastroenteritis, from drinking water reservoirs in Hong Kong. *J Appl Microbiol* 2007, **103**:507-515.
8. Ni XP, Ren SH, Sun JR, Xiang HQ, Gao Y, Kong QX, Cha J, Pan JC, Yu H, Li HM: *Laribacter hongkongensis* isolated from a patient with community-acquired gastroenteritis in Hangzhou City. *J Clin Microbiol* 2007, **45**:255-256.
9. Lau SK, Lee LC, Fan RY, Teng JL, Tse CW, Woo PC, Yuen KY: Isolation of *Laribacter hongkongensis*, a novel bacterium associated with gastroenteritis, from Chinese tiger frog. *Int J Food Microbiol* 2009, **129**:78-82.
10. Tsang N, Macnab R, Koshland DE Jr: Common mechanism for repellents and attractants in bacterial chemotaxis. *Science* 1973, **181**:60-63.
11. Falke JJ, Bass RB, Butler SL, Chervitz SA, Danielson MA: The two-component signaling pathway of bacterial chemotaxis: a molecular view of signal transduction by receptors, kinases, and adaptation enzymes. *Annu Rev Cell Dev Biol* 1997, **13**:457-512.
12. Blair DF: How bacteria sense and swim. *Annu Rev Microbiol* 1995, **49**:489-522.
13. Stock AM, Mowbray SL: Bacterial chemotaxis—a field in motion. *Curr Opin Struct Biol* 1995, **5**:744-751.
14. Woo PC, Lau SK, Tse H, Teng JL, Curream SO, Tsang AK, Fan RY, Wong GK, Huang Y, Loman NJ, Snyder LA, Cai JJ, Huang JD, Mak W, Pallen MJ, Lok S, Yuen KY: The complete genome and proteome of *Laribacter hongkongensis* reveal potential mechanisms for adaptations to different temperatures and habitats. *PLoS Genet* 2009, **5**:e1000416.
15. Jap BK, Walian PJ: Structure and functional mechanism of porins. *Physiol Rev* 1996, **76**:1073-1088.
16. Benz R, Bauer K: Permeation of hydrophilic molecules through the outer membrane of gram-negative bacteria. Review on bacterial porins. *Eur J Biochem* 1988, **176**:1-19.
17. Booth IR, Edwards MD, Black S, Schumann U, Miller S: Mechanosensitive channels in bacteria: signs of closure? *Nat Rev Microbiol* 2007, **5**:431-440.
18. Pao SS, Paulsen IT, Saier MH Jr: Major facilitator superfamily. *Microbiol Mol Biol Rev* 1998, **62**:1-34.
19. Tseng TT, Gratwick KS, Kollman J, Park D, Nies DH, Goffeau A, Saier MH Jr: The RND permease superfamily: an ancient, ubiquitous and diverse family that includes human disease and development proteins. *J Mol Microbiol Biotechnol* 1999, **1**:107-125.
20. Blair JM, Piddock LJ: Structure, function and inhibition of RND efflux pumps in Gram-negative bacteria: an update. *Curr Opin Microbiol* 2009, **12**:512-519.
21. Janasch IG, Zientz E, Tran QH, Kroger A, Unden G: C4-dicarboxylate carriers and sensors in bacteria. *Biochim Biophys Acta* 2002, **1553**:39-56.
22. Dimroth P, Schink B: Energy conservation in the decarboxylation of dicarboxylic acids by fermenting bacteria. *Arch Microbiol* 1998, **170**:69-77.
23. Mulligan C, Fischer M, Thomas GH: Tripartite ATP-independent periplasmic (TRAP) transporters in bacteria and archaea. *FEMS Microbiol Rev* 2011, **35**:68-86.
24. Kelly DJ, Thomas GH: The tripartite ATP-independent periplasmic (TRAP) transporters of bacteria and archaea. *FEMS Microbiol Rev* 2001, **25**:405-424.
25. Fischer M, Zhang QY, Hubbard RE, Thomas GH: Caught in a TRAP: substrate-binding proteins in secondary transport. *Trends Microbiol* 2010, **18**:471-478.
26. Forward JA, Behrendt MC, Wyborn NR, Cross R, Kelly DJ: TRAP transporters: a new family of periplasmic solute transport systems encoded by the dctPQM genes of *Rhodobacter capsulatus* and by homologs in diverse gram-negative bacteria. *J Bacteriol* 1997, **179**:5482-5493.
27. Mulligan C, Geertsma ER, Severi E, Kelly DJ, Poolman B, Thomas GH: The substrate-binding protein imposes directionality on an electrochemical sodium gradient-driven TRAP transporter. *Proc Natl Acad Sci USA* 2009, **106**:1778-1783.
28. Johnston JW, Coussens NP, Allen S, Houtman JC, Turner KH, Zaleski A, Ramaswamy S, Gibson BW, Apicella MA: Characterization of the N-acetyl-5-neuraminic acid-binding site of the extracytoplasmic solute receptor (SiaP) of nontypeable *Haemophilus influenzae* strain 2019. *J Biol Chem* 2008, **283**:855-865.
29. Grangeiro TB, Jorge DM, Bezerra WM, Vasconcelos AT, Simpson AJ: Transport genes of *Chromobacterium violaceum*: an overview. *Genet Mol Res* 2004, **3**:117-133.
30. Kos V, Ford RC: The ATP-binding cassette family: a structural perspective. *Cell Mol Life Sci* 2009, **66**:3111-3126.
31. Higgins CF: ABC transporters: physiology, structure and mechanism—an overview. *Res Microbiol* 2001, **152**:205-210.
32. Dimroth P: Primary sodium ion translocating enzymes. *Biochim Biophys Acta* 1997, **1318**:11-51.
33. Barabote RD, Saier MH Jr: Comparative genomic analyses of the bacterial phosphotransferase system. *Microbiol Mol Biol Rev* 2005, **69**:608-634.
34. Fischbach MA, Lin H, Liu DR, Walsh CT: How pathogenic bacteria evade mammalian sabotage in the battle for iron. *Nat Chem Biol* 2006, **2**:132-138.
35. Carniel E: The *Yersinia* high-pathogenicity island: an iron-uptake island. *Microbes Infect* 2001, **3**:561-569.
36. Braun V: Energy-coupled transport and signal transduction through the gram-negative outer membrane via TonB-ExbB-ExbD-dependent receptor proteins. *FEMS Microbiol Rev* 1995, **16**:295-307.
37. Simon N, Coulanges V, Andre P, Vidon DJ: Utilization of exogenous siderophores and natural catechols by *Listeria monocytogenes*. *Appl Environ Microbiol* 1995, **61**:1643-1645.
38. Wandersman C, Stojiljkovic I: Bacterial heme sources: the role of heme, hemoprotein receptors and hemophores. *Curr Opin Microbiol* 2000, **3**:215-220.
39. Cartron ML, Maddocks S, Gillingham P, Craven CJ, Andrews SC: Feo-transport of ferrous iron into bacteria. *Biometals* 2006, **19**:143-157.

40. Harrison PM, Arosio P: **The ferritins: molecular properties, iron storage function and cellular regulation.** *Biochim Biophys Acta* 1996, **1275**:161-203.
41. Gibson TJ, Koonin EV, Musco G, Pastore A, Bork P: **Friedreich's ataxia protein: phylogenetic evidence for mitochondrial dysfunction.** *Trends Neurosci* 1996, **19**:465-468.
42. Cavadini P, O'Neill HA, Benada O, Isaya G: **Assembly and iron-binding properties of human frataxin, the protein deficient in Friedreich ataxia.** *Hum Mol Genet* 2002, **11**:217-227.
43. Taylor BL, Zhulin IB, Johnson MS: **Aerotaxis and other energy-sensing behavior in bacteria.** *Annu Rev Microbiol* 1999, **53**:103-128.
44. Aravind L, Ponting CP: **The cytoplasmic helical linker domain of receptor histidine kinase and methyl-accepting proteins is common to many prokaryotic signalling proteins.** *FEMS Microbiol Lett* 1999, **176**:111-116.
45. Egger LA, Park H, Inouye M: **Signal transduction via the histidyl-aspartyl phosphorelay.** *Genes Cells* 1997, **2**:167-184.
46. Porter SL, Armitage JP: **Phosphotransfer in *Rhodobacter sphaeroides* chemotaxis.** *J Mol Biol* 2002, **324**:35-45.
47. Welch M, Oosawa K, Aizawa S, Eisenbach M: **Phosphorylation-dependent binding of a signal molecule to the flagellar switch of bacteria.** *Proc Natl Acad Sci USA* 1993, **90**:8787-8791.
48. Pereira M, Parente JA, Bataus LA, Cardoso DD, Soares RB, Soares CM: **Chemotaxis and flagellar genes of *Chromobacterium violaceum*.** *Genet Mol Res* 2004, **3**:92-101.
49. Russell CB, Stewart RC, Dahlquist FW: **Control of transducer methylation levels in *Escherichia coli*: investigation of components essential for modulation of methylation and demethylation reactions.** *J Bacteriol* 1989, **171**:3609-3618.
50. Sherris D, Parkinson JS: **Posttranslational processing of methyl-accepting chemotaxis proteins in *Escherichia coli*.** *Proc Natl Acad Sci USA* 1981, **78**:6051-6055.
51. West AH, Martinez-Hackert E, Stock AM: **Crystal structure of the catalytic domain of the chemotaxis receptor methylesterase, CheB.** *J Mol Biol* 1995, **250**:276-290.
52. Porter SL, Warren AV, Martin AC, Armitage JP: **The third chemotaxis locus of *Rhodobacter sphaeroides* is essential for chemotaxis.** *Mol Microbiol* 2002, **46**:1081-1094.
53. DePamphilis ML, Adler J: **Purification of intact flagella from *Escherichia coli* and *Bacillus subtilis*.** *J Bacteriol* 1971, **105**:376-383.
54. Kendall MM, Sperandio V: **Quorum sensing by enteric pathogens.** *Curr Opin Gastroenterol* 2007, **23**:10-15.
55. Nealson KH, Platt T, Hastings JW: **Cellular control of the synthesis and activity of the bacterial luminescent system.** *J Bacteriol* 1970, **104**:313-322.
56. Sperandio V, Torres AG, Kaper JB: **Quorum sensing *Escherichia coli* regulators B and C (QseBC): a novel two-component regulatory system involved in the regulation of flagella and motility by quorum sensing in *E. coli*.** *Mol Microbiol* 2002, **43**:809-821.
57. Sperandio V, Mellies JL, Nguyen W, Shin S, Kaper JB: **Quorum sensing controls expression of the type III secretion gene transcription and protein secretion in enterohemorrhagic and enteropathogenic *Escherichia coli*.** *Proc Natl Acad Sci USA* 1999, **96**:15196-15201.
58. Moreira CG, Palmer K, Whiteley M, Sircili MP, Trabulsi LR, Castro AF, Sperandio V: **Bundle-forming pili and EspA are involved in biofilm formation by enteropathogenic *Escherichia coli*.** *J Bacteriol* 2006, **188**:3952-3961.
59. Clarke MB, Hughes DT, Zhu C, Boedeker EC, Sperandio V: **The QseC sensor kinase: a bacterial adrenergic receptor.** *Proc Natl Acad Sci USA* 2006, **103**:10420-10425.
60. Clarke MB, Sperandio V: **Transcriptional regulation of flhDC by QseBC and sigma FliA) in enterohaemorrhagic *Escherichia coli*.** *Mol Microbiol* 2005, **57**:1734-1749.
61. Bateman A, Birney E, Ceruti L, Durbin R, Eddy SR, Griffiths-Jones S, Howe KL, Marshall M, Sonnhammer EL: **The Pfam protein families database.** *Nucleic Acids Res* 2002, **30**:276-280.
62. Project BNG: **The complete genome sequence of *Chromobacterium violaceum* reveals remarkable and exploitable bacterial adaptability.** *Proc Natl Acad Sci* 2003, **100**:11660-11665.
63. Tettelin H, Saunders NJ, Heidelberg J, Jeffries AC, Nelson KE, Eisen JA, Ketchum KA, Hood DW, Peden JF, Dodson RJ, Nelson WC, Gwinn ML, DeBoy R, Peterson JD, Hickey EK, Haft DH, Salzberg SL, White O, Fleischmann RD, Dougherty BA, Mason T, Ciecko A, Parksey DS, Blair E, Cittone H, Clark EB, Cotton MD, Utterback TR, Khouri H, Qin H, *et al*: **Complete genome sequence of *Neisseria meningitidis* serogroup B strain MC58.** *Science* 2000, **287**:1809-1815.
64. Parkhill J, Achtman M, James KD, Bentley SD, Churcher C, Klee SR, Morelli G, Basham D, Brown D, Chillingworth T, Davies RM, Davis P, Devlin K, Feltwell T, Hamlin N, Holroyd S, Jagels K, Leather S, Moule S, Mungall K, Quail MA, Rajandream MA, Rutherford KM, Simmonds M, Skelton J, Whitehead S, Spratt BG, Barrell BG: **Complete DNA sequence of a serogroup A strain of *Neisseria meningitidis* Z2491.** *Nature* 2000, **404**:502-506.
65. Bentley SD, Vernikos GS, Snyder LA, Churcher C, Arrowsmith C, Chillingworth T, Cronin A, Davis PH, Holroyd NE, Jagels K, Maddison M, Moule S, Rabinowitsch E, Sharp S, Unwin L, Whitehead S, Quail MA, Achtman M, Barrell B, Saunders NJ, Parkhill J: **Meningococcal genetic variation mechanisms viewed through comparative analysis of serogroup C strain FAM18.** *PLoS Genet* 2007, **3**:e23.
66. Chung GT, Yoo JS, Oh HB, Lee YS, Cha SH, Kim SJ, Yoo CK: **Complete genome sequence of *Neisseria gonorrhoeae* NCCP11945.** *J Bacteriol* 2008, **190**:6035-6036.
67. Peng J, Yang L, Yang F, Yang J, Yan Y, Nie H, Zhang X, Xiong Z, Jiang Y, Cheng F, Xu X, Chen S, Sun L, Li W, Shen Y, Shao Z, Liang X, Xu J, Jin Q: **Characterization of ST-4821 complex, a unique *Neisseria meningitidis* clone.** *Genomics* 2008, **91**:78-87.
68. Schoen C, Blom J, Claus H, Schramm-Gluck A, Brandt P, Müller T, Goesmann A, Joseph B, Konietzny S, Kurzai O, Schmitt C, Friedrich T, Linke B, Vogel U, Frosch M: **Whole-genome comparison of disease and carriage strains provides insights into virulence evolution in *Neisseria meningitidis*.** *Proc Natl Acad Sci USA* 2008, **105**:3473-3478.
69. Rusniok C, Vallenet D, Floquet S, Ewles H, Mouzè-Soulama C, Brown D, Lajus A, Buchrieser C, Médigue C, Glaser P, Pelicic V: **NeMeSys: a biological resource for narrowing the gap between sequence and function in the human pathogen *Neisseria meningitidis*.** *Genome Biol* 2009, **10**:R110.
70. Joseph B, Schneiker-Bekel S, Schramm-Gluck A, Blom J, Claus H, Linke B, Schwarz RF, Becker A, Goesmann A, Frosch M, Schoen C: **Comparative genome biology of a serogroup B carriage and disease strain supports a polygenic nature of meningococcal virulence.** *J Bacteriol* 2010, **192**:5363-5377.
71. Emanuelsson O, Brunak S, von Heijne G, Nielsen H: **Locating proteins in the cell using TargetP, SignalP and related tools.** *Nat Protoc* 2007, **2**:953-971.
72. Krogh A, Larsson B, von Heijne G, Sonnhammer EL: **Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes.** *J Mol Biol* 2001, **305**:567-580.
73. Roy A, Kucukural A, Zhang Y: **I-TASSER: a unified platform for automated protein structure and function prediction.** *Nat Protoc* 2010, **5**:725-738.
74. Laskowski RA, MacArthur MW, Moss DS, Thornton JM: **PROCHECK - a program to check the stereochemical quality of protein structures.** *J App Cryst* 1993, **26**:283-291.
75. Wiederstein M, Sippl MJ: **ProSA-web: interactive web service for the recognition of errors in three-dimensional structures of proteins.** *Nucleic Acids Res* 2007, **35**:W407-410.
76. Ferre F, Clote P: **DiANNA 1.1: an extension of the DiANNA web server for ternary cysteine classification.** *Nucleic Acids Res* 2006, **34**:W182-185.
77. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE: **UCSF Chimera—a visualization system for exploratory research and analysis.** *J Comput Chem* 2004, **25**:1605-1612.

doi:10.1186/2045-3701-1-28

Cite this article as: Lau *et al*: Transport genes and chemotaxis in *Laribacter hongkongensis*: a genome-wide analysis. *Cell & Bioscience* 2011 1:28.