

Speculations on the activation of ROS generation in *C. elegans* innate immune signaling

Ransome van der Hoeven, Katie C. McCallum and Danielle A. Garsin*

Department of Microbiology and Molecular Genetics; The University of Texas Health Science Center at Houston; Houston, TX USA

We recently published work demonstrating that ROS (reactive oxygen species) generated by the dual oxidase, Ce-Duox1/BLI-3, in response to infection in *Caenorhabditis elegans* activates the transcription factor SKN-1, initiating a protective response. Moreover, we showed that the crucial innate immune pathway, p38 MAPK signaling, was responsible for relaying the activating signal. In this commentary, we speculate on the signaling pathway upstream of Ce-Duox1/BLI-3 that triggers its activity. Specifically, we hypothesize that a G-protein signaling pathway comprising Gαq - PLCβ - TPA-1 - DKF-2 activates Ce-Duox1/BLI-3. Our rationale is based on work showing that these components are connected to p38 MAPK signaling and innate immunity in the worm, and investigations in other organisms demonstrating that some of these components are involved in dual oxidase activation.

Keywords: dual oxidase, reactive oxygen species, innate immunity, *C. elegans*, signaling

Abbreviations: ROS, reactive oxygen species; NOX, NADPH oxidase; DUOX, dual oxidase

Submitted: 02/03/12

Accepted: 02/20/12

<http://dx.doi.org/10.4161/worm.19767>

*Correspondence to: Danielle A. Garsin;
Email: Danielle.A.Garsin@uth.tmc.edu

Commentary to: van der Hoeven R, McCallum KC, Cruz MR, Garsin DA. Ce-Duox1/BLI-3 generated reactive oxygen species trigger protective SKN-1 activity via p38 MAPK signaling during infection in *C. elegans*. PLoS Pathog 2011; 7:e1002453; PMID:22216003; <http://dx.doi.org/10.1371/journal.ppat.1002453>

Coordinated regulation of immunity is not only crucial for fighting invading pathogens, but also vital for safeguarding host tissues from injury due to excessive reactions. Unchecked immune responses can lead to tissue damage, disease and sometimes death of the host. An example of an immune response that can potentially harm host tissue is the purposeful generation of reactive oxygen species (ROS) by NADPH oxidase enzymes.

The first characterized NADPH oxidase was the phagocytic gp91phox/Nox2. It produces ROS in the phagolysosomes of neutrophils and other immune cells, contributing to the destruction of invading microbes.^{1,2} The robust consumption of

molecular oxygen and the generation of superoxide anions by this enzyme was termed the oxidative burst. Six other homologs of Nox2 (Nox1, Nox3, Nox4, Nox5, Duox1 and Duox2) were subsequently identified in the human genome and found to be expressed in a wide range of tissues, giving rise to the Nox/Duox (NADPH oxidase/dual oxidase) family of proteins. All members of this family of proteins retain a catalytic C-terminal domain comprising NADPH and FAD binding sites and two, membrane-bound heme moieties. In addition to the C-terminal catalytic domain, calcium binding EF hand motifs are found in dual oxidases and Nox5. Dual oxidases also possess an N-terminal heme-containing peroxidase domain. Nox and Duox enzymes occur in plants, algae, fungi, amoeba, nematodes, echinoderms, urochordates, insects, fish, reptiles, birds and mammals, but are absent in prokaryotes.¹ Therefore, a variety of systems can be used to study these proteins.

The genome of *C. elegans* contains two genes that encode for dual oxidases, Ce-Duox1/BLI-3 and Ce-Duox2, but lacks genes encoding for Nox enzymes.³ However, Ce-Duox2 may be a pseudogene because it does not appear to be expressed, and a deletion mutant has no phenotype.^{4,5} In initial studies, a functional role for Ce-Duox1/BLI-3 in the biogenesis of the cuticle was described, and it was localized to the hypodermis.³ Ce-Duox1/BLI-3 was postulated to generate hydrogen peroxide by the C-terminal catalytic domain for use by the N-terminal peroxidase domain. Specifically, the model proposes that the peroxidase domain uses hydrogen peroxide as an electron donor to

generate radical tyrosine molecules that react with one another, creating protein crosslinks that stabilize and strengthen the cuticle. Consistent with this model, loss of Ce-Duox1/BLI-3 results in blistering and bubbling of the cuticle.³

Another role for Ce-Duox1/BLI-3, which others and we have established, is the protective generation of ROS in the intestine in response to infection.^{4,6} Though overall this response was beneficial, there was evidence that the elevated levels of host-generated ROS caused cellular damage.^{7,8} We speculated that to maintain redox homeostasis during infection, the worm might simultaneously engage oxidative stress response programs, such as the phase II detoxification response, to scavenge free radicals and other reactive molecules using glutathione.⁹ Phase II detoxification of xenobiotic and chemically induced oxidative stress in *C. elegans* has been extensively studied and shown to be regulated by SKN-1, which is distantly related to the human Nuclear factor erythroid related factor (Nrf).¹⁰⁻¹² In mice, Nrf2 mediated regulation of redox status was shown to modulate immune responses during inflammation.¹³ However, a role for SKN-1 in responding to infection in the worm had not been demonstrated. Additionally, a link between ROS produced by NOX/DUOX enzymes and the activation of the Nrf2/SKN-1 family of transcription factors had not been established in any organism.

In van der Hoeven et al., we demonstrated, using a variety of techniques, that SKN-1 is activated in the intestine of the worm in response to the human pathogens *Enterococcus faecalis* and *Pseudomonas aeruginosa*.⁹ For example, we observed increased transcription of several SKN-1 dependent genes, such as *gcs-1*, *gst-4*, *gst-5*, *gst-7* and *gst-10*, by qRT-PCR and promoter fusions to *gfp*, in some cases. Moreover, we established that ROS produced by Ce-Duox1/BLI-3 activates SKN-1 through the p38 MAPK signaling pathway, similarly to that previously shown for chemically induced oxidative stress.¹⁴ The p38 MAPK signaling pathway is central to *C. elegans* innate immune response and is comprised of the Toll/IL-1 receptor domain protein, TIR-1, the MAPKKK, NSY-1, the MAPKK, SEK-1,

and the MAPK, PMK-1.^{15,16} Our analysis of the p38 MAPK pathway revealed that components NSY-1, SEK-1 and PMK-1 are required for activation of SKN-1, while TIR-1, which is essential in responding to pathogens, is not required.⁹ Finally and most importantly, we showed that SKN-1 positively impacts survival during infection. Loss of *skn-1* decreased resistance to the pathogens, whereas overexpression resulted in enhanced survival. Overall, SKN-1 is activated via signaling through the p38 MAPK pathway in response to the oxidative burst generated by Ce-Duox1/BLI-3 during infection of the worm intestine.⁹

Currently, we are focused on understanding how Ce-Duox1/BLI-3 is triggered to produce ROS in response to pathogen. Interestingly, the p38 MAPK pathway was shown to upregulate *DUOX* gene expression in *Drosophila*.¹⁷ However by western analysis, we see no increase in Ce-Duox1/BLI-3 protein levels in infected, as compared with uninfected, *C. elegans* (Garsin lab, unpublished data), leading us to speculate that activation of Ce-Duox1/BLI-3 happens post-translationally. Genetic evidence in *Drosophila* has shown the G α q-phospholipase C β (PLC β) pathway utilizes secondary messengers, Inositol triphosphate (Ins(1,4,5) P_3) and Ca²⁺, to regulate *DUOX* activity.¹⁸ It was proposed that a microbe-derived ligand triggers a G-protein coupled receptor, leading to the release of G α q, which subsequently activates PLC β to hydrolyze phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5) P_2) into inositol 1,4,5-trisphosphate (Ins(1,4,5) P_3) and diacylglycerol (DAG). Next, Ins(1,4,5) P_3 binds to the Ins(1,4,5) P_3 receptor (Ins(1,4,5) P_3 R), located on the endoplasmic reticulum (ER) membrane, causing the induction and release of intracellular calcium. The released calcium is thought to bind to the EF hands, modulating the activity of *DUOX*.^{17,18} In another study using mammalian cell lines derived from the intestinal epithelium, Duox activity was shown to be dependent on intracellular calcium levels and to be regulated by protein kinase C and protein kinase A.¹⁹ DAG released by the hydrolysis of PtdIns(4,5) P_2 binds to protein kinase C, while protein kinase A is activated by the binding of cAMP synthesized by adeny-

cyclase. The study highlights that phosphorylation of Duox at specific serine residues enhances its sensitivity to calcium, further modulating its activity.¹⁹ Boots et al. demonstrated activation of Duox1 in immortalized human bronchial epithelial (HBE1) cells by extracellular ATP and purinergic receptor stimulation.²⁰ Based on this information, we postulate that these components also may regulate Ce-Duox1/BLI-3.

Though no evidence currently supports a role for the above-mentioned components in ROS generation in the worm, many do appear to affect susceptibility to pathogen, as would be predicted if they did regulate Ce-Duox1/BLI-3. For example, work by Kawli et al. showed that G α q (EGL-30) and PLC β (EGL-8), in *C. elegans*, regulated both pathogen immune responses and oxidative stress responses through the p38 MAPK pathway.²¹ DAG, released by PLC β hydrolysis of membrane lipids, also regulates p38 MAPK activity in the intestine. DAG can additionally interact with C1 domain containing proteins TPA-1 (protein kinase C δ) and DKF2 (protein kinase D2), which are expressed in the *C. elegans* intestine.²²⁻²⁴ Interestingly, the TPA-1 - DKF2 module induces the production of immune effectors and the oxidative stress response, via the p38 MAPK pathway. Furthermore, worms lacking DFK-2 and TPA-1 were hypersensitive to killing by pathogenic bacteria.²³ Taken together, there is good evidence that the G α q - PLC β - TPA-1 - DKF-2 pathway activates the p38 MAPK pathway by DAG signaling. Considering that we have established ROS generated by Ce-Duox1 works through the p38 MAPK pathway to activate the SKN-1-dependent oxidative stress response, it is plausible that a G α q - PLC β - TPA-1 - DKF-2 pathway could regulate Ce-Duox1/BLI-3 activity in the worm, and that the ROS generated by BLI-3 acts as a signaling molecule to link the upstream G-protein signaling pathway to the downstream p38 MAPK signaling module (Fig. 1). Validating this hypothesis would enable us to explore the possibility of identifying G-protein coupled receptors that regulate this response by screening for differences in pathogen-triggered ROS production.




Figure 1. Hypothetical model depicting the activation of Ce-Duox1/BLI-3 and the p38 MAPK pathway in *C. elegans* during infection.

In conclusion, we have established a connection between ROS produced by Ce-Duox1/BLI-3 and the activation of the oxidative stress transcription factor SKN-1, in a p38 MAPK dependent

manner, during *C. elegans* infection. The next challenge is to identify the regulatory network that controls Ce-Duox1/BLI-3 activity in response to pathogen invasion.

Acknowledgments

This work was supported by the National Institutes of Health grant R01AI076406 to D.A.G.

References

- Aguirre J, Lambeth JD. Nox enzymes from fungus to fly to fish and what they tell us about Nox function in mammals. *Free Radic Biol Med* 2010; 49:1342-53; PMID:20696238; <http://dx.doi.org/10.1016/j.freeradbiomed.2010.07.027>
- Bedard K, Krause KH. The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev* 2007; 87:245-313; PMID:17237347; <http://dx.doi.org/10.1152/physrev.00044.2005>
- Edens WA, Sharling L, Cheng G, Shapira R, Kinkade JM, Lee T, et al. Tyrosine cross-linking of extracellular matrix is catalyzed by Duox, a multidomain oxidase/peroxidase with homology to the phagocyte oxidase subunit gp91phox. *J Cell Biol* 2001; 154:879-91; PMID:11514595; <http://dx.doi.org/10.1083/jcb.200103132>
- Chavez V, Mohri-Shiomi A, Garsin DA. Ce-Duox1/BLI-3 generates reactive oxygen species as a protective innate immune mechanism in *Caenorhabditis elegans*. *Infect Immun* 2009; 77:4983-9.
- Hill AA, Hunter CP, Tsung BT, Tucker-Kellogg G, Brown EL. Genomic analysis of gene expression in *C. elegans*. *Science* 2000; 290:809-12; PMID:11052945; <http://dx.doi.org/10.1126/science.290.5492.809>
- Jain C, Yun M, Politz SM, Rao RP. A pathogenesis assay using *Saccharomyces cerevisiae* and *Caenorhabditis elegans* reveals novel roles for yeast AP-1, Yap1, and host dual oxidase BLI-3 in fungal pathogenesis. *Eukaryot Cell* 2009; 8:1218-27; PMID:19502579; <http://dx.doi.org/10.1128/EC.00367-08>
- Chávez V, Mohri-Shiomi A, Maadani A, Vega LA, Garsin DA. Oxidative stress enzymes are required for DAF-16-mediated immunity due to generation of reactive oxygen species by *Caenorhabditis elegans*. *Genetics* 2007; 176:1567-77; PMID:17483415; <http://dx.doi.org/10.1534/genetics.107.072587>
- Mohri-Shiomi A, Garsin DA. Insulin signaling and the heat shock response modulate protein homeostasis in the *Caenorhabditis elegans* intestine during infection. *J Biol Chem* 2008; 283:194-201; PMID:17951251; <http://dx.doi.org/10.1074/jbc.M707956200>
- van der Hoeven R, McCallum KC, Cruz MR, Garsin DA. Ce-Duox1/BLI-3 generated reactive oxygen species trigger protective SKN-1 activity via p38 MAPK signaling during infection in *C. elegans*. *PLoS Pathog* 2011; 7:e1002453; PMID:22216003; <http://dx.doi.org/10.1371/journal.ppat.1002453>
- An JH, Blackwell TK. SKN-1 links *C. elegans* mesodermal specification to a conserved oxidative stress response. *Genes Dev* 2003; 17:1882-93; PMID:12869585; <http://dx.doi.org/10.1101/gad.1107803>
- An JH, Vranas K, Lucke M, Inoue H, Hisamoto N, Matsumoto K, et al. Regulation of the *Caenorhabditis elegans* oxidative stress defense protein SKN-1 by glycogen synthase kinase-3. *Proc Natl Acad Sci U S A* 2005; 102:16275-80; PMID:16251270; <http://dx.doi.org/10.1073/pnas.0508105102>
- Tullet JM, Hertweck M, An JH, Baker J, Hwang JY, Liu S, et al. Direct inhibition of the longevity-promoting factor SKN-1 by insulin-like signaling in *C. elegans*. *Cell* 2008; 132:1025-38; PMID:18358814; <http://dx.doi.org/10.1016/j.cell.2008.01.030>
- Cho HY, Kleeberger SR. Nrf2 protects against airway disorders. *Toxicol Appl Pharmacol* 2010; 244:43-56; PMID:19646463; <http://dx.doi.org/10.1016/j.taap.2009.07.024>
- Inoue H, Hisamoto N, An JH, Oliveira RP, Nishida E, Blackwell TK, et al. The *C. elegans* p38 MAPK pathway regulates nuclear localization of the transcription factor SKN-1 in oxidative stress response. *Genes Dev* 2005; 19:2278-83; PMID:16166371; <http://dx.doi.org/10.1101/gad.1324805>
- Kim DH, Feinbaum R, Allosing G, Emerson FE, Garsin DA, Inoue H, et al. A conserved p38 MAP kinase pathway in *Caenorhabditis elegans* innate immunity. *Science* 2002; 297:623-6; PMID:12142542; <http://dx.doi.org/10.1126/science.1073759>

16. Liberati NT, Fitzgerald KA, Kim DH, Feinbaum R, Golenbock DT, Ausubel FM. Requirement for a conserved Toll/interleukin-1 resistance domain protein in the *Caenorhabditis elegans* immune response. *Proc Natl Acad Sci U S A* 2004; 101:6593-8; PMID: 15123841; <http://dx.doi.org/10.1073/pnas.0308625101>
17. Ha EM, Lee KA, Seo YY, Kim SH, Lim JH, Oh BH, et al. Coordination of multiple dual oxidase-regulatory pathways in responses to commensal and infectious microbes in drosophila gut. *Nat Immunol* 2009; 10:949-57; PMID:19668222; <http://dx.doi.org/10.1038/ni.1765>
18. Ha EM, Lee KA, Park SH, Kim SH, Nam HJ, Lee HY, et al. Regulation of DUOX by the Galphaq-phospholipase Cbeta-Ca2+ pathway in Drosophila gut immunity. *Dev Cell* 2009; 16:386-97; PMID:19289084; <http://dx.doi.org/10.1016/j.devcel.2008.12.015>
19. Rigutto S, Hoste C, Grasberger H, Milenkovic M, Communi D, Dumont JE, et al. Activation of dual oxidases Duox1 and Duox2: differential regulation mediated by camp-dependent protein kinase and protein kinase C-dependent phosphorylation. *J Biol Chem* 2009; 284:6725-34; PMID:19144650; <http://dx.doi.org/10.1074/jbc.M806893200>
20. Boots AW, Hristova M, Kasahara DI, Haenen GR, Bast A, van der Vliet A. ATP-mediated activation of the NADPH oxidase DUOX1 mediates airway epithelial responses to bacterial stimuli. *J Biol Chem* 2009; 284:17858-67; PMID:19386603; <http://dx.doi.org/10.1074/jbc.M809761200>
21. Kawli T, Wu C, Tan MW. Systemic and cell intrinsic roles of Gqalpha signaling in the regulation of innate immunity, oxidative stress, and longevity in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A* 2010; 107:13788-93; PMID:20647387; <http://dx.doi.org/10.1073/pnas.0914715107>
22. Feng H, Ren M, Chen L, Rubin CS. Properties, regulation, and in vivo functions of a novel protein kinase D: *Caenorhabditis elegans* DKF-2 links diacylglycerol second messenger to the regulation of stress responses and life span. *J Biol Chem* 2007; 282:31273-88; PMID:17728253; <http://dx.doi.org/10.1074/jbc.M701532200>
23. Ren M, Feng H, Fu Y, Land M, Rubin CS. Protein kinase D is an essential regulator of *C. elegans* innate immunity. *Immunity* 2009; 30:521-32; PMID: 19371715; <http://dx.doi.org/10.1016/j.immuni.2009.03.007>
24. Ziegler K, Kurz CL, Cypowyj S, Couillault C, Pophillat M, Pujol N, et al. Antifungal innate immunity in *C. elegans*: PKCdelta links G protein signaling and a conserved p38 MAPK cascade. *Cell Host Microbe* 2009; 5:341-52; PMID:19380113; <http://dx.doi.org/10.1016/j.chom.2009.03.006>