



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

- 8 Patrick DM, Sbihi H, Dai DLY, et al. Decreasing antibiotic use, the gut microbiota, and asthma incidence in children: a population-based study and a prospective cohort study *Lancet Respir Med* 2020; published online March 24. [https://doi.org/10.1016/S2213-2600\(20\)30052-7](https://doi.org/10.1016/S2213-2600(20)30052-7).
- 9 Zven SE, Susi A, Mitre E, Nylund CM. association between use of multiple classes of antibiotic in infancy and allergic disease in childhood. *JAMA Pediatr* 2019; published online Dec 20. DOI:10.1001/jamapediatrics.2019.4794.
- 10 Stokholm J, Sevelsted A, Bønnelykke K, Bisgaard H. Maternal propensity for infections and risk of childhood asthma: a registry-based cohort study. *Lancet Respir Med* 2014; 2: 631–37.

Inhaled corticosteroids and COVID-19-related mortality: confounding or clarifying?

Inhaled corticosteroids (ICSs) are the mainstay of anti-inflammatory therapy for asthma and chronic obstructive pulmonary disease (COPD).^{1,2} Studies have shown worse outcomes in patients with COVID-19 who have been admitted to hospital and have comorbidities including chronic lung diseases.^{3–5} Whether ICSs protect against COVID-19 or contribute to worse outcomes from COVID-19 has been debated.^{6,7} ICS use might reduce antiviral immunity and increase the frequency of pneumonia in patients with COPD.⁸ However, ICS use reduces the frequency of exacerbations¹ and might even reduce replication of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2),⁹ supporting the case for protection against COVID-19.


In *The Lancet Respiratory Medicine*, Anna Schultze and colleagues¹⁰ use UK electronic primary care records to retrospectively interrogate associations between current ICS use (defined as a prescription within 4 months) and COVID-19-related deaths in patients with asthma and COPD. The COPD cohort (n=148 557) consisted of patients currently using ICSs and a long-acting β -agonists (LABA), with or without a long-acting muscarinic antagonist (LAMA), compared with patients using a LABA and LAMA. Although comorbidities were similar between groups, exacerbations in the previous year were more frequent in the ICS combination group, reflecting that ICSs are used in patients with exacerbations. In the asthma cohort (n=818 490), current ICS users were compared with those using short-acting β agonists (SABAs) only. The mean age, number of comorbidities, and number of exacerbations in the past year were lower in non-ICS users, indicating that between-group analyses might be confounded by baseline differences.

In the COPD cohort, ICS use was associated with an increased risk of COVID-19-related death after adjusting

for relevant factors, including age and comorbidities (hazard ratio [HR] 1.39 [95% CI 1.10–1.76]). A sensitivity analysis showed that the risk of death in the COPD population was highest with ICS–LABA–LAMA (triple therapy; adjusted HR 1.43 [1.12–1.83]), and lower and non-significant with ICS–LABA (1.29 [0.96–1.74]), a difference that cannot be explained by ICS use. These sensitivity analyses suggest factors driving increased risk in the ICS combination group that are not attributable to ICS itself, including confounding by treatment indication (ie, patients prescribed triple therapy have an increased disease burden and worse prognosis). A negative control analysis supported this hypothesis, with increased risk of non-COVID-19-related death in ICS users.

In the asthma cohort, no increased risk of COVID-19-related death was found in low-to-medium dose ICS users compared with non-ICS users (adjusted HR 1.14 [95% CI 0.85–1.54]), suggesting ICS use had no bearing on COVID-19-related mortality. Schultze and colleagues found increased risk of COVID-19-related mortality in high-dose ICS users compared with non-ICS users (1.55 [1.10–2.18]), but no increase in non-COVID-19-related deaths. This result might relate to underlying disease characteristics beyond those captured in the health records, such as increased susceptibility to viral infections with more severe asthma.

Patients with asthma and COPD are understandably concerned about developing COVID-19. They are also concerned whether their medication affects their risk of becoming infected or their prognosis if they do develop COVID-19, and they look to their clinicians for answers to these questions. Does this analysis help answer their questions? We think it provides some insights, but not conclusive answers. The negative results for use of low-to-medium dose ICS in asthma and ICS–LABA in COPD argue against the hypothesis that ICS use



Published Online
September 24, 2020
[https://doi.org/10.1016/S2213-2600\(20\)30447-1](https://doi.org/10.1016/S2213-2600(20)30447-1)

See [Articles](#) page 1106

For the [Global Initiative for Asthma](#) see <http://www.ginasthma.org>

increases the risk of COVID-19-related mortality. For the hypothesis that ICS use protects against COVID-19-related mortality, the results rule out a benefit large enough to overcome the effects of confounding factors, but do not completely exclude a smaller benefit.

Overall, the analysis is confounded and does not provide definitive answers that patients and clinicians need, although it hints that ICS use does not provide a strong protective effect. Similar to Schultze and colleagues, we believe that had the analysis taken into account clinical factors, such as disease severity and history of exacerbations, which might have influenced the choice of maintenance therapy, it might have reached different conclusions about possible harms. ICSs are used to reduce future risk of events including exacerbations and mortality,^{1,2} therefore, ICS use inevitably identifies individuals with an increased disease burden associated with increased future risk. Analyses of associations between ICS use and COVID-19-related outcomes in real-life datasets cannot escape this issue, but the comprehensive analysis reported by Schultze and colleagues in a large sample of almost 1 million people is a valiant attempt to provide some clarity despite the confounding by treatment indication observed.

The analysis does not completely resolve whether regular ICS therapy for asthma or COPD either decreases or increases risk of death from COVID-19. This finding is in contrast with the very real harm patients requiring ICS therapy for their asthma or COPD might be at risk if they stop treatment because of unfounded concerns related to their effects in COVID-19. Until more information is available, patients with asthma and COPD who are stable while using ICS must continue on their treatment during the ongoing COVID-19 pandemic.

DS reports personal fees from AstraZeneca, Boehringer Ingelheim, Chiesi, Cipla, Genentech, GlaxoSmithKline, Glenmark, Gossamerbio, Menarini, Mundipharma, Novartis, Peptinnovate, Pfizer, Pulmatrix, Theravance, and Verona outside of the submitted work. DMGH reports personal fees from AstraZeneca, Chiesi, GlaxoSmithKline, Pfizer, and Sanofi, and personal fees and non-financial support from Boehringer Ingelheim and Novartis outside of the submitted work.

Copyright © 2020 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

*Dave Singh, David M G Halpin
dsingh@meu.org.uk

University of Manchester, Manchester University NHS Foundation Trust, Manchester, M23 9QZ, UK (DS); and University of Exeter Medical School, College of Medicine and Health, University of Exeter, Exeter, Devon, UK (DMGH)

- 1 Singh D, Agusti A, Anzueto A, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive lung disease: the GOLD science committee report 2019. *Eur Respir J* 2019; **53**: 1900164.
- 2 Reddel HK, FitzGerald JM, Bateman ED, et al. GINA 2019: a fundamental change in asthma management: treatment of asthma with short-acting bronchodilators alone is no longer recommended for adults and adolescents. *Eur Respir J* 2019; **53**: 190104.
- 3 Alqahtani JS, Oyelade T, Aldhahir AM, et al. Prevalence, severity and mortality associated with COPD and smoking in patients with COVID-19: a rapid systematic review and meta-analysis. *PLoS One* 2020; **15**: e0233147.
- 4 Argenziano MG, Bruce SL, Slater CL, et al. Characterization and clinical course of 1000 patients with coronavirus disease 2019 in New York: retrospective case series. *BMJ* 2020; **369**: m1996.
- 5 Docherty AB, Harrison EM, Green CA, et al. Features of 20133 UK patients in hospital with covid-19 using the ISARIC WHO clinical characterisation protocol: prospective observational cohort study. *BMJ* 2020; **369**: m1985.
- 6 Halpin DMG, Singh D, Hadfield RM. Inhaled corticosteroids and COVID-19: a systematic review and clinical perspective. *Eur Respir J* 2020; **55**: 2001009.
- 7 Halpin DMG, Faner R, Sibila O, Badia JR, Agusti A. Do chronic respiratory diseases or their treatment affect the risk of SARS-CoV-2 infection? *Lancet Respir Med* 2020; **8**: 436–38.
- 8 Singanayagam A, Johnston SL. Long-term impact of inhaled corticosteroid use in asthma and chronic obstructive pulmonary disease (COPD): review of mechanisms that underlie risks. *J Allergy Clin Immunol* 2020; published online Jan 14. <https://doi.org/10.1016/j.jaci.2019.12.907>.
- 9 Jeon S, Ko M, Lee J, et al. Identification of antiviral drug candidates against SARS-CoV-2 from FDA-approved drugs. *Antimicrob Agents Chemother* 2020; **64**: e00819-20.
- 10 Schultze A, Walker AJ, MacKenna B, et al. Risk of COVID-19 related death among patients with chronic obstructive pulmonary disease or asthma prescribed inhaled corticosteroids: an observational cohort study using the OpenSAFELY platform. *Lancet Respir Med* 2020; published online Sept 24. [https://doi.org/10.1016/S2213-2600\(20\)30415-X](https://doi.org/10.1016/S2213-2600(20)30415-X).



ECMO for severe ARDS associated with COVID-19: now we know we can, but should we?

Published Online
August 13, 2020
[https://doi.org/10.1016/S2213-2600\(20\)30357-X](https://doi.org/10.1016/S2213-2600(20)30357-X)
See [Articles](#) page 1121

The initial months of the COVID-19 pandemic were dominated by studies reporting poor and varied outcomes in patients who developed severe acute respiratory distress syndrome (ARDS) associated with the disease. Variable mortality could have been related to heterogeneity in patient populations and pre-pandemic intensive care infrastructure, resource constraints imposed during the

pandemic, and variability in duration of follow-up. As the pandemic has evolved, lower mortality attributable to the disease has been reported. For instance, in a cohort of 742 patients with COVID-19-associated ARDS from Spain, mortality for severe ARDS was 39%,¹ similar to findings of a large epidemiological study of patients with severe ARDS who did not have COVID-19.²