



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Letter to the Editor Regarding “Pulmonary Embolism Prophylaxis in Patients With COVID-19: an Emerging Issue”, Heart Lung Circ. 2021;30(10):1435–41.



To the Editor,

I read the article by Sanidas and colleagues [1] with great interest and congratulate the authors on their excellent work. However, I would like to draw attention to several points. Recent findings suggest that SARS-CoV-2 controls the expression of host regulatory molecules such as miRNAs and transcription factors (TFs), often regulating gene expression at transcriptional and post-transcriptional levels and forming highly complex network interactions [2]. Research has identified four miRNAs (hsa-mir-9-5p, hsa-mir-324-3p, hsa-mir-1827, and hsa-mir-1277-5p) and TFs (EP300, MYC, E2F1, and SP1) that co-regulate important target genes responsible for SARS-CoV-2 host invasion [2]. Studies have found evidence supporting common molecular trajectories of SARS-CoV-2 with clotting, pulmonary embolism, pulmonary oedema, and systematic inflammation, showing both fibrinolysis as a major pathway leading to D-dimer increase and cytokines TGF- β and TNF- α as major regulators of fibrinolysis controlling proteins PAI-1 and plasminogen activators also highlighting the role of hsa-mir-9-5p, hsa-mir-324-3p, EP300, and SP1 on pulmonary embolism [2]. It has also been reported that in the circulating exosomes of high D-dimer COVID-19 patients (cut-off 3 $\mu\text{g}/\text{mL}$) there is significant miR-424 upregulation and miR-103a, miR-145, and miR-885 downregulation. Exosomal miR-424 is an independent thromboembolic event predictor in COVID-19 patients [3,4], whereas miR-103a independently regulates D-dimer levels [4]. Furthermore, miR-885 targets von Willebrand factor and miR-145 targets tissue factor [3,4], showing negative correlations with their respective targets [3]. The findings of Sanidas and colleagues [1] add significant information to previously published data; however, also evaluating these

aspects would be useful for better understanding of the interplay between pulmonary embolism in patients with COVID-19 and its complex regulatory network.

Conflict of Interest

None declared.

Salvatore Patanè, MD*

Cardiologia Ospedale San Vincenzo Taormina (Me) Azienda Sanitaria Provinciale di Messina, Contrada Sirina, Taormina (Messina), Italy

*Corresponding author at: Cardiologia Ospedale San Vincenzo Taormina (Me) Azienda Sanitaria Provinciale di Messina, Contrada Sirina, 98039 Taormina (Messina), Italy.

Email: patane@libero.it

Received 18 September 2021; received in revised form 9 October 2021; accepted 12 October 2021; online published-ahead-of-print 4 November 2021

References

- [1] Sanidas E, Grassos C, Papadopoulos D, Velliou M, Barbetseas J. Pulmonary embolism prophylaxis in patients with COVID-19: an emerging issue. *Heart Lung Circ.* 2021;30(10):1435–41.
- [2] Khurana P, Gupta A, Sugadev R, Sharma YK, Varshney R, Ganju L. nSARS-Cov-2, pulmonary edema and thrombosis: possible molecular insights using miRNA-gene circuits in regulatory networks. *ExRNA.* 2020;2(1):16.
- [3] Plowman T, Lagos D. Non-coding RNAs in COVID-19: emerging insights and current questions. *Non-coding RNA.* 2021;7(3):54.
- [4] Gambardella J, Sardu C, Morelli MB, Messina V, Marfella R, Maggi P. Exosomal microRNAs drive thromboembolism in COVID-19. *Circulation.* 2020;142(Suppl_4):A221.