
Restoration cannot be scaled up globally to save reefs from loss and degradation

In the format provided by the
authors and unedited

Supplementary Information

Restoration cannot be scaled up globally to save reefs from loss and degradation

Clelia Mula¹, Corey J. A. Bradshaw^{2,3,4}, Mar Cabeza¹, Federica Manca¹, Simone Montano^{5,6}, Giovanni Strona^{7*}

*giovanni.strona@ec.europa.eu

¹Faculty of Biological and Environmental Sciences, Organismal and Evolutionary Biology Research Programme, University of Helsinki, PO Box 65 Viikinkaari 1, Helsinki, 00014 Finland ²Global Ecology | *Partuyarta Ngadluku Wardli Kuu*, College of Science and Engineering, Flinders University, GPO Box 2100, Adelaide, South Australia 5001, Australia

²Australian Research Council Centre of Excellence for Australian Biodiversity and Heritage, Wollongong, New South Wales, Australia

³Australian Research Council Centre of Excellence for Indigenous and Environmental Histories and Futures, Cairns, Queensland, Australia

⁴Global Ecology | *Partuyarta Ngadluku Wardli Kuu*, College of Science and Engineering, Flinders University, GPO Box 2100, Adelaide, South Australia 5001, Australia

⁵Department of Earth and Environmental Sciences (DISAT), University of Milano-Bicocca, Piazza Della Scienza 1, 20126, Milan, Italy

⁶MaRHE Centre (Marine Research and High Education Center), Magoodhoo Island, Faafu Atoll, 12030, Republic of Maldives

⁷European Commission, Joint Research Centre (JRC), Ispra, Italy




Figure S1 | Pairwise correlation (Pearson’s R²) between the independent variables used to predict the choice of restoration sites. Total n observations = 3324.




Figure S2 | Pairwise correlation (Pearson's R2) between the independent variables used to predict coral restoration success. Total n observations = 134.