

Endovascular revascularisation of acute tandem vertebrobasilar artery occlusion: seven case series with literature reviews

Haihua Yang,^{1,2} Ning Ma,¹ Shiyong Zhang,³ Xiaochuan Huo,¹ Feng Gao,¹ Xuan Sun,¹ Dapeng Mo,¹ Zhongrong Miao¹

To cite: Yang H, Ma N, Zhang S, *et al.* Endovascular revascularisation of acute tandem vertebrobasilar artery occlusion: seven case series with literature reviews. *Stroke and Vascular Neurology* 2018;**3**: e000125. doi:10.1136/svn-2017-000125

Received 14 November 2017

Revised 24 December 2017

Accepted 4 January 2018

Published Online First

24 January 2018

ABSTRACT

Background The outcome of acute ischaemic stroke due to tandem vertebrobasilar artery occlusion was poor. Endovascular revascularisation may be a positive approach for acute basilar artery occlusion combined with vertebral ostium stenosis or occlusion. We reported seven patients with acute vertebrobasilar tandem occlusion by using angioplasty or stenting for proximal lesion and thrombectomy for distal occlusion.

Materials and methods Consecutive patients with acute tandem vertebrobasilar artery occlusion at two centres were included in this study. We retrospectively analysed the clinical, technical and functional outcomes of the patients.

Results From March 2016 to March 2017, seven patients were included. The mean age and National Institute of Health Stroke Scale score at admission was 57 years and 22, respectively. A reverse approach was used in five patients. The Thrombolysis in Cerebral Infarction score 2b-3 was acquired in all patients. There were no procedural complications. The modified Rankin Scale score was 1–2 for three patients and 4 for one patient at 3 months. Three patients died at 3 months of follow-up.

Conclusions Endovascular revascularisation may be feasible for acute tandem vertebrobasilar artery occlusion, and more researches are needed.

INTRODUCTION

In patients with isolated acute basilar artery (BA) occlusion, the use of modern thrombectomy devices for recanalisation was a better approach to improve recanalisation rates and clinical outcomes.^{1–3} At the same time, the clinical outcome of acute occlusion of the ostial vertebral artery (VA) concomitant occlusion of BA was generally extreme poor.^{4–6} The benefit of the second thrombectomy device has been verified in patients with acute tandem lesions in anterior circulation in Endovascular Treatment for Small Core and Anterior Circulation Proximal Occlusion with Emphasis on Minimizing CT to Recanalization Times (ESCAPE) and Revascularization with Solitaire FR device VS. Best Medical Therapy in the Treatment of Acute Stroke due to Anterior Circulation Large Vessel Occlusion Presenting within

Eight hours of Symptom onset (REVASCAT) trials.^{7,8} However, the efficacy of these devices for patients with tandem vertebrobasilar arterial occlusion remains uncertain.

In this paper, we reported seven patients with acute vertebrobasilar artery tandem lesions treated with endovascular revascularisation techniques to achieve successful recanalisation.

MATERIALS AND METHODS

Patients

This is a retrospective study from two stroke centres in Beijing, China. The time interval of this study ranged from March 2016 to March 2017. The clinical data of patients undergoing endovascular revascularisation of acute vertebrobasilar artery tandem occlusion were collected and analysed. The local ethics committee approved the study and informed consent of the patient was obtained.

Endovascular treatment procedure

The procedure of endovascular treatment was performed under general anaesthesia in all patients. A 6 Fr guiding catheter was placed at the proximal subclavian artery. The procedure was performed in all patients using the endovascular revascularisation approach through the occlusive or stenotic vertebral artery (dirty-road path). The microguidewire crossed the lesion of VA occlusion or stenosis. Predilation of the VA lesion was performed using a small balloon and then digital subtraction angiography (DSA) was performed to understand the diameter of the VA. Then reverse approach (mechanical thrombectomy of BA occlusion followed by proximal VA stenting angioplasty) or antegrade approach (proximal VA stenting angioplasty followed by mechanical thrombectomy of BA occlusion)⁴ was used to perform endovascular treatment. Before stent implantation, 300 mg aspirin and 300 mg clopidogrel were given via a nasal



¹Department of Interventional Neurology, Beijing Tiantan Hospital, Capital Medical University, Beijing, China

²Department of Neurology, Beijing Daxing Hospital, Capital Medical University, Beijing, China

³Department of Interventional Neurology, Beijing Fengtai Youanmen Hospital, Beijing, China

Correspondence to

Dr Zhongrong Miao;
zhongrongm@163.com




Figure 1 Illustration of the reverse (A) and the antegrade (B) technique for the tandem vertebrobasilar artery occlusion through the occlusive or severe stenotic vertebral artery (dirty-road path). (A) A guiding catheter was advanced to the distal left vertebral artery (LVA) V2 segment through the stenosis segment over the partially reinflated balloon. A stent retriever was deployed at the segment of the basilar artery (BA). After successful recanalisation was achieved, the microguidewire was sent to the V2 segment and then the guiding catheter was gently pulled back to the proximal subclavian artery. A balloon-expandable stent was then implanted at the ostial vertebral artery. (B) A balloon-expandable stent was sent to the ostial vertebral artery over the microguidewire and was implanted at the lesion exactly. The guiding catheter was then navigated to the distal V2 segment through the partially reinflated balloon across the implanted stent gently. Subsequently, the stent-assisted thrombectomy was performed as described above.

gastric tube immediately. Then tirofiban (glycoprotein IIb/IIIa inhibitor) was given according to the operator's discretion. Final angiography was performed to exclude reocclusion or thrombosis in the stent after the stent was deployed for 10–20 min. Patients received double antiplatelet agents for 3 months, and then aspirin or clopidogrel was continued alone.

Reverse technique

The guiding catheter was then sent to distal of V2 segment through the stenosis segment over the partially reinflated balloon (similar to the triaxial system) under the road map. The microcatheter was then advanced to the distal BA or the largest posterior cerebral artery across the BA occlusion over a microguidewire. Then angiography was performed through the microcatheter to identify the distal thrombus. A stent retriever was deployed at the segment of the BA occlusion through the microcatheter. The stent retriever was unsheathed to fully expand through the clot for 3–7 min. Subsequently, the stent retriever was gently pulled back with aspiration through the guiding catheter. If recanalisation was insufficient, further procedure would be attempted to achieve successful reperfusion of the BA. Thrombolysis in Cerebral Infarction (TICI) 2b-3 was defined as successful recanalisation after mechanical thrombectomy. After successful recanalisation was achieved, the microguidewire was sent to the V2 segment and then the guiding catheter was gently pulled back to the proximal subclavian artery. A balloon-expandable stent was then implanted at the ostial vertebral artery over the microguidewire (figure 1A).

Antegrade technique

A balloon-expandable stent was sent to the ostial vertebral artery over the microguidewire and was implanted at the lesion exactly. The guiding catheter was then navigated to the distal V2 segment through the partially reinflated balloon across the implanted stent gently. Subsequently, the stent-assisted thrombectomy was performed as described above (figure 1B).

RESULTS

Seven patients were identified to have acute tandem vertebrobasilar artery occlusions (seven men, mean age 57 years, range 48–66). Mean National Institute of Health Stroke Scale (NIHSS) score at admission was 22 (range 12–28). One patient received tissue plasminogen activator prior to endovascular treatment.

The reverse approach was used in five patients, while antegrade approach was used in two patients. TICI score 2b-3 was acquired in all patients. There were no technical complications. The modified Rankin Scale (mRS) score was 1–2 for three patients and 4 for one patient at 3 months. Three patients died at 3 months of follow-up. The clinical data, treatment strategy and patient clinical outcomes are presented in table 1.

Case example 1

A 57-year-old patient with a history of coronary heart disease and diabetes mellitus was transferred to our emergency room due to quadriplegia. The NIHSS score was 12 and posterior circulation Acute Stroke Prognosis Early CT Score (ASPECTS) on diffusion-weighted imaging (DWI) was 8. DWI showed pontine infarction, and MRA revealed BA occlusion. The time from the onset to femoral artery puncture was 402 min.

Cerebral DSA showed the left dominant VA and BA tandem occlusion (figure 2A, D) and the right contralateral non-dominant VA ending in the posterior inferior cerebellar artery (figure 2C). So the left occlusive VA (dirty-road path) was the only access to perform mechanical thrombectomy for BA occlusion.

A 6 Fr Envoy guiding catheter (Cordis, Miami Lakes, Florida, USA) was placed at the proximal subclavian artery, and the Transcend microguidewire (Boston Scientific, Fremont, California, USA) crossed the lesion of the VA occlusion. Predilation of the VA ostial occlusion was performed using a 2 mm x 15 mm balloon (Boston Scientific), and then the balloon was exchanged with a 4 mm x 18 mm balloon-expandable Apollo stent (MicroPort, Shanghai, China). After the stent was placed, the balloon of the stent was partially reinflated and the 6 Fr guiding catheter was advanced into V2 segment through the stent over the partially reinflated balloon (figure 2B). A microcatheter (Stryker Neurovascular, Fremont, California, USA) was placed at the right posterior cerebral artery over the microguidewire, and then a Trevo stent retriever (Stryker Neurovascular) was used to perform mechanical thrombectomy. Follow-up angiography

Table 1 Summary of patient clinical data

Case no	Sex/age (years)	NIHSS at admission	Time to puncture (min)	Time to recanalisation (min)	Treatment strategy	Occlusion	Contralateral vertebral artery	TICI score	Haemorrhage	mRS at 3 months
1	M/57	12	207	90	AT	LV1+BA	Non-dominant ending in PICA	3	N	0
2	M/57	28	125	105	RT	LV1+BA	Non-dominant ending in PICA	3	N	2
3	M/57	28	600	110	AT	LV1+BA	RV1 95% stenosis	3	N	6
4	M/58	16	855	77	RT	LV1+BA	RV1 95% stenosis	3	N	2
5	M/59	20	695	118	RT	LV1+BA	Non-dominant ending in PICA	2b	N	4
6	M/48	25	960	175	RT	RV1+BA	Non-dominant (<2mm)	3	N	6
7	M/66	25	390	194	RT	LV1+BA	Non-dominant ending in PICA	2b	N	6

*AT/RT, antegrade technique/reverse technique; BA, basilar artery; L, left; M, male; mRS, modified Rankin Scale; N, no; NIHSS, National Institute of Health Stroke Scale; N, no; NIHSS, National Institute of Health Stroke Scale; PICA, posterior inferior cerebellar artery; R, right; TICI, Thrombolysis in Cerebral Infarction; V1, first segment of the vertebral artery.




Figure 2 Case example 1. (A) Left vertebral artery occlusion. (B) A 6 Fr catheter crossing the left vertebral artery through the stent over the partially reinflated balloon. (C) Right non-dominant ending in posterior inferior cerebellar artery. (D) Basilar artery occlusion. (E) Recanalisation of the basilar artery. (F) Final angiography of the left vertebral artery.

showed complete BA reperfusion (figure 2E) and the final status of the left ostium VA stent (figure 2F). The time from femoral puncture to reperfusion was 90 min. Follow-up CT scan revealed no cerebral haemorrhagic transformation. The mRS was 1 at 3 months.

Case example 2

A 58-year-old patient with a history of diabetes mellitus and hypertension was transferred to our hospital due to vertigo and consciousness disturbance. The time from onset to femoral artery puncture was 855 min, and the NIHSS score was 16. The posterior circulation ASPECTS on DWI was 7 and MRA showed distal BA occlusion.




Figure 3 Case example 2. (A) Left vertebral artery occlusion. (B) Right vertebral artery ostial severe stenosis. (C) Predilation of the right vertebral artery stenosis. (D) Distal basilar artery clot. (E) Recanalisation of the basilar artery. (F) A stent-assisted angioplasty at the right vertebral artery.

Cerebral DSA showed left VA ostium occlusion (figure 3A), right vertebral artery ostium severe stenosis (figure 3B) and distal BA clot. A 6 Fr guiding catheter (Cordis) was placed at the right subclavian artery and a microguide-wire (Abbott, USA) crossed the stenosis segment to the V2 segment. Predilation was performed using a 3.0×20 mm balloon (Boston Scientific) (figure 3C). The guiding catheter was sent to V2 segment through the stenosis segment over the partially reinflated balloon. Angiography showed that the distal BA occluded (figure 3D). A 4 mm × 20 mm Solitaire stentriever (Irvine, California, USA) was used to perform mechanical thrombectomy. Follow-up angiography showed successful recanalisation of the distal BA occlusion (figure 3E). A 4 mm × 15 mm balloon-expandable stent (Boston Scientific) was implanted at the right vertebral artery stenosis (figure 3F). The time from puncture to recanalisation was 77 min. Follow-up CT scan showed no cerebral haemorrhage. The mRS was 2 at 3 months.

DISCUSSION

In this case series, we reported the endovascular revascularisation for the acute tandem occlusion of vertebrobasilar artery through the occlusive or stenotic road path (dirty-road path).

Ecker *et al*⁶ reported six patients with acute vertebral occlusion below a basilar thrombus were treated with endovascular reconstruction of VA occlusion prior to basilar thrombectomy, and one-third of the patients died due to progressive infarction.

Cohen *et al*⁴ detailed seven patients who suffered from acute tandem vertebrobasilar occlusion and underwent emergent endovascular revascularisation, and two patients died. Though the device and approach for endovascular revascularisation were different, the clinical outcome was similar.

It was more complicated and difficult to perform endovascular revascularisation in the patients with acute tandem vertebrobasilar occlusion than that in the patients with an isolated acute BA occlusion. Generally, in tandem vertebrobasilar artery occlusion, the contralateral VA (clean-road path) was the priority access to perform mechanical thrombectomy of the BA occlusion.⁴ Cohen *et al*⁴ reported only two patients who underwent thrombectomy through the clean-road path.

In this case series, the contralateral VA of five patients was non-dominant artery and another two patients' contralateral dominant VA was severe stenosis. So endovascular revascularisation was performed through the occlusive VA in five patients and the contralateral stenotic VA in another two patients.

There was usually no need for the predilation of the stenotic VA prior to implanting the stent for reducing the risk of artery embolism.^{4,9} However, in the case of VA occlusion or extremely severe stenosis, we predilate the lesions with a small balloon, and then angiography was performed to understand the extent and width of the VA before the stent was implanted.

Cohen *et al*⁴ described two techniques (antegrade technique and reverse technique) that were used to perform endovascular revascularisation when the access was dirty-road path. In Cohen *et al*'s report, five patients underwent endovascular recanalisation through occlusive VA; reverse technique was used for four patients and antegrade technique was used for one patient. They thought there were risks of delaying reperfusion, stent slipping towards the subclavian artery and increasing the difficulty to advance the guiding catheter through the stent.⁴

However, for the reverse technique, the guiding catheter could not cross the occlusion of VA with severe residual stenosis after primary VA angioplasty or the blood flow was very slow when the guiding catheter advanced into the V2 segment. Finally, there was the risk of re-embolisation when the stent of the ostial VA was implanted.⁹ In this case series, antegrade technique was used for two patients without stent slipping and deformation; reverse technique was used for five patients without re-embolisation.

For antegrade strategy, how to deliver the guiding catheter to V2 segment through the stent was a technological challenge. A partially reinflated balloon within the stent can help deliver the guiding catheter to the VA.⁹ In this procedure, the guiding catheter passed through the stent over a partially reinflated balloon when we used the antegrade technique. Because each of these methods has its advantages and disadvantages, it needs careful consideration to choose antegrade or reverse revascularisation technique.

The limitations of this study were the small number of patients, retrospective analysis and selection bias. In addition, this study was not a randomised controlled design. The long-term outcomes of these patients were uncertain.

CONCLUSIONS

In this case series, endovascular revascularisation of acute vertebrobasilar artery tandem occlusion through dirty-road path can be performed successfully without technical complications and may be feasible, while more studies are needed.

Contributors All authors provided substantial contributions to the conception and design of the study and drafted the work or revised it and provided the final approval of the manuscript for submission. All authors are agreeable to be accountable for all aspects of the work, including its accuracy or integrity.

Funding This work was supported by the National Natural Science Foundation of China (Contract grant number: 81471390 to N.M., 81371290 to Z.R.M.), Beijing High-level Personnel Funds (Contract grant number: 2013-2-19 to Z.R.M.).

Competing interests None declared.

Patient consent Obtained.

Ethics approval Tiantan Hospital, Capital Medical University.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

1. Mokin M, Sonig A, Sivakanthan S, *et al.* Clinical and procedural predictors of outcomes from the endovascular treatment of posterior circulation strokes. *Stroke* 2016;47:782–788.
2. Yoon W, Kim SK, Heo TW, *et al.* Predictors of good outcome after stent-retriever thrombectomy in acute basilar artery occlusion. *Stroke* 2015;46:2972–5.
3. Singer OC, Berkefeld J, Nolte CH, *et al.* Mechanical recanalization in basilar artery occlusion: the ENDOSTROKE study. *Ann Neurol* 2015;77:415–24.
4. Cohen JE, Leker RR, Gomori JM, *et al.* Emergent revascularization of acute tandem vertebrobasilar occlusions: endovascular approaches and technical considerations—confirming the role of vertebral artery ostium stenosis as a cause of vertebrobasilar stroke. *J Clin Neurosci* 2016;34:70–6.
5. Savitz SI, Caplan LR. Vertebrobasilar disease. *N Engl J Med* 2005;352:2618–26.
6. Ecker RD, Tsujiura CA, Baker CB, *et al.* Endovascular reconstruction of vertebral artery occlusion prior to basilar thrombectomy in a series of six patients presenting with acute symptomatic basilar thrombosis. *J Neurointerv Surg* 2014;6:379–83.
7. Goyal M, Demchuk AM, Menon BK, *et al.* Randomized assessment of rapid endovascular treatment of ischemic stroke. *N Engl J Med* 2015;372:1019–30.
8. Jovin TG, Chamorro A, Cobo E, *et al.* Thrombectomy within 8 hours after symptom onset in ischemic stroke. *N Engl J Med* 2015;372:2296–306.
9. Du B, Wong EH, Jiang WJ. Long-term outcome of tandem stenting for stenoses of the intracranial vertebrobasilar artery and vertebral ostium. *AJNR Am J Neuroradiol* 2009;30:840–4.