

Phthalocyanines | Hot Paper |

Iron-Catalyzed Oxidative C–C Cross-Coupling Reaction of Tertiary Anilines with Hydroxyarenes by Using Air as Sole Oxidant**

Alexander Purtsas,^[a] Olga Kataeva,^[b] and Hans-Joachim Knölker^{*,[a]}

Abstract: A mild procedure for the oxidative C–C cross-coupling of tertiary anilines with phenols is described which provides the products generally in high yields and with excellent selectivity. The reaction is catalyzed by the hexadeca-

fluorinated iron–phthalocyanine complex FePcF₁₆ in the presence of substoichiometric amounts of methanesulfonic acid and ambient air as sole oxidant.


Introduction

The development of transition metal-catalyzed cross-coupling reactions represents a useful and powerful tool for C–C bond formation. While noble metals have been used as catalysts for classical cross-coupling reactions (e.g., Heck, Negishi, Suzuki–Miyaura),^[1] first-row transition metals have been extensively studied over the past two decades due to their low cost, large abundance in the earth’s crust and lower toxicity.^[2] Thus for economic and ecological reasons, iron-catalyzed organic transformations have received most attention in recent years.^[3] Moreover, the pre-functionalized substrates required for classical cross-coupling reactions lead to a low atom economy and the formation of by-products as unavoidable drawbacks. Oxidative (or dehydrogenative) cross-coupling reactions for C–C bond formation by activation of two C–H bonds are a highly useful synthetic alternative and represent a perfect example of a green and economical chemical process.^[4,5] However, most of the iron-catalyzed oxidative coupling reactions require peroxides,^[6–8] peracids,^[9] or quinones^[10] as stoichiometric oxidants to re-oxidize iron(II) to iron(III). Reactions using molecular oxygen or air as sole oxidant have rarely been explored.^[11,12]


Recently, Shindo and co-workers described a heterogeneous Rh-catalyzed aerobic oxidative cross-coupling for the synthesis of unsymmetrical biaryl compounds (Scheme 1).^[13]

Iron as first-row transition metal is environmentally safe and thus has become a highly promising catalyst.^[3] The groups of Li,^[6] Chandrasekharam,^[7] and Pappo^[8] reported iron-catalyzed oxidative cross-coupling reactions which were achieved by re-oxidation of iron with different peroxides. Despite the recent advances, this process still remains a challenge because of the formation of homocoupling products and oxidized by-products

a) Heterogeneous rhodium-catalyzed oxidative cross-coupling by Shindo




b) Iron-catalyzed oxidative cross-coupling by Chandrasekharam



c) Our previous work: iron-catalyzed oxidative homocoupling



d) This work: iron-catalyzed oxidative cross-coupling with air as sole oxidant



Scheme 1. Biaryl bond formation by transition metal-catalyzed oxidative cross-coupling reactions. TFA = trifluoroacetic acid, TBHP = *tert*-butyl hydroperoxide, TIPS = Triisopropylsilyl, MsOH = methanesulfonic acid.

[a] A. Purtsas, Prof. Dr. H.-J. Knölker
Fakultät Chemie, Technische Universität Dresden
Bergstraße 66, 01069 Dresden (Germany)
E-mail: hans-joachim.knoelker@tu-dresden.de

[b] Dr. O. Kataeva
A. E. Arbusov Institute of Organic and Physical Chemistry
Russian Academy of Sciences
Arbusov Str. 8, Kazan 420088 (Russia)

[**] *Transition Metals in Organic Synthesis, Part 143. For Part 142, see: J. R. Suresh, G. Whitener, D. J. Bröcher, G. Theumer, I. Bauer, W. Massa, H.-J. Knölker, Chem. Eur. J.* **2019**, *25*, 13759–13765.

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/chem.201905595>.

© 2019 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of Creative Commons Attribution NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.




Figure 1. Structure of iron(II)-hexadecafluorophthalocyanine (FePcF₁₆).

resulting from stoichiometric amounts of a strong oxidant (e.g., peroxide).^[6–8] Recently, we have developed an iron-catalyzed oxidative homocoupling by using the hexadecafluorinated iron–phthalocyanine complex FePcF₁₆ as catalyst (Figure 1) and air as re-oxidant.^[14]


This reaction has been applied to the homocoupling of diarylamines and 1- and 2-hydroxycarbazoles.^[14,15] In contrast to most of the alternative iron-catalyzed oxidative cross-coupling reactions,^[6–10] our procedure is using ambient air as sole oxidant.

Biaryl compounds represent key structural motifs in many biologically active natural products and pharmaceutical compounds and therefore, diverse methods for their synthesis have been established.^[116] Herein, we report a mild and efficient synthesis of unsymmetrical biaryl compounds by an iron-catalyzed oxidative cross-coupling of tertiary anilines and hydroxyarenes in the presence of air.

Results and Discussion

The oxidative homocoupling of diarylamines and hydroxycarbazoles has been achieved under mild reaction conditions and in high yields with catalytic amounts of iron(II)–hexadecafluorophthalocyanine (FePcF₁₆)^[14,17] in the presence of methanesulfonic acid or acetic acid and ambient air as final oxidant.^[14,15] Oxidative cross-coupling reactions with iron–phthalocyanine (FePc) or iron–porphyrin complexes as catalysts have been reported previously.^[18,19] However, in order to regenerate the catalytically active iron(III) complex, an oxidant (e.g., peroxide) had to be added in stoichiometric amounts.^[8c,d] Based on our previous studies,^[14,15,20] we envisaged to develop an iron-catalyzed cross-coupling of anilines with phenols by using air as sole oxidant. Moreover, we wanted to avoid the need for a transient connection between the two reactants for cross-coupling.^[21] According to previous reports,^[7,13,14,22] we hypothesized that the homocoupling of anilines could be suppressed by substituents causing a steric hindrance at the amino group. Reaction of a radical cation generated by SET oxidation of a tertiary aniline with a sterically less hindered phenol should preferentially form a C–C bond either via radical or electrophilic aromatic substitution (Scheme 2). A second SET and final proton loss with concomitant re-aromatization would lead to the cross-coupling product.

Thus, we selected *N,N*-dimethyl-*p*-toluidine (**1a**) and 2-naphthol (**2a**) as model system for optimizing the reaction conditions of the iron-catalyzed oxidative cross-coupling. Various temperatures and iron catalysts have been tested for the reaction of **1a** with two equivalents of **2a** in dichloromethane under air (Table 1). Using catalytic amounts (1 mol%) of FePcF₁₆ and substoichiometric amounts (40 mol%) of methanesulfonic acid as additive at room temperature afforded the cross-coupling product **3a** in 64% yield along with the homo-



Scheme 2. Proposed mechanism for the iron-catalyzed oxidative cross-coupling of tertiary anilines and phenols.

Table 1. Optimization of the iron-catalyzed oxidative cross-coupling reaction.^[a]

Entry	T [°C]	[Fe]	Yield 3a [%]	Yield 4a [%]
1	20	FePcF ₁₆	64	9
2 ^[b]	70	FePcF ₁₆	trace	–
3	0	FePcF ₁₆	84	18
4	0	–	–	–
5 ^[c]	0	FePcF ₁₆	–	–
6 ^[d]	0	FePcF ₁₆	17	27
7 ^[e]	0	FePcF ₁₆	37	26
8	0	FePc	66	20
9	0	FePc(SO ₃) ₄ H ₃ Na	14	trace
10 ^[f]	0	FePc(SO ₃) ₄ H ₃ Na	20	–
11	0	FePc(CO ₂ C ₅ H ₁₁) ₈	33	trace
12	0	FePcCl ₁₆	–	–
13	0	FePpz	–	–
14	0	FePpzCl ₈	–	–
15	0	FePpz(Ph) ₈	–	–
16	0	ClFeTPPF ₂₀	–	–
17	0	FeCl ₂ (99.99%)	–	–

[a] **1a** (1.0 equiv), **2a** (2.0 equiv). [b] Solvent: (CH₂Cl)₂, 4 h. [c] Reaction under argon. [d] Reaction without MsOH. [e] Reaction with AcOH (3.0 equiv) instead of MsOH. [f] Solvent: H₂O. FePcF₁₆ = iron(II)-hexadecafluorophthalocyanine, FePc = iron(II)-phthalocyanine, FePc(SO₃)₄H₃Na = iron(II)-phthalocyaninetetrasulfonic acid monosodium salt · x H₂O, FePpz = iron(II)-pyrazinoporphyrazine, ClFeTTPP = tetraphenylporphyrin-iron(III) chloride (structures shown in the Supporting Information).

coupling product 1,1'-binaphthyl-2,2'-diol (BINOL) (**4a**)^[23] in 9% yield (entry 1). The structure of **3a** was confirmed by an X-ray analysis (Figure 2). Variation of the reaction temperature (entries 2 and 3) showed that the cross-coupling reaction proceeds even better at 0 °C and provides the biaryl compound **3a** in 84% yield.^[7a] In the following, we have performed several blank experiments in order to support our hypothesis of an iron(III)-catalyzed oxidative coupling which is promoted by the Brønsted acid additive. Reaction in the absence of iron catalyst gave no product at all (entry 4). The same result was obtained when the reaction was performed in the presence of 1 mol% of FePcF₁₆ but under argon instead of air (entry 5). These experiments confirm that FePcF₁₆ as well as its oxidation by air to




Figure 2. Molecular structure of the biaryl compound **3a** in the crystal (thermal ellipsoids are shown at the 50% probability level).

the μ -oxo-diiron(III) complex are essential for the catalytic $[\text{Fe}^{\text{III}}]/[\text{Fe}^{\text{I}}]$ cycle leading to the oxidative biaryl coupling.^[14, 18–20] The iron-catalyzed oxidative cross-coupling in the absence of methanesulfonic acid resulted in a significantly lower yield (17%) of the biaryl compound **3a** (entry 6) and an excess of a weaker acid (acetic acid) as additive led to **3a** in only 37% yield (entry 7).

The unsubstituted iron(II)–phthalocyanine complex (FePc) afforded the cross-coupling product **3a** in 66% yield (entry 8), whereas other iron complexes led to either low yields of **3a** (entries 9–11) or no product formation at all (entries 12–17). The present results confirm previous findings that FePcF₁₆ has a higher activity compared to the parent complex FePc which is ascribed to the electronic effect of the fluorine atoms.^[14, 20, 24] It is noteworthy that by using the tetrasulfonato-substituted iron-phthalocyanine complex as catalyst, the oxidative cross-coupling to **3a** could successfully be achieved in water as solvent (entry 10), albeit yet in low yield (20%).

Using the optimized reaction conditions, we have studied the synthetic scope of this cross-coupling reaction between various tertiary arylamines **1** and hydroxyarenes **2** (Table 2). As the simple monocyclic phenols we have tested gave no cross-coupling products, we focused on polycyclic hydroxyarenes as substrates. The iron-catalyzed oxidative coupling of *N,N*-dimethyl-*p*-toluidine (**1a**) with the substituted 2-naphthols **2a–d** provided selectively the corresponding cross-coupling products **3a–d** in 64–87% yield. Interestingly, homocoupling of the 2-naphthols was observed only for the parent compound **2a**. The results emphasize the functional group tolerance of this reaction. The cross-coupling of 2-hydroxydibenzofuran (**2e**) with **1a** appeared to be more difficult and afforded compound **3e** only in 42% yield. The structure of the 1-aryl-coupled dibenzofuran-2-ol **3e**, which was reported to show anti-TB activity and previously obtained as a red liquid,^[7b] was confirmed by an X-ray analysis (Figure 3). The iron-catalyzed cross-coupling of a range of tertiary anilines **1** with 2-naphthols **2** proceeded well and led to the corresponding products **3f–j** in high yields (65–88%). Whereas the reaction between *p*-chloro-*N,N*-dimethylaniline and 2-naphthol (**2a**) afforded the coupling product **3k** only in moderate yield.

Table 2. Synthesis of unsymmetrical biaryls **3** by iron-catalyzed oxidative cross-coupling of *N,N*-dialkylanilines **1** with various hydroxyarenes **2**.

<p>3a, 84%</p>	<p>3b, 87%</p>	<p>3c, 76%</p>
<p>3d, 64% [a]</p>	<p>3e, 42% [b]</p>	
<p>3f, 65% [c]</p>	<p>3g, 69% [d]</p>	<p>3h, 71% [e]</p>
<p>3i, 82%</p>	<p>3j, 88%</p>	<p>3k, 36% [f]</p>
<p>[a] 0 °C to RT. [b] 2 mol% of FePcF₁₆. [c] Reaction time: 4 h. [d] 6.0 equiv of 2a, 48 h. [e] 6.0 equiv of 2a. [f] 4.0 equiv of 2a.</p>		






Figure 3. Molecular structure of the biaryl compound **3e** in the crystal (thermal ellipsoids are shown at the 50% probability level).

The *meta*-silyloxy-substituted *N,N*-dimethylaniline (**5**) is very electron-rich and highly reactive at the positions *ortho/para* to the dimethylamino group (Scheme 3). Thus, the iron-catalyzed reaction led to an oxidative coupling of two molecules of **5** at their sterically least hindered reactive position (C-6) with C-1 of 2-naphthol (**2a**) and provided compound **6** by dearomatization of one benzo ring.



Scheme 3. Iron-catalyzed oxidative cross-coupling of 2-naphthol (**2a**) with *N,N*-dimethyl-3-(triisopropylsilyloxy)aniline (**5**). Reaction conditions: CH_2Cl_2 , air, 0°C , 24 h; **6**: 37%, **4a**: 19% yield.

The application of our iron-catalyzed C–C cross-coupling to the reaction of 2-(dimethylamino)naphthalene (**7**) with **2a** provided *N,N*-dimethyl-NOBIN (2-(dimethylamino)-2'-hydroxy-1,1'-binaphthyl) (**8**) (Scheme 4).^[25] Compound **8** was obtained in 62%



Scheme 4. Iron-catalyzed oxidative cross-coupling of 2-naphthol (**2a**) with 2-(dimethylamino)naphthalene (**7**). Reaction conditions: CH_2Cl_2 , air, 0°C , 24 h; **8**: 62%, **9**: 11%, **4a**: 20% yield.

yield and confirmed by an X-ray crystal structure determination (Figure 4). NOBIN derivatives were shown to be highly useful as axially chiral biaryl ligands in asymmetric catalysis and effi-




Figure 4. Molecular structure of *N,N*-dimethyl-NOBIN (**8**) in the crystal (thermal ellipsoids are shown at the 50% probability level).

cient routes for their synthesis are still being searched.^[25,26] Remarkably, this reaction afforded the C–O cross-coupled compound **9** as a by-product, thus indicating that ethers may be accessible by iron-catalyzed C–O cross-coupling reactions using air as oxidant.

Conclusions

We have developed an efficient and environmentally friendly method for the synthesis of unsymmetrical biaryl compounds by iron-catalyzed oxidative cross-coupling of tertiary anilines and hydroxyarenes. Our procedure requires 1–2 mol% of the hexadecafluorinated iron-phthalocyanine complex FePcF_{16} as catalyst, substoichiometric amounts of methanesulfonic acid as additive, and air as the sole oxidant. The unsymmetrical biaryls are obtained with excellent selectivity. Further studies of this process and applications are in progress.

Experimental Section

General methods: All iron-catalyzed reactions were performed with non-dried solvents in an atmosphere of air unless stated otherwise. All other reactions were performed in oven-dried glassware with dry solvents. Dichloromethane and THF were dried by using a solvent-purification system (MBraun-SPS). Chemicals were used as received from commercial sources. Reaction mixtures were cooled by using a Lauda Compact PI 1 cryostat. Automated flash column chromatography was performed with silica gel (Acros Organics; 0.035–0.070 mm) on a Büchi Sepacore system equipped with an UV monitor. TLC analysis was performed on TLC plates from Merck (60 F254) with UV light for visualization. Melting points were measured on a Gallenkamp MPD 350 melting-point apparatus. UV spectra were recorded on a PerkinElmer 25 UV/Vis spectrometer. Fluorescence spectra were measured on a Varian Cary Eclipse spectrophotometer. Wavelengths are reported in nm. IR spectra were recorded on a Thermo Nicolet Avatar 360 FTIR spectrometer using the attenuated total reflectance (ATR) method. Wavenumbers are reported in cm^{-1} . NMR spectra were recorded on Bruker DRX 500 and Avance III 600 spectrometers. Chemical shifts δ are reported in ppm with the solvent signal as an internal standard. The following abbreviations have been used: s=singlet, d=doublet, t=triplet, q=quartet, quin=quintet, spt=septet, m=multiplet, and br=broad. EI mass spectra were recorded by GC-MS coupling on an Agilent Technologies 6890 N GC system equipped with a 5973 mass-selective detector (70 eV). ESI mass spectra were recorded on a Bruker Esquire LC mass spectrometer with an ion-trap detector. Positive and negative ions were detected. ESI-HRMS were recorded on a Waters Xevo G2-XS QTOF mass spectrometer. Elemental analyses were measured on a EuroVector EuroEA3000 elemental analyzer. Weight portions are given in percent. X-ray crystal structure analyses were performed with a Bruker Kappa Apex II CCD X-ray diffractometer equipped with a 700 series Cryostream low temperature device from Oxford Cryosystems. SHELXS-97,^[27] SADABS version 2.10,^[28] SHELXL-97,^[29] POV-Ray for Windows version 3.7.0.msvc10.win64, and ORTEP-3 for Windows^[30] were used as software.

2-Acetyldibenzofuran:^[7b] A solution of dibenzofuran (3.11 g, 18.5 mmol) in chloroform (30 mL) was added over a period of 10 min via a syringe pump to a mixture of aluminum trichloride (2.99 g, 22.4 mmol) and freshly distilled acetyl chloride (1.6 mL, 1.8 g, 23 mmol) in chloroform (30 mL) at room temperature under

argon. After 45 min of stirring, the reaction mixture was poured into a mixture of ice water (100 mL) and 1 N HCl (50 mL). The aqueous layer was extracted three times with dichloromethane. The combined organic layers were dried over MgSO₄ and the solvent was removed in vacuo. Purification of the crude product by column chromatography on silica gel with isohexane/ethyl acetate (10:1) afforded 2-acetyldibenzofuran (3.47 g, 16.5 mmol, 89%) as a colorless solid. M.p. 66.5 °C; UV (MeOH): $\lambda = 216, 238, 253, 287$ nm; fluorescence (MeOH): $\lambda_{\text{ex}} = 253$ nm, $\lambda_{\text{em}} = 415$ nm; IR (ATR): $\tilde{\nu} = 3319, 3061, 3041, 2991, 2962, 2912, 2056, 2030, 2009, 1907, 1844, 1794, 1772, 1735, 1717, 1699, 1667, 1633, 1583, 1553, 1488, 1476, 1453, 1421, 1364, 1345, 1327, 1304, 1278, 1215, 1199, 1185, 1119, 1103, 1064, 1022, 955, 907, 893, 871, 841, 820, 769, 748, 729, 711, 652, 632$ cm⁻¹; ¹H NMR (500 MHz, CDCl₃): $\delta = 2.72$ (s, 3H), 7.40 (td, $J = 7.5, 0.9$ Hz, 1H), 7.51 (m, 1H), 7.60 (m, 2H), 8.01 (m, 1H), 8.11 (dd, $J = 8.5, 1.9$ Hz, 1H), 8.59 (dd, $J = 1.8, 0.5$ Hz, 1H) ppm; ¹³C NMR and DEPT (125 MHz, CDCl₃): $\delta = 26.94$ (CH₃), 111.72 (CH), 112.09 (CH), 121.09 (CH), 121.75 (CH), 123.53 (CH), 123.87 (C), 124.71 (C), 128.11 (CH), 128.13 (CH), 132.65 (C), 157.00 (C), 159.06 (C), 197.46 (C=O) ppm; MS (EI): m/z (%) = 210 (62, [M]⁺), 195 (100), 167 (35), 139 (68), 113 (10); elemental analysis (%) calcd for C₁₄H₁₀O₂: C 79.98, H 4.79; found: C 79.62, H 4.80.

2-Hydroxydibenzofuran (2e):^[7b] Trifluoroacetic acid (1.8 mL, 2.7 g, 23 mmol) and then *m*-chloroperoxybenzoic acid (2.49 g, 14.4 mmol) were slowly added to a solution of 2-acetyldibenzofuran (1.16 g, 5.52 mmol) in dry dichloromethane (20 mL) at 0 °C. The reaction mixture was stirred at room temperature for 3 days and quenched with a sat. solution of aqueous iron(II) sulfate. The aqueous layer was extracted three times with dichloromethane. The combined organic layers were washed with a sat. aqueous solution of Na₂CO₃, then with water and dried over MgSO₄. After evaporation of the solvent, the crude product was suspended in MeOH (25 mL) and sodium methoxide (1.26 g, 23.2 mmol) was added. The mixture was stirred at room temperature for 1 h and quenched with 2 N HCl (15 mL). The aqueous layer was extracted three times with dichloromethane, the combined organic layers were dried over MgSO₄, and the solvent was removed in vacuo. Purification of the crude product by column chromatography on silica gel with isohexane/ethyl acetate (10:1) provided 2-hydroxydibenzofuran (2e) (779 mg, 4.23 mmol, 77% over two steps) as a colorless solid. M.p. 129 °C; UV (MeOH): $\lambda = 236, 242, 252, 290, 308$ nm; fluorescence (MeOH): $\lambda_{\text{ex}} = 252$ nm, $\lambda_{\text{em}} = 354$ nm; IR (ATR): $\tilde{\nu} = 3193$ (br), 3061, 3035, 2011, 1974, 1934, 1894, 1845, 1740, 1635, 1598, 1574, 1475, 1440, 1420, 1397, 1361, 1332, 1306, 1279, 1210, 1188, 1165, 1148, 1117, 1100, 1015, 933, 898, 867, 840, 798, 740, 717, 621 cm⁻¹; ¹H NMR (500 MHz, CDCl₃): $\delta = 4.80$ (br s, 1H), 6.96 (dd, $J = 8.8, 2.5$ Hz, 1H), 7.33 (td, $J = 7.5, 1.1$ Hz, 1H), 7.38 (d, $J = 2.5$ Hz, 1H), 7.43 (d, $J = 8.8$ Hz, 1H), 7.45 (td, $J = 7.8, 1.4$ Hz, 1H), 7.54 (d, $J = 8.2$ Hz, 1H), 7.89 (d, $J = 7.9$ Hz, 1H) ppm; ¹³C NMR and DEPT (125 MHz, CDCl₃): $\delta = 106.40$ (CH), 111.89 (CH), 112.25 (CH), 115.39 (CH), 120.84 (CH), 122.59 (CH), 124.35 (C), 125.19 (C), 127.44 (CH), 151.13 (C), 151.55 (C), 157.14 (C) ppm; MS (EI): m/z (%) = 184 (100, [M]⁺), 155 (10), 128 (16); elemental analysis (%) calcd for C₁₂H₈O₂: C 78.25, H 4.38; found: C 78.59, H 4.21.

1-(4-Tolyl)pyrrolidine: 1,4-Dibromobutane (475 mg, 2.20 mmol) was added to a mixture of K₂CO₃ (692 mg, 5.01 mmol) and *para*-toluidine (214 mg, 2.00 mmol) in anhydrous DMF (10 mL) and the reaction mixture was heated to 80 °C for 24 h under argon. Subsequently, the reaction mixture was cooled to room temperature and diluted with ethyl acetate and water. The layers were separated and the organic layer was extracted three times with 1 N HCl. The aqueous layers were combined, adjusted to pH 8 with 1 N NaOH, and then extracted three times with ethyl acetate. The combined

organic layers were washed with brine and dried over MgSO₄. Evaporation of the solvent in vacuo and purification of the crude product by column chromatography on silica gel with isohexane/ethyl acetate (20:1) afforded 1-(4-tolyl)pyrrolidine (251 mg, 1.56 mmol, 78%) as a pale yellow solid. ¹H NMR (500 MHz, CDCl₃): $\delta = 2.00$ (m, 4H), 2.27 (s, 3H), 3.27 (m, 4H), 6.53 (d, $J = 7.2$ Hz, 2H), 7.06 (d, $J = 8.3$ Hz, 2H) ppm; ¹³C NMR and DEPT (125 MHz, CDCl₃): $\delta = 20.41$ (CH₃), 25.53 (2 CH₂), 47.98 (2 CH₂), 111.92 (2 CH), 124.52 (C), 129.76 (2 CH), 146.23 (C) ppm; MS (EI): m/z (%) = 161 (72, [M]⁺), 160 (100), 118 (20), 105 (40); elemental analysis (%) calcd for C₁₁H₁₅N: C 81.94, H 9.38, N 8.69; found: C 82.24, H 9.39, N 8.66. For further analytical data, see ref. [31].

1-(4-Tolyl)piperidine and 4-(4-tolyl)morpholine (by adaptation of a literature procedure^[32]): A mixture of 4-bromotoluene (180 mg, 1.05 mmol), piperidine (86.1 mg, 1.01 mmol) or morpholine (87.2 mg, 1.00 mmol), Pd(OAc)₂ (2.7 mg, 12 μ mol), RuPhos (9.5 mg, 20 μ mol), and powdered NaOtBu (124 mg, 1.29 mmol) was stirred at 110 °C overnight. After cooling to room temperature, the mixture was dissolved in CH₂Cl₂/H₂O (1:1), and the aqueous layer was extracted twice with CH₂Cl₂ (5 mL). The combined organic layers were dried over MgSO₄, the solvent was evaporated in vacuo, and the crude product was purified by column chromatography on silica gel with isohexane/ethyl acetate (20:1).

1-(4-Tolyl)piperidine (118 mg, 0.672 mmol, 67%) was obtained as a colorless oil. ¹H NMR (500 MHz, CDCl₃): $\delta = 1.56$ (m, 2H), 1.73 (m, 4H), 2.27 (s, 3H), 3.10 (t, $J = 5.4$ Hz, 4H), 6.88 (d, $J = 8.2$ Hz, 2H), 7.07 (d, $J = 8.2$ Hz, 2H) ppm; ¹³C NMR and DEPT (125 MHz, CDCl₃): $\delta = 20.58$ (CH₃), 24.39 (CH₂), 26.04 (2 CH₂), 51.55 (2 CH₂), 117.17 (2 CH), 129.05 (C), 129.69 (2 CH), 150.36 (C) ppm; MS (EI): m/z (%) = 175 (66, [M]⁺), 174 (100), 146 (10), 119 (43), 91 (48), 65 (14); elemental analysis (%) calcd for C₁₂H₁₇N: C 82.23, H 9.78, N 7.99; found: C 82.22, H 9.50, N 8.14. For further analytical data, see ref. [31].

4-(4-Tolyl)morpholine (163 mg, 0.921 mmol, 92%) was obtained as a colorless solid. ¹H NMR (500 MHz, CDCl₃): $\delta = 2.29$ (s, 3H), 3.11 (m, 4H), 3.87 (m, 4H), 6.85 (d, $J = 8.5$ Hz, 2H), 7.10 (d, $J = 8.2$ Hz, 2H) ppm; ¹³C NMR and DEPT (125 MHz, CDCl₃): $\delta = 20.55$ (CH₃), 50.10 (2 CH₂), 67.08 (2 CH₂), 116.21 (2 CH), 129.72 (C), 129.84 (2 CH), 149.24 (C) ppm; MS (EI): m/z (%) = 177 (45, [M]⁺), 119 (100), 91 (40), 65 (13); HRMS (ESI) calcd for C₁₁H₁₆NO⁺ ([M+H]⁺): 178.1232; found: 178.1230. For further analytical data, see ref. [31].

4-Methoxy-*N,N*-dimethylaniline, 4-chloro-*N,N*-dimethylaniline, and 2-*N,N*-dimethylaminonaphthalene (7): Aqueous formaldehyde (37%, 12.2 mL, 4.92 g, 164 mmol) was slowly added to a solution of *p*-anisidine (1.28 g, 10.4 mmol) or *p*-chloroaniline (1.32 g, 10.4 mmol) or naphthalen-2-amine (1.49 g, 10.4 mmol) in dry THF (110 mL) and the solution was stirred at room temperature. After 15 min, sodium cyanoborohydride (4.63 g, 73.7 mmol) was added and the mixture was stirred at room temperature for 15 min. Subsequently, acetic acid (1.7 mL) was added dropwise while keeping the reaction at room temperature. The reaction mixture was stirred at room temperature for 2–4 h until all starting material was consumed as indicated by TLC and quenched with 2 N NaOH (pH > 7). The mixture was extracted three times with ethyl acetate and the combined organic layers were dried over MgSO₄. After evaporation of the solvent in vacuo, the crude product was purified by column chromatography on silica gel with isohexane/ethyl acetate (gradient elution from 15:1 to 10:1).

4-Methoxy-*N,N*-dimethylaniline (1.21 g, 8.01 mmol, 77%) was obtained as a grey solid. M.p. 35 °C; UV (MeOH): $\lambda = 246, 306$ nm; fluorescence (MeOH): $\lambda_{\text{ex}} = 246$ nm, $\lambda_{\text{em}} = 373$ nm; IR (ATR): $\tilde{\nu} = 3071, 3038, 2993, 2949, 2884, 2831, 2798, 1851, 1613, 1574, 1558, 1539, 1513, 1441, 1341, 1300, 1244, 1225, 1180, 1129, 1064, 1034,$

945, 816, 801, 681 cm^{-1} ; $^1\text{H NMR}$ (600 MHz, CDCl_3): δ = 2.88 (s, 6H), 3.77 (s, 3H), 6.78 (d, J = 8.7 Hz, 2H), 6.86 (d, J = 8.7 Hz, 2H) ppm; $^{13}\text{C NMR}$ and DEPT (151 MHz, CDCl_3): δ = 42.03 (2 CH_3), 55.90 (CH_3), 114.80 (2 CH), 115.15 (2 CH), 145.81 (C), 152.26 (C) ppm; MS (EI): m/z (%) = 151 (58, $[\text{M}]^+$), 136 (100), 120 (11), 108 (22), 92 (10), 65 (12); elemental analysis (%) calcd for $\text{C}_9\text{H}_{13}\text{NO}$: C 71.49, H 8.67, N 9.26; found: C 71.45, H 8.64, N 9.34.

4-Chloro-*N,N*-dimethylaniline (1.47 g, 9.45 mmol, 91%) was obtained as a colorless solid. M.p. 43.5 °C; UV (MeOH): λ = 259, 313 nm; fluorescence (MeOH): λ_{ex} = 259 nm, λ_{em} = 364 nm; IR (ATR): $\tilde{\nu}$ = 3091, 2989, 2886, 2853, 2803, 2109, 1956, 1865, 1755, 1613, 1590, 1559, 1490, 1442, 1340, 1222, 1188, 1164, 1124, 1093, 1060, 994, 941, 806, 758, 697 cm^{-1} ; $^1\text{H NMR}$ (500 MHz, CDCl_3): δ = 2.93 (s, 6H), 6.65 (d, J = 8.5 Hz, 2H), 7.18 (d, J = 8.8 Hz, 2H) ppm; $^{13}\text{C NMR}$ and DEPT (125 MHz, CDCl_3): δ = 40.86 (2 CH_3), 113.83 (2 CH), 121.60 (C), 128.95 (2 CH), 149.26 (C) ppm; MS (EI): m/z (%) = 155 (78, $[\text{M}]^+$), 154 (100), 139 (23), 118 (12), 111 (17), 77 (14), 75 (19), 50 (11), 42 (13); elemental analysis (%) calcd for $\text{C}_8\text{H}_{10}\text{ClN}$: C 61.74, H 6.48, N 9.00; found: C 61.82, H 6.47, N 9.06.

2-(Dimethylamino)naphthalene (**7**) (1.65 g, 9.62 mmol, 93%) was obtained as a pale yellow solid. M.p. 43.5 °C; UV (MeOH): λ = 214, 247, 287, 297, 352 nm; fluorescence (MeOH): λ_{ex} = 247 nm, λ_{em} = 409 nm; IR (ATR): $\tilde{\nu}$ = 3048, 3021, 2975, 2889, 2869, 2801, 1736, 1699, 1624, 1595, 1559, 1540, 1507, 1484, 1440, 1365, 1328, 1277, 1241, 1198, 1145, 1062, 1014, 956, 865, 824, 803, 740, 614 cm^{-1} ; $^1\text{H NMR}$ (500 MHz, CDCl_3): δ = 3.06 (s, 6H), 6.97 (br s, 1H), 7.19 (dd, J = 8.8, 2.5 Hz, 1H), 7.23 (m, 1H), 7.38 (td, J = 7.6, 1 Hz, 1H), 7.67 (d, J = 8.2 Hz, 1H), 7.70 (d, J = 8.4 Hz, 1H), 7.72 (d, J = 8.5 Hz, 1H) ppm; $^{13}\text{C NMR}$ and DEPT (125 MHz, CDCl_3): δ = 41.21 (2 CH_3), 107.02 (CH), 116.63 (CH), 122.36 (CH), 126.38 (2 CH), 127.17 (C), 127.58 (CH), 128.87 (CH), 135.03 (C), 148.58 (C) ppm; MS (EI): m/z (%) = 171 (100, $[\text{M}]^+$), 170 (97), 155 (22), 154 (11), 129 (12), 128 (36), 127 (30), 126 (11), 115 (12), 77 (11); elemental analysis (%) calcd for $\text{C}_{12}\text{H}_{13}\text{N}$: C 84.17, H 7.65, N 8.18; found: C 84.46, H 7.73, N 8.24.

N,N-Dimethyl-3-(triisopropylsilyloxy)aniline (**5**): Triisopropylsilyl triflate (2.70 mL, 3.08 g, 10.1 mmol) was added dropwise to a solution of *N,N*-dimethyl-3-aminophenol (1.00 g, 7.29 mmol) and triethylamine (2.70 mL, 1.97 g, 19.5 mmol) in dichloromethane (62 mL) at 0 °C under an argon atmosphere. After stirring of the solution for 22 h at room temperature, water (20 mL) was added and the aqueous layer was extracted three times with dichloromethane. The combined organic layers were dried over MgSO_4 and the solvent was removed. Purification of the crude product by column chromatography on silica gel with isohexane/ethyl acetate (15:1) provided compound **5** (2.08 g, 7.09 mmol, 97%) as a reddish oil. UV (MeOH): λ = 212, 253, 296 nm; fluorescence (MeOH): λ_{ex} = 212 nm, λ_{em} = 348 nm; IR (ATR): $\tilde{\nu}$ = 2943, 2889, 2865, 2800, 1604, 1573, 1499, 1452, 1396, 1353, 1248, 1168, 1151, 1062, 1002, 925, 881, 827, 783, 756, 681 cm^{-1} ; $^1\text{H NMR}$ (500 MHz, CDCl_3): δ = 1.11 (d, J = 7.6 Hz, 18H), 1.26 (m, 3H), 2.91 (s, 6H), 6.29 (m, 2H), 6.35 (d, J = 7.9 Hz, 1H), 7.06 (t, J = 8.0 Hz, 1H) ppm; $^{13}\text{C NMR}$ and DEPT (125 MHz, CDCl_3): δ = 12.88 (6 CH_3), 18.13 (3 CH), 40.91 (2 CH_3), 104.95 (CH), 106.08 (CH), 108.64 (C), 129.64 (2 CH), 157.11 (C) ppm; ESI-MS (+25 V): m/z = 294.3 $[\text{M}+\text{H}]^+$; elemental analysis (%) calcd for $\text{C}_{17}\text{H}_{31}\text{NOSi}$: C 69.56, H 10.65, N 4.77; found: C 69.40, H 10.87, N 5.00.

1,2,3,4,8,9,10,11,15,16,17,18,22,23,24,25-Hexadecafluorophthalocyaninato-iron(II) (for experimental details, see refs. [14,17,20]): FePcF_{16} (70% yield) was obtained as a dark blue to violet powder. IR (ATR): $\tilde{\nu}$ = 3636, 3180, 2645, 2436, 2043, 1772, 1733, 1682, 1618, 1559, 1520, 1484, 1459, 1399, 1318, 1266, 1147, 958, 944, 837, 751, 655 cm^{-1} ; ESI-MS (+100 V): m/z = 857.1 $[\text{M}+\text{H}]^+$; ESI-MS (−50 V):

m/z = 914.7 $[\text{M}+\text{OAc}]^-$, 886.7 $[\text{M}+\text{OMe}]^-$; elemental analysis (%) calcd for $\text{C}_{32}\text{F}_{16}\text{FeN}_8$: C 44.89, N 13.09; found: C 44.33, N 13.18.

General procedure for the iron-catalyzed oxidative cross-coupling: A 0.1 M solution of the tertiary aniline **1** (1.0 equiv) in dichloromethane (2–6 mL) was added to a mixture of the hydroxyarene **2** (2.0–6.0 equiv), methanesulfonic acid (40 mol%), and FePcF_{16} (1.0 mol%). The reaction mixture was vigorously stirred at 0 °C for 4–48 h under air and then quenched with a sat. aqueous solution of Na_2CO_3 (5 mL). The aqueous layer was extracted three times with dichloromethane. The combined organic layers were dried over MgSO_4 and evaporated in vacuo. Purification of the crude product by automated column chromatography on silica gel provided the biaryl compound **3**.

1-(2-(Dimethylamino)-5-methylphenyl)naphthalen-2-ol (3a): *N,N*-Dimethyl-*p*-toluidine (**1a**) (83.1 mg, 615 μmol), dichloromethane (6 mL), 2-naphthol (**2a**) (173 mg, 1.20 mmol), methanesulfonic acid (23.2 mg, 241 μmol), FePcF_{16} (5.1 mg, 6.0 μmol). Reaction time: 24 h, column chromatography (isohexane/ethyl acetate, 1–3%, 1 h, 3–15%, 0.5 h, 15%, 20 min). **3a** (144 mg, 519 μmol , 84%) was obtained as a pale yellow solid. M.p. 166.5 °C; UV (MeOH): λ = 231, 286, 333 nm; fluorescence (MeOH): λ_{ex} = 231 nm, λ_{em} = 422 nm; IR (ATR): $\tilde{\nu}$ = 3028, 2995, 2955, 2916, 2873, 2842, 2795, 1892, 1619, 1590, 1558, 1497, 1458, 1397, 1358, 1334, 1296, 1231, 1177, 1156, 1130, 1093, 1037, 1015, 955, 925, 891, 868, 816, 751, 720, 688, 662, 634 cm^{-1} ; $^1\text{H NMR}$ (600 MHz, CDCl_3): δ = 2.33 (s, 3H), 2.68 (s, 6H), 7.19 (m, 1H), 7.22 (m, 2H), 7.28 (d, J = 8.7 Hz, 1H), 7.34 (m, 1H), 7.38 (m, 1H), 7.78 (d, J = 8.8 Hz, 1H), 7.80 (d, J = 8.5 Hz, 1H), 7.82 (dd, J = 8.4, 1.1 Hz, 1H), 10.15 (br s, 1H) ppm; $^{13}\text{C NMR}$ and DEPT (151 MHz, CDCl_3): δ = 20.80 (CH_3), 43.95 (2 CH_3), 118.23 (CH), 120.82 (CH), 121.07 (C), 123.20 (CH), 125.48 (CH), 126.20 (CH), 128.28 (CH), 129.19 (CH), 129.45 (CH), 130.05 (C), 130.66 (C), 132.82 (C), 133.51 (C), 135.87 (CH), 147.50 (C), 152.27 (C) ppm; ESI-MS (+10 V): m/z = 278.1 $[\text{M}+\text{H}]^+$; elemental analysis (%) calcd for $\text{C}_{19}\text{H}_{19}\text{NO}$: C 82.28, H 6.90, N 5.05; found: C 81.90, H 7.00, N 4.89.

1,1'-Binaphthyl-2,2'-diol (BINOL) (4a): (30.5 mg, 107 μmol , 18%) was obtained as a brown solid. $^1\text{H NMR}$ (600 MHz, CDCl_3): δ = 5.06 (br s, 2H), 7.16 (d, J = 8.3 Hz, 2H), 7.31 (td, J = 7.7, 1.1 Hz, 2H), 7.38 (m, 4H), 7.90 (d, J = 8.3 Hz, 2H), 7.98 (d, J = 9.0 Hz, 2H) ppm; $^{13}\text{C NMR}$ and DEPT (151 MHz, CDCl_3): δ = 110.94 (2 C), 117.90 (2 CH), 124.19 (2 CH), 124.35 (2 CH), 127.64 (2 CH), 128.57 (2 CH), 129.61 (2 C), 131.60 (2 CH), 133.54 (2 C), 152.89 (2 C) ppm; ESI-MS (+25 V): m/z = 287.2 $[\text{M}+\text{H}]^+$. For further analytical data, see ref. [23].

Crystallographic data for compound 3a: $\text{C}_{19}\text{H}_{19}\text{NO}$, M = 277.35 g mol^{-1} , crystal size 0.250 × 0.300 × 0.330 mm^3 , monoclinic, space group $P2_1/c$, a = 9.2180(17), b = 8.8167(17), c = 18.835(4) Å, V = 1491.1(5) Å³, Z = 4, ρ_{calcd} = 1.235 g cm^{-3} , μ = 0.076 mm^{-1} , λ = 0.71073 Å, T = 120(2) K, θ range = 2.22–28.49°, reflections collected 17 145, independent reflections 3766 (R_{int} = 0.0477), 197 parameters. The structure was solved by direct methods and refined by full-matrix least-squares on F^2 ; final R indices [$I > 2\sigma(I)$] $R1$ = 0.0462 and $wR2$ = 0.1105 maximal residual electron density 0.282 e Å^{-3} . CCDC 1963408 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.

6-Bromo-1-(2-(dimethylamino)-5-methylphenyl)naphthalen-2-ol (3b): *N,N*-Dimethyl-*p*-toluidine (**1a**) (81.0 mg, 599 μmol), dichloromethane (6 mL), 6-bromo-2-naphthol (268 mg, 1.20 mmol), methanesulfonic acid (23.1 mg, 240 μmol), FePcF_{16} (5.1 mg, 6.0 μmol). Reaction time: 24 h, column chromatography (isohexane/ethyl acetate, 2–6%, 1 h, 6–15%, 0.5 h, 15–20%, 20 min). Biaryl **3b** (186 mg, 522 μmol , 87%) was obtained as a brownish yellow solid. M.p. 189.5 °C; UV (MeOH): λ = 236, 291, 343 nm; fluorescence (MeOH):

$\lambda_{\text{ex}}=343$ nm, $\lambda_{\text{em}}=423$ nm; IR (ATR): $\tilde{\nu}=3064, 3041, 3025, 2988, 2949, 2912, 2876, 2840, 2797, 1908, 1585, 1490, 1449, 1417, 1358, 1334, 1295, 1278, 1233, 1175, 1134, 1098, 1074, 1036, 952, 923, 897, 816, 755, 694, 670$ cm^{-1} ; $^1\text{H NMR}$ (600 MHz, CDCl_3): $\delta=2.32$ (s, 3H), 2.66 (s, 6H), 7.17 (s, 1H), 7.18 (d, $J=5.6$ Hz, 1H), 7.23 (dd, $J=8.3, 1.9$ Hz, 1H), 7.28 (d, $J=8.7$ Hz, 1H), 7.43 (dd, $J=9.0, 2.3$ Hz, 1H), 7.68 (d, $J=3.8$ Hz, 1H), 7.69 (d, $J=4.1$ Hz, 1H), 7.96 (d, $J=2.3$ Hz, 1H), 10.27 (br s, 1H) ppm; $^{13}\text{C NMR}$ and DEPT (151 MHz, CDCl_3): $\delta=20.78$ (CH_3), 43.87 (2 CH_3), 116.88 (C), 118.32 (CH), 121.44 (C), 121.96 (CH), 127.44 (CH), 128.42 (CH), 129.35 (CH), 129.44 (CH), 130.12 (CH), 130.22 (C), 131.23 (C), 132.11 (C), 132.82 (C), 135.72 (CH), 147.73 (C), 152.66 (C) ppm; ESI-MS (+10 V): $m/z=356.1$ $[\text{M}+\text{H}]^+$; ESI-MS (−10 V): $m/z=354.2$ $[\text{M}-\text{H}]^-$; elemental analysis (%) calcd for $\text{C}_{19}\text{H}_{18}\text{BrNO}$: C 64.06, H 5.09, N 3.93; found: C 63.75, H 5.08, N 3.91.

1-(2-(Dimethylamino)-5-methylphenyl)-7-methoxynaphthalen-2-ol (3c): *N,N*-Dimethyl-*p*-toluidine (**1a**) (81.1 mg, 600 μmol), dichloromethane (6 mL), 7-methoxy-2-naphthol (209 mg, 1.20 mmol), methanesulfonic acid (23.1 mg, 240 μmol), FePcF_{16} (5.1 mg, 6.0 μmol). Reaction time: 24 h, column chromatography (isohexane/ethyl acetate, 2–8%, 1 h, 8–15%, 0.5 h, 15–20%, 20 min). Biaryl **3c** (141 mg, 459 μmol , 76%) was obtained as a beige solid. M.p. 116 °C; UV (MeOH): $\lambda=238, 301$ nm; fluorescence (MeOH): $\lambda_{\text{ex}}=301$ nm, $\lambda_{\text{em}}=412$ nm; IR (ATR): $\tilde{\nu}=2995, 2958, 2932, 2916, 2873, 2840, 2797, 1988, 1904, 1733, 1697, 1619, 1559, 1493, 1454, 1397, 1373, 1358, 1327, 1297, 1263, 1230, 1214, 1175, 1160, 1136, 1092, 1038, 1018, 980, 933, 877, 863, 841, 817, 751, 679, 662, 640, 615$ cm^{-1} ; $^1\text{H NMR}$ (600 MHz, CDCl_3): $\delta=2.33$ (s, 3H), 2.70 (s, 6H), 3.76 (s, 3H), 7.01 (dd, $J=2.6, 1.0$ Hz, 1H), 7.10 (m, 1H), 7.14 (d, $J=8.7$ Hz, 1H), 7.22 (m, 2H), 7.30 (s, 1H), 7.70 (d, $J=9.0$ Hz, 1H), 7.72 (d, $J=8.7$ Hz, 1H), 10.09 (br s, 1H) ppm; $^{13}\text{C NMR}$ and DEPT (151 MHz, CDCl_3): $\delta=20.75$ (CH_3), 44.08 (2 CH_3), 55.38 (CH_3), 104.91 (CH), 115.00 (CH), 118.29 (CH), 118.54 (CH), 119.94 (C), 125.39 (C), 129.24 (CH), 129.37 (CH), 129.88 (CH), 130.72 (C), 133.34 (C, HMBC), 134.76 (C), 135.44 (CH), 146.93 (C), 152.86 (C), 158.16 (C) ppm; ESI-MS (+10 V): $m/z=308.1$ $[\text{M}+\text{H}]^+$; ESI-MS (−10 V): $m/z=305.8$ $[\text{M}-\text{H}]^-$; elemental analysis (%) calcd for $\text{C}_{20}\text{H}_{21}\text{NO}_2$: C 78.15, H 6.89, N 4.56; found: C 77.89, H 6.92, N 4.53.

Methyl 5-(2-(dimethylamino)-5-methylphenyl)-6-hydroxy-2-naphthoate (3d): *N,N*-Dimethyl-*p*-toluidine (**1a**) (81.0 mg, 599 μmol), dichloromethane (6 mL), methyl 6-hydroxy-2-naphthoate (245 mg, 1.21 mmol), methanesulfonic acid (23.1 mg, 240 μmol), FePcF_{16} (5.1 mg, 6.0 μmol). The reaction mixture was warmed up slowly from 0 °C to room temperature. Reaction time: 24 h, column chromatography (isohexane/ethyl acetate, 3–7%, 1.5 h, 7–15%, 0.5 h). Biaryl **3d** (128 mg, 382 μmol , 64%) was obtained as a reddish solid. M.p. 112 °C; UV (MeOH): $\lambda=246, 303$ nm; fluorescence (MeOH): $\lambda_{\text{ex}}=246$ nm, $\lambda_{\text{em}}=441$ nm; IR (ATR): $\tilde{\nu}=3025, 2953, 2909, 2876, 2839, 2794, 1917, 1844, 1792, 1771, 1733, 1709, 1653, 1635, 1618, 1593, 1495, 1472, 1436, 1366, 1341, 1291, 1230, 1196, 1176, 1154, 1135, 1106, 1092, 1038, 975, 957, 929, 909, 819, 790, 753, 694, 674$ cm^{-1} ; $^1\text{H NMR}$ (600 MHz, CDCl_3): $\delta=2.33$ (s, 3H), 2.67 (s, 6H), 3.97 (s, 3H), 7.18 (m, 2H), 7.24 (dd, $J=8.3, 1.3$ Hz, 1H), 7.32 (d, $J=8.7$ Hz, 1H), 7.84 (m, 1H), 7.88 (d, $J=8.7$ Hz, 1H), 7.94 (dd, $J=9.0, 1.5$ Hz, 1H), 8.57 (d, $J=1.5$ Hz, 1H) ppm; $^{13}\text{C NMR}$ and DEPT (151 MHz, CDCl_3): $\delta=20.78$ (CH_3), 43.89 (2 CH_3), 52.23 (CH_3), 118.34 (CH), 121.30 (C), 121.67 (CH), 124.72 (C), 125.62 (CH), 125.68 (CH), 128.99 (C), 129.47 (CH), 130.23 (C), 130.86 (CH), 131.43 (CH), 132.94 (C), 135.81 (CH), 136.06 (C), 147.59 (C), 154.58 (C), 167.57 (C=O) ppm; ESI-MS (+10 V): $m/z=336.2$ $[\text{M}+\text{H}]^+$; ESI-MS (−10 V): $m/z=333.9$ $[\text{M}-\text{H}]^-$; elemental analysis (%) calcd for $\text{C}_{21}\text{H}_{21}\text{NO}_3$: C 75.20, H 6.31, N 4.18; found: C 74.92, H 6.51, N 4.09.

1-(2-(Dimethylamino)-5-methylphenyl)dibenzo[*b,d*]furan-2-ol (3e): *N,N*-Dimethyl-*p*-toluidine (**1a**) (81.3 mg, 601 μmol), dichloromethane (6 mL), 2-hydroxydibenzofuran (**2e**) (223 mg, 1.21 mmol), methanesulfonic acid (23.2 mg, 241 μmol), FePcF_{16} (10.3 mg, 12.0 μmol , 2 mol%). Reaction time: 24 h, column chromatography (isohexane/ethyl acetate, 1–5%, 1 h, 5–10%, 45 min, 10–15%, 15 min). Biaryl **3e** (80.8 mg, 255 μmol , 42%) was obtained as a light pink solid. M.p. 115.5 °C; UV (MeOH): $\lambda=218, 292$ nm; fluorescence (MeOH): $\lambda_{\text{ex}}=292$ nm, $\lambda_{\text{em}}=456$ nm; IR (ATR): $\tilde{\nu}=3002, 2959, 2919, 2873, 2839, 2793, 1907, 1845, 1727, 1613, 1559, 1542, 1499, 1469, 1440, 1423, 1397, 1339, 1299, 1259, 1213, 1185, 1156, 1130, 1112, 1083, 1064, 1041, 1017, 963, 932, 903, 887, 860, 840, 826, 805, 783, 751, 694, 644, 629$ cm^{-1} ; $^1\text{H NMR}$ (600 MHz, CDCl_3): $\delta=2.33$ (s, 3H), 2.71 (s, 6H), 7.09 (m, 1H), 7.17 (d, $J=8.7$ Hz, 1H), 7.21 (d, $J=8.3$ Hz, 1H), 7.28 (dd, $J=8.3, 1.5$ Hz, 1H), 7.39 (m, 2H), 7.48 (m, 2H), 7.55 (d, $J=8.3$ Hz, 1H), 9.89 (br s, 1H) ppm; $^{13}\text{C NMR}$ and DEPT (151 MHz, CDCl_3): $\delta=20.68$ (CH_3), 44.14 (2 CH_3), 111.68 (CH), 111.80 (CH), 118.19 (CH), 118.75 (CH), 121.94 (CH), 122.25 (CH), 122.35 (C), 122.42 (C), 124.76 (C), 126.72 (CH), 129.85 (CH), 130.14 (C), 133.22 (C), 134.72 (CH), 147.09 (C), 150.73 (C), 151.54 (C), 157.13 (C) ppm; ESI-MS (+10 V): $m/z=318.1$ $[\text{M}+\text{H}]^+$; elemental analysis (%) calcd for $\text{C}_{21}\text{H}_{19}\text{NO}_2$: C 79.47, H 6.03, N 4.41; found: C 79.18, H 6.31, N 4.35.

Crystallographic data for compound 3e: $\text{C}_{21}\text{H}_{19}\text{NO}_2$, $M=317.37$ g mol^{-1} , crystal size $0.486 \times 0.628 \times 0.673$ mm^3 , orthorhombic, space group $P2_12_12_1$, $a=7.9372(6)$, $b=12.2884(9)$, $c=17.0584(13)$ Å, $V=1663.8(2)$ Å³, $Z=4$, $\rho_{\text{calcd}}=1.267$ g cm^{-3} , $\mu=0.081$ mm^{-1} , $\lambda=0.71073$ Å, $T=150(2)$ K, θ range = 2.91–28.00°, reflections collected 12725, independent reflections 4017 ($R_{\text{int}}=0.0879$), 224 parameters. The structure was solved by direct methods and refined by full-matrix least-squares on F^2 ; final R indices [$I > 2\sigma(I)$] $R1=0.0472$ and $wR2=0.0876$ maximal residual electron density 0.199 e Å^{-3} . CCDC 1963409 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.

1-(5-Methyl-2-(pyrrolidin-1-yl)phenyl)naphthalen-2-ol (3f): 1-(4-Tolyl)pyrrolidine (64.0 mg, 397 μmol), dichloromethane (4 mL), 2-naphthol (**2a**) (116 mg, 805 μmol), methanesulfonic acid (15.5 mg, 161 μmol), FePcF_{16} (3.4 mg, 4.0 μmol). Reaction time: 4 h, column chromatography (isohexane/ethyl acetate, 1–5%, 1 h, 5–15%, 20 min). Biaryl **3f** (79.0 mg, 65%) was obtained as a pale reddish solid. M.p. 145 °C; UV (MeOH): $\lambda=231, 323$ nm; fluorescence (MeOH): $\lambda_{\text{ex}}=323$ nm, $\lambda_{\text{em}}=480$ nm; IR (ATR): $\tilde{\nu}=3051, 3032, 2970, 2922, 2876, 2853, 1740, 1621, 1591, 1499, 1466, 1396, 1335, 1297, 1276, 1221, 1186, 1154, 1110, 1091, 1041, 1015, 956, 933, 890, 815, 787, 753, 674, 633$ cm^{-1} ; $^1\text{H NMR}$ (600 MHz, CDCl_3): $\delta=1.79$ (m, 4H), 2.32 (s, 3H), 2.83 (m, 2H), 3.08 (m, 2H), 7.08 (d, $J=8.3$ Hz, 1H), 7.18 (m, 2H), 7.25 (d, $J=8.7$ Hz, 1H), 7.35 (m, 1H), 7.40 (m, 1H), 7.78 (d, $J=9.0$ Hz, 1H), 7.82 (d, $J=7.5$ Hz, 1H), 7.85 (d, $J=8.7$ Hz, 1H), 8.73 (br s, 1H) ppm; $^{13}\text{C NMR}$ and DEPT (151 MHz, CDCl_3): $\delta=20.64$ (CH_3), 24.64 (2 CH_2), 51.07 (2 CH_2), 116.59 (CH), 119.68 (CH), 122.13 (C), 123.19 (CH), 125.83 (CH), 126.17 (CH), 127.22 (C), 128.16 (CH), 129.15 (CH), 129.20 (CH), 129.75 (C), 130.57 (C), 133.45 (C), 135.44 (CH), 145.75 (C), 151.96 (C) ppm; ESI-MS (+10 V): $m/z=304.1$ $[\text{M}+\text{H}]^+$; ESI-MS (−10 V): $m/z=301.8$ $[\text{M}-\text{H}]^-$; elemental analysis (%) calcd for $\text{C}_{21}\text{H}_{21}\text{NO}$: C 83.13, H 6.98, N 4.62; found: C 83.05, H 7.20, N 4.57.

1-(5-Methyl-2-(piperidin-1-yl)phenyl)naphthalen-2-ol (3g): 1-(4-Tolyl)piperidine (35.5 mg, 203 μmol), dichloromethane (2 mL), 2-naphthol (**2a**) (116 mg, 806 μmol), methanesulfonic acid (7.8 mg, 81 μmol), FePcF_{16} (1.8 mg, 2.1 μmol), after 24 h, addition of further **2a** (57.9 mg, 402 μmol). Reaction time: 48 h, column chromatography (isohexane/ethyl acetate, 1–3%, 1 h, 3–8%, 0.5 h, 8–15%,

20 min). Biaryl **3g** (44.2 mg, 139 μmol , 69%) was obtained as a colorless solid. M.p. 126.5 °C; UV (MeOH): $\lambda = 231, 286, 335$ nm; fluorescence (MeOH): $\lambda_{\text{ex}} = 231$ nm, $\lambda_{\text{em}} = 411$ nm; IR (ATR): $\tilde{\nu} = 3058, 3028, 2936, 2920, 2844, 1590, 1495, 1465, 1438, 1378, 1332, 1276, 1227, 1147, 1099, 1063, 1030, 953, 918, 903, 867, 817, 751, 726, 676, 635$ cm^{-1} ; $^1\text{H NMR}$ (600 MHz, CDCl_3): $\delta = 1.55$ (m, 6H), 2.32 (s, 3H), 2.89 (m, 4H), 7.13 (d, $J = 8.3$ Hz, 1H), 7.21 (m, 2H), 7.30 (d, $J = 8.7$ Hz, 1H), 7.34 (m, 1H), 7.38 (m, 1H), 7.77 (d, $J = 9.0$ Hz, 1H), 7.82 (m, 2H) ppm; $^{13}\text{C NMR}$ and DEPT (151 MHz, CDCl_3): $\delta = 20.81$ (CH_3), 23.93 (CH_2), 26.22 (2 CH_2), 53.72 (2 CH_2), 118.69 (CH), 121.02 (CH), 121.31 (C), 123.14 (CH), 125.60 (CH), 126.06 (CH), 128.26 (CH), 129.15 (CH), 129.34 (CH), 130.16 (C), 131.19 (C), 132.63 (C), 133.63 (C), 135.87 (CH), 148.09 (C), 151.83 (C) ppm; ESI-MS (+10 V): $m/z = 318.2$ [$M+H$] $^+$; elemental analysis (%) calcd for $\text{C}_{22}\text{H}_{23}\text{NO}$: C 83.24, H 7.30, N 4.41; found: C 83.44, H 7.42, N 4.35.

4a (49.0 mg, 171 μmol , 28%).

1-(5-Methyl-2-morpholinophenyl)naphthalen-2-ol (3h): 4-(4-Tolyl)morpholine (70.9 mg, 400 μmol), dichloromethane (4 mL), 2-naphthol (**2a**) (346 mg, 2.40 mmol), methanesulfonic acid (15.5 mg, 161 μmol), FePcF_{16} (3.4 mg, 4.0 μmol). Reaction time: 24 h, column chromatography (isohexane/ethyl acetate, 2–5%, 1 h, 5–8%, 45 min, 8–15%, 0.5 h). Biaryl **3h** (90.3 mg, 283 μmol , 71%) was obtained as a brown solid. M.p. 94 °C; UV (MeOH): $\lambda = 230, 288, 334$ nm; fluorescence (MeOH): $\lambda_{\text{ex}} = 334$ nm, $\lambda_{\text{em}} = 412$ nm; IR (ATR): $\tilde{\nu} = 3055, 2960, 2912, 2851, 1726, 1618, 1593, 1497, 1450, 1362, 1334, 1298, 1275, 1227, 1154, 1113, 1068, 1045, 954, 916, 877, 815, 749, 730, 676, 662, 635, 613$ cm^{-1} ; $^1\text{H NMR}$ (600 MHz, CDCl_3): $\delta = 2.34$ (s, 3H), 2.99 (m, 4H), 3.68 (m, 4H), 7.18 (d, $J = 8.3$ Hz, 1H), 7.22 (d, $J = 1.5$ Hz, 1H), 7.25 (d, $J = 1.5$ Hz, 1H), 7.32 (d, $J = 8.7$ Hz, 1H), 7.36 (m, 1H), 7.38 (m, 1H), 7.73 (d, $J = 8.3$ Hz, 1H), 7.79 (d, $J = 8.7$ Hz, 1H), 7.83 (dd, $J = 7.9, 1.5$ Hz, 1H), 9.69 (br s, 1H) ppm; $^{13}\text{C NMR}$ and DEPT (151 MHz, CDCl_3): $\delta = 20.84$ (CH_3), 52.49 (2 CH_2), 66.84 (2 CH_2), 118.52 (CH), 120.60 (CH), 120.64 (C), 123.45 (CH), 125.35 (CH), 126.40 (CH), 128.34 (CH), 129.58 (CH), 129.77 (CH), 130.12 (C), 130.44 (C), 133.40 (C), 133.73 (C), 135.97 (CH), 146.26 (C), 151.42 (C) ppm; ESI-MS (+10 V): $m/z = 320.2$ [$M+H$] $^+$; ESI-MS (–10 V): $m/z = 318.1$ [$M-H$] $^-$, HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{22}\text{NO}_2^+$ ([$M+H$] $^+$): 320.1651; found: 320.1654.

4a (116 mg, 405 μmol , 34%).

1-(2-(Dimethylamino)-5-methoxyphenyl)naphthalen-2-ol (3i): 4-Methoxy-*N,N*-dimethylaniline (90.7 mg, 600 μmol), dichloromethane (6 mL), 2-naphthol (**2a**) (177 mg, 1.23 mmol), methanesulfonic acid (23.3 mg, 241 μmol), FePcF_{16} (5.1 mg, 6.0 μmol). Reaction time: 24 h, column chromatography (isohexane/ethyl acetate, 2–8%, 1.5 h, 8–15%, 20 min). Biaryl **3i** (144 mg, 491 μmol , 82%) was obtained as a colorless solid. M.p. 141.5 °C; UV (MeOH): $\lambda = 233, 287, 336$ nm; fluorescence (MeOH): $\lambda_{\text{ex}} = 233$ nm, $\lambda_{\text{em}} = 412$ nm; IR (ATR): $\tilde{\nu} = 3045, 3004, 2972, 2954, 2919, 2873, 2829, 2790, 1619, 1594, 1558, 1542, 1500, 1460, 1408, 1361, 1337, 1308, 1292, 1238, 1192, 1168, 1037, 1005, 954, 922, 879, 861, 816, 754, 724, 698, 675, 636, 606$ cm^{-1} ; $^1\text{H NMR}$ (600 MHz, CDCl_3): $\delta = 2.67$ (s, 6H), 3.75 (s, 3H), 6.98 (m, 2H), 7.24 (d, $J = 8.3$ Hz, 1H), 7.29 (d, $J = 8.7$ Hz, 1H), 7.34 (m, 1H), 7.38 (m, 1H), 7.80 (d, $J = 8.7$ Hz, 1H), 7.83 (m, 2H), 10.49 (br s, 1H) ppm; $^{13}\text{C NMR}$ and DEPT (151 MHz, CDCl_3): $\delta = 44.26$ (2 CH_3), 55.78 (CH_3), 114.33 (CH), 119.40 (CH), 120.11 (CH), 120.65 (C), 120.94 (CH), 123.25 (CH), 125.30 (CH), 126.31 (CH), 128.33 (CH), 129.70 (CH), 129.99 (C), 132.30 (C), 133.49 (C), 143.20 (C), 152.39 (C), 155.44 (C) ppm; ESI-MS (+10 V): $m/z = 294.1$ [$M+H$] $^+$; elemental analysis (%) calcd for $\text{C}_{19}\text{H}_{19}\text{NO}_2$: C 77.79, H 6.53, N 4.77; found: C 77.76, H 6.81, N 4.77.

4a (34.2 mg, 120 μmol , 20%).

6-Bromo-1-(2-(dimethylamino)-5-methoxyphenyl)naphthalen-2-ol (3j): 4-Methoxy-*N,N*-dimethylaniline (90.7 mg, 600 μmol), dichloromethane (6 mL), 6-bromo-2-naphthol (268 mg, 1.20 mmol), methanesulfonic acid (23.2 mg, 241 μmol), FePcF_{16} (5.1 mg, 6.0 μmol). Reaction time: 24 h, column chromatography (isohexane/ethyl acetate, 2–8%, 1.5 h, 8–15%, 20 min). Biaryl **3j** (196 mg, 526 μmol , 88%) was obtained as a dark yellow solid. M.p. 169 °C; UV (MeOH): $\lambda = 235, 282, 348$ nm; fluorescence (MeOH): $\lambda_{\text{ex}} = 235$ nm, $\lambda_{\text{em}} = 422$ nm; IR (ATR): $\tilde{\nu} = 3058, 3021, 2985, 2949, 2924, 2850, 2830, 2790, 1736, 1679, 1630, 1574, 1490, 1451, 1415, 1360, 1331, 1290, 1233, 1203, 1163, 1068, 1038, 1003, 953, 924, 877, 813, 769, 672, 615$ cm^{-1} ; $^1\text{H NMR}$ (600 MHz, CDCl_3): $\delta = 2.67$ (s, 6H), 3.75 (s, 3H), 6.91 (d, $J = 3.0$ Hz, 1H), 6.98 (dd, $J = 8.7, 3.0$ Hz, 1H), 7.23 (d, $J = 8.7$ Hz, 1H), 7.30 (d, $J = 9.0$ Hz, 1H), 7.42 (dd, $J = 9.0, 2.3$ Hz, 1H), 7.69 (d, $J = 6.0$ Hz, 1H), 7.70 (d, $J = 6.4$ Hz, 1H), 7.96 (d, $J = 2.3$ Hz, 1H), 10.51 (br s, 1H) ppm; $^{13}\text{C NMR}$ and DEPT (151 MHz, CDCl_3): $\delta = 44.30$ (2 CH_3), 55.78 (CH_3), 114.54 (CH), 116.91 (C), 119.56 (CH), 119.98 (CH), 120.83 (C), 122.09 (CH), 127.18 (CH), 128.74 (CH), 129.45 (CH), 130.20 (CH), 131.16 (C), 131.83 (C), 132.07 (C), 143.11 (C), 152.79 (C), 155.56 (C) ppm; ESI-MS (+10 V): $m/z = 372.1$ [$M+H$] $^+$; ESI-MS (–10 V): $m/z = 369.9$ [$M-H$] $^-$, elemental analysis (%) calcd for $\text{C}_{19}\text{H}_{18}\text{BrNO}_2$: C 61.30, H 4.87, N 3.76; found: C 60.97, H 4.83, N 3.52.

1-(5-Chloro-2-(dimethylamino)phenyl)naphthalen-2-ol (3k): 4-Chloro-*N,N*-dimethylaniline (93.4 mg, 600 μmol), dichloromethane (6 mL), 2-naphthol (**2a**) (347 mg, 2.41 mmol), methanesulfonic acid (23.1 mg, 240 μmol), FePcF_{16} (5.1 mg, 6.0 μmol). Reaction time: 24 h, column chromatography (isohexane/ethyl acetate, 1–3%, 0.5 h, 3–7%, 1.5 h, 7–15%, 20 min). Biaryl **3k** (63.8 mg, 214 μmol , 36%) was obtained as a brownish yellow solid. M.p. 128 °C; UV (MeOH): $\lambda = 230, 282, 333$ nm; fluorescence (MeOH): $\lambda_{\text{ex}} = 230$ nm, $\lambda_{\text{em}} = 364, 422$ nm; IR (ATR): $\tilde{\nu} = 3051, 2959, 2916, 2876, 2845, 2797, 2056, 1701, 1632, 1592, 1510, 1474, 1455, 1391, 1358, 1336, 1299, 1272, 1233, 1176, 1150, 1127, 1112, 1038, 997, 952, 924, 890, 849, 815, 787, 746, 721, 686, 640, 624$ cm^{-1} ; $^1\text{H NMR}$ (600 MHz, CDCl_3): $\delta = 2.67$ (s, 6H), 7.20 (d, $J = 8.7$ Hz, 1H), 7.27 (d, $J = 8.8$ Hz, 1H), 7.37 (m, 2H), 7.40 (d, $J = 2.4$ Hz, 1H), 7.42 (m, 1H), 7.76 (d, $J = 8.5$ Hz, 1H), 7.81 (d, $J = 8.8$ Hz, 1H), 7.83 (m, 1H), 9.66 (br s, 1H) ppm; $^{13}\text{C NMR}$ and DEPT (151 MHz, CDCl_3): $\delta = 43.72$ (2 CH_3), 119.62 (CH), 119.85 (C), 120.70 (CH), 123.51 (CH), 124.97 (CH), 126.72 (CH), 128.33 (C), 128.39 (CH), 128.40 (CH), 130.04 (C), 130.09 (CH), 132.50 (C), 133.13 (C), 134.81 (CH), 148.72 (C), 152.26 (C) ppm; ESI-MS (+10 V): $m/z = 298.1$ [$M+H$] $^+$; elemental analysis (%) calcd for $\text{C}_{18}\text{H}_{16}\text{ClNO}$: C 72.60, H 5.42, N 4.70; found: C 72.63, H 5.50, N 4.39.

4a (90.5 mg, 316 μmol , 26%).

1,1-Bis(2-(dimethylamino)-4-(triisopropylsilyloxy)phenyl)naphthalen-2(1H)-one (6): *N,N*-Dimethyl-3-(triisopropylsilyloxy)aniline (**5**) (173 mg, 589 μmol), dichloromethane (6 mL), 2-naphthol (**2a**) (173 mg, 1.20 mmol), methanesulfonic acid (23.1 mg, 240 μmol), FePcF_{16} (10.5 mg, 12.3 μmol , 2 mol%). Reaction time: 24 h, column chromatography (isohexane/ethyl acetate, 1–3%, 1 h, 3–15%, 0.5 h, 15%, 20 min). Compound **6** was obtained as a pale yellow solid (78.2 mg, 108 μmol , 37%). M.p. 261 °C; UV (MeOH): $\lambda = 219, 259, 293$ nm; fluorescence (MeOH): $\lambda_{\text{ex}} = 259$ nm, $\lambda_{\text{em}} = 443$ nm; IR (ATR): $\tilde{\nu} = 2942, 2886, 2864, 2790, 1738, 1690, 1651, 1606, 1557, 1542, 1508, 1453, 1423, 1356, 1255, 1185, 1122, 1091, 998, 922, 882, 798, 752, 680, 641$ cm^{-1} ; $^1\text{H NMR}$ (500 MHz, CDCl_3): $\delta = 0.83$ (d, $J = 7.6$ Hz, 9H), 0.88 (d, $J = 7.6$ Hz, 9H), 0.94 (d, $J = 7.6$ Hz, 9H), 1.01 (d, $J = 7.6$ Hz, 9H), 1.06 (m, 3H), 1.33 (m, 3H), 2.86 (s, 6H), 2.87 (s, 6H), 5.94 (d, $J = 10.1$ Hz, 1H), 6.04 (m, 2H), 6.13 (d, $J = 2.2$ Hz, 1H), 6.21 (br d, $J = 9.1$ Hz, 2H), 6.75 (m, 2H), 6.97 (d, $J = 10.1$ Hz, 1H), 7.09 (m, 1H), 7.19 (m, 2H) ppm; $^{13}\text{C NMR}$ and DEPT (125 MHz, CDCl_3): $\delta = 13.23$ (3 CH), 13.61 (3 CH), 18.09 (3 CH_3), 18.19 (3 CH_3),

18.29 (3 CH₃), 18.42 (3 CH₃), 40.49 (2 CH₃), 41.50 (2 CH₃), 63.91 (C), 103.24 (CH), 104.23 (CH), 104.86 (CH), 105.03 (CH), 116.07 (C), 119.06 (C), 126.57 (CH), 127.37 (CH), 128.28 (CH), 128.76 (CH), 129.52 (CH), 130.43 (CH), 132.33 (C), 133.29 (CH), 139.09 (CH), 146.27 (C), 151.21 (C), 154.33 (C), 157.57 (C), 194.74 (C=O) ppm (the signal for one C is missing due to overlapping); ESI-MS (+25 V): m/z = 727.8 [M+H]⁺; elemental analysis (%) calcd for C₄₄H₆₆N₂O₃Si₂: C 72.67, H 9.15, N 3.85; found: C 72.97, H 9.19, N 4.10.

4a (33.4 mg, 117 μmol, 19%).

2'-Dimethylamino-2-hydroxy-1,1'-binaphthyl (N,N-dimethyl-NOBIN) (8) and N,N-dimethyl-1-(naphthalen-2-yloxy)naphthalen-2-amine (9): A solution of 2-N,N-dimethylaminonaphthalene (7) (413 mg, 2.41 mmol) in dichloromethane (26 mL) was added to a mixture of 2-naphthol (**2a**) (1.41 g, 9.80 mmol), methanesulfonic acid (93.2 mg, 967 μmol), and FePcF₁₆ (41.1 mg, 48.0 μmol, 2 mol%). The reaction mixture was vigorously stirred at 0 °C for 24 h under air and then quenched with a saturated aqueous solution of Na₂CO₃ (25 mL). The aqueous layer was extracted three times with dichloromethane (25 mL) and the combined organic layers were dried over MgSO₄. Evaporation of the solvent in vacuo and purification of the crude product by automated column chromatography on silica gel (isohexane/ethyl acetate, 0–2%, 0.5 h, 2–6%, 2 h, 6–18%, 0.5 h) afforded the coupling products **8**, **9**, and **4a**. *N,N*-Dimethyl-NOBIN (**8**) (470 mg, 1.50 mmol, 62%) was obtained as a yellow solid. M.p. 151.5 °C; UV (MeOH): λ = 219, 254, 290, 335 nm; fluorescence (MeOH): λ_{ex} = 254 nm, λ_{em} = 423 nm; IR (ATR): $\tilde{\nu}$ = 3061, 2975, 2953, 2866, 2835, 2789, 1737, 1618, 1590, 1542, 1507, 1458, 1397, 1353, 1325, 1293, 1271, 1223, 1195, 1142, 1092, 1068, 1039, 988, 952, 935, 857, 816, 748, 697, 655, 631 cm⁻¹; ¹H NMR (600 MHz, CDCl₃): δ = 2.66 (s, 6H), 7.04 (d, J = 8.7 Hz, 1H), 7.09 (d, J = 8.5 Hz, 1H), 7.16 (ddd, J = 8.5, 7.0, 1.1 Hz, 1H), 7.21 (ddd, J = 8.5, 7.0, 1.1 Hz, 1H), 7.32 (m, 2H), 7.38 (d, J = 8.8 Hz, 1H), 7.52 (d, J = 9.0 Hz, 1H), 7.84 (d, J = 8.3 Hz, 1H), 7.87 (d, J = 8.1 Hz, 1H), 7.90 (d, J = 8.8 Hz, 1H), 7.96 (d, J = 9.0 Hz, 1H) ppm; ¹³C NMR and DEPT (151 MHz, CDCl₃): δ = 43.85 (2 CH₃), 118.41 (C), 118.45 (CH), 119.60 (CH), 122.48 (C), 123.38 (CH), 124.46 (CH), 125.84 (CH), 126.10 (CH), 126.48 (CH), 126.71 (CH), 128.07 (CH), 128.31 (CH), 129.42 (C), 129.90 (CH), 130.12 (CH), 130.34 (C), 134.07 (C), 134.27 (C), 149.24 (C), 151.79 (C) ppm; ESI-MS (+25 V): m/z = 314.1 [M+H]⁺; ESI-MS (–25 V): m/z = 311.8 [M–H][–]; elemental analysis (%) calcd for C₂₂H₁₉NO: C 84.31, H 6.11, N 4.47; found: C 84.44, H 6.12, N 4.43.

Crystallographic data for compound 8: C₂₂H₁₉NO, M = 313.38 g mol⁻¹, crystal size 0.210 × 0.320 × 0.670 mm³, orthorhombic, space group *P*2₁2₁1, a = 8.3318(7), b = 10.8815(9), c = 18.0791(15) Å, V = 1639.1(2) Å³, Z = 4, ρ_{calcd} = 1.270 g cm⁻³, μ = 0.077 mm⁻¹, λ = 0.71073 Å, T = 100(2) K, θ range = 2.25–28.98°, reflections collected 30910, independent reflections 4356 (R_{int} = 0.0578), 223 parameters. The structure was solved by direct methods and refined by full-matrix least-squares on F^2 ; final R indices [$I > 2\sigma(I)$] $R1$ = 0.0450 and $wR2$ = 0.1027; maximal residual electron density 0.290 e Å⁻³. CCDC 1963410 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.

N,N-Dimethyl-1-(naphthalen-2-yloxy)naphthalen-2-amine (**9**) (81.1 mg, 25.9 μmol, 11%) was obtained as a beige solid. M.p. 102 °C; UV (MeOH): λ = 215, 226, 254, 303, 328 nm; fluorescence (MeOH): λ_{ex} = 226 nm, λ_{em} = 416 nm; IR (ATR): $\tilde{\nu}$ = 3058, 2948, 2843, 2800, 1734, 1697, 1653, 1624, 1596, 1559, 1540, 1504, 1457, 1428, 1375, 1320, 1249, 1231, 1212, 1158, 1115, 1070, 991, 957, 910, 844, 801, 747, 713, 685, 634 cm⁻¹; ¹H NMR (500 MHz, CDCl₃): δ = 2.91 (s, 6H), 6.91 (d, J = 2.5 Hz, 1H), 7.28 (dd, J = 8.8, 2.5 Hz, 1H), 7.34 (m,

5H), 7.54 (d, J = 8.2 Hz, 1H), 7.74 (d, J = 8.8 Hz, 1H), 7.78 (m, 2H), 7.82 (m, 1H), 7.86 (m, 1H) ppm; ¹³C NMR and DEPT (125 MHz, CDCl₃): δ = 42.89 (2 CH₃), 109.58 (CH), 117.80 (CH), 119.46 (CH), 121.31 (CH), 123.84 (CH), 123.87 (CH), 125.78 (CH), 126.35 (CH), 126.58 (CH), 127.04 (C), 127.74 (CH), 127.76 (CH), 129.38 (CH), 129.60 (CH), 129.88 (C), 134.59 (C), 138.41 (C), 142.12 (C), 156.35 (C) ppm (the signal for one C is missing due to overlapping); MS (EI): m/z = 313 (70, [M]⁺), 186 (26), 184 (10), 173 (13), 172 (100), 170 (12), 142 (15), 141 (23), 128 (14), 127 (49), 126 (12), 115 (40), 102 (10), 77 (11), 42 (42); elemental analysis (%) calcd for C₂₂H₁₉NO: C 84.31, H 6.11, N 4.47; found: C 84.67, H 6.38, N 4.54.

4a (281 mg, 981 μmol, 20%).

Acknowledgements

We are grateful to the Deutsche Forschungsgemeinschaft (DFG) for the financial support of our project „Green and Sustainable Catalysts for Synthesis of Organic Building Blocks“ (DFG grant KN 240/19-2). We thank Stefan König for experimental support.

Conflict of interest

The authors declare no conflict of interest.

Keywords: C–H bond activation · homogeneous catalysis · iron · oxidative cross-coupling · phthalocyanines

- [1] For reviews, see: a) I. P. Beletskaya, A. V. Cheprakov, *Chem. Rev.* **2000**, *100*, 3009–3066; b) N. J. Whitcombe, K. K. Hii, S. E. Gibson, *Tetrahedron* **2001**, *57*, 7449–7476; c) N. Miyaura, *Top. Curr. Chem.* **2002**, *219*, 11–59; d) *Metal-Catalyzed Cross-Coupling Reactions, Vol. 1 and 2*, (Eds.: A. de Meijere, F. Diederich), Wiley-VCH, Weinheim, 2nd ed., **2004**; e) A. Suzuki, Y. Yamamoto, *Chem. Lett.* **2011**, *40*, 894–901; f) C. C. C. Johansson Seechurn, M. O. Kitching, T. J. Colacot, V. Snieckus, *Angew. Chem. Int. Ed.* **2012**, *51*, 5062–5085; *Angew. Chem.* **2012**, *124*, 5150–5174.
- [2] a) *Catalysis without Precious Metals* (Ed.: R. M. Bullock), Wiley-VCH, Weinheim, **2010**, pp. 1–290; b) N. V. Tzouras, I. K. Stamatopoulos, A. T. Papatavrou, A. A. Liori, G. C. Vougioukalakis, *Coord. Chem. Rev.* **2017**, *343*, 25–138; c) *Non-Noble Metal Catalysis: Molecular Approaches and Reactions* (Eds.: R. J. M. Klein Gebbink, M.-E. Moret), Wiley-VCH, Weinheim, **2019**, pp. 1–590.
- [3] a) C. Bolm, J. Legros, J. Le Paih, L. Zani, *Chem. Rev.* **2004**, *104*, 6217–6254; b) *Iron Catalysis in Organic Chemistry: Reactions and Applications* (Ed.: B. Plietker), Wiley-VCH, Weinheim, **2008**, pp. 1–279; c) B. D. Sherry, A. Fürstner, *Acc. Chem. Res.* **2008**, *41*, 1500–1511; d) S. Enthaler, K. Junge, M. Beller, *Angew. Chem. Int. Ed.* **2008**, *47*, 3317–3321; *Angew. Chem.* **2008**, *120*, 3363–3367; e) K. Junge, K. Schröder, M. Beller, *Chem. Commun.* **2011**, *47*, 4849–4859; f) C.-L. Sun, B.-J. Li, Z.-J. Shi, *Chem. Rev.* **2011**, *111*, 1293–1314; g) K. Gopalaiah, *Chem. Rev.* **2013**, *113*, 3248–3296; h) E. Bauer, *Top. Organomet. Chem.* **2015**, *50*, 1–356; i) I. Bauer, H.-J. Knölker, *Chem. Rev.* **2015**, *115*, 3170–3387; j) A. Fürstner, *ACS Cent. Sci.* **2016**, *2*, 778–789; k) R. Shang, L. Ilies, E. Nakamura, *Chem. Rev.* **2017**, *117*, 9086–9139.
- [4] a) C.-J. Li, *Acc. Chem. Res.* **2009**, *42*, 335–344; b) C. S. Yeung, V. M. Dong, *Chem. Rev.* **2011**, *111*, 1215–1292; c) S. A. Girard, T. Knauber, C.-J. Li, *Angew. Chem. Int. Ed.* **2014**, *53*, 74–100; *Angew. Chem.* **2014**, *126*, 76–103.
- [5] For the Scholl reaction, see: a) A. A. O. Sarhan, C. Bolm, *Chem. Soc. Rev.* **2009**, *38*, 2730–2744; b) M. Grzybowski, K. Skonieczny, H. Butenschön, D. T. Gryko, *Angew. Chem. Int. Ed.* **2013**, *52*, 9900–9930; *Angew. Chem.* **2013**, *125*, 10084–10115; c) M. Grzybowski, B. Sadowski, H. Butenschön, D. T. Gryko, *Angew. Chem. Int. Ed.* **2020**, *59*, 2998–3027; *Angew. Chem.* **2020**, *132*, 3020–3050; for electrosynthetic coupling reactions, see:

- d) B. R. Rosen, E. W. Werner, A. G. O'Brien, P. S. Baran, *J. Am. Chem. Soc.* **2014**, *136*, 5571–5574; e) S. R. Waldvogel, B. Janza, *Angew. Chem. Int. Ed.* **2014**, *53*, 7122–7123; *Angew. Chem.* **2014**, *126*, 7248–7249; f) B. Elsler, D. Schollmeyer, K. M. Dyballa, R. Franke, S. R. Waldvogel, *Angew. Chem. Int. Ed.* **2014**, *53*, 4979; *Angew. Chem.* **2014**, *126*, 5311–5314; g) T. Gieshoff, D. Schollmeyer, S. R. Waldvogel, *Angew. Chem. Int. Ed.* **2016**, *55*, 9437–9440; *Angew. Chem.* **2016**, *128*, 9587–9590; h) A. Wiebe, D. Schollmeyer, K. M. Dyballa, R. Franke, S. R. Waldvogel, *Angew. Chem. Int. Ed.* **2016**, *55*, 11801–11805; *Angew. Chem.* **2016**, *128*, 11979–11983.
- [6] a) Z. Li, L. Cao, C.-J. Li, *Angew. Chem. Int. Ed.* **2007**, *46*, 6505–6507; *Angew. Chem.* **2007**, *119*, 6625–6627; b) Y. Zhang, C.-J. Li, *Eur. J. Org. Chem.* **2007**, 4654–4657.
- [7] a) M. Chandrasekharam, B. Chiranjeevi, K. S. V. Gupta, B. Sridhar, *J. Org. Chem.* **2011**, *76*, 10229–10235; b) B. Chiranjeevi, G. Koyyada, S. Prabuseenivasan, V. Kumar, P. Sujitha, C. G. Kumar, B. Sridhar, S. Shaik, M. Chandrasekharam, *RSC Adv.* **2013**, *3*, 16475–16485; c) B. Chiranjeevi, B. Vinayak, T. Parsharamulu, V. S. PhaniBabu, B. Jagadeesh, B. Sridhar, M. Chandrasekharam, *Eur. J. Org. Chem.* **2014**, 7839–7849.
- [8] a) E. Gaster, Y. Vainer, A. Regev, S. Narute, K. Sudheendran, A. Werbeloff, H. Shalit, D. Pappo, *Angew. Chem. Int. Ed.* **2015**, *54*, 4198–4202; *Angew. Chem.* **2015**, *127*, 4272–4276; b) V. Vershinin, A. Dyadyuk, D. Pappo, *Tetrahedron* **2017**, *73*, 3660–3668; c) H. Shalit, A. Libman, D. Pappo, *J. Am. Chem. Soc.* **2017**, *139*, 13404–13413; d) H. Shalit, A. Dyadyuk, D. Pappo, *J. Org. Chem.* **2019**, *84*, 1677–1686.
- [9] K. Wang, M. Lü, A. Yu, X. Zhu, Q. Wang, *J. Org. Chem.* **2009**, *74*, 935–938.
- [10] a) S. Guo, Y. Li, Y. Wang, X. Guo, X. Meng, B. Chen, *Adv. Synth. Catal.* **2015**, *357*, 950–954; b) K. Liang, J. Yang, X. Tong, W. Shang, Z. Pan, C. Xia, *Org. Lett.* **2016**, *18*, 1474–1477.
- [11] a) H. Egami, T. Katsuki, *J. Am. Chem. Soc.* **2009**, *131*, 6082–6083; b) H. Egami, K. Matsumoto, T. Oguma, T. Kunisu, T. Katsuki, *J. Am. Chem. Soc.* **2010**, *132*, 13633–13635.
- [12] T. Niu, Y. Zhang, *Tetrahedron Lett.* **2010**, *51*, 6847–6851.
- [13] K. Matsumoto, M. Yoshida, M. Shindo, *Angew. Chem. Int. Ed.* **2016**, *55*, 5272–5276; *Angew. Chem.* **2016**, *128*, 5358–5362.
- [14] R. F. Fritsche, G. Theumer, O. Kataeva, H.-J. Knölker, *Angew. Chem. Int. Ed.* **2017**, *56*, 549–553; *Angew. Chem.* **2017**, *129*, 564–568.
- [15] C. Brütting, R. F. Fritsche, S. K. Kutz, C. Börger, A. W. Schmidt, O. Kataeva, H.-J. Knölker, *Chem. Eur. J.* **2017**, *23*, 458–470.
- [16] a) G. Bringmann, A. J. P. Mortimer, P. A. Keller, M. J. Gresser, J. Garner, M. Breuning, *Angew. Chem. Int. Ed.* **2005**, *44*, 5384–5427; *Angew. Chem.* **2005**, *117*, 5518–5563; b) D. Alberico, M. E. Scott, M. Lautens, *Chem. Rev.* **2007**, *107*, 174–238; c) M. C. Kozlowski, B. J. Morgan, E. C. Linton, *Chem. Soc. Rev.* **2009**, *38*, 3193–3207; d) G. Bringmann, T. Gulder, T. A. M. Gulder, M. Breuning, *Chem. Rev.* **2011**, *111*, 563–639; e) A. W. Schmidt, K. R. Reddy, H.-J. Knölker, *Chem. Rev.* **2012**, *112*, 3193–3328; f) J. E. Smyth, N. M. Butler, P. A. Keller, *Nat. Prod. Rep.* **2015**, *32*, 1562–1583; g) P. Loxq, E. Manoury, R. Poli, E. Deydier, A. Labande, *Coord. Chem. Rev.* **2016**, *308*, 131–190.
- [17] J. G. Jones, M. V. Twigg, *Inorg. Chem.* **1969**, *8*, 2018–2019.
- [18] For synthesis and reactivity of iron-phthalocyanine complexes, see: a) D. Vonderschmitt, K. Bernauer, S. Fallab, *Helv. Chim. Acta* **1965**, *48*, 951–954; b) I. Collamati, *Inorg. Chim. Acta* **1979**, *35*, L303–L304; c) C. Ercolani, F. Monacelli, G. Rossi, *Inorg. Chem.* **1979**, *18*, 712–716; d) C. Ercolani, G. Rossi, F. Monacelli, *Inorg. Chim. Acta* **1980**, *44*, L215–L216; e) C. Ercolani, M. Gardini, F. Monacelli, G. Pennesi, G. Rossi, *Inorg. Chem.* **1983**, *22*, 2584–2589; f) A. L. Balch, Y. W. Chan, R. J. Cheng, G. N. La Mar, L. Latos-Grazynski, M. W. Renner, *J. Am. Chem. Soc.* **1984**, *106*, 7779–7785; g) R. Dieing, G. Schmid, E. Witke, C. Feucht, M. Dreßen, J. Pohmer, M. Hanack, *Chem. Ber.* **1995**, *128*, 589–598.
- [19] a) A. B. Sorokin, *Chem. Rev.* **2013**, *113*, 8152–8191; b) A. B. Sorokin, *Coord. Chem. Rev.* **2019**, *389*, 141–160.
- [20] F. Puls, H.-J. Knölker, *Angew. Chem. Int. Ed.* **2018**, *57*, 1222–1226; *Angew. Chem.* **2018**, *130*, 1236–1240.
- [21] T. Doba, T. Matsubara, L. Iliès, R. Shang, E. Nakamura, *Nat. Catal.* **2019**, *2*, 400–406.
- [22] a) X. Ling, Y. Xiong, R. Huang, X. Zhang, S. Zhang, C. Chen, *J. Org. Chem.* **2013**, *78*, 5218–5226; b) J. Pan, J. Li, R. Huang, X. Zhang, H. Shen, Y. Xiong, X. Zhu, *Tetrahedron* **2015**, *71*, 5341–5346.
- [23] a) E. Yashima, C. Yamamoto, Y. Okamoto, *J. Am. Chem. Soc.* **1996**, *118*, 4036–4048; b) K. Ding, Y. Wang, L. Zhang, Y. Wu, T. Matsuura, *Tetrahedron* **1996**, *52*, 1005–1010; et loc. cit.
- [24] a) M.-S. Liao, T. Kar, S. M. Gorun, S. Scheiner, *Inorg. Chem.* **2004**, *43*, 7151–7161; b) M.-S. Liao, J. D. Watts, M.-J. Huang, S. M. Gorun, T. Kar, S. Scheiner, *J. Chem. Theory Comput.* **2005**, *1*, 1201–1210.
- [25] a) M. Smrčina, Š. Vyskočil, J. Polívková, J. Polávková, J. Sejbal, V. Hanuš, M. Poláček, H. Verrier, P. Kočovský, *Tetrahedron: Asymmetry* **1997**, *8*, 537–546; b) Š. Vyskočil, S. Jaracz, M. Smrčina, M. Štícha, V. Hanuš, M. Poláček, P. Kočovský, *J. Org. Chem.* **1998**, *63*, 7727–7737; c) K. Ding, H. Guo, X. Li, Y. Yuan, Y. Wang, *Top. Catal.* **2005**, *35*, 105–116.
- [26] a) K. Körber, W. Tang, X. Hu, X. Zhang, *Tetrahedron Lett.* **2002**, *43*, 7163–7165; b) D. C. Patel, Z. S. Breitbach, R. M. Woods, Y. Lim, A. Wang, F. W. Foss Jr., D. W. Armstrong, *J. Org. Chem.* **2016**, *81*, 1295–1299.
- [27] G. M. Sheldrick, SHELXS-97, Programs for Crystal Structure Solution, University of Göttingen, Germany, **1997**.
- [28] G. M. Sheldrick, SADABS, v. 2.10, Bruker/Siemens Area Detector Absorption Correction Program, Bruker AXS Inc., Madison, WI, USA, **2002**.
- [29] G. M. Sheldrick, SHELXL-97, Programs for Crystal Structure Refinement, University of Göttingen, Germany, **1997**.
- [30] L. J. Farrugia, *J. Appl. Crystallogr.* **1997**, *30*, 565.
- [31] M. J. Cawley, F. G. N. Cloke, R. J. Fitzmaurice, S. E. Pearson, J. S. Scott, S. Caddick, *Org. Biomol. Chem.* **2008**, *6*, 2820–2825.
- [32] M. A. Topchiy, A. F. Asachenko, M. S. Nechaev, *Eur. J. Org. Chem.* **2014**, 3319–3322.

Manuscript received: December 11, 2019

Accepted manuscript online: December 20, 2019

Version of record online: February 4, 2020