

## Diversity of desert rangelands of Tunisia

Mouldi Gamoun\*, Azaiez Ouled Belgacem, Mounir Louhaichi

International Center for Agricultural Research in the Dry Areas (ICARDA)



### ARTICLE INFO

#### Article history:

Received 15 February 2018

Received in revised form

19 June 2018

Accepted 20 June 2018

Available online 20 August 2018

(Editor: Richard Corlett)

#### Keywords:

Vegetation

Dryland

Species richness

### ABSTRACT

Plants are important components of any rangeland. However, the importance of desert rangeland plant diversity has often been underestimated. It has been argued that desert rangelands of Tunisia in good ecological condition provide more services than those in poor ecological condition. This is because rangelands in good condition support a more diverse mixture of vegetation with many benefits, such as forage for livestock and medicinal plants.

Nearly one-quarter of Tunisia, covering about 5.5 million hectares, are rangelands, of which 87% are located in the arid and desert areas (45% and 42%, respectively). Here, we provide a brief review of the floristic richness of desert rangelands of Tunisia. Approximately 135 species are specific to desert rangelands. The predominant families are Asteraceae, Poaceae, Brassicaceae, Chenopodiaceae, and Fabaceae. These represent approximately 50% of Tunisian desert flora.

Copyright © 2018 Kunming Institute of Botany, Chinese Academy of Sciences. Publishing services by Elsevier B.V. on behalf of KeAi Communications Co., Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### 1. Introduction

Rangelands cover about 40% of the world's land area (White et al., 2000) and are as ecologically important as rain forests (Casper, 2009). Arid ecosystems comprise one-third of the global land surface, support 14% of the world's inhabitants, and provide a significant share of the world's agriculture (Nicholson, 2011). Arid environments may not be the most hospitable places on Earth, but the 30% or more of the global land surface that they cover does support an ever-growing human population and has fascinated explorers and scientists for centuries (Thomas, 2011).

Desert rangelands, like other arid rangelands, suffer from severe natural disturbances such as high rates of soil degradation and extremely low rainfall distribution, which, in part, may be caused by the climate due to their geographical location (Le Houérou, 2009). These problems are generally compounded by anthropogenic factors such as overgrazing and wood harvesting (Le Houérou, 2009). Taken together, these factors contribute to decreased biological diversity and rangeland productivity, as well as high rates of erosion, all of which are widespread problems in Tunisia (Gamoun, 2012; Tarhouni et al., 2014).

Tunisia is a small country, yet has highly diverse climatic and edaphic conditions (Floret and Pontanier, 1982). Rangelands

constitute the largest use of land in Tunisia, where pastoralism remains vital. Furthermore, rangelands not only provide important goods and services but also represent a tremendous source of biodiversity. Nearly one quarter of Tunisia is rangeland, occupying about 5.5 million hectares, 87% of which are located in the arid and desert areas (45% and 42%, respectively). In Northern Africa, the term "desert" is most properly applied to zones that receive less than 100 mm of average annual rainfall, are little affected by human activity, and possess a very low production potential or likelihood of future evolution (Floret and Pontanier, 1982). The majority of the rangelands of south Tunisia exhibit moderate or severe desertification, and are being damaged and made less productive by mismanagement (Nefzaoui et al., 2011). Desertification includes deterioration of ecosystems and degradation of various forms of vegetation (Le Houérou, 1969).

The causes of desertification are myriad and often interconnected. They include overgrazing and overcutting for firewood and timber, inappropriate farming, poor irrigation, and poor management, which has led to salinity problems, mining, construction of highways and utility corridors, air pollution, recreational activities, particularly off-road vehicle recreation, and climate change (Bainbridge, 2007). The underlying causes of dryland degradation are commonly economic and cultural, rather than ecological (Hallsworth, 1987; Carney and Farrington, 1998; Chambers et al., 1991). The primary causes of current degradation in some arid areas have been identified as intensified and irrational human activities and climate variability (Ouled Belgacem and Louhaichi,

\* Corresponding author.

E-mail address: [m.gamoun@cgiar.org](mailto:m.gamoun@cgiar.org) (M. Gamoun).

Peer review under responsibility of Editorial Office of Plant Diversity.

2013). Overgrazing is the main cause of rangeland degradation and desertification (Le Houérou, 1996; Ibáñez et al., 2007; Gamoun, 2014a, 204b). It may be driven by economic pressure, greed, desperation, and sometimes ignorance (Bainbridge, 2007). For example, Africa as a whole contributes 36% of the world's total land degraded by overgrazing and 49%–90% of the continent's rangelands are believed to already be in the process of long-term degradation (Hudak, 1999).

The primary effects of overgrazing have been reported to be mainly of a biotic nature, resulting in a decrease in plant cover, and in particular, a loss of perennials, which constitutes one of the most significant indicators of desertification (Verstraete and Schwartz, 1991; Aronson et al., 1993). The main consequences of overloading rangelands are summarized as follows: (i) an increase in cultivated land decreases rangeland area; (ii) an increase in livestock number decreases rangeland area; (iii) rangeland is degraded, decreasing both forage yield and carrying capacity (Zhao et al., 1994).

Desertification is largely a response to climatic trends and fluctuations in the availability of resources (Wang et al., 2005). Climate variability (high temperatures and low rainfall distribution) negatively affects rangeland productivity (Peters et al., 2013), and is most severe in dryland areas (Le Houérou, 1996; Darkoh, 1998; Harris, 2010; Reynolds, 2013). Sequences of dry years have been a major climatic force in the degradation of rangelands (Verner, 2013). The unreliable distribution of rainfall is an important component which contributes significantly towards the deterioration of rangeland productivity in the dry areas (Floret et al., 1978; Gamoun, 2013, 2014a, 2014b, 2016a, 2016b). Evidence of this is provided through long term annual rainfall figures in Tunisia, which reveal that desert rangelands are characterized by a dry climate, with hot dry summers and winters and very low levels of precipitation—about 75 mm—from 1997 to 2016 (Figs. 1 and 2).

The flora of desert areas has always attracted ecologists from around the world (Ward, 2016). As a result, the arid zones of Tunisia have been extensively studied over the past fifty years; indeed, the arid zone flora, climate, ecology, hydrogeology, and soils of Tunisia are among the best known in the world.

By any measure, the world's arid rangelands qualify as modest repositories of biodiversity (Shachak et al., 2005). The biological diversity of many desert and semi-desert areas continues to be threatened by many different human activities, such as overharvesting of firewood and overgrazing (Bainbridge, 2007). In these areas, where overgrazing and wood collection drive changes in the structure and functioning of rangelands, any decrease in perennial plant cover and plant species diversity leads to changes in floristic composition, soil erosion, and modifications in nutrient, water, and energy flow (Floret et al., 1978; Jauffret and Lavorel, 2003).

In this paper, we provide a brief review of the floristic richness of part of a desert rangeland in Tunisia.

## 2. Rangeland diversity

Researchers in Tunisia have focused a lot of attention on rangelands as well as on plant diversity and community ecology (Ouled Belgacem et al., 2011; Tarhouni et al., 2015). Tunisia's rangelands span a wide variety of environments from the arid steppe to desert rangelands of southern Tunisia. As in all Mediterranean countries, these steppe zones were previously used mainly for grazing and have been subjected to severe pressure from an increasing population, which has slowly stabilized (Floret and Hadje, 1977). In response to climatic conditions and grazing, the species composition, diversity and abundance, community structure, and plant life forms also changed (Ouled Belgacem et al., 2011; Tarhouni et al., 2015; Gamoun et al., 2016a, 2016b). Heavy grazing has left




Fig. 1. Isohyetal map of Tunisia (curves with mm values represent mean annual rainfall).

Tunisian ecosystems with a homogenized flora consisting only of species that are unpalatable and highly tolerant to herbivory and other forms of disturbance (Gondard et al., 2003; Jauffret and Lavorel, 2003). Sand deposits due to strong desert winds also further decrease vegetation growth by burying some plant parts, gradually decreasing the floristic variety (Bendali et al., 1990).

Sustainable rangeland management approaches have now been used to protect some sections of these desert rangelands in an effort to reduce the effects of degradation by increasing the vegetation composition, spatial distribution, and structure (Floret, 1981; Ouled Belgacem et al., 2008; Gamoun et al., 2010; Tarhouni et al., 2015). As a result, such sustainable practices have yielded diverse rangeland resources, with an abundant and rich diversity of plant species (Figs. 3 and 4 and Table 1).

Tunisia's rangelands support 2162 species (Le Floc'h et al., 2010). In addition, rangelands in pre-Saharan Tunisia harbor a rich flora which includes about 836 species (Ferchichi, 2000). Approximately 135 of these species are specific to desert rangelands (Gamoun, 2012). Five predominant plant families, Asteraceae, Poaceae, Brassicaceae, Chenopodiaceae, and Fabaceae, represent approximately 50% of Tunisian desert flora (Fig. 5).

In the Tunisian desert rangelands the abundance of plant life forms includes 75 species of Therophyte (49%), 38 species of Chamaephyte (25%), 20 species of Hemicryptophyte (13%), 11 species of Nanophanerophyte (7%), 7 species of Geophytes (5%) and 2 species of Phanerophyte (1%). Anthropogenic disturbances in dry habitats have been shown to generate an increase in micro-scale species richness (Holzaphel et al., 1992). As for Therophytes, their




Fig. 2. Rainfall variation between 1997 and 2016 in Southern Tunisia (Remada).





Fig. 3. Photographs showing different rangelands types: (a) Psammophytes of *Stipagrostis pungens* on sand accumulation being grazed by camels from the typical desert vegetation site where protecting the vegetation promotes infiltration and minimizes runoff. (b) Example of an almost flat, sandy area unit, where the vegetation is dominated by *Haloxylon schmittianum*. (c) Good stand of *Anthyllis henoniana* community type in stony terrain showing the spacing of the shrubs. (d) *Retama raetam* growing on sand dunes.

presence may be explained in particular by the large number of micro-habitats available to annual plants that have rapid germination and growth, thus increasing their abundance (Gamoun et al., 2011, 2012). Chamaephytes are over-represented in arid rangelands because they are highly adapted to arid conditions (Raunkiaer, 1934; Orshan et al., 1984; Floret et al., 1990; Jauffret and Visser, 2003; Gamoun et al., 2012). The low abundance of Hemicryptophytes reduces competition for soil moisture to the benefit of Chamaephytes (Gamoun et al., 2011) (Fig. 6). On the other hand, the low presence of Geophytes mainly reflects the long period of drought in these arid rangelands (Gamoun et al., 2011, 2012).

Grazing is not the only driver in arid ecosystems. Variable and unpredictable precipitation regimes, as well as limited soil nutrient concentrations are also major factors that determine arid ecosystem features (Noy-Meir, 1985; Walker, 1987). Gamoun et al. (2012) reported changes in the plant community composition in response to grazing and soil texture. In arid rangelands, edaphic factor availability strongly determines vegetation, and soil type has been found to limit arid plant dynamics within rangelands by affecting grazing regimes practices (Gamoun, 2012). Each soil type is characterized by specific vegetation. Range production, cover, and species richness are low and highly irregular in arid zones, and



**Fig. 4.** Some plant species of desert rangelands of Tunisia (photos by Mouldi Gamoun); (a) *Rhanterium suaveolens* Desf., (b) *Echiochilon fruticosum* Desf., (c) *Plantago albicans* L., (d) *Stipa tenacissima* L., (e) *Artemisa herba-alba* Asso, (f) *Periploca angustifolia* Labill., (g) *Gymnocarpus decander* Forssk., (h) *Anthyllis henoniana* Batt., (i) *Stipagrostis ciliata* (Desf.) de Winter (j) *Ziziphus lotus* (L.) Lam., (k) *Helianthemum kahircum* Delile, (l) *Retama raetam* (Forssk.) Webb & Berthel., (m) *Haloxylon scoparium* Pomel., (n) *Haloxylon schmittianum* Pomel., (o) *Stipagrostis pungens* (Desf.) de Winter, (p) *Stipa lagascae* Roem. & Schult.

always spatially limited to sandy and gravelly soils (Gamoun, 2012). The introduction of herbivores is generally believed to reduce plant diversity. Selective grazing by livestock has been shown to negatively affect plant diversity and species composition in arid ecosystems. Heavy grazing of natural rangelands leads to loss of several highly palatable species (Louhaichi et al., 2009). However, the soil type influences the response to grazing. For example, vegetation response on sandy and gravelly soils is more diversified and more productive than on limestone and loamy soil, whereas the latter is

more adapted to grazing pressure (Gamoun et al., 2011). On the whole, grazing causes a spatial homogenization of the plant community in areas dominated by Chamaephytes. Under grazed conditions, the vegetation is dominated by perennial species which are generally present in the ungrazed rangelands. These perennial species may benefit from an adaptation caused by drought and grazing (Fensham et al., 2010). The absence of perennial species such as *Echiochilon fruticosum* Desf., *Stipa lagascae* Roem. & Schult., and *Stipagrostis ciliata* (Desf.) de Winter. is due to their high




Fig. 4. (continued).

palatability. Alternatively, perennial species may have decreased because of intensive grazing and possibly from a trampling effect from livestock in the sandy and gravelly soils of such rangelands (Gamoun et al., 2011). Otherwise, on ideal grazing land, there is a greater variety of plant species available for selective grazing and grazing animals are highly selective when given the opportunity. Thus, positive manipulation of the soil-forage plant-grazing animal complex should play a central role in any grazing management strategy for arid rangelands (Valentine, 2001).

Few studies have evaluated the effects of controlled grazing on plant diversity in arid areas. Gamoun (2014) reported that vegetation diversity was higher in protected areas than in heavily grazed

areas in the desert rangeland of southern Tunisia. In contrast, moderate grazing did not significantly affect species richness, diversity index, or species composition. In another study in the same area, Gamoun and Hanchi (2014) found that plant diversity increased as grazing intensity decreased. Ward et al. (2000) and Jauffret and Lavorel (2003), among others, have shown that palatability may play an important role in determining the effects of grazing on arid ecosystems. For these studies, however, the differences in plant diversity between the ungrazed and lightly grazed areas were small. The largest difference between the ungrazed and heavily-grazed sites was the disappearance of very palatable species under a heavy grazing treatment, including *Anabasis*

**Table 1**  
Family, life form, and livestock acceptability index of main desert species of Tunisia. 0: Refusal or Toxic; 1: Occasionally palatable; 2: Few palatable; 3: Palatable; 4: Very palatable; 5: Extremely palatable.

Species	Family	Life form	Acceptability index
<i>Anthyllis henoniana</i> Batt.	Fabaceae	Chamaephyte	4
<i>Ajuga iva</i> (L.) Schreb.	Lamiaceae	Therophyte	3
<i>Allium roseum</i> L.	Alliaceae	Geophyte	2
<i>Anabasis oropedium</i> Maire.	Chenopodiaceae	Chamaephyte	5
<i>Anacyclus clavatus</i> (Desf.) Pers.	Asteraceae	Therophyte	4
<i>Anacyclus monanthos</i> ssp <i>cyrtolopidioides</i> (Pomel)	Asteraceae	Therophyte	4
<i>Anarrhinum fruticosum</i> Desf. subsp. <i>Brevifolium</i>	Scrophulariaceae	Chamaephyte	1
<i>Argyrolobium uniflorum</i> (Deene.) Jaub. & Spach.	Fabaceae	Chamaephyte	5
<i>Arnebia decumbens</i> (Vent.) Coss. & Kralik	Boraginaceae	Therophyte	2
<i>Artemisia campestris</i> L.	Asteraceae	Chamaephyte	2
<i>Artemisia herba-alba</i> Asso	Asteraceae	Chamaephyte	2
<i>Asphodelus refractus</i> Boiss.	Asphodelaceae	Therophyte	1
<i>Asphodelus tenuifolius</i> Cav.	Asphodelaceae	Therophyte	0
<i>Astragalus armatus</i> Willd.	Fabaceae	Chamaephyte	2
<i>Astragalus asterias</i> Steven.	Fabaceae	Therophyte	2
<i>Astragalus corrugatus</i> Bertol.	Fabaceae	Therophyte	2
<i>Atractylis cancellata</i> L.	Asteraceae	Therophyte	1
<i>Atractylis carduus</i> (Forssk.) C. Chr.	Asteraceae	Chamaephyte	0
<i>Atractylis prolifera</i> Boiss.	Asteraceae	Therophyte	1
<i>Atractylis serratuloides</i> Sieber ex Cass.	Asteraceae	Hemicryptophyte	2
<i>Atriplex halimus</i> L.	Chenopodiaceae	Nanophanerophyte	3
<i>Bassia muricata</i> (L.) Asc.	Chenopodiaceae	Therophyte	2
<i>Brassica tournefortii</i> Gouan.	Brassicaceae	Therophyte	3
<i>Calendula tripterocarpa</i> Rupr.	Asteraceae	Therophyte	1
<i>Calicotome villosa</i> (Poir.) Link.	Fabaceae	Nanophanerophyte	3
<i>Calligonum azel</i> Maire	Polygonaceae	Nanophanerophyte	3
<i>Calligonum polygonoides</i> L.	Polygonaceae	Nanophanerophyte	4
<i>Carthamus eriocephalus</i> (Boiss.) Greuter	Asteraceae	Therophyte	1
<i>Centaurea furfuracea</i> Coss. & Durieu.	Asteraceae	Therophyte	3
<i>Citrullus colocynthis</i> (L.) Schrad.	Cucurbitaceae	Hemicryptophyte	0
<i>Cleome amblyocarpa</i> Barratte & Murb.	Cleomeaceae	Hemicryptophyte	0
<i>Convolvulus supinus</i> Coss. & Kralik.	Convolvulaceae	Therophyte	2
<i>Cuscuta epithimum</i> (L.) L.	Cuscutaceae	Therophyte	0
<i>Cutandia dichotoma</i> (Forssk.) Batt. & Trab.	Poaceae	Therophyte	4
<i>Cynara cardunculus</i> L. subsp. <i>Cardunculus</i>	Asteraceae	Hemicryptophyte	0
<i>Cynodon dactylon</i> (L.) Pers.	Poaceae	Geophyte	5
<i>Cynomorium coccineum</i> L.	Cynomoriaceae	Geophyte	0
<i>Daucus sahariensis</i> Murb.	Apiaceae	Therophyte	3
<i>Deverra denudata</i> (Viv.) R. Pfisterer & Podlech.	Apiaceae	Chamaephyte	2
<i>Deverra tortuosa</i> (Desf.) DC.	Apiaceae	Chamaephyte	2
<i>Dipcadi serotinum</i> (L.) Medik.	Hyacinthaceae	Geophyte	0
<i>Diplotaxis harra</i> (Forssk.) Boiss.	Brassicaceae	Therophyte	2
<i>Diplotaxis simplex</i> Spreng.	Brassicaceae	Therophyte	3
<i>Echinops spinosus</i> L.	Asteraceae	Therophyte	0
<i>Echiochilon fruticosum</i> Desf.	Boraginaceae	Chamaephyte	5
<i>Enarthrocarpus clavatus</i> Godr.	Brassicaceae	Therophyte	2
<i>Ephedra altissima</i> Desf.	Ephedraceae	Nanophanerophyte	2
<i>Erodium crassifolium</i> L'Hér.	Geraniaceae	Therophyte	3
<i>Erodium glaucophyllum</i> (L.) L'Hér.	Geraniaceae	Therophyte	1
<i>Erucaria pinnata</i> (Viv.) Täckh. & Boulos	Brassicaceae	Therophyte	3
<i>Eryngium ilicifolium</i> Lam.	Apiaceae	Therophyte	0
<i>Euphorbia retusa</i> Forssk.	Euphorbiaceae	Therophyte	0
<i>Euphorbia terracina</i> L.	Euphorbiaceae	Therophyte	0
<i>Fagonia cretica</i> L.	Zygophyllaceae	Therophyte	0
<i>Fagonia glutinosa</i> Delile.	Zygophyllaceae	Therophyte	0
<i>Farsetia aegyptia</i> Turra.	Brassicaceae	Chamaephyte	3
<i>Filago germanica</i> L.	Asteraceae	Therophyte	1
<i>Gagea fibrosa</i> (Desf.) Schult. & Schult. f.	Liliaceae	Geophyte	0
<i>Gymnarrhena micrantha</i> Desf.	Asteraceae	Therophyte	2
<i>Gymnocarpos decander</i> Forssk.	Caryophyllaceae	Chamaephyte	5
<i>Halocnemum strobilaceum</i> (Pall.) M. Bieb.	Plumbaginaceae	Hemicryptophyte	2
<i>Haloxylon schmittianum</i> Pomel.	Chenopodiaceae	Chamaephyte	1
<i>Haloxylon scoparium</i> Pomel.	Chenopodiaceae	Chamaephyte	1
<i>Haplophyllum tuberculatum</i> (Forssk.) Juss.	Rutaceae	Chamaephyte	0
<i>Helianthemum kahiricum</i> Delile.	Cistaceae	Chamaephyte	4
<i>Helianthemum sessiliflorum</i> (Desf.)	Cistaceae	Chamaephyte	5
<i>Herniaria fontanesii</i> J. Gay.	Caryophyllaceae	Chamaephyte	3
<i>Hippocrepis areolata</i> Desv.	Fabaceae	Therophyte	4
<i>Ifloga spicata</i> (Forssk.) Sch. Bip.,	Asteraceae	Therophyte	1
<i>Kickxia aegyptiaca</i> (L.) Nábelek.	Scrophulariaceae	Chamaephyte	2
<i>Koelpinia linearis</i> Pall.	Asteraceae	Therophyte	4
<i>Launaea angustifolia</i> (Desf.) Muschl.	Asteraceae	Therophyte	4
<i>Launaea capitata</i> (Spreng.) Dandy	Asteraceae	Therophyte	3

Table 1 (continued)

Species	Family	Life form	Acceptability index
<i>Launaea fragilis</i> (Asso) Pau.	Asteraceae	Therophyte	2
<i>Launaea nudicaulis</i> (Linn.) Hook. f.	Asteraceae	Therophyte	4
<i>Lavandula multifida</i> L.	Lamiaceae	Chamaephyte	0
<i>Limoniastrum guyonianum</i> Boiss.	Plumbaginaceae	Hemicryptophyte	3
<i>Limoniastrum monopetalum</i> (L.) Boiss.	Plumbaginaceae	Hemicryptophyte	3
<i>Limonium pruinatum</i> (L.) Chaz.	Plumbaginaceae	Hemicryptophyte	1
<i>Linaria laxiflora</i> Desf.	Scrophulariaceae	Therophyte	1
<i>Lobularia libyca</i> (Viv.) Meissn.	Brassicaceae	Therophyte	3
<i>Lotus halophilus</i> Boiss. & Spruner	Asteraceae	Therophyte	3
<i>Lycium shawii</i> Roem. & Schult.	Solanaceae	Nanophanerophyte	2
<i>Lygeum spartum</i> Loefl. ex L.	Poaceae	Hemicryptophyte	2
<i>Matthiola longipetala</i> (Vent.) DC.	Brassicaceae	Therophyte	2
<i>Medicago minima</i> (L.) L.	Fabaceae	Therophyte	3
<i>Moricandia arvensis</i> (L.) DC.	Brassicaceae	Chamaephyte	2
<i>Muricaria prostrata</i> (Desf.) Desv.	Brassicaceae	Therophyte	3
<i>Neurada procumbens</i> L.	Neuradaceae	Therophyte	2
<i>Nitraria retusa</i> (Forssk.) Asch.	Nitrariaceae	Nanophanerophyte	2
<i>Nolletia chrysocomoides</i> (Desf.) Cass. ex Less.	Asteraceae	Chamaephyte	2
<i>Nonea calycina</i> (Roem. & Schult.) Selvi	Boraginaceae	Therophyte	2
<i>Pallenis hierochuntica</i> (Michon) Greuter.	Asteraceae	Therophyte	2
<i>Paronychia arabica</i> (L.) DC.	Caryophyllaceae	Therophyte	1
<i>Peganum harmala</i> L.	Zygophyllaceae	Chamaephyte	0
<i>Pennisetum divisum</i> (Forssk. ex J.F. Gmel.) Henrard	Poaceae	Hemicryptophyte	3
<i>Periploca angustifolia</i> subsp. <i>angustifolia</i> (Labill.)	Asclepiadaceae	Nanophanerophyte	3
<i>Plantago albicans</i> L.	Plantaginaceae	Hemicryptophyte	5
<i>Plantago ovata</i> Forssk.	Plantaginaceae	Therophyte	3
<i>Polygonum equisetiforme</i> Sibth. et Sm.	Polygonaceae	Hemicryptophyte	4
<i>Reaumuria vermiculata</i> L.	Tamaricaceae	Chamaephyte	0
<i>Reichardia tingitana</i> (L.) Roth	Asteraceae	Therophyte	2
<i>Reseda alba</i> L. subsp. <i>Alba</i>	Resedaceae	Therophyte	1
<i>Retama raetam</i> (Forssk.) Webb & Berthel.	Fabaceae	Nanophanerophyte	3
<i>Rhanterium suaveolens</i> Desf.	Asteraceae	Chamaephyte	2
<i>Rhus tripartita</i> (Ucria) Grande	Anacardiaceae	Nanophanerophyte	2
<i>Salsola tetragona</i> Delile	Chenopodiaceae	Phanerophyte	2
<i>Salsola tetrandra</i> Forssk.	Chenopodiaceae	Phanerophyte	2
<i>Salsola vermiculata</i> L.	Chenopodiaceae	Chamaephyte	3
<i>Salvia aegyptiaca</i> L.	Lamiaceae	Chamaephyte	3
<i>Salvia verbenaca</i> L.	Lamiaceae	Chamaephyte	3
<i>Savignya parviflora</i> (Delile) Webb.	Brassicaceae	Therophyte	3
<i>Scabiosa arenaria</i> Forssk.	Dipsacaceae	Therophyte	2
<i>Schismus barbatus</i> (L.) Thell.	Poaceae	Therophyte	4
<i>Scorzonera undulata</i> Vahl.	Asteraceae	Geophyte	4
<i>Senecio gallicus</i> L.	Asteraceae	Therophyte	1
<i>Stipa capensis</i> Thunb.	Poaceae	Therophyte	2
<i>Stipa lagascae</i> Roem. & Schult.	Poaceae	Hemicryptophyte	4
<i>Stipa parviflora</i> Desf.	Poaceae	Hemicryptophyte	4
<i>Stipa tenacissima</i> L.	Poaceae	Hemicryptophyte	3
<i>Stipagrostis ciliata</i> (Desf.) de Winter.	Poaceae	Hemicryptophyte	4
<i>Stipagrostis obtusa</i> (Delile) Nees.	Poaceae	Hemicryptophyte	4
<i>Stipagrostis plumosa</i> (L.) Munro.	Poaceae	Hemicryptophyte	4
<i>Stipagrostis pungens</i> (Desf.) de Winter.	Poaceae	Hemicryptophyte	3
<i>Suaeda vermiculata</i> Forssk. ex J.F. Gmel.	Chenopodiaceae	Chamaephyte	3
<i>Tamarix gallica</i> L.	Tamaricaceae	Nanophanerophyte	2
<i>Teucrium alopecurus</i> De Noé	Primulaceae	Chamaephyte	1
<i>Teucrium polium</i> L.	Lamiaceae	Chamaephyte	2
<i>Thesium humile</i> Vahl, Symb.	Santalaceae	Therophyte	1
<i>Thymelaea hirsuta</i> (L.) Endl.	Thymelaeaceae	Chamaephyte	0
<i>Thymelaea microphylla</i> Coss. & Durieu.	Thymelaeaceae	Chamaephyte	2
<i>Traganum nudatum</i> Delile.	Chenopodiaceae	Chamaephyte	3
<i>Tribulus terrestris</i> L.	Zygophyllaceae	Therophyte	1
<i>Ziziphus lotus</i> (L.) Lam.	Rhamnaceae	Nanophanerophyte	2
<i>Zygophyllum album</i> L. f.	Zygophyllaceae	Chamaephyte	0

*oropediorum* (Maire), *Cutandia dichotoma* (Forssk.) Batt. & Trab., *E. fruticosum* (Desf.), *Helianthemum kahiricum* (Delile), *Helianthemum sessiliflorum* (Desf.), *Hippocrepis areolata* Desv., *Launaea nudicaulis* (Linn.) Hook. f., *Launaea angustifolia* (Desf.) Muschl., *Koelpinia linearis* Pall., *Polygonum equisetiforme* S. & Sm., *Scorzonera undulata* Vahl. and *S. lagascae* Roem. & Schult., (Gamoun, 2014a, 2014b). Jauffret and Lavorel (2003) note that long-spine species such as *Astragalus armatus* Willd., and unpalatable, highly fibrous species such as *Thymelaea hirsuta* (L.) Endl., are dominant in arid Tunisian

rangelands, and suggest this is a consequence of long grazing history. Similarly, unpalatable shrubs such as *Haloxylon scoparium* Pomel., *T. hirsuta* (L.) Endl., and *Anabasis articulata* (Forssk.) Moq., are often dominant in heavily grazed arid regions of the Middle East (Ward, 2004).

Deserts are biologically stressful environments and plants have acquired two principal strategies to cope with the harsh arid conditions and herbivory: avoidance and tolerance (Laity, 2008). These strategies allow plants to cope with heat and drought to ensure that




Fig. 5. Family distribution of plant species of desert rangelands of southern Tunisia (%).




Fig. 6. Life form distribution of plant species of desert rangelands of southern Tunisia (%). The: Therophyte; Cha: Chamaephyte; Hem: Hemicryptophyte; Npha: Nanophanerophyte; Geo: Geophyte.

neither internal temperatures nor tissue dehydration reach deadly levels. The majority of the flora in deserts are evaders, surviving stressful periods by living permanently or temporarily in cooler and/or moister microhabitats (Laity, 2008).

### 3. Conclusion

Many plants live in deserts, but it is only xerophytes that can withstand and live under long-term dry conditions. Asteraceae, Poaceae, Brassicaceae, Chenopodiaceae, and Fabaceae represent approximately 50% of Tunisian desert flora. This diversity provides many benefits that can meet the demands of both the local people for medicinal plants and fruits, as well as livestock requirements through providing forage. Mismanagement and climate change have gradually destroyed previously productive ecosystems. To reverse the degradation and desertification of these natural resources, restoration and improved management is essential.

### Acknowledgements

This work was supported by the International Center for Agricultural Research in the Dry Areas (ICARDA), the Arid Regions Institute (IRA – Medenine, Tunisia) and the CGIAR Research Program on Livestock (CRP Livestock).

### Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.pld.2018.06.004>.

### References

- Aronson, J., Floret, C., Le Floch, E., Oualle, C., Pontanier, R., 1993. Restoration and rehabilitation of degraded ecosystems in arid and semi-arid lands. II. Case studies in Southern Tunisia, Central Chile and Northern Cameroon. *Restor. Ecol.* 1, 168–187.
- Bainbridge, D.A., 2007. *Guide for Desert and Dryland Restoration: New Hope for Arid Lands*. Island Press, Washington, DC, USA, p. 416.
- Bendali, F., Floret, C., Le Floch, E., Pontanier, R., 1990. The dynamics of vegetation and sand mobility in arid regions of Tunisia. *J. Arid Environ.* 18, 21–32.
- Carney, D., Farrington, J., 1998. *Natural Resource Management and Institutional Change*. Routledge, London, p. 136.
- Casper, J.K., 2009. *Changing Ecosystems: Effects of Global Warming*. Facts On File, Inc, p. 273.
- Chambers, R., Saxena, N.C., Shah, T., 1991. *To the Hand of the Poor: Water and Trees*. Intermediate Technology Publication, London, p. 273p.
- Darkoh, M.B.K., 1998. The nature, causes and consequences of Desertification in the drylands of Africa. *Land Degrad. Dev.* 9, 1–20.
- Fensham, R.J., Fairfax, R.J., Dwyer, J.M., 2010. Vegetation responses to the first 20 years of cattle grazing in an Australian desert. *Ecology* 91, 681–692.
- Ferchichi, A., 2000. Rangelands biodiversity in presaharian Tunisia. *Cah. Options Mediterr.* 45, 69–72.
- Floret, C., 1981. The effects of protection on steppic vegetation in the Mediterranean arid zone of Southern Tunisia. *Vegetatio* 46, 117–129.
- Floret, C., Hadjei, M.S., 1977. An attempt to combat desertification in Tunisia. *Ambio* 6, 366–368.
- Floret, C., Le Floch, E., Pontanier, R., Romane, F., 1978. *Modèle écologique régional en vue de la planification et de l'aménagement agro-pastoral des régions arides*. Application à la région de Zougrata. Inst. Reg. Arides-Medenine, Dir. Ress. Eau et Sols Tunis, CEPE/CNRS Montpellier & ORSTOM, Paris, FR, p. 73.
- Floret, C., Pontanier, R., 1982. L'aridité en Tunisie présaharienne : Climat – sol – végétation et aménagement, p. 544. *Trav. et Doc. ORSTOM*, n° 150, Paris.
- Floret, C., Galan, M.J., Le floch, E., Orshan, G., Romane, F., 1990. Growth forms and phenomorphology traits along an environmental gradient: tools for studying vegetation. *J. Veg. Sci.* 1, 71–80.
- Gamoun, M., 2012. *The Impact of Rest from Grazing on Vegetation Dynamics: Application to Sustainable Management of Saharan Rangelands in Southern Tunisia*. PhD. thesis. University of Tunis El Manar, Tunisia, p. 202.
- Gamoun, M., 2013. *Management and resilience of Saharan rangelands: south Tunisia*. *Fourrages* 216, 321–328.
- Gamoun, M., 2014. *Grazing intensity effects on the vegetation in desert rangelands of Southern Tunisia*. *J. Arid Land* 6, 324–333.
- Gamoun, M., 2016. *Rain use efficiency, primary production and rainfall relationships in desert rangelands of Tunisia*. *Land Degrad. Dev.* 27, 738–747.
- Gamoun, M., Chaieb, M., Ouled Belgacem, A., 2010. Evolution of ecological characteristics along a gradient of soil degradation in the South Tunisian rangelands. *Ecol. Mediterr.* 36, 5–16.
- Gamoun, M., Tarhouni, M., Ouled Belgacem, A., Hanchi, B., Neffati, M., 2011. Response of different arid rangelands to protection and drought. *Arid Land Res. Manag.* 25, 372–378.



- Gamoun, M., Ouled Belgacem, A., Hanchi, B., Neffati, M., Gillet, F., 2012. Impact of grazing on the floristic diversity of arid rangelands in South Tunisia. *Revue d'Écologie (Terre Vie)* 67, 271–282.
- Gamoun, M., Hanchi, B., 2014. Natural vegetation cover dynamic under grazing-rotation managements in desert rangelands of Tunisia. *Arid Ecosyst.* 4, 277–284.
- Gamoun, M., Essifi, B., Dickens, C., Hanchi, B., 2016. Interactive effects of grazing and drought on desert rangelands of Tunisia. *Biologija* 62, 105–115.
- Gondard, H., Jauffret, S., Aronson, J., Lavorel, S., 2003. Plant functional types: a promising tool for management and restoration of degraded lands. *Appl. Veg. Sci.* 6, 223–234.
- Hallsworth, E.G., 1987. *Anatomy, Physiology and Psychology of Erosion*. Wiley, New York, p. 188p.
- Harris, R.B., 2010. Rangeland degradation on the Qinghai-Tibetan plateau: a review of the evidence of its magnitude and causes. *J. Arid Environ.* 74, 1–12.
- Holzaphel, C., Schmidt, W., Shmida, A., 1992. Effects of human-caused distribution on the flora along a Mediterranean-desert gradient. *Flora* 186, 261–270.
- Hudak, A.T., 1999. Rangeland mismanagement in South Africa: failure to apply ecological knowledge. *Hum. Ecol.* 27, 55–78.
- Ibáñez, J., Martínez, J., Schnabel, S., 2007. Desertification due to overgrazing in a dynamic commercial livestock-grass-soil system. *Ecol. Model.* 205, 277–288.
- Jauffret, S., Lavorel, S., 2003. Are plant functional types relevant to describe degradation in arid, southern Tunisian steppes? *J. Veg. Sci.* 14, 399–408.
- Jauffret, S., Visser, M., 2003. Assigning life history traits to plant species to better qualify arid land degradation in Presaharian Tunisia. *J. Arid Environ.* 55, 1–28.
- Laity, J., 2008. *Deserts and Desert Environments*. A John Wiley & Sons, Inc., Publication, p. 357.
- Le Floc'h, E., Boulos, L., Vêla, E., 2010. *Flore de Tunisie, Catalogue synonymique commenté*. Tunis, p. 500.
- Le Houérou, H.N., 1969. Vegetation of the Tunisian steppe (with the references to Morocco, Algeria and Libya). *Ann. Inst. Natl. Rech. Agron. Tunis. (Tunisia)* 42 (5), 622.
- Le Houérou, H.N., 1996. Climate change, drought and desertification. *J. Arid Environ.* 34, 133–185.
- Le Houérou, H.N., 2009. *Bioclimatology and Biogeography of Africa*. Springer-Verlag Berlin Heidelberg, p. 241.
- Louhaichi, M., Salkini, A.K., Petersen, S.L., 2009. Effect of small ruminant grazing on the plant community characteristics of semi-arid Mediterranean ecosystems. *Int. J. Agric. Biol.* 11, 681–689.
- Nefzaoui, A., Ben Salem, H., El Mourid, M., 2011. Innovations in small ruminants feeding systems in arid Mediterranean areas. In: Bouche, R., Derkimba, A., Casabianca, F. (Eds.), *New Trends for Innovation in the Mediterranean Animal Production*. Wageningen Academic Publishers, pp. 99–116.
- Nicholson, E.S., 2011. *Dryland Climatology*. Cambridge University Press, New York, p. 528p.
- Noy-Meir, I., 1985. Desert ecosystem structure and function. In: Evenari, M., Noy-Meir, I., Goodall, D.W. (Eds.), *Hot Deserts and Arid Shrublands*. Elsevier, Amsterdam, p. 458.
- Orshan, G., Montenegro, G., Avila, G., Aljaro, M.E., Walckowiak, A., Mujica, A.M., 1984. Plant growth forms of Chilean matorral. A monocharacter growth form analysis along an altitudinal transect from sea level to 2000 m a.s.l. *Bulletin de la Société Botanique de France. Actualités Botaniques* 131, 411–425.
- Ouled Belgacem, A., Ben Salem, H., Bouaicha, A., El Mourid, M., 2008. Communal rangeland rest in arid area, a tool for facing animal feed costs and drought mitigation: the case of Chenini Community, Southern Tunisia. *J. Biol. Sci.* 8, 822–825.
- Ouled Belgacem, A., Tarhouni, M., Louhaichi, M., 2011. Effect of protection on plant community dynamics in the Mediterranean arid zone of southern Tunisia: a case study from Bou Hedma National Park. *Land Degrad. Dev.* 24, 57–62.
- Ouled Belgacem, A., Louhaichi, M., 2013. The vulnerability of native rangeland plant species to global climate change in the West Asia and North African regions. *Climatic Change* 119, 451–463.
- Peters, D.P.C., Bestelmeyer, B.T., Havstad, K.M., Rango, A., Archer, S.R., Comrie, A.C., Gimblett, H.R., López-Hoffman, L., Sala, O.E., Vivoni, E.R., Brooks, M.L., Brown, J., Monger, H.C., Goldstein, J.H., Okin, G.S., Tweedie, C.E., 2013. Desertification of rangelands. *Clim. Vulnerability* 4, 230–259.
- Raunkiaer, C., 1934. *The Life Form of Plants and Statistical Plant Geography*. Collected Papers. Clarendon Press, Oxford, p. 632p.
- Reynolds, J.F., 2013. Desertification. *Encycl. Biodivers.* 2, 479–494.
- Shachak, M., Gosz, J.R., Pickett, S.T.A., Perevolotsky, A., 2005. Biodiversity in Drylands: toward a Unified Framework. Oxford University Press, Inc., p. p366
- Tarhouni, M., Ben Salem, F., Ouled Belgacem, A., Neffati, M., 2014. Impact of livestock exclusion on Sidi Toui National park vegetation communities, Tunisia. *Int. J. Biodivers.* <https://doi.org/10.1155/2014/620405>.
- Tarhouni, M., Ben Hmida, W., Neffati, M., 2015. Long-term changes in plant life forms as a consequence of grazing exclusion under arid climatic conditions. *Land Degrad. Dev.* 28, 1199–1211.
- Thomas, D.S.G., 2011. *Arid Zone Geomorphology: Process, Form and Change in Drylands*, third ed. A John Wiley & Sons, Ltd., Publication, p. 648.
- Vallentine, J.F., 2001. *Grazing. Management*, second ed. Academic Press, London.
- Verner, D., 2013. Tunisia in a Changing Climate: Assessment and Actions for Increased Resilience and Development. The world bank, Washington, D.C, p. 184.
- Verstraete, M.M., Schwartz, S.A., 1991. Desertification and global change. In: Henderson-Sellers, A., Pitman, A.J. (Eds.), *Vegetation and Climate Interactions in Semi-arid Regions*. Kluwer Academic Publishers, Belgium, pp. 3–13.
- Walker, B.H., 1987. A general model of savanna structure and function. In: Walker, B.H. (Ed.), *Determinants of Tropical Savannas*. IUBS Monograph Series No. 3. IRL Press, Oxford, pp. 1–12.
- Wang, X., Chen, F.H., Dong, Z., Xia, D., 2005. Evolution of the southern Mu Us desert in north China over the past 50 years: an analysis using proxies of human activity and climate parameters. *Land Degrad. Dev.* 16, 351–366.
- Ward, D., Saltz, D., Olsvig-Whittaker, L., 2000. Distinguishing signal from noise: long-term studies of vegetation in Makhtesh Ramon erosion cirque, Negev desert, Israel. *Plant Ecol.* 150, 27–36.
- Ward, D., 2004. The effects of grazing on plant biodiversity in arid ecosystems. In: Shachak, M., Pickett, S.T.A., Gosz, J.R., Perevolotsky, A. (Eds.), *Biodiversity in Drylands: towards a Unified Framework*. Oxford University Press, Oxford, pp. 233–249.
- Ward, D., 2016. *The Biology of Deserts*. Oxford University Press, p. 395.
- White, R., Murray, S., Rohweder, M., 2000. *Pilot Analysis of Global Ecosystems Grassland Ecosystems*. World Resources Institute, Washington D.C.
- Zhao, H., Li, S.G., Zhang, T.H., Okhuro, T., Zhou, R.L., 1994. Sheep gain and species diversity in sandy grassland, Inner Mongolia. *Rangel. Ecol. Manag.* 57 (2), 187–190.