



Crystal structure of 5-(1,3-dithian-2-yl)-2H-1,3-benzodioxole

Julio Zukerman-Schpector,^{a*} Ignez Caracelli,^b Hélio A. Stefani,^c Olga Gozhina^c and Edward R. T. Tiekink^d

^aDepartamento de Química, Universidade Federal de São Carlos, 13565-905 São Carlos, SP, Brazil, ^bDepartamento de Física, Universidade Federal de São Carlos, 13565-905 São Carlos, SP, Brazil, ^cDepartamento de Farmácia, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, 05508-900 São Paulo-SP, Brazil, and ^dDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia. *Correspondence e-mail: julio@power.ufscar.br

Received 4 February 2015; accepted 5 February 2015

Edited by W. T. A. Harrison, University of Aberdeen, Scotland


In the title compound, C₁₁H₁₂O₂S₂, two independent but virtually superimposable molecules, *A* and *B*, comprise the asymmetric unit. In each molecule, the 1,3-dithiane ring has a chair conformation with the 1,4-disposed C atoms being above and below the plane through the remaining four atoms. The substituted benzene ring occupies an equatorial position in each case and forms dihedral angles of 85.62 (9) (molecule *A*) and 85.69 (8)° (molecule *B*) with the least-squares plane through the 1,3-dithiane ring. The difference between the molecules rests in the conformation of the five-membered 1,3-dioxole ring which is an envelope in molecule *A* (the methylene C atom is the flap) and almost planar in molecule *B* (r.m.s. deviation = 0.046 Å). In the crystal, molecules of *A* self-associate into supramolecular zigzag chains (generated by glide symmetry along the *c* axis) *via* methylene C—H... π interactions. Molecules of *B* form similar chains. The chains pack with no specific directional intermolecular interactions between them.

Keywords: crystal structure; 1,3-dithiane; conformation; 1,3-benzodioxole; C—H... π interactions.

CCDC reference: 1047484

1. Related literature

The title compound has been prepared previously, see: Ballesteros *et al.* (2005). For the structure of a related compound containing the same molecular skeleton as in the title compound, *i.e.* (19*R*,21*R*,25*S*)-(–)-2-(2-menthyloxy-carbonyl-3,4-methylenedioxyphenyl)1,3-dithiane, see: Ratajczak-Sitarz *et al.* (1996).



2. Experimental

2.1. Crystal data

C ₁₁ H ₁₂ O ₂ S ₂	<i>V</i> = 2272.05 (9) Å ³
<i>M_r</i> = 240.33	<i>Z</i> = 8
Monoclinic, <i>P</i> 2 ₁ / <i>c</i>	Mo <i>K</i> α radiation
<i>a</i> = 11.4765 (3) Å	μ = 0.45 mm ^{−1}
<i>b</i> = 17.5504 (4) Å	<i>T</i> = 296 K
<i>c</i> = 11.6397 (2) Å	0.59 × 0.40 × 0.26 mm
β = 104.275 (1)°	

2.2. Data collection

Bruker APEXII CCD diffractometer	14839 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	4164 independent reflections
<i>T</i> _{min} = 0.702, <i>T</i> _{max} = 0.745	3759 reflections with <i>I</i> > 2σ(<i>I</i>)
	<i>R</i> _{int} = 0.020

2.3. Refinement

<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)] = 0.035	271 parameters
<i>wR</i> (<i>F</i> ²) = 0.092	H-atom parameters constrained
<i>S</i> = 1.03	$\Delta\rho_{\text{max}}$ = 0.29 e Å ^{−3}
4164 reflections	$\Delta\rho_{\text{min}}$ = −0.44 e Å ^{−3}

Table 1

Hydrogen-bond geometry (Å, °).

*Cg*1 and *Cg*2 are the centroids of the C5–C11 and C16–C21 rings, respectively.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C4—H4b... <i>Cg</i> 1 ⁱ	0.97	2.77	3.731 (2)	170
C13—H13a... <i>Cg</i> 2 ⁱⁱ	0.97	2.54	3.4841 (19)	165

Symmetry codes: (i) *x*, −*y* + $\frac{1}{2}$, *z* − $\frac{1}{2}$; (ii) *x*, −*y* − $\frac{1}{2}$, *z* − $\frac{3}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SIR2014* (Burla *et al.*, 2015); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012), *QMOL* (Gans & Shalloway, 2001) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *MarvinSketch* (ChemAxon, 2010) and *publCIF* (Westrip, 2010).

Acknowledgements

We thank Professor Regina H. A. Santos from IQSC-USP for the X-ray data collection. The Brazilian agencies CNPq

(305626/2013-2 to JZS, 306121/2013-2 to IC and 308320/2010-7 to HAS), FAPESP (2012/00424-2 and 2013/21925-2) and CAPES are acknowledged for financial support.

Supporting information for this paper is available from the IUCr electronic archives (Reference: HB7361).

References

- Ballesteros, L., Noguez, O., Arroyo, G., Velasco, B., Delgado, F. & Miranda, R. (2005). *J. Mex. Chem. Soc.* **49**, 302–306.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burla, M. C., Caliendo, R., Carrozzini, B., Cascarano, G. L., Cuocci, C., Giacovazzo, C., Mallamo, M., Mazzzone, A. & Polidori, G. (2015). *J. Appl. Cryst.* **48**, 306–309.
- ChemAxon (2010). *Marvinsketch*. <http://www.chemaxon.com>.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Gans, J. & Shalloway, D. (2001). *J. Mol. Graph. Model.* **19**, 557–559.
- Ratajczak-Sitarz, M., Narusiewicz, M., Kosturkiewicz, Z., Jozkowiak, J. & Rozwadowska, M. D. (1996). *Pol. J. Chem.* **70**, 310–319.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2015). E71, o167–o168 [doi:10.1107/S2056989015002455]

Crystal structure of 5-(1,3-dithian-2-yl)-2*H*-1,3-benzodioxole


Julio Zukerman-Schpector, Ignez Caracelli, Hélio A. Stefani, Olga Gozhina and Edward R. T. Tiekink

S1. Experimental

A solution of the corresponding 2*H*-1,3-benzodioxole-5-carbaldehyde (0.037 mol, 1 equiv.) in chloroform (20 ml) was combined with an equimolar amount of propane-1,3-dithiol (3.7 ml, 0.037 mol) at room temperature. The solution was stirred for 1 h at this temperature, then cooled to -20 °C after which BF₃ etherate (0.46 ml, 0.0037 mol, 0.1 equiv.) was added drop-wise. The reaction solution was allowed to warm to room temperature and stirred overnight. After this time, the solution was washed three times each with water, 10% aqueous KOH, then water followed by drying over MgSO₄. Evaporation of the solvent furnishes a pure product as colourless crystals in 97% yield. To obtain crystals suitable for X-ray analysis, the product was crystallized from CH₃OH. The spectroscopic data matched those reported in the literature (Ballesteros *et al.*, 2005).

S2. Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H = 0.93–0.98 Å) and were included in the refinement in the riding model approximation, with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$.

**Figure 1**

The molecular structures of the two independent molecules in title compound showing the atom-labelling scheme and displacement ellipsoids at the 35% probability level.






Figure 2

Superimposition of the two independent molecules. Molecule *A* is shown in red and *B* in blue. The molecules have been superimposed such that the benzene rings are overlapped.

**Figure 3**

A view of the zigzag supramolecular chain comprising molecules of *A* along the *c* axis (glide symmetry) mediated by C—H... π interactions are shown as purple dashed lines.

**Figure 4**

A view in projection down the *c* axis of the unit-cell contents. The C—H... π interactions are shown as purple dashed lines.

5-(1,3-Dithian-2-yl)-2*H*-1,3-benzodioxole

Crystal data

$C_{11}H_{12}O_2S_2$
 $M_r = 240.33$

Monoclinic, $P2_1/c$
 $a = 11.4765$ (3) Å

$b = 17.5504$ (4) Å
 $c = 11.6397$ (2) Å
 $\beta = 104.275$ (1)°
 $V = 2272.05$ (9) Å³
 $Z = 8$
 $F(000) = 1008$
 $D_x = 1.405$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 9068 reflections
 $\theta = 2.9\text{--}25.4^\circ$
 $\mu = 0.45$ mm⁻¹
 $T = 296$ K
 Prism, colourless
 $0.59 \times 0.40 \times 0.26$ mm

Data collection

Bruker APEXII CCD
 diffractometer
 φ and ω scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.702$, $T_{\max} = 0.745$
 14839 measured reflections

4164 independent reflections
 3759 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$
 $\theta_{\max} = 25.4^\circ$, $\theta_{\min} = 1.8^\circ$
 $h = -13 \rightarrow 13$
 $k = -21 \rightarrow 19$
 $l = -14 \rightarrow 8$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.092$
 $S = 1.03$
 4164 reflections
 271 parameters
 0 restraints

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0459P)^2 + 0.9363P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.29$ e Å⁻³
 $\Delta\rho_{\min} = -0.44$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.36505 (5)	0.56934 (3)	0.82081 (4)	0.05661 (16)
S2	0.50542 (5)	0.71160 (3)	0.79700 (4)	0.05188 (15)
O1	0.23644 (16)	0.62085 (13)	0.26070 (13)	0.0865 (6)
O2	0.40467 (15)	0.55784 (9)	0.36797 (12)	0.0660 (4)
C1	0.36233 (15)	0.66240 (10)	0.75248 (15)	0.0432 (4)
H1	0.3013	0.6935	0.7762	0.052*
C2	0.4034 (2)	0.59674 (13)	0.97523 (16)	0.0553 (5)
H2A	0.3401	0.6291	0.9899	0.066*
H2B	0.4066	0.5513	1.0234	0.066*
C3	0.52212 (18)	0.63848 (12)	1.01417 (16)	0.0524 (5)
H3A	0.5408	0.6455	1.0994	0.063*
H3B	0.5849	0.6071	0.9962	0.063*
C4	0.5228 (2)	0.71522 (12)	0.95595 (17)	0.0567 (5)
H4A	0.5980	0.7406	0.9919	0.068*
H4B	0.4583	0.7459	0.9720	0.068*
C5	0.32654 (15)	0.65344 (11)	0.62010 (15)	0.0428 (4)

C6	0.22667 (18)	0.69163 (15)	0.55466 (19)	0.0640 (6)
H6	0.1835	0.7230	0.5937	0.077*
C7	0.1889 (2)	0.68440 (18)	0.4319 (2)	0.0779 (7)
H7	0.1215	0.7100	0.3882	0.094*
C8	0.25501 (19)	0.63842 (14)	0.37937 (16)	0.0602 (6)
C9	0.35527 (17)	0.60054 (11)	0.44322 (15)	0.0464 (4)
C10	0.39414 (16)	0.60659 (10)	0.56338 (15)	0.0426 (4)
H10	0.4621	0.5809	0.6056	0.051*
C11	0.3389 (3)	0.57881 (16)	0.25229 (18)	0.0757 (7)
H11A	0.3891	0.6094	0.2143	0.091*
H11B	0.3145	0.5335	0.2047	0.091*
S3	0.09983 (5)	0.28175 (2)	0.03890 (4)	0.04927 (14)
S4	0.16401 (4)	0.44631 (2)	0.00810 (4)	0.04651 (14)
O3	0.33099 (13)	0.39219 (10)	0.48171 (12)	0.0623 (4)
O4	0.17354 (14)	0.42693 (9)	0.56131 (12)	0.0626 (4)
C12	0.07298 (15)	0.38075 (9)	0.06991 (15)	0.0393 (4)
H12	−0.0118	0.3922	0.0342	0.047*
C13	0.05738 (19)	0.28499 (10)	−0.12132 (16)	0.0496 (4)
H13A	0.0628	0.2340	−0.1517	0.060*
H13B	−0.0258	0.3011	−0.1470	0.060*
C14	0.13363 (18)	0.33772 (11)	−0.17442 (16)	0.0484 (4)
H14A	0.1138	0.3306	−0.2597	0.058*
H14B	0.2176	0.3245	−0.1438	0.058*
C15	0.11586 (19)	0.42070 (10)	−0.14730 (16)	0.0504 (5)
H15A	0.0312	0.4330	−0.1756	0.060*
H15B	0.1596	0.4518	−0.1914	0.060*
C16	0.09597 (16)	0.39253 (9)	0.20158 (15)	0.0401 (4)
C17	0.00141 (17)	0.41118 (10)	0.25062 (18)	0.0460 (4)
H17	−0.0757	0.4149	0.2014	0.055*
C18	0.01841 (18)	0.42465 (11)	0.37213 (18)	0.0521 (5)
H18	−0.0452	0.4381	0.4044	0.063*
C19	0.13232 (18)	0.41716 (10)	0.44076 (16)	0.0451 (4)
C20	0.22716 (16)	0.39691 (10)	0.39325 (16)	0.0435 (4)
C21	0.21260 (16)	0.38462 (11)	0.27471 (15)	0.0441 (4)
H21	0.2772	0.3716	0.2437	0.053*
C22	0.2965 (2)	0.40546 (14)	0.58912 (18)	0.0616 (5)
H22A	0.3083	0.3596	0.6371	0.074*
H22B	0.3455	0.4457	0.6338	0.074*

Atomic displacement parameters ($\text{\AA}^2$)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0779 (4)	0.0577 (3)	0.0369 (3)	−0.0258 (3)	0.0194 (2)	−0.0058 (2)
S2	0.0583 (3)	0.0509 (3)	0.0473 (3)	−0.0141 (2)	0.0146 (2)	−0.0003 (2)
O1	0.0774 (11)	0.1433 (18)	0.0318 (7)	−0.0069 (11)	0.0003 (7)	−0.0008 (9)
O2	0.0837 (11)	0.0812 (11)	0.0364 (7)	0.0008 (8)	0.0209 (7)	−0.0108 (7)
C1	0.0412 (9)	0.0521 (10)	0.0391 (9)	0.0008 (8)	0.0155 (7)	−0.0036 (8)
C2	0.0696 (13)	0.0651 (13)	0.0352 (9)	−0.0103 (10)	0.0208 (9)	−0.0019 (9)

C3	0.0580 (12)	0.0621 (12)	0.0359 (9)	−0.0002 (9)	0.0092 (8)	−0.0045 (8)
C4	0.0662 (13)	0.0545 (12)	0.0459 (11)	−0.0100 (10)	0.0068 (9)	−0.0122 (9)
C5	0.0380 (9)	0.0554 (11)	0.0370 (9)	−0.0024 (8)	0.0130 (7)	0.0002 (8)
C6	0.0459 (11)	0.0932 (17)	0.0539 (12)	0.0180 (11)	0.0142 (9)	0.0020 (11)
C7	0.0512 (12)	0.124 (2)	0.0516 (12)	0.0217 (14)	−0.0001 (10)	0.0118 (13)
C8	0.0507 (11)	0.0919 (16)	0.0343 (10)	−0.0106 (11)	0.0032 (8)	0.0042 (10)
C9	0.0516 (10)	0.0543 (11)	0.0358 (9)	−0.0091 (8)	0.0155 (8)	−0.0023 (8)
C10	0.0439 (9)	0.0509 (10)	0.0335 (8)	0.0003 (8)	0.0104 (7)	0.0015 (7)
C11	0.0981 (19)	0.0934 (18)	0.0356 (11)	−0.0223 (15)	0.0166 (11)	−0.0127 (11)
S3	0.0712 (3)	0.0321 (2)	0.0429 (3)	−0.0009 (2)	0.0108 (2)	0.00570 (17)
S4	0.0588 (3)	0.0368 (2)	0.0442 (3)	−0.01144 (19)	0.0132 (2)	−0.00229 (18)
O3	0.0544 (8)	0.0930 (11)	0.0390 (7)	0.0080 (7)	0.0104 (6)	−0.0048 (7)
O4	0.0798 (10)	0.0687 (9)	0.0452 (8)	0.0103 (8)	0.0265 (7)	−0.0080 (7)
C12	0.0401 (9)	0.0347 (8)	0.0426 (9)	0.0011 (7)	0.0091 (7)	0.0036 (7)
C13	0.0643 (12)	0.0356 (9)	0.0443 (10)	−0.0069 (8)	0.0046 (9)	−0.0028 (7)
C14	0.0598 (11)	0.0477 (10)	0.0362 (9)	−0.0050 (8)	0.0092 (8)	−0.0039 (8)
C15	0.0677 (12)	0.0404 (10)	0.0420 (10)	−0.0090 (9)	0.0111 (9)	0.0061 (8)
C16	0.0447 (9)	0.0330 (8)	0.0448 (9)	0.0036 (7)	0.0155 (8)	0.0024 (7)
C17	0.0434 (9)	0.0388 (9)	0.0578 (11)	0.0092 (7)	0.0161 (8)	0.0032 (8)
C18	0.0550 (11)	0.0453 (10)	0.0654 (13)	0.0117 (8)	0.0325 (10)	−0.0021 (9)
C19	0.0623 (12)	0.0348 (9)	0.0440 (10)	0.0037 (8)	0.0240 (9)	−0.0034 (7)
C20	0.0460 (10)	0.0436 (10)	0.0424 (9)	0.0032 (8)	0.0136 (8)	0.0000 (7)
C21	0.0411 (9)	0.0535 (11)	0.0412 (9)	0.0062 (8)	0.0169 (8)	−0.0004 (8)
C22	0.0757 (15)	0.0689 (14)	0.0411 (10)	−0.0015 (11)	0.0162 (10)	−0.0090 (10)

Geometric parameters (Å, °)

S1—C2	1.8071 (18)	S3—C13	1.8084 (18)
S1—C1	1.8136 (19)	S3—C12	1.8163 (17)
S2—C4	1.812 (2)	S4—C15	1.8128 (19)
S2—C1	1.8142 (18)	S4—C12	1.8176 (17)
O1—C8	1.380 (2)	O3—C20	1.372 (2)
O1—C11	1.412 (3)	O3—C22	1.420 (2)
O2—C9	1.377 (2)	O4—C19	1.377 (2)
O2—C11	1.420 (3)	O4—C22	1.419 (3)
C1—C5	1.502 (2)	C12—C16	1.504 (2)
C1—H1	0.9800	C12—H12	0.9800
C2—C3	1.515 (3)	C13—C14	1.507 (3)
C2—H2A	0.9700	C13—H13A	0.9700
C2—H2B	0.9700	C13—H13B	0.9700
C3—C4	1.509 (3)	C14—C15	1.515 (3)
C3—H3A	0.9700	C14—H14A	0.9700
C3—H3B	0.9700	C14—H14B	0.9700
C4—H4A	0.9700	C15—H15A	0.9700
C4—H4B	0.9700	C15—H15B	0.9700
C5—C6	1.383 (3)	C16—C17	1.384 (2)
C5—C10	1.402 (2)	C16—C21	1.405 (2)
C6—C7	1.393 (3)	C17—C18	1.399 (3)

C6—H6	0.9300	C17—H17	0.9300
C7—C8	1.353 (3)	C18—C19	1.360 (3)
C7—H7	0.9300	C18—H18	0.9300
C8—C9	1.377 (3)	C19—C20	1.383 (2)
C9—C10	1.363 (2)	C20—C21	1.365 (2)
C10—H10	0.9300	C21—H21	0.9300
C11—H11A	0.9700	C22—H22A	0.9700
C11—H11B	0.9700	C22—H22B	0.9700
C2—S1—C1	99.80 (9)	C13—S3—C12	99.16 (8)
C4—S2—C1	99.93 (9)	C15—S4—C12	100.05 (8)
C8—O1—C11	105.10 (17)	C20—O3—C22	105.77 (15)
C9—O2—C11	104.86 (18)	C19—O4—C22	105.78 (14)
C5—C1—S1	109.05 (13)	C16—C12—S3	109.78 (11)
C5—C1—S2	110.02 (12)	C16—C12—S4	110.05 (12)
S1—C1—S2	112.76 (10)	S3—C12—S4	112.62 (9)
C5—C1—H1	108.3	C16—C12—H12	108.1
S1—C1—H1	108.3	S3—C12—H12	108.1
S2—C1—H1	108.3	S4—C12—H12	108.1
C3—C2—S1	114.00 (13)	C14—C13—S3	113.97 (13)
C3—C2—H2A	108.8	C14—C13—H13A	108.8
S1—C2—H2A	108.8	S3—C13—H13A	108.8
C3—C2—H2B	108.8	C14—C13—H13B	108.8
S1—C2—H2B	108.8	S3—C13—H13B	108.8
H2A—C2—H2B	107.6	H13A—C13—H13B	107.7
C4—C3—C2	113.67 (17)	C13—C14—C15	112.54 (16)
C4—C3—H3A	108.8	C13—C14—H14A	109.1
C2—C3—H3A	108.8	C15—C14—H14A	109.1
C4—C3—H3B	108.8	C13—C14—H14B	109.1
C2—C3—H3B	108.8	C15—C14—H14B	109.1
H3A—C3—H3B	107.7	H14A—C14—H14B	107.8
C3—C4—S2	114.57 (13)	C14—C15—S4	115.01 (13)
C3—C4—H4A	108.6	C14—C15—H15A	108.5
S2—C4—H4A	108.6	S4—C15—H15A	108.5
C3—C4—H4B	108.6	C14—C15—H15B	108.5
S2—C4—H4B	108.6	S4—C15—H15B	108.5
H4A—C4—H4B	107.6	H15A—C15—H15B	107.5
C6—C5—C10	120.29 (17)	C17—C16—C21	120.01 (16)
C6—C5—C1	119.61 (17)	C17—C16—C12	119.63 (16)
C10—C5—C1	120.09 (16)	C21—C16—C12	120.36 (15)
C5—C6—C7	121.8 (2)	C16—C17—C18	121.92 (18)
C5—C6—H6	119.1	C16—C17—H17	119.0
C7—C6—H6	119.1	C18—C17—H17	119.0
C8—C7—C6	116.8 (2)	C19—C18—C17	116.85 (16)
C8—C7—H7	121.6	C19—C18—H18	121.6
C6—C7—H7	121.6	C17—C18—H18	121.6
C7—C8—C9	122.08 (18)	C18—C19—O4	128.74 (17)
C7—C8—O1	128.4 (2)	C18—C19—C20	121.81 (17)

C9—C8—O1	109.6 (2)	O4—C19—C20	109.45 (17)
C10—C9—O2	127.83 (18)	C21—C20—O3	128.03 (16)
C10—C9—C8	122.31 (18)	C21—C20—C19	122.16 (17)
O2—C9—C8	109.86 (16)	O3—C20—C19	109.81 (16)
C9—C10—C5	116.72 (17)	C20—C21—C16	117.23 (16)
C9—C10—H10	121.6	C20—C21—H21	121.4
C5—C10—H10	121.6	C16—C21—H21	121.4
O1—C11—O2	109.06 (18)	O4—C22—O3	108.66 (16)
O1—C11—H11A	109.9	O4—C22—H22A	110.0
O2—C11—H11A	109.9	O3—C22—H22A	110.0
O1—C11—H11B	109.9	O4—C22—H22B	110.0
O2—C11—H11B	109.9	O3—C22—H22B	110.0
H11A—C11—H11B	108.3	H22A—C22—H22B	108.3
C2—S1—C1—C5	−177.74 (12)	C13—S3—C12—C16	176.91 (13)
C2—S1—C1—S2	59.74 (11)	C13—S3—C12—S4	−60.10 (11)
C4—S2—C1—C5	179.08 (13)	C15—S4—C12—C16	−178.87 (12)
C4—S2—C1—S1	−58.95 (11)	C15—S4—C12—S3	58.30 (11)
C1—S1—C2—C3	−59.52 (17)	C12—S3—C13—C14	61.79 (16)
S1—C2—C3—C4	65.6 (2)	S3—C13—C14—C15	−67.4 (2)
C2—C3—C4—S2	−64.9 (2)	C13—C14—C15—S4	65.0 (2)
C1—S2—C4—C3	58.13 (18)	C12—S4—C15—C14	−57.47 (16)
S1—C1—C5—C6	122.30 (18)	S3—C12—C16—C17	−113.76 (16)
S2—C1—C5—C6	−113.55 (18)	S4—C12—C16—C17	121.74 (15)
S1—C1—C5—C10	−57.71 (19)	S3—C12—C16—C21	66.16 (19)
S2—C1—C5—C10	66.4 (2)	S4—C12—C16—C21	−58.33 (19)
C10—C5—C6—C7	0.9 (3)	C21—C16—C17—C18	1.7 (3)
C1—C5—C6—C7	−179.2 (2)	C12—C16—C17—C18	−178.35 (16)
C5—C6—C7—C8	−0.3 (4)	C16—C17—C18—C19	−1.2 (3)
C6—C7—C8—C9	−0.3 (4)	C17—C18—C19—O4	−179.94 (18)
C6—C7—C8—O1	179.1 (2)	C17—C18—C19—C20	−0.2 (3)
C11—O1—C8—C7	172.7 (3)	C22—O4—C19—C18	174.8 (2)
C11—O1—C8—C9	−7.8 (3)	C22—O4—C19—C20	−5.0 (2)
C11—O2—C9—C10	−173.3 (2)	C22—O3—C20—C21	−176.4 (2)
C11—O2—C9—C8	7.3 (2)	C22—O3—C20—C19	3.8 (2)
C7—C8—C9—C10	0.3 (3)	C18—C19—C20—C21	1.2 (3)
O1—C8—C9—C10	−179.17 (18)	O4—C19—C20—C21	−179.04 (17)
C7—C8—C9—O2	179.8 (2)	C18—C19—C20—O3	−178.97 (17)
O1—C8—C9—O2	0.3 (2)	O4—C19—C20—O3	0.8 (2)
O2—C9—C10—C5	−179.15 (18)	O3—C20—C21—C16	179.54 (18)
C8—C9—C10—C5	0.2 (3)	C19—C20—C21—C16	−0.7 (3)
C6—C5—C10—C9	−0.8 (3)	C17—C16—C21—C20	−0.7 (3)
C1—C5—C10—C9	179.21 (16)	C12—C16—C21—C20	179.33 (16)
C8—O1—C11—O2	12.4 (3)	C19—O4—C22—O3	7.3 (2)
C9—O2—C11—O1	−12.2 (2)	C20—O3—C22—O4	−6.8 (2)

Hydrogen-bond geometry (Å, °)

Cg1 and Cg2 are the centroids of the C5–C11 and C16–C21 rings, respectively.

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C4—H4b $\cdots$ Cg1 ⁱ	0.97	2.77	3.731 (2)	170
C13—H13a $\cdots$ Cg2 ⁱⁱ	0.97	2.54	3.4841 (19)	165

Symmetry codes: (i) $x, -y+1/2, z-1/2$; (ii) $x, -y-1/2, z-3/2$.