



Relating cardiorespiratory responses to work rate during incremental ramp exercise on treadmill in children and adolescents: sex and age differences

Nicola Lai^{1,2,3} · Justin J. Fiutem^{2,4} · Nora Pfaff⁵ · Desy Salvadego⁶ · James Strainic^{2,4}

Received: 22 January 2021 / Accepted: 8 June 2021 / Published online: 18 June 2021
© The Author(s) 2021

Abstract

Purpose Evaluation of cardiopulmonary exercise testing (CPET) slopes such as dHR/dWR_{tot} (cardiac/skeletal muscle function) and $d\dot{V}O_2/dWR_{tot}$ (O_2 delivery/utilization), using treadmill protocols is limited because the difficulties in measuring the total work rate (WR_{tot}). To overcome this limitation, we proposed a new method in quantifying WR_{tot} to determine CPET slopes.

Methods CPET's were performed by healthy patients, ($n = 674$, 9–18 year) 300 female (F) and 374 male (M), using an incremental ramp protocol on a treadmill. For this protocol, a quantitative relationship based on biomechanical principles of human locomotion, was used to quantify the WR_{tot} of the subject. CPET slopes were determined by linear regression of the data recorded until the gas exchange threshold occurred.

Results The method to estimate WR_{tot} was substantiated by verifying that: $d\dot{V}O_2/dWR_{tot}$ for treadmill exercise corresponded to an efficiency of muscular work similar to that of cycle ergometer; $d\dot{V}O_2/dWR_{tot}$ ($mL\ min^{-1}\ W^{-1}$) was invariant with age and greater in M than F older than 12 years old (13–14 years: 9.6 ± 1.5 (F) vs. 10.5 ± 1.8 (M); 15–16 years: 9.7 ± 1.7 (F) vs. 10.6 ± 2.2 (M); 17–18 years: 9.6 ± 1.7 (F) vs. 11.0 ± 2.3 (M), $p < 0.05$); similar to cycle ergometer exercise, dHR/dWR_{tot} was inversely related to body weight (BW) ($r = 0.71$) or $\dot{V}O_{2, peak}$ ($r = 0.66$) and $d\dot{V}O_2/dWR_{tot}$ was not related to BW ($r = -0.01$), but had a weak relationship with $\dot{V}O_{2, peak}$ ($r = 0.28$).

Conclusion The proposed approach can be used to estimate WR_{tot} and quantify CPET slopes derived from incremental ramp protocols at submaximal exercise intensities using the treadmill, like the cycle ergometer, to infer cardiovascular and metabolic function in both healthy and diseased states.

Keywords Kinetics · Muscular efficiency · CPET slopes · Oxygen delivery · External work rate

Communicated by I. Markolfert.

✉ Nicola Lai
nicola.lai@unica.it

¹ Department of Mechanical, Chemical and Materials Engineering, University of Cagliari, Cagliari, Italy

² Departments of Pediatrics, Case Western Reserve University, Cleveland, OH, USA

³ Biomedical Engineering, Case Western Reserve University, Cleveland, OH, USA

⁴ Rainbow Babies and Children's Hospital, Cleveland, OH, USA

⁵ Department of Pediatrics, University of California, San Francisco, CA, USA

⁶ Department of Automation, Biocybernetics and Robotics, Jožef Stefan Institute, Ljubljana, Slovenia

List of symbols

dHR/dWR_{tot}	Slope: variation of HR per variation of WR_{tot}
$d\dot{V}O_2/dHR$	Slope: variation of $\dot{V}O_2$ per variation of HR
$d\dot{V}O_2/dWR_{tot}$	Slope: variation of $\dot{V}O_2$ per variation of WR_{tot}
f_{ext}	Fraction of the external work rate
g	Acceleration of gravity
GET	Gas exchange threshold
HR	Heart beat
i	Inclination
m	Body mass
s	Speed of the subject
t	Time
$\dot{V}\dot{C}O_2$	Rate of carbon dioxide release
\dot{V}_e	Ventilation rate

$\dot{V}O_2$	Pulmonary oxygen uptake
$\dot{V}O_{2,GET}$	Pulmonary oxygen uptake at GET
$\dot{V}O_{2,peak}$	Pulmonary oxygen uptake at peak
WR_{ext}	External work rate
WR_{tot}	Total work rate

Introduction

Cardiopulmonary exercise testing (CPET) continues to be the ideal method in assessing cardiovascular and respiratory function in children and adolescents (Guazzi et al. 2017). CPET is performed using a cycle ergometer or a treadmill (Paridon et al. 2006; Armstrong and Welsman 2019a) with a progressive increase in workload to challenge the subject to the limit of tolerance. The work rate (WR) can be imposed and measured in protocols with the cycle ergometer in a precise manner, whereas in treadmill CPET, the WR depends on body weight, inclination, running speed, and other factors that cannot be easily quantified.

This cycle ergometer feature allows clinicians and researchers to relate pulmonary gas-exchange and heart rate adjustments to work rate by $d\dot{V}O_2/dWR$ and dHR/dWR (i.e., CPET slopes) to evaluate cardiorespiratory function and the efficiency of O_2 delivery and utilization processes (Cooper et al. 2014). CPET slopes quantify the integrated responses of the cardiovascular, respiratory, and skeletal muscle systems to an increase in ATP demand and provide complementary diagnostic information to the maximal pulmonary O_2 uptake ($\dot{V}O_{2,max}$). Because CPET slopes are obtained from data recorded at submaximal exercise intensities (Cooper et al. 2014), they represent a suitable alternative to $\dot{V}O_{2,max}$ measurement in assessing cardiorespiratory function in pediatric populations for those subjects who cannot reliably reach $\dot{V}O_{2,max}$ (Shaibi et al. 2006; Poole and Jones 2017).

The CPET slopes are clinically relevant in both pediatric (Groen et al. 2010) and adult populations (Troutman et al. 1998; Guazzi et al. 2012, 2017; Ashish et al. 2015; Barron et al. 2016; Popovic et al. 2019). Their determinations rely on the knowledge of WR also referred to as total work rate (WR_{tot}) in this manuscript and can only be determined when using cycle ergometer protocols. In treadmill protocols, especially in clinical settings, it is difficult to quantify WR_{tot} . The possibility of relating CPET variables to WR could enable investigators to analyze CPET slopes obtained from treadmill protocols.

Computational and experimental studies provide robust information on quantifying the external (WR_{ext}) and internal work rate as major components of the work rate for walking and running (Cavagna and Kaneko 1977; Willems et al. 1995; Minetti 2000). Specifically, studies on kinetic

movements showed that the WR_{ext} fraction was approximately 40–50% and 65–75% of the WR_{tot} (internal and external) for walking and running, respectively (Minetti et al. 1994b, 2000; Willems et al. 1995). This relationship can quantify WR_{tot} from WR_{ext} which could be estimated from biomechanical principles of human locomotion. This approach to estimate WR_{tot} for treadmill protocols is promising but has not been tested yet.

Physiological functions change over the course of growth and development (Cooper et al. 1984, 2014) and CPET slopes are a valuable tool to evaluate these changes. In a cycle ergometer study (Cooper et al. 2014) for children and adolescents, dWR/dHR and $d\dot{V}O_2/dHR$ slopes were linearly related to body size whereas $d\dot{V}O_2/dWR$ slope was body weight independent. In the same study, these slopes have been reported to be lower in females than males. These CPET slope characteristics (muscular efficiency, sex differences, dependence on body weight) observed for cycle ergometer are expected to be similar for treadmill protocols and can be used to verify the consistency of the WR_{tot} estimate for treadmill exercise.

The aim of this study was to develop an equation to quantify WR_{tot} for incremental ramp protocols using a treadmill and provide evidence that WR_{tot} estimates are consistent with CPET slope characteristics. The equation is based on the relationship between WR_{tot} and WR_{ext} . WR_{ext} is estimated from the work required to move an object in a plane at a specific inclination and speed (Margaria 1978; Porszasz et al. 2003). To determine the efficacy of the method proposed to estimate WR_{tot} , we evaluate whether (a) $d\dot{V}O_2/dWR_{tot}$ for treadmill was similar to the efficiency of muscular work for walking, running and cycling (Whipp and Wasserman 1969; Brooks 2012) and (b) the CPET slopes (dHR/dWR_{tot} and $d\dot{V}O_2/dWR_{tot}$) are related to body weight or $\dot{V}O_{2,peak}$. Also, we substantiate the versatility of the WR_{tot} equation by investigating gender and age differences in the CPET slopes obtained in children and adolescents exercising on a treadmill. We hypothesize that in adolescents: (1) males have a higher $d\dot{V}O_2/dWR_{tot}$ and $d\dot{V}O_2/dHR$ than females; (2) females have higher dHR/dWR_{tot} than males; and (3) dHR/dWR_{tot} decrease and $d\dot{V}O_2/dHR$ increase with age; (4) CPET slopes are related to body weight except $d\dot{V}O_2/dWR_{tot}$.

Methods

This was a retrospective study based on data from 1934 patients, mainly children and adolescents who had undergone CPET on a treadmill at Rainbow Babies and Children's Hospital between the years 2000 and 2013. The study initially included healthy subjects and patients with congenital heart disease (ages 9–18 years); the analysis focused on the healthy

patients. Patients were excluded if they were underweight with a BMI less than the 5th percentile, overweight with a BMI greater than the 85th percentile, had cardiopulmonary disease, had an RER < 1.0 or an exhaustion time less than 350 s. The final number of included patients in the study was 674 (300 females and 374 males).

The subjects underwent a modified Bruce protocol with an incremental ramp (DiBella et al. 2002) with a linear change in both speed and slope of the treadmill (Fig. 1). During the test, a 12-lead EKG was used to measure heart rate (HR), while the metabolic cart was used to determine pulmonary oxygen uptake ($\dot{V}O_2$), carbon dioxide release ($\dot{V}CO_2$), and minute ventilation (\dot{V}_e).

The protocol consisted of three phases: warmup, exercise, and recovery. The test started with 3 min of walking at 26.8 m min⁻¹ (warm-up) and 0% incline and continued until speed and inclination were increased to 45.6 m min⁻¹ and 10%, respectively. After 3 min, both speed (*s*) and inclination (*i*) were increased every minute by 7.4 m min⁻¹ and 0.66%, respectively. Thus, *s* and *i* change with time (*t*) according to the relationships:

$$s(t) = 45.6 + 7.4 t. \tag{1}$$

$$i(t) = 10 + 0.66 t. \tag{2}$$

The subjects continued exercise until they reached the limit of tolerance.

The external work rate (WR_{ext}) generated by the subject running on the treadmill with an angle of inclination $\alpha(t)$ and speed *s*(*t*) was estimated as:

$$WR_{ext}(t) = m g s(t) \sin [\alpha(t)]. \tag{3}$$




Fig. 1 Comparison between incremental ramp protocol used in this study and Bruce protocol: speed and inclination changes with time

where *m* and *g* represent the body mass and the acceleration of gravity, respectively. The angle of inclination is related to the inclination by the trigonometric relationship:

$$\sin [\alpha(t)] = i(t) / \sqrt{1 + i(t)^2}. \tag{4}$$

It was assumed that the fraction of the external work rate (f_{ext}) was the same at different speeds and inclinations. The external work rate was 70% of the total work rate ($f_{ext} = 0.7$) (Gosseye et al. 2010). With this assumption, the total work rate can be computed as a function of WR_{ext} , and f_{ext} :

$$WR_{tot}(t) = WR_{ext}(t) / f_{ext}. \tag{5}$$

The estimate of WR_{tot} by Eq. 5 accounts for inclination, speed, and body weight of the subject running on the treadmill. The data were processed using excel macros. Outliers sporadically observed in HR and breath-by-breath CPET data were identified and excluded for the determination of peak (i.e., $\dot{V}O_2$ and WR_{tot}) or slopes.

A facemask (8900 Series, Hans Rudolph, Inc.; Kansas, MO) was carefully fitted and sealed with a gel (Hans Rudolph, Inc.) to obviate any gas leaks during exercise. The subjects were given several minutes to familiarize themselves with the breathing apparatus to minimize unusual breathing patterns. To measure gas exchange, subjects breathed through a mass flow sensor (hot-wire anemometer) connected to a metabolic cart system (VMax 29, Sensor-Medics, Yorba Linda, CA). Before each exercise test, the volume sensor was calibrated using a 3-L syringe while the O₂ and CO₂ analyzers were calibrated as previously reported (Lai et al. 2008).

Gas exchange threshold (GET, $\dot{V}O_{2,GET}$) was determined by the V-slope method in combination with ventilatory equivalents and the end-tidal O₂ responses (Beaver et al. 1986). Peak values for HR, $\dot{V}O_2$, $\dot{V}CO_2$, O₂ pulse calculated as $\dot{V}O_2/HR$, and \dot{V}_e were determined as the average of the last 20 s prior to exhaustion. Data were processed with a moving average of 10 s.

For each subject's data, a linear regression between WR_{tot} and time, $\dot{V}O_2$ and time, HR and WR_{tot} , $\dot{V}O_2$ and HR, $\dot{V}O_2$ and WR_{tot} was obtained to determine the slopes for dWR_{tot}/dt , $d\dot{V}O_2/dt$, dHR/dWR_{tot} , $d\dot{V}O_2/dHR$, and $d\dot{V}O_2/dWR_{tot}$. The linear regression was performed with the recorded data in the interval of time between 60 s after the onset of exercise until GET occurred.

Statistical analysis

The results are reported as means ± standard deviation. Linear regression was used to evaluate the relationship between CPET slopes and body weight and between CPET slopes and $\dot{V}O_{2,peak}$. The correlation between the variable was quantified

with Pearson's coefficient (r). The comparisons of the CPET variables and slopes obtained from female and male tests for different age groups were evaluated with a two-way ANOVA with Bonferroni for multiple comparisons (Origin software). The interaction effect between age and gender was investigated for the CPET slopes. Student t -test was used to compare regression lines between female and male groups and determine whether the parameters of the regression lines were significantly different. Difference of $p < 0.05$ was considered significant.

Results

Anthropometric measurements were grouped by sex and age and reported in Table 1. Both height and weight increased with age and there were no significant differences in height, weight, or BMI percentiles between male and female patients. For both female and male groups, BMI percentile was similar at 50–60%.

Cardiopulmonary response to peak exercise: sex and age

CPET variables for males and females measured among different age groups are reported in Table 2. Maximal HR was not affected by sex, whereas $\dot{V}O_{2,GET}$, O_2 pulse, peak of WR_{tot} , \dot{V}_e , and $\dot{V}O_2$ were higher in male than in female patients > 12 years of age. The peak WR_{tot} and O_2 pulse increased with age in both sexes.

CPET slopes at submaximal exercise: relating the estimate of WR_{tot} to $\dot{V}O_2$ and HR

The $d\dot{V}O_2/dWR_{tot}$, $d\dot{V}O_2/dHR$, and dHR/dWR_{tot} individual slopes were determined using data recorded before the gas exchange threshold, as reported in Fig. 2. The range of $d\dot{V}O_2/dWR_{tot}$ slopes (Fig. 3a, 9.5–11 mL O_2 min⁻¹ W⁻¹) based on the estimate of $WR_{tot}(t)$ across different age groups corresponded to a range of typical values of muscular efficiency from 30 to 26% (assuming an energy equivalent for oxygen of 20.9 J mL O_2 ⁻¹). The $d\dot{V}O_2/dWR_{tot}$ slope was greater in male than in female patients > 12 years of age (Fig. 3a), and the effect of age was absent in both female and male groups. For both $d\dot{V}O_2/dHR$ and dHR/dWR_{tot} slopes, there was a significant difference between females and males among different age groups (Fig. 3b, c): $d\dot{V}O_2/dHR$ was lower and dHR/dWR_{tot} was higher in females than males. Furthermore, for both males and females $d\dot{V}O_2/dHR$ increased, whereas dHR/dWR_{tot} decreased in older subjects. The interaction between sex and age was also statistically significant for both $d\dot{V}O_2/dHR$ and dHR/dWR_{tot} slopes.

In Table 3 was reported the total work rate variation with time during the incremental ramp (dWR_{tot}/dt). The dWR_{tot}/dt slope increased with age in both females and males and was higher in males than females in groups older than 12 years old.

To further investigate the sex differences observed in $d\dot{V}O_2/dWR_{tot}$, its components dWR_{tot}/dt and $d\dot{V}O_2/dt$ were analyzed separately. Specifically, dWR_{tot}/dt (Fig. 4a) and $d\dot{V}O_2/dt$ (Fig. 4b) were plotted against body weight for

Table 1 Summary of the anthropometric measurement of the population

	9–10	11–12	13–14	15–16	17–18
N. Subjects (–)					
Female	19	39	86	121	35
Male	26	66	106	107	69
Height (m)					
Female	1.42 ± 0.08	1.52 ± 0.07 [§]	1.60 ± 0.07 [†]	1.64 ± 0.06 [†]	1.64 ± 0.07 [†]
Male	1.39 ± 0.08	1.41 ± 0.08 [§]	1.68 ± 0.09 ^{*,†}	1.74 ± 0.08 ^{*,†}	1.76 ± 0.06 ^{*,†}
Weight (kg)					
Female	36.1 ± 6.1	45.7 ± 7.5 [§]	52.6 ± 8.3 [†]	59.0 ± 8.7 [†]	61.2 ± 9.6 [†]
Male	35.2 ± 5.8	42.5 ± 7.5 [§]	59.0 ± 11.1 ^{*,†}	66.6 ± 9.8 ^{*,†}	71.8 ± 9.5 ^{*,†}
BMI (kg m ⁻²)					
Female	17.9 ± 2.1	19.7 ± 2.2	20.3 ± 2.5 [§]	21.7 ± 2.7 [†]	22.7 ± 3.2 [†]
Male	18.0 ± 2.0	18.4 ± 2.1	20.7 ± 2.6 [†]	21.8 ± 2.4 [†]	23.1 ± 2.4 [†]
BMI (Percentile)					
Female	58.2 ± 24.9	64.5 ± 20.7	55.4 ± 24.8	58.8 ± 24.5	59.0 ± 26.0
Male	62.1 ± 26.4	53.5 ± 25.6	61.9 ± 25.6	60.7 ± 24.2	60.3 ± 23.8

*Influence of gender: statistically different from female ($p < 10^{-3}$)

[†]Influence of age within the group: statistically different from 9- to 10- and 11–12-year-old groups ($p < 10^{-2}$)

[§]Influence of age within the group: statistically different from 9- to 10-year-old group ($p < 10^{-4}$)

Table 2 Summary of the CPET variables measured during the incremental ramp protocol

	9–10	11–12	13–14	15–16	17–18
Exercise time (min)					
Female	7.3 ± 1.1	7.4 ± 1.2	7.9 ± 1.1	8.1 ± 1.4	7.9 ± 1.6
Male	7.8 ± 0.9	8.5 ± 1.4*	9.2 ± 1.2*, †	9.7 ± 1.4*, †	9.7 ± 1.3*, †
Peak HR (beat min ⁻¹)					
Female	192 ± 10	193 ± 11	193 ± 11	190 ± 10	190 ± 12
Male	192 ± 10	191 ± 9	194 ± 9	191 ± 10	188 ± 11
Peak $\dot{V}O_2$ (mL kg ⁻¹ min ⁻¹)					
Female	41.5 ± 6	40.0 ± 7	41.7 ± 6	41.3 ± 7	39.8 ± 7
Male	47.9 ± 8	48.5 ± 6*	51.0 ± 7*, †	52.5 ± 8*, †	52.7 ± 8*, †
Peak $\dot{V}CO_2$ (mL kg ⁻¹ min ⁻¹)					
Female	45.8 ± 7	45.1 ± 9	47.7 ± 7	48.3 ± 8	47.0 ± 9
Male	50.2 ± 8	52.6 ± 7*	56.7 ± 7*, †	59.3 ± 8*, †	59.8 ± 8*, †
Peak <i>RER</i> (-)					
Female	1.10 ± 0.05	1.12 ± 0.07	1.14 ± 0.06	1.17 ± 0.08	1.18 ± 0.07
Male	1.05 ± 0.04	1.08 ± 0.06	1.11 ± 0.06	1.13 ± 0.07*	1.14 ± 0.08
Peak \dot{V}_e (mL kg ⁻¹ min ⁻¹)					
Female	1.60 ± 0.4	1.48 ± 0.3	1.47 ± 0.3	1.45 ± 0.3	1.38 ± 0.2
Male	1.62 ± 0.3	1.62 ± 0.4	1.74 ± 0.3*	1.73 ± 0.3*	1.69 ± 0.3*
Peak $\dot{V}O_2/HR$ (mL beat ⁻¹)					
Female	7.8 ± 1.7	9.4 ± 1.9	11.3 ± 2.2 [§]	12.8 ± 2.7 [†]	12.7 ± 2.6 [†]
Male	8.7 ± 1.3	10.7 ± 2.2	15.5 ± 3.5*, †	18.3 ± 3.7*, †	20.1 ± 3.9*, †
$\dot{V}O_{2,GET}$ (mL kg ⁻¹ min ⁻¹)					
Female	29.4 ± 4	28.9 ± 5	29.7 ± 5	29.8 ± 5	28.6 ± 5
Male	33.6 ± 5	34.9 ± 5*	35.3 ± 5*	35.9 ± 6*	35.9 ± 6*
Peak WR_{tot} (W)					
Female	126 ± 28	160 ± 30	194 ± 38 [†]	224 ± 46 [†]	225 ± 44 [†]
Male	129 ± 22	167 ± 40 [§]	252 ± 60*, †	299 ± 57*, †	322 ± 57*, †

*Influence of gender: statistically different from female ($p < 10^{-3}$)

[†]Influence of age within the group: statistically different from 9- to 10- and 11–12-year-old groups ($p < 10^{-2}$)

[§]Influence of age within the group: statistically different from 9- to 10-year-old group ($p < 10^{-4}$)




Fig. 2 Representative case to determine the slope of $d\dot{V}O_2/dWR_{tot}$ (a), $d\dot{V}O_2/dHR$ (b), and dHR/dWR_{tot} (c) by linear regression

both male and female groups. Both WR_{tot}/dt and $d\dot{V}O_2/dt$ have a strong linear relation to body weight. In males, the dWR_{tot}/dt change with body weight was only 6% greater

than that observed for females, whereas the $d\dot{V}O_2/dt$ change with body weight was 41% greater than that observed for females. Thus, the higher $d\dot{V}O_2/dWR_{tot}$ slopes observed




Fig. 3 Effects of gender and age on the slope $d\dot{V}O_2/dWR_{tot}$ (a), $d\dot{V}O_2/dHR$ (b), dHR/dWR_{tot} (c). *Influence of gender: statistically different from female for $d\dot{V}O_2/dWR_{tot}$ ($p < 0.05$) and for $d\dot{V}O_2/dHR$ and dHR/dWR_{tot} ($p < 10^{-4}$). †Influence of age: within

the group: statistically different from 9–10 and 11–12 years old groups ($p < 10^{-3}$). §Influence of age within the group: statistically different from 9–10 years old ($p < 10^{-3}$)

Table 3 Summary of the CPET variables measured during the incremental ramp protocol

	9–10	11–12	13–14	15–16	17–18
dWR_{tot}/dt ($W \text{ min}^{-1}$)					
Female	11.4 ± 1.0	14.4 ± 2.3§	16.7 ± 2.6†	18.8 ± 2.8†	19.4 ± 2.8†
Male	11.2 ± 1.8	13.7 ± 2.3§	19.5 ± 3.7*,†	22.1 ± 3.1*,†	23.8 ± 3.0*,†
$d\dot{V}O_2/dWR_{tot}$ ($\text{mL min}^{-1} W^{-1}$)					
Female	9.6 ± 1.6	9.5 ± 1.5	9.6 ± 1.5	9.7 ± 1.7	9.6 ± 1.7
Male	10.6 ± 1.8	10.6 ± 1.2	10.5 ± 1.8*	10.6 ± 2.2*	11.0 ± 2.3*

*Influence of gender: statistically different from female ($p < 10^{-2}$)

†Influence of age within the group: statistically different from 9- to 10- and 11–12-year-old groups ($p < 10^{-3}$)

§Influence of age within the group: statistically different from 9- to 10-year-old group ($p < 10^{-2}$)




Fig. 4 Comparison of the linear relationship dWR_{tot}/dt vs. body weight (BW) (a) between male and female groups, slope is significantly different from zero for male ($dWR_{tot}/dt = 0.33 BW - 0.14$, $p < 10^{-8}$, solid line) and female ($dWR_{tot}/dt = 0.31 BW - 0.37$, $p < 10^{-8}$, dot line). Slopes are statistically different $p < 10^{-4}$, com-

parison of the linear relationship $d\dot{V}O_2/dt$ vs. BW between male and female groups, slope is significantly different from zero for male ($d\dot{V}O_2/dt = 3.6 BW - 2.4$, $p < 10^{-8}$, solid line) and female ($d\dot{V}O_2/dt = 2.5 BW + 29$, $p < 10^{-8}$, dot line), slopes are statistically different $p < 10^{-5}$

in males is mainly due to the higher $d\dot{V}O_2/dt$ because dWR_{tot}/dt (i.e., workload changes) in females and males with the same body weight was similar (Fig. 4a).

To further evaluate the reliability of $WR_{tot}(t)$ estimates, we tested whether any of the $d\dot{V}O_2/dWR_{tot}$, $d\dot{V}O_2/dHR$, and dHR/dWR_{tot} slopes were related to body weight (Fig. 5) or $\dot{V}O_{2, peak}$ (Fig. 6). The $d\dot{V}O_2/dWR_{tot}$ was not related to body weight (Fig. 5a) but it was significantly related to $\dot{V}O_{2, peak}$ (Fig. 6a). The $d\dot{V}O_2/dHR$ was linearly related to body weight (Fig. 5b) and $\dot{V}O_{2, peak}$ (Fig. 6b), whereas dHR/dWR_{tot} was inversely related to body weight (Fig. 5c) and $\dot{V}O_{2, peak}$ (Fig. 6c).

Discussion

This study focused on a method to quantify WR_{tot} during incremental ramp exercise on a treadmill in children and adolescents and to evaluate whether the CPET

slope characteristics determined with WR_{tot} expression are like those obtained with the cycle ergometer. The estimates of WR_{tot} for treadmill exercise were consistent with typical values of muscular efficiency equivalent to the $d\dot{V}O_2/dWR_{tot}$ range values (9.5–11 mL O_2 $min^{-1} W^{-1}$) observed for both female and male groups. The $d\dot{V}O_2/dWR_{tot}$ slope is greater in males than females and unchanged with age as previously reported for cycling exercise (Cooper et al. 2014). The proposed equation is also substantiated by the linear relationship between dHR/dWR_{tot} and body weight or $\dot{V}O_{2, peak}$, and between $d\dot{V}O_2/dWR_{tot}$ and $\dot{V}O_{2, peak}$ which is similar to that observed using the cycle ergometer (Cooper et al. 2014). Males rely on a higher stroke volume (i.e., $d\dot{V}O_2/dHR$) than females, whereas females compensate for this difference with larger changes in heart rate for the same workload increase (i.e., dHR/dWR_{tot}). In both males and females, the dHR/dWR_{tot} decrease and $d\dot{V}O_2/dHR$ increase with age indicates a compensatory mechanism that was likely due to an increase of stroke volume.




Fig. 5 Assessment of the linear relationship between $d\dot{V}O_2/dWR_{tot}$ and body weight (BW) (a), slope is not significantly different from zero ($p > 0.8$); $d\dot{V}O_2/dHR$ and BW (b), slope significantly different from zero ($p < 10^{-10}$); and dHR/dWR_{tot} and BW (c), slope significantly different from zero ($p < 10^{-10}$)




Fig. 6 Assessment of the linear regression between $d\dot{V}O_2/dWR_{tot}$ and $\dot{V}O_{2, peak}$ (a); $d\dot{V}O_2/dHR$ and $\dot{V}O_{2, peak}$ (b); dHR/dWR_{tot} and $\dot{V}O_{2, peak}$ (c), slope is significantly different from zero for each case ($p < 10^{-10}$)

Maximal work rate

The estimate of the peak WR_{tot} observed in this study increases with age as does $\dot{V}O_{2, peak}$ (Table 2). In this study, $\dot{V}O_{2, peak}$ and peak WR_{tot} obtained during incremental exercise on a treadmill are 10% and 25% higher than those obtained by (Cooper et al. 2014) incremental exercise on a cycle ergometer in populations with comparable age and body weight. Although these comparisons are limited by the fact that the cycle ergometer and treadmill data are not obtained from the same adolescent groups, the treadmill data are consistently higher than those of the cycle ergometer. The cycle ergometer involves less muscle mass recruitment when compared to the exercise on a treadmill (Krahenbuhl et al. 1985; Armstrong and Welsman 1994) which involves weight bearing muscle groups. In a study with children under the age of 10, $\dot{V}O_{2, peak}$ on a treadmill was 11% higher than that obtained on the cycle ergometer for both females and males (Turley et al. 1995). In another study on children, adolescents, and adults, a similar $\dot{V}O_{2, peak}$ difference was observed in female adolescents (Bar-Yoseph et al. 2019). Thus, a higher peak WR_{tot} for treadmill exercise than that for a cycle ergometer is consistent with the $\dot{V}O_{2, peak}$ differences observed between the two exercise modalities. The WR_{tot} expression allows for the detection of maximal WR_{tot} differences related to sex and age like those for $\dot{V}O_{2, peak}$.

External and total work rate in relation to $\dot{V}O_2$

The equation used to estimate WR_{tot} is based on biomechanical principles to study internal, external, and total work rates in human movement, (Cavagna et al. 1964; Cavagna and Kaneko 1977; Willems et al. 1995; Minetti et al. 2000) and used to optimize treadmill protocols (Porszasz et al. 2003). The external work rate quantifies the power required to move the body on a treadmill at specific speeds and inclines, but it does not include the work made by the joints to move the body's center of mass (i.e., internal) (Herman 2008). Thus, we related the external to the total work rate for running based on the analysis of the kinetic and gravitational potential energy of the body's center of mass (Cavagna et al. 1964; Cavagna and Kaneko 1977; Willems et al. 1995; Gosseye et al. 2010). Specifically, it is assumed that the fraction of the external work rate (f_{ext}) is 70% of the total work rate during the incremental ramp protocol. The value of f_{ext} is close to the range of values (60–75%) previously determined on a treadmill with a plate force to measure the force exerted under the foot during running (Schepens et al. 2001; Gosseye et al. 2010). A similar range was also observed in other studies (Minetti et al. 1994b, 2000; Willems et al. 1995; Saibene and Minetti 2003). Although the assumption of considering $f_{ext}=70\%$ is consistent with the CPET slope characteristics, it might not fully reflect the individual

body movement characteristics at different running speeds. In pediatric and adult studies on locomotion, f_{ext} decreased from 75 to 70% (Minetti et al. 1994b; Schepens et al. 2001; Gosseye et al. 2010) for an increase in running speed in the range (50–85 m min⁻¹) used in our protocol. If these f_{ext} changes with speed are considered, $d\dot{V}O_2/dWR_{tot}$ slope decreases by 10% of the values obtained with a constant f_{ext} (Figs. 2, 3). This potential effect on $d\dot{V}O_2/dWR_{tot}$ is the same in female and male because the workload was the same for subjects of both sexes with the same body weight (Fig. 4a), thus sex differences observed in our work were not affected. Also, age-related changes in walking characteristics have been reported in children (Malloggi et al. 2019). Potential differences of body movement associated to age, sex, speed and inclination (Minetti et al. 1994a) are not considered in the expression proposed to quantify WR_{tot} . Nevertheless, the $d\dot{V}O_2/dWR_{tot}$ (9.5–11 mL min⁻¹ W⁻¹) values obtained in our study are similar to those obtained for cycle ergometer exercise (Lai et al. 2012; Jones and Poole 2013; Cooper et al. 2014), thus indirectly supporting the validity of the proposed equation to estimate the total work rate. The $d\dot{V}O_2/dWR_{tot}$ values observed among age groups correspond to the range of muscular work efficiency (30–26%) determined for cycling exercise (Whipp and Wasserman 1969). This evidence indicates that the WR_{tot} estimated for males and females at different age groups is consistent with the thermodynamic constraints of the biochemical processes in contracting muscle.

To further substantiate the approach proposed in calculating WR_{tot} and estimating CPET slopes at submaximal treadmill exercise, we verify that the relationship between CPET slopes and body weight or $\dot{V}O_{2, peak}$ for the treadmill is similar to that observed for the cycle ergometer (Figs. 5 and 6). The inverse correlation between dHR/dWR_{tot} and body weight and between dHR/dWR_{tot} and $\dot{V}O_{2, peak}$ are consistent with the linear relationship between dWR_{tot}/dHR and body weight and between dWR_{tot}/dHR and $\dot{V}O_{2, peak}$ observed in cycle ergometer exercise (Cooper et al. 2014). Although $d\dot{V}O_2/dWR_{tot}$ was not related to body weight, it was significantly related to $\dot{V}O_{2, peak}$ (Fig. 5). Both findings echoed those reported for cycle ergometer exercise in the same age groups. Thus, the WR_{tot} estimated using the treadmill protocol is body size dependent and enables the analysis of size-dependent and size-independent properties of the CPET slopes similarly to the analysis obtained using cycle ergometer (Cooper et al. 2014).

dHR/dWR_{tot} and $d\dot{V}O_2/dHR$: sex and age differences

The higher dHR/dWR_{tot} and lower $d\dot{V}O_2/dHR$ slopes in females suggests a sex difference in the cardiovascular response to submaximal exercise (Fig. 3). Whereas the first

slope quantifies HR change with increasing power output, the second slope quantifies the product of stroke volume and arterio-venous difference. Maximal cardiac output differences between females and males have been reported at different ages (Miyamura and Honda 1973). Specifically, in pediatric populations, males appear to have higher maximal cardiac outputs than females and this difference in part is related to a higher stroke volume in males (Rowland et al. 2000). Also, at submaximal exercise intensities, males relied on higher stroke volume with a lower HR (Turley and Wilmore 1997a, b; Armstrong et al. 2007) whereas arterio-venous oxygen difference was similar between genders. In these studies, the greater stroke volume observed in males was mainly attributed to a larger left ventricular muscle size (Turley and Wilmore 1997b). Thus, in our study, under the assumption that the arterio-venous oxygen difference in male is like that in female, the greater $d\dot{V}O_2/dHR$ in male patients was consistent with the finding of a higher stroke volume in males as compared to female patients. Furthermore, the higher dHR/dWR_{tot} in females (Fig. 3c) was consistent with a higher amplitude of HR per WR observed at moderate and heavy exercise intensities in females (Lai et al. 2016). In another study relating CPET variables to work rate for treadmill exercise (Bar-Yoseph et al. 2019), the dHR/dWR slope in children and adolescents was similar to that found in our study. In this study, WR was determined imposing metabolic efficiency, whereas in our work, WR was independently derived from a biomechanical relationship between external and total work rate.

An age effect on the slopes was also observed in our study. The increase of $d\dot{V}O_2/dHR$ with age is in part related to the growth of left ventricular mass contributing to higher stroke volume (Daniels et al. 1995). It has been previously reported that the stroke volume of older children is higher than that of younger children for a given exercise intensity (Bar-Or 1983). Thus, the decrease of dHR/dWR_{tot} with age observed in our study is consistent with a compensatory mechanism of an increase in stroke volume (indicated by an increase in the $d\dot{V}O_2/dHR$ slope). This cardiovascular adaptation in exercise is known in the pediatric population (Bar-Or 1983). Sex differences for $d\dot{V}O_2/dHR$ during growth as observed in our study in subjects older than 12 years old, can be partially explained by a greater growth in left ventricular mass in males when compared to females undergoing pubertal development (Pelà et al. 2016).

Our analysis was based on retrospective data obtained from exercise tests in which lean body mass was not systematically recorded. The analysis was mainly focused on corroborating the method proposed to estimate WR_{tot} . Although we cannot quantify the effects of body composition on CPET slopes, we can exclude any effect of BMI on sex differences because BMI was comparable in males and females at each age group (Cooper et al. 2016). To study

sex differences in $\dot{V}O_{2,peak}$ during development it is crucial to account for lean body mass (Armstrong and Welsman 2019b). Our work focused on CPET slopes, which in the case of dHR/dWR_{tot} and $d\dot{V}O_2/dHR$, should also be affected by body composition.

In conclusion, the results of this work indicate that the methodology proposed has the potential to be used as a tool in clinical and research settings to estimate the workload to determine the CPET slopes derived from an incremental ramp protocol using the treadmill. The estimates of the WR_{tot} for treadmill were substantiated by verifying thermodynamic constraint and CPET slope characteristics. Nevertheless, future work should focus on further validating the proposed approach. A direct comparison of treadmill and cycle ergometer exercises as well as ways to change and measure work rate from individual locomotion characteristics for treadmill exercise can contribute to the design of validation strategies. As for the cycle ergometer, CPET slopes depending on WR_{tot} , can be used to infer cardiovascular and metabolic function at submaximal exercise intensities in healthy and diseased states.

Author contributions NL contributed to the study conception. NL, JF and JS contributed to the design. Data collection was performed by NL, JF and NP. All authors contributed to the formal analysis and interpretation of the data. The first draft of the manuscript was written by NL and all authors commented on previous versions of the manuscript. All authors have read and approved the final version of the manuscript.

Funding Open access funding provided by Università degli Studi di Cagliari within the CRUI-CARE Agreement. This research was supported in part by National Institutes of Health Grants GM6630901 and K25AR057206.

Declarations

Conflict of interest The authors have no conflicts of interest to declare that are relevant to the content of this article.

Ethical approval The investigation procedures were approved by the University Hospitals of Cleveland Institutional Review Board.

Consent to participate A written consent was obtained from both subjects and their guardians/parents.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Armstrong N, Welsman J (1994) Assessment and interpretation of aerobic fitness in children and adolescents. *Exerc Sport Sci Rev* 22:435–476
- Armstrong N, Welsman J (2019a) Development of peak oxygen uptake from 11–16 years determined using both treadmill and cycle ergometry. *Eur J Appl Physiol* 119:801–812. <https://doi.org/10.1007/s00421-019-04071-3>
- Armstrong N, Welsman J (2019b) Clarity and confusion in the development of youth aerobic fitness. *Front Physiol* 10:1–7. <https://doi.org/10.3389/fphys.2019.00979>
- Armstrong N, Baxter-Jones ADG, Chia M et al (2007) Paediatric exercise physiology. Elsevier, Amsterdam
- Ashish N, Bammam MM, Cerny FJ et al (2015) The clinical translation gap in child health exercise research: a call for disruptive innovation. *Clin Transl Sci* 8:67–76. <https://doi.org/10.1111/cts.12194>
- Bar-Or O (1983) Pediatric sports medicine for the practitioner from physiologic principles to clinical applications. Springer, Berlin
- Barron A, Francis DP, Mayet J et al (2016) Oxygen uptake efficiency slope and breathing reserve, not anaerobic threshold, discriminate between patients with cardiovascular disease over chronic obstructive pulmonary disease. *J Am Coll Cardiol HF* 4:252–261. <https://doi.org/10.1016/j.jchf.2015.11.003>
- Bar-Yoseph R, Porszasz J, Radom-Aizik S et al (2019) The effect of test modality on dynamic exercise biomarkers in children, adolescents, and young adults. *Physiol Rep* 7:1–15. <https://doi.org/10.14814/phy2.14178>
- Beaver WL, Wasserman K, Whipp BJ (1986) A new method for detecting threshold by gas exchange anaerobic. *J Appl Physiol* 60:2020–2027
- Brooks GA (2012) Bioenergetics of Exercising Humans. *Compr Physiol* 2:537–562. <https://doi.org/10.1002/cphy.c110007>
- Cavagna GA, Kaneko M (1977) Mechanical work and efficiency in level walking and running. *J Physiol* 268:467–481
- Cavagna GA, Saibene FP, Margaria R (1964) Mechanical work in running. *J Appl Physiol* 19:249–256
- Cooper DM, Weiler-Ravell D, Whipp BJ, Wasserman K (1984) Aerobic parameters of exercise as a function of body size during growth in children. *J Appl Physiol Respir Environ Exerc Physiol* 56:628–634. <https://doi.org/10.1152/jappl.1984.56.3.628>
- Cooper DM, Leu SY, Galassetti P, Radom-Aizik S (2014) Dynamic interactions of gas exchange, body mass, and progressive exercise in children. *Med Sci Sports Exerc* 46:877–886. <https://doi.org/10.1249/MSS.0000000000000180>
- Cooper DM, Leu SY, Taylor-Lucas C et al (2016) Cardiopulmonary exercise testing in children and adolescents with high body mass index. *Pediatr Exerc Sci* 28:98–108. <https://doi.org/10.1123/pes.2015-0107>
- Daniels SR, Kimball TR, Morrison JA et al (1995) Effect of lean body mass, fat mass, blood pressure, and sexual maturation on left ventricular mass in children and adolescents: statistical, biological, and clinical significance. *Circulation* 92:3249–3254. <https://doi.org/10.1161/01.CIR.92.11.3249>
- DiBella JA, Johnson EM, Cabrera ME (2002) Ramped vs. standard bruce protocol in children: a comparison of exercise responses. *Pediatr Exerc Sci* 14:391–400
- Gosseye TP, Willems PA, Heglund NC (2010) Biomechanical analysis of running in weightlessness on a treadmill equipped with a subject loading system. *Eur J Appl Physiol* 110:709–728. <https://doi.org/10.1007/s00421-010-1549-9>
- Groen WG, Hulzebos HJ, Helders PJ, Takken T (2010) Oxygen uptake to work rate slope in children with heart, lung or muscle disease. *Int J Sport Med* 31:202–206
- Guazzi EM, Adams V, Conraads V et al (2012) EACPR/AHA scientific statement clinical recommendations for cardiopulmonary exercise. *Circulation* 126:2261–2274. <https://doi.org/10.1161/CIR.0b013e31826fb946>
- Guazzi M, Bandera F, Ozemek C et al (2017) Cardiopulmonary exercise Testing. *J Am Coll Cardiol* 70:1618–1636. <https://doi.org/10.1016/j.jacc.2017.08.012>
- Herman IP (2008) Physics of the human body: biological and medical physics, biomedical engineering. Metabolism: energy, heat, work, and power of the body. Springer, Berlin, p 857
- Jones AM, Poole DC (2013) Oxygen uptake kinetics in sport, exercise and medicine. Routledge, England
- Krahenbuhl G, Skinner J, Kohrt W (1985) Developmental aspects of maximal aerobic power in children. *Exerc Sport Sci Rev* 13:503–538
- Lai N, Nasca MM, Silva MA et al (2008) Influence of exercise intensity on pulmonary oxygen uptake kinetics at the onset of exercise and recovery in male adolescents. *Appl Physiol Nutr Metab* 33:107–117. <https://doi.org/10.1139/H07-154>
- Lai N, Tolentino-Silva F, Nasca MM et al (2012) Exercise intensity and oxygen uptake kinetics in African-American and Caucasian women. *Eur J Appl Physiol* 112:973–982. <https://doi.org/10.1007/s00421-011-2054-5>
- Lai N, Martis A, Belfiori A et al (2016) Gender differences in VO₂ and HR kinetics at the onset of moderate and heavy exercise intensity in adolescents. *Physiol Rep* 4(18):1–12. <https://doi.org/10.14814/phy2.12970>
- Malloggi C, Rota V, Catino L et al (2019) Three-dimensional path of the body centre of mass during walking in children: an index of neural maturation. *Int J Rehabil Res* 42:112–119. <https://doi.org/10.1097/MRR.0000000000000345>
- Margaria R (1978) Biomechanics and energetics of muscular exercise. Biomechanics and energetics of muscular exercise. Oxford University Press, Oxford, pp 172–172
- Minetti AE, Ardigo LP, Saibene F (1994a) The transition between walking and running in humans: metabolic and mechanical aspects at different gradients. *Acta Physiol Scand* 150:315–323
- Minetti AE, Saibene F, Ardigo LP et al (1994b) Pygmy locomotion. *Eur J Appl Physiol* 68:285–290
- Minetti AE, Ardigo LP, Saibene F et al (2000) Mechanical and metabolic profile of locomotion in adults with childhood-onset GH deficiency. *Eur J Endocrinol* 142:35–41
- Miyamura M, Honda Y (1973) Maximum cardiac output related to sex and age. *Jpn J Physiol* 23:645–656
- Paridon SM, Alpert BS, Boas SR et al (2006) Clinical stress testing in the pediatric age group: a statement from the American Heart Association council on cardiovascular disease in the young, committee on atherosclerosis, hypertension, and obesity in youth. *Circulation* 113:1905–1920. <https://doi.org/10.1161/CIRCULATIONAHA.106.174375>
- Pelà G, Crocimo A, Li Calzi M et al (2016) Sex-related differences in left ventricular structure in early adolescent non-professional athletes. *Eur J Prev Cardiol* 23:777–784. <https://doi.org/10.1177/2047487315608826>
- Poole DC, Jones AM (2017) Measurement of the maximum oxygen uptake VO₂max: VO₂peak is no longer acceptable. *J Appl Physiol* 122:997–1002. <https://doi.org/10.1152/japplphysiol.01063.2016>
- Popovic D, Guazzi M, Jakovljevic DG et al (2019) Quantification of coronary artery disease using different modalities of cardiopulmonary exercise testing. *Int J Cardiol* 285:11–13. <https://doi.org/10.1016/j.ijcard.2019.03.012>
- Porszasz J, Casaburi R, Somfay A et al (2003) A treadmill ramp protocol using simultaneous changes in speed and grade. *Med Sci Sports Exerc* 35:1596–1603. <https://doi.org/10.1249/01.MSS.0000084593.56786.DA>

- Rowland T, Goff D, Martel L, Ferrone L (2000) Influence of cardiac functional capacity on gender differences in maximal oxygen uptake in children. *Chest* 117:629–635
- Saibene F, Minetti AE (2003) Biomechanical and physiological aspects of legged locomotion in humans. *Eur J Appl Physiol* 88:297–316. <https://doi.org/10.1007/s00421-002-0654-9>
- Schepens B, Willems PA, Cavagna GA, Heglund NC (2001) Mechanical power and efficiency in running children. *Pflug Arch Eur J Physiol* 442:107–116. <https://doi.org/10.1007/s004240000511>
- Shaibi GQ, Ball GDC, Goran MI (2006) Aerobic fitness among Caucasian, African–American, and Latino youth. *Ethn Dis* 16:120–125
- Troutman WB, Barstow TJ, Galindo AJ, Cooper DM (1998) Abnormal dynamic cardiorespiratory responses to exercise in pediatric patients after fontan procedure. *J Am Coll Cardiol* 31:668–673. [https://doi.org/10.1016/S0735-1097\(97\)00545-7](https://doi.org/10.1016/S0735-1097(97)00545-7)
- Turley KR, Wilmore JH (1997a) Cardiovascular responses to treadmill and cycle ergometer exercise in children and adults. *J Appl Physiol* 83:948–957. <https://doi.org/10.1152/jappl.1997.83.3.948>
- Turley KR, Wilmore JH (1997b) Cardiovascular responses to submaximal exercise in 7- to 9-yr-old boys and girls. *Med Sci Sports Exerc* 29:824–832. <https://doi.org/10.1097/00005768-199706000-00014>
- Turley KR, Rogers DM, Harper KM et al (1995) Maximal treadmill versus cycle ergometry testing in children: differences, reliability, and variability of responses. *Pediatr Exerc Sci* 7:49–60
- Whipp BJ, Wasserman K (1969) Efficiency of muscular work. *J Appl Physiol* 26:644–648. <https://doi.org/10.1152/jappl.1969.26.5.644>
- Willems PA, Cavagna GA, Heglund NC (1995) External, internal and total work in human locomotion. *J Exp Biol* 198:379–393

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.