

# De Novo Whole-Genome Sequence of *Micromonospora carbonacea* JXNU-1 with Broad-Spectrum Antimicrobial Activity, Isolated from Soil Samples

Yun Jiang, Yun-hong Huang, Zhong-er Long

College of Life Science, Jiangxi Normal University, Nanchang, Jiangxi, Zhōngguó, China

***Micromonospora carbonacea* JXNU-1 is an actinomycete with broad-spectrum antimicrobial activity, isolated from soil samples from the farmland in the area of Yaohu Lake in Nanchang, China. Here, we report the whole-genome sequence of *M. carbonacea* JXNU-1.**

Received 8 February 2015 Accepted 23 February 2015 Published 9 April 2015

**Citation** Jiang Y, Huang Y-H, Long Z-E. 2015. *De novo* whole-genome sequence of *Micromonospora carbonacea* JXNU-1 with broad-spectrum antimicrobial activity, isolated from soil samples. *Genome Announc* 3(2):e00174-15. doi:10.1128/genomeA.00174-15.

**Copyright** © 2015 Jiang et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Zhong-er Long, longzhonger@163.com.

*Micromonospora carbonacea* JXNU-1 is an actinomycete with broad-spectrum antimicrobial activity. The strain was initially isolated from soil samples from farmland in the area of Yaohu Lake in Nanchang, China (1). The single antibiotic component from the strain showed broad-spectrum antimicrobial activity and featured nucleosides (2), unlike the other strain of *M. carbonacea* (3–7).

The genomic DNA from *M. carbonacea* JXNU-1 was obtained from mycelia cultured in liquid supplemented minimal medium (ISP<sub>2</sub> medium) under shaken conditions at 200 rpm at 28°C for 5 days. The Illumina HiSeq 2500 and Illumina MiSeq platforms were used to construct 2 different genomic DNA libraries according to the manufacturer's instructions. Long-insert (about 5-kb) libraries were sequenced by the paired-end mode using the Illumina HiSeq 2500, and the short-insert (500-bp) libraries were sequenced by the paired-end mode with Illumina MiSeq. DNA sequencing resulted in 3,487 Mb raw reads, of which 2,393 Mb reads passed stringent quality filters and were used to create the final assembly using the SOAPdenovo alignment tool (version 2.0) (8), with multiplex PCR used to close the gaps (9, 10). The assembly consists of 8 scaffolds, including 9 large contigs (sum, 7.63 Mb; *N*<sub>50</sub>, 1.11 Mb; maximum length, 2.31 Mb).

The whole genome of *M. carbonacea* JXNU-1 contains a single circular chromosome of 7,635,725 bp, with an average G+C content of 73.85%. A total of 6,444 coding sequences (CDs) were identified by GeneMarkS (11, 12). In addition, there are 27 interspersed repeats (IRs) with 2,731 bp and 4,100 tandem repeats (TRs) with 196,933 bp in the genome, respectively. We also used rRNAmmer (13), tRNAscan (14), and Rfam to identify noncoding RNAs; 6 rRNAs, 51 tRNAs, and 23 sRNAs were found in genome, respectively. In addition, 16 genomic islands (GIs) and 22 clustered regularly interspaced short palindromic repeats (CRISPR) were predicted using IslandPath-DIOMB software and CRISPRFinder (<http://crispr.u-psud.fr/>), respectively.

**Nucleotide sequence accession numbers.** This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank un-

der the accession number [JX SX00000000](https://doi.org/10.1101/000000). The version described in this paper is the first version, JX SX01000000.

## ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (grants 31160029 and 31360018) and the Natural Science Foundation of Jiangxi Province of China (grant 20132BAB204007).

## REFERENCES

- Long ZE, Zhu YJ, Huang YH, Fu XQ. 2008. Isolation and identification of a strain of *Micromonospora* with broad-spectrum antimicrobial activity. *Microbiol China* 35:378–384. <http://dx.doi.org/10.3969/j.issn.0253-2654.2008.03.012>.
- Long ZE, Zhu YJ, Huang YH, Fu XQ. 2008. Studies on the isolation and purification of antibiotic from the fermentation broth of *Micromonospora carbonacea* JXNU-1 and its physical-chemical properties. *Microbiol China* 35:1450–1454. <http://dx.doi.org/10.3969/j.issn.0253-2654.2008.09.019>.
- Weinstein MJ, Luedemann GM, Oden EM, Wagman GH. 1964. Everninomicin, a new antibiotic complex from *Micromonospora carbonacea*. *Antimicrob Agents Chemother* 10:24–32.
- Luedemann GM, Brodsky B. 1964. *Micromonospora carbonacea* sp. n., an everninomicin producing organism. *Antimicrob Agents Chemother* 10:47–52.
- Herzog HL, Meseck E, Delorenzo S, Murawski A, Charney W, Rosselet JP. 1965. Chemistry of antibiotics from *Micromonospora*: III. Isolation and characterization of everninomicin D and everninomicin B. *Appl Microbiol* 13:515–520.
- Sanders WE, Sanders CC. 1974. Microbiological characterization of everninomicins B and D. *Antimicrob Agents Chemother* 6:232–238. <http://dx.doi.org/10.1128/AAC.6.3.232>.
- Puar MS, Chan TM, Hegde V, Patel M, Bartner P, Ng KJ, Pramanik BN, MacFarlane RD. 1998. A novel thiostrepton from *Micromonospora carbonacea*. *J Antibiot* 51:221–224. <http://dx.doi.org/10.7164/antibiotics.51.221>.
- Bolotin A, Quinquis B, Ehrlich SD, Sorokin A. 2012. Complete genome sequence of *Lactococcus lactis* subsp. cremoris A76. *J Bacteriol* 194:1241–1242. <http://dx.doi.org/10.1128/JB.06629-11>.
- Fukao M, Oshima K, Morita H, Toh H, Suda W, Kim SW, Suzuki S, Yakabe T, Hattori M, Yajima N. 2013. Genomic analysis by deep sequencing of the probiotic *Lactobacillus brevis* KB290 harboring nine plasmids reveals genomic stability. *PLoS One* 8:e60521. <http://dx.doi.org/10.1371/journal.pone.0060521>.
- Sorokin A, Lapidus A, Capuano V, Galleron N, Pujic P, Ehrlich SD.

1996. A new approach using multiplex long accurate PCR and yeast artificial chromosomes for bacterial chromosome mapping and sequencing. *Genome Res* 6:448–453. <http://dx.doi.org/10.1101/gr.6.5.448>.
11. Stanke M, Diekhans M, Baertsch R, Haussler D. 2008. Using native and syntenically mapped cDNA alignments to improve *de novo* gene finding. *Bioinformatics* 24:637–644. <http://dx.doi.org/10.1093/bioinformatics/btn013>.
  12. Besemer J, Lomsadze A, Borodovsky M. 2001. GeneMarkS: a self-training method for prediction of gene starts in microbial genomes. Implications for finding sequence motifs in regulatory regions. *Nucleic Acids Res* 29:2607–2618. <http://dx.doi.org/10.1093/nar/29.12.2607>.
  13. Lagesen K, Hallin P, Rødland EA, Stærfeldt H-H, Rognes T, Ussery DW. 2007. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 35:3100–3108. <http://dx.doi.org/10.1093/nar/gkm160>.
  14. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 25:955–964. <http://dx.doi.org/10.1093/nar/25.5.0955>.