


Pueraria mirifica, an estrogenic tropical herb, unveiled the severity of Type 1 LQTS caused by *KCNQ1-T587M*

Asami Kashiwa MD^{1,2}  | Yukio Hosaka MD, PhD² | Kazuyoshi Takahashi MD² | Seiko Ohno MD, PhD³ | Yuko Wada MD, PhD^{4,5} | Takeru Makiyama MD, PhD¹ | Hirotaka Oda MD, PhD² | Minoru Horie MD, PhD⁵

¹Department of Cardiovascular Medicine, Kyoto University Graduate School of Medicine, Kyoto, Japan

²Department of Cardiovascular Medicine, Niigata City General Hospital, Niigata, Japan

³Department of Bioscience and Genetics, National Cerebral and Cardiovascular Center, Osaka, Japan

⁴Vanderbilt Center for Arrhythmia Research and Therapeutics, Division of Clinical Pharmacology, Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA

⁵Department of Cardiovascular Medicine, Shiga University of Medical Science, Shiga, Japan

Correspondence

Asami Kashiwa, Department of Cardiovascular Medicine, Kyoto University Graduate School of Medicine, 54, Shogoin Kawahara-cho, Sakyo-ku, Kyoto 606-8507, Japan.

Email: asamika@kuhp.kyoto-u.ac.jp

Keywords: acquired long QT syndrome, congenital long QT syndrome, drug-induced long QT syndrome, estrogens, genetics

1 | INTRODUCTION

One-third of patients with apparent acquired long QT syndrome (LQTS) carry a mutation in, at least, one of the three major LQTS-related genes including *KCNQ1*, which is responsible for type 1 LQTS.¹ We here report a case of acquired LQTS carrying a *KCNQ1-T587M* mutation,²⁻⁴ which presented a very severe phenotype triggered by *Pueraria mirifica*. The compound is derived from a tropical plant containing estrogen-like substances and potentially modifies cardiac repolarization.⁵⁻⁷

2 | CASE REPORT

A 24-year-old woman developed sudden onset of pre-syncope while she was resting at home during the daytime and called emergency medical services. In the ambulance, she collapsed and ventricular fibrillation (VF) was detected. Immediate electrical shock terminated VF. Upon arrival at the hospital, her ECG showed a marked QT prolongation (corrected QT [QTc] = 674 ms) and notched T waves (Figure 1A). Her laboratory examination

showed no electrolyte abnormality (K: 3.8 mEq/L, Ca: 8.9 mg/dL, Mg: 1.7 mg/dL). Soon after her arrival, we injected magnesium sulfate. There was neither medical history of syncope nor family history of cardiac disease or sudden death. Further history taking revealed that she had been taking *Pueraria mirifica*, an estrogenic herb containing miroestrol and deoxymiroestrol, for a week before hospitalization without any other QT prolonging agents. Her menstrual cycle was regular, and the event occurred 7 days before her menstruation (luteal phase). After admission, the estrogen supplement was discontinued.

On day 2, her ECG still presented prolonged QTc interval (QTc = 764 ms) (Figure 1B) and she developed repetitive torsade de pointes (TdP) on emotional agitation (Figure 2A). After oral mexiletine (300 mg/kg/d) was started, QTc interval shortened to 631 ms, but TdP was not completely suppressed. A total of 14 electrical shocks were required. Mild conscious sedation finally suppressed the TdP storms. QTc gradually shortened, and on day 4, mexiletine and sedation were discontinued (QTc = 544 ms). On day 8, QTc interval further shortened to 439 ms (Figure 2B). Epinephrine stress test⁸ was conducted and revealed drastic QTc prolongation from 491 to 683 ms with marked T-wave abnormality (Figure 3A,B).

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Journal of Arrhythmia* published by John Wiley & Sons Australia, Ltd on behalf of Japanese Heart Rhythm Society.

After obtaining the informed consent for genetic analysis approved by our institutional review board, we performed screening for *KCNQ1* and *KCNH2* by denatured high-performance liquid chromatography (WAVE system; Transgenomic Omaha) in addition to the analysis including all LQTS-related genes using HaloPlex HS custom panel (Agilent Technology)⁹ and identified a heterozygous *KCNQ1* variant: c.1760C>T;p.Thr587Met (T587M). There were no

pathogenic variants in other genes. Her relatives did not consent for genetic testing. After taking bisoprolol (0.06 mg/kg/d), she still presented marked QT prolongation during exercise. Considering her high risk of arrhythmia, we implanted an ICD after a careful discussion with the patient.

Five years after the first event, the patient has been free from recurrence of arrhythmic events.

FIGURE 1 Twelve-lead surface ECGs of the patient. A, ECG on admission showing long QT interval. The corrected QT interval (QTc) = 674 ms. (B) ECG on day 2 after admission. QTc = 764 ms




FIGURE 2 Twelve-lead surface ECGs of the patient on day 2 and day 8. A, ECG demonstrating torsade de pointes (TdP) on day 2. B, ECG on day 8 after admission showing normalized QT interval. QTc = 439 ms





FIGURE 3 Twelve-lead ECGs at epinephrine stress test. A, Baseline (QTc = 491 ms). B, Steady-state (QTc = 683 ms)



3 | DISCUSSION

We report a concealed type 1 LQTS case (*KCNQ1-T587M*) that was unmasked by the ingestion of *Pueraria mirifica*, a commercially available rejuvenating supplement, containing estrogen-like substances.^{10,11} Estrogen was reported to prolong action potential duration and QT interval.⁵⁻⁷ One of the proposed mechanisms is that 17 β -estradiol (E2) can suppress I_{Kr} in a receptor-independent manner.⁷ However, the suppression level was so small that its impact on baseline QTc interval was not prominent in the presence of repolarization reserve.¹² In fact, to the best of our knowledge, there are no reports of *Pueraria mirifica*-induced ventricular arrhythmias, and it is unlikely to cause dramatic changes in ECGs in healthy individuals.

Meanwhile, its estrogenic activity played a critical role in our patient by unveiling the most malignant phenotype of *KCNQ1-T587M* mutation, leading to refractory and repetitive TdP. The mutation was first reported to cause trafficking defects of $K_v7.1$ encoded by *KCNQ1*.² Later, it has been shown that the *KCNQ1-T587M* failed to function as a chaperone that transports hERG proteins (responsible for I_{Kr}) to the plasma membranes.^{13,14} These multiple mechanisms that modulate both I_{Kr} and I_{Ks} may underlie malignant phenotypes which were often seen in the *KCNQ1-T587M* carriers. Therefore, estrogenic substances predisposed the patient to life-threatening arrhythmias by the decrease of repolarization reserve caused by *KCNQ1-T587M*.

4 | CONCLUSION

We report the first case of severe phenotype of type 1 LQTS unveiled by *Pueraria mirifica*.

CONFLICT OF INTEREST

The authors declare no conflict of interests for this article.

ORCID

Asami Kashiwa  <https://orcid.org/0000-0001-9562-8981>

REFERENCES

- Itoh H, Crotti L, Aiba T, Spazzolini C, Denjoy I, Fressart V, et al. The genetics underlying acquired long QT syndrome: impact for genetic screening. *Eur Heart J*. 2016;37(18):1456–64.
- Yamashita F, Horie M, Kubota T, Yoshida H, Yumoto Y, Kobori A, et al. Characterization and subcellular localization of *KCNQ1* with a heterozygous mutation in the C terminus. *J Mol Cell Cardiol*. 2001;33(2):197–207.
- Furushima H, Chinushi M, Sato A, Aizawa Y, Kikuchi A, Takakuwa K, et al. Fetal atrioventricular block and postpartum augmentative QT prolongation in a patient with long-QT syndrome with *KCNQ1* mutation. *J Cardiovasc Electrophysiol*. 2010;21(10):1170–3.
- Chen S, Zhang L, Bryant RM, et al. *KCNQ1* mutations in patients with a family history of lethal cardiac arrhythmias and sudden death. *Clin Genet*. 2003;63(4):273–82.
- Hara M, Danilo P Jr, Rosen MR. Effects of gonadal steroids on ventricular repolarization and on the response to E4031. *J Pharmacol Exp Ther*. 1998;285(3):1068–72.
- Saito T, Ciobotaru A, Bopassa JC, Toro L, Stefani E, Eghbali M. Estrogen contributes to gender differences in mouse ventricular repolarization. *Circ Res*. 2009;105(4):343–52.
- Kurokawa J, Tamagawa M, Harada N, et al. Acute effects of oestrogen on the guinea pig and human *IKr* channels and drug-induced prolongation of cardiac repolarization. *J Physiol*. 2008;586(12):2961–73.
- Shimizu W, Noda T, Takaki H, Kurita T, Nagaya N, Satomi K, et al. Epinephrine unmasks latent mutation carriers with LQT1 form of congenital long-QT syndrome. *J Am Coll Cardiol*. 2003;41(4):633–42.
- Ohno S, Ozawa J, Fukuyama M, Makiyama T, Horie M. An NGS-based genotyping in LQTS; minor genes are no longer minor. *J Hum Genet*. 2020;65(12):1083–91.
- Cain JC. Miroestrol: an oestrogen from the plant *Pueraria mirifica*. *Nature*. 1960;188:774–7.
- Malavijitnond S. Medical applications of phytoestrogens from the Thai herb *Pueraria mirifica*. *Front Med*. 2012;6:8–21.
- Kurokawa J, Kodama M, Clancy CE, Furukawa T. Sex hormonal regulation of cardiac ion channels in drug-induced QT syndromes. *Pharmacol Ther*. 2016;168:23–8.
- Biliczki P, Girmatsion Z, Brandes RP, et al. Trafficking-deficient long QT syndrome mutation *KCNQ1-T587M* confers severe clinical phenotype by impairment of *KCNH2* membrane localization: evidence for clinically significant *IKr-IKs* alpha-subunit interaction. *Heart Rhythm*. 2009;6(12):1792–801.
- Wu J, Sakaguchi T, Takenaka K, Toyoda F, Tsuji K, Matsuura H, et al. A trafficking-deficient *KCNQ1* mutation, T587M, causes a severe phenotype of long QT syndrome by interfering with intracellular hERG transport. *J Cardiol*. 2019;73(5):343–50.

How to cite this article: Kashiwa A, Hosaka Y, Takahashi K, et al. *Pueraria mirifica*, an estrogenic tropical herb, unveiled the severity of Type 1 LQTS caused by *KCNQ1-T587M*. *J Arrhythmia*. 2021;37:1114–1116. <https://doi.org/10.1002/joa3.12576>