

OPEN

Neurologic Manifestations of Severe Respiratory Viral Contagions

Christopher P. Robinson, DO, MS and Katharina M. Busl, MD, MS

Objectives: Endemic and pandemic viral respiratory infections have recently emerged as a critical topic of investigation given the recent severe acute respiratory syndrome coronavirus-2 outbreak. Data from such outbreaks indicate that severe systemic comorbidities including acute neurologic illness are associated with illness and lead to significant outcome differences. Herein, we will discuss the neurologic manifestations of severe viral respiratory infections including coronavirus, influenza, respiratory syncytial virus, metapneumovirus, and enterovirus.

Data Sources: PubMed and EMBASE were searched by two independent investigators up to March 2020.

Study Selection: Data selection included preclinical and clinical studies detailing neurologic manifestations of viral respiratory infections.

Data Extraction and Synthesis: Two independent investigators reviewed and extracted the data.

Conclusions: Neurologic manifestations including seizures, status epilepticus, encephalitis, critical illness neuromyopathy, acute disseminated encephalomyelitis, acute necrotizing encephalitis, Guillan-Barré syndrome, transverse myelitis, and acute flaccid myelitis have all been associated with severe viral respiratory infections. Having an understanding of the direct neurotropism of such viruses is imperative to understanding pathogenesis, clinical presentation, and potential treatment paradigms aimed at improving morbidity and mortality.

Key Words: encephalitis; meningitis; neurology; respiratory; seizure; virus

Novel viral respiratory infections are an emerging global public health emergency with high rates of morbidity and mortality. Some of the most common viral pathogens known to effect the human population with respiratory

infections include coronavirus, influenza, human metapneumovirus (hMPV), respiratory syncytial virus (RSV), and enterovirus (1). Over the past 20 years, pandemics such as severe acute respiratory syndrome coronavirus (SARS Co-V), Middle Eastern respiratory syndrome (MERS), influenza (H1N1 and H5N1), and currently the coronavirus SARS Co-V2 with coronavirus disease 2019 (COVID-19) have placed a strain on the healthcare systems and societies. In addition to the costs associated with clinical care and the severe respiratory syndromes that such diseases may manifest with, severe and sometimes irreversible neurologic complications can arise. Although manifestation such as mild upper respiratory symptoms are most common, more severe manifestations such as bronchitis, pneumonia and, acute respiratory distress syndrome (ARDS) can occur as well as can neurologic manifestations varying from self-limited aseptic meningitis to fulminant status epilepticus (2–7). Severe neurologic complications can include encephalitis, acute disseminated encephalomyelitis (ADEM), cerebellitis, Guillan-Barré syndrome (GBS), acute flaccid myelitis, and necrotizing encephalopathy as well as symptomatic manifestation with status epilepticus (Table 1) (8). In this review, we will discuss the neural invasion of respiratory viruses and the typical neurologic complications of each individual respiratory infection as well as the routes by which CNS transmission occurs.

NEURAL INVASION OF RESPIRATORY VIRUSES

Invasion of viral respiratory pathogens into the CNS requires at least a degree of neurotropism; that is, the ability to invade and survive within neural tissue. The blood-brain barrier (BBB), the essential homeostatic bridge to neural function, plays a vital role in neural protection and molecular passage of such microorganisms. Composed of endothelial cells, pericytes, astrocytes, microglia, and extracellular matrix, the BBB controls permeability and acts to control quiescence of the immune system (9). Disruption of such factors through either inflammation or vascular breakdown leads to increased BBB permeability and potentiates unwanted CNS effects. Several modes of CNS transmission by viral pathogens exist including transcellular, paracellular, and retrograde axonal transport along sensory and olfactory nerves (10). Hematogenous spread of viral pathogens, or viremia, is the common mechanism allowing for cellular migration of the virus. Transcellular migration occurs due to

Both authors: Departments of Neurology and Neurosurgery, University of Florida, Gainesville, FL.

Copyright © 2020 The Authors. Published by Wolters Kluwer Health, Inc. on behalf of the Society of Critical Care Medicine. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Crit Care Expl 2020; 2:e0107

DOI: 10.1097/CCE.000000000000107

TABLE 1. Associated Neurologic Manifestations of Severe Viral Respiratory Infections

Respiratory Viral Pathogens	Associated Neurologic Manifestations
Human coronavirus (human variants coronavirus A/B, SARS, Middle Eastern respiratory syndrome, SARS coronavirus 2)	Encephalitis
	Seizures
	Refractory status epilepticus
	ADEM
	GBS
	Leukoencephalopathy
	Critical illness neuromyopathy
Influenza virus (flu A, flu B, H1N1, H5N1)	Febrile seizures
	Acute necrotizing encephalitis
	Encephalitis/encephalopathy
	Reye's syndrome
	ADEM
	GBS
	Seizures
	Myelitis
	Mutism
Respiratory syncytial virus/orthopneumovirus	Encephalitis
	Seizures
	Status epilepticus
	Central apnea
Human metapneumovirus	Encephalitis
	Focal seizures
	Status epilepticus
Enterovirus (D68, D71)	Acute flaccid myelitis
	Aseptic meningitis
	Viral encephalitis

ADEM = acute disseminated encephalomyelitis, GBS = Guillan-Barré syndrome, SARS = severe acute respiratory syndrome.

invasion of either host cells or phagocytic macrophages to overcome the BBB or the blood-cerebrospinal fluid (CSF) barrier, whereas paracellular migration occurs with pathogen invasion of BBB tight junctions (10). Axonal transport of viral pathogens can occur via sensory, motor, or olfactory nerves. Axonal transport occurs via adherence of motor proteins responsible for retrograde neuronal transport in the periphery as well as in the olfactory nerve which has direct communication with the nasal endothelium where respiratory viral pathogens preferably infect (11). Although each individual virus may have its own predilection for neuroinvasion, there is likely a combination or

additive pathognomonic mode of transmission that occurs during viremia (12).

CORONAVIRUS

Coronaviridae are a family of RNA viruses that were first isolated in the 1960s from patients with upper respiratory symptoms (13). The virus infects both birds and mammals. Human variants of coronavirus (HCoV) isolated as alphacoronavirus and betacoronavirus are endemic worldwide and generally produce the classical symptoms of HCoV including rhinitis, pharyngitis, laryngitis, bronchitis, and otitis (14). In 2002, a novel coronavirus identified as SARS Co-V was isolated, propagated by zoonotic transmission from bats through and intermediate reservoir (15). Unlike typical HCoV, SARS Co-V was extremely virulent and complicated by high rates of ARDS, multiple organ dysfunction syndrome, and mortality in up to 10% of cases (16, 17). In 2012, a second novel coronavirus, MERS Co-V evolved from zoonotic transmission causing severe respiratory symptoms with a mortality rate of 35% (18). In 2019, a third novel coronavirus, SARS Co-V2, has emerged. This virus has rapidly disseminated to become a pandemic outbreak in early 2020. It shares highly homologous sequence with SARS Co-V and causes acute, highly lethal pneumonia (COVID-19) with clinical symptoms similar to that reported for SARS Co-V and MERS Co-V (19). One difference is that upper respiratory tract signs and symptoms are rare, indicating that the target cells of SARS CoV-2 may be located in the lower airway (20).

The innate virility of Co-V has tropism for both the respiratory tract and the CNS. The neuroinvasive and neurotropic potential of the virus has been well described in studies isolating viral RNA in brain parenchyma outside the blood vessels and is accepted as a vector leading to acute or chronic neurologic disease (5, 21). Viral particles are almost exclusively discovered in neuronal cells, and hematologic or lymphatic routes of viral dissemination are postulated in addition to direct invasion of peripheral nerve terminals and subsequent trans-synaptic transfer (21–23). For acute neurologic disease, the neuropathogenicity of Co-V has been described with associated cases of encephalitis, refractory status epilepticus, GBS, and ADEM during active infection with confirmatory CSF results (24–28). In addition to acute virulence, chronic and persistent infection of human oligodendrocytes and glial cells by Co-V has been demonstrated and proposed to be involved in neuropathologic changes (29, 30). Specifically, an association with development of multiple sclerosis or other demyelinating diseases has been posed (5). Similarly, several acute neurologic manifestations have been reported for SARS Co-V and MERS Co-V, including critical illness neuromyopathy, considered secondary to direct nerve invasion, ADEM, and leukoencephalopathy (31, 32). To date, no rigorous evidence exists describing neurologic involvement of the novel SARS Co-V2. Extrapolating knowledge of neuroinvasion from Co-V in general, it has been proposed that SARS Co-V's neurotropism may play a role in the severe respiratory symptoms, by invasion of medullary neurons and interference with functions of nucleus ambiguus and nucleus tractus solitarius, potentially causing central apnea, and lack of feedback from lung

and respiratory tract receptors (19). Furthermore, headaches were reported by 8% in one of the published series from China (19). In a retrospective case series (not yet peer reviewed) of patients hospitalized with COVID-19 in Wuhan, China, the authors describe that of 214 cases, 36% had neurologic manifestations, with higher prevalence in more severe cases (33). CNS symptoms were present in 25%, with dizziness and headache being the most common, whereas peripheral neuromuscular symptoms were present in 19%. Severe manifestations with impaired consciousness or cerebrovascular disease was also reported, but much less common (33). Acute management strategies for such presentations should be individualized for each underlying manifestation, but overall remains largely supportive and symptomatic. Several drugs have been proposed to be repurposed for the new coronavirus (34). Effectiveness of corticosteroids remains unclear and likely dependent on the phase of illness during which steroids are given (35). Currently, several medications are being investigated for their efficacy and safety for COVID-19 pneumonia, including complement mediators and nucleotide analogues (36), but no specific treatment or prevention of neurologic manifestations exist, other than primary prevention of exposure and infection.

INFLUENZA

The influenza viruses, formally classified as orthomyxoviridae, are a family of RNA viruses commonly implicated in severe upper respiratory syndromes. There are three genera: influenza virus A, influenza virus B, and influenza virus C, of which, type A is the most virulent human pathogen and causes the most severe form of the disease (37). Diverse phenotypic symptoms are common, from pyrexia, headache, pharyngitis/laryngitis, to ARDS. Influenza infects upwards of 10% of the population annually and has a mortality rate of 4–8.8 deaths per 100,000 persons (38, 39). Individual subtypes of the influenza virus are determined by the structural proteins hemagglutinin (H) and neuraminidase (N). Antigenic shifts that occur within these proteins during zoonotic transmission have led to several pandemics including H1N1 and H5N1 (40, 41). Phenotypic expression within these subtypes tends to be more severe with higher rates of severe lower respiratory infection, ARDS, and death. In addition to the classical respiratory symptoms associated with influenza, a variety of extrapulmonary complications including neurologic, cardiovascular, ocular, hematologic, renal, and hepatic can occur (42).

The neurologic manifestations of influenza have been extensively described (43), vary widely, and have important clinical and prognostic implications. Although generally more prevalent in children, encephalitis, GBS, ADEM, Reye's syndrome, seizures, myelitis, mutism, and Parkinsonian symptoms have been reported in both adult and pediatric populations (42). Historically, febrile seizure is the most frequently encountered influenza-associated CNS complication, with one in five children hospitalized with influenza affected (44). Thirty percent of pediatric deaths related to H1N1 in the United Kingdom presented with encephalopathy and seizures, and neurologic sequelae are among the most severe consequences of the pandemic influenza A H1N1 in survivors (45). The epidemic influenza A (H3N2) strain was associated with a surge in cases of encephalitis/encephalopathy in Japan, Europe,

and the United States (43). One of the best characterized and severe neurologic complications of influenza viremia is acute necrotizing encephalopathy (ANE). Although originally described in the pediatric population, ANE also can affect adults and presents with rapid deterioration in consciousness, symmetric bithalamic, cortical, or infratentorial lesions on MRI and often absence of CSF pleocytosis (see case example in Fig. 1) (46–50). Presence of hemorrhage and tissue loss on imaging in conjunction with CSF are poor prognostic indicators (51). Secondary complications include seizures and status epilepticus. The mortality rate of ANE is upwards of 30% with a significant proportion of patients having long-term neurologic sequelae following recovery (52). Progression to brain death is also a documented complication of ANE (53). In a study of influenza A and B strains in Japan, elevated serum aspartate aminotransferase and creatine phosphokinase and thrombocytopenia ($<50,000$ platelets/ μL) appeared to correlate with unfavorable outcome in influenza-associated encephalopathy (54). Another commonly recognized complication of influenza is GBS, a para- or postinfectious demyelinating polyneuropathy resulting in flaccid quadriplegia and cranial nerve deficits (55, 56). In addition to active infection, the influenza vaccine has also been associated with GBS, although a recent large case-series refutes this evidence (57). Reye's syndrome has also been highly implicated as a parainfectious complication of the influenza virus, especially in children (58). Defined as acute encephalopathy and fatty liver infiltration, Reye's patients generally have a poor prognosis with high mortality. Additional but rare neurologic complications resulting from viral infiltration of specific neuronal structures (basal ganglia, hypothalamus, myelinated axons) include Klein-Levin syndrome, Parkinsonism, transverse myelitis, and ADEM (44).

The exact pathogenesis of the neurologic manifestations of influenza are not entirely clear. Data showing ANE but absence of the influenza virus in the CNS have led to conclude that the inflammatory insult outside the CNS is the trigger for ANE (59), whereas others have hypothesized that disruption of the BBB in the presence of systemic hypercytokinemia could be responsible for inducing necrotic brain lesions (44). Historically, occurrence of Parkinsonian symptoms and "encephalitis lethargica" was reported after the 1918 influenza pandemic (60). However, although there was documented presence of influenza A antigens in patients with "encephalitis lethargica" (61), viral RNA could not be detected in brains of postencephalitic Parkinsonian patients, questioning the exact role of the influenza virus in the genesis of Parkinsonism (62), and calling for long-term follow-up studies among survivors of H5N1-infected individuals to further elucidate the possible direct destructive effect of influenza to the neurons of the substantia nigra.

The mainstay of treatment for influenza infection remains supportive. Patients in high-risk groups are recommended to receive antiviral therapy to shorten illness duration and decrease the occurrence of secondary complications (63), but not all data are affirming strong efficacy for complication prevention (64, 65). For ANE, steroids given within 24 hours after the onset were associated with better outcome of children without brainstem lesions (51) in one series. The use of salicylates should be avoided in children and adolescents due to the association with Reye syndrome.




Figure 1. MRI of a 56-yr-old female who presented with 2 d of shortness of breath and upper respiratory congestion, followed by onset of rapidly progressive confusion culminating in seizures and then obtundation. She tested positive for influenza A antigen. Cerebrospinal fluid analysis showed a mild pleocytosis of 13 (82% neutrophils), and polymerase chain reaction testing for other viral encephalitides remained negative. She was treated with oseltamivir immediately upon confirmation of positive influenza testing. MRI at admission showed restricted diffusion and susceptibility along with Fluid-attenuated inversion recovery (FLAIR) hyperintensity, without enhancement, in bilateral thalami (**A–C**) and bilateral mesial temporal lobes (**D–F**) (A+D, FLAIR, B+E, diffusion weighted imaging, C+F, susceptibility weighted imaging sequences).

RESPIRATORY SYNCYTIAL VIRUS

Human RSV is an RNA virus of the pneumoviridae family. Accordingly, the virus has been renamed orthopneumovirus, but for the clinical context of this article, we will use “RSV” (66). Bronchiolitis, pneumonia, and bronchopneumonia from RSV largely infects the pediatric population and is responsible for a majority of PICU admissions related to pulmonary disease (67). Immunocompromised individuals are also susceptible to infection, and emerging literature suggests a large infection burden within adult populations (68). Extrapulmonary manifestations of RSV are well reported and organ systems typically involved include neurologic, cardiovascular, and hepatic (69).

Neurologic manifestations of acute RSV infection include encephalitis, seizures, status epilepticus, and central apnea and are reported in up to 39% of patients (70, 71). Application of very stringent criteria for diagnosis of neurologic disease in the largest available series yielded, however, only a prevalence of 1.2% (71). Seizure frequency among pediatric patients with RSV was 6.6% in one of the pediatric series (70) but only made for 0.7% of all

patients and represented the majority of neurologic manifestations in other large series (6, 71). Another important neurologic complication following RSV infection is central apnea, which is found in 16%–21% of patients admitted to the ICU (72). Central apnea at admission is a strong predictor for recurrent apneic events and clinical predictor for mechanical ventilation (72). Studies have shown an abnormal laryngeal chemoreflex in patients infected with RSV, implicating CNS pathology as a culprit (73). The exact mechanism of CNS penetration and pathologic response of RSV are unknown; however, CSF samples were positive for the virus in 50% of patients with seizures in the above series (70). With reference to the adult population, scant evidence exists in the literature implicating RSV as a neuropathogen. Treatment for the neurologic sequelae of RSV is generally supportive and symptomatic.

HUMAN METAPNEUMOVIRUS

Human metapneumovirus (hMPV) is an RNA virus of the paramyxovirus family responsible for upper respiratory infection in all age groups, with more severe infections such as bronchiolitis

and pneumonia affecting the pediatric and immunocompromised populations (74). Since its discovery in 2001, extrapulmonary neurologic manifestations of hMPV have been reported including encephalitis, focal seizures, and status epilepticus or even severe fatal encephalitis mostly in the pediatric but also in the adult population (75–77). The overall prevalence of neurologic complications seems rare, with exact prevalence unknown and most data coming from small case series or single case reports (75). Among 205 pediatric patients referred to the California encephalitis project between 2004 and 2006, hMPV was detected in nasopharyngeal swabs of five patients (6). Compared with infection with RSV, the frequency of seizures is reported to be higher (6.3% vs 0.7%) (6). In a literature review of pediatric cases of hMPV infection complicated by encephalitis, seizures occurred in 10 of 13 cases (75). Infection likely occurs due to direct CNS invasion, as hMPV has been found in brain tissue after autopsy (78). MRI findings can range from normal parenchymal structures to having various degrees of white matter involvement. CSF analysis can range from normal to showing varying degrees of pleocytosis and inconsistent polymerase chain reaction positivity (6). Treatment for the secondary neurologic complications of hMPV remains largely supportive.

ENTEROVIRUS D68 AND 71

Enteroviruses (EV) D68 and 71 (EV68, EV71) are RNA viruses belonging to the picornaviridae family. Initially discovered in 1968, EV68 was described as a nonpolio enterovirus that infected the upper respiratory tract (79). Isolated in 1969, EV71 was a novel enterovirus found causes hand foot and mouth syndrome (80). A majority of the enterovirus family cause mild symptoms; however, certain strains with specific neurotropism such as EV68 and EV71 have been implicated in severe secondary neurologic sequelae of infection (80–82). EV71 specifically has been demonstrated to cause both aseptic meningitis and viral encephalitis (83, 84). More notably, however, are the recent spikes in cases of acute flaccid paralysis or acute flaccid myelitis (AFM) affecting the pediatric population worldwide (85). Both EV68 and EV71 can result in nonpolio enterovirus paralysis, with EV68 being causative during the most recent outbreak (82, 86, 87). A diagnostic classification for AFM has been proposed for epidemiologic purposes and describes the condition as the acute onset of flaccid paralysis with neuroimaging evidence of spinal cord grey matter lesions (85, 88). High levels of CSF EV-specific antibodies are frequently identified in AFM cases, providing evidence for a causal role of nonpolio EV in AFM (89). Treatment is usually supportive and rehabilitative with outcomes varying from complete clinical recovery to those having persistent residual neurologic deficits (85) and slow recovery (90).

OTHER COMMON RESPIRATORY VIRUSES

Several other well-known respiratory viruses including rhinovirus, adenovirus, and parainfluenza virus commonly infect the general population. The respiratory symptoms related to infection are generally mild and include upper respiratory manifestations including rhinitis, pharyngitis, and tonsillitis. although usually self-limited, mild neurologic sequelae including aseptic meningitis, encephalopathy, and febrile seizures have been associated in

up to 3% of patients with such infections (91–95). Aseptic or viral meningitis specifically can have varying clinical presentations ranging from fatigue or headache to alterations in consciousness and seizures (96). CSF analysis typically shows a lymphocytic pleocytosis with hyperproteinemia, although patients can present with normal CSF findings (96, 97). Neurologic symptoms with such viruses tend to be mild, improve with supportive care, and have low-cause morbidity and mortality (95, 98).

CONCLUSIONS

The neurologic manifestations of severe respiratory viruses including Co-V, influenza, RSV, hMPV, and enterovirus D68 and D71 are a diagnostic and therapeutic challenge for many physicians. Clinical syndromes including status epilepticus, encephalitis, critical illness neuromyopathy, ADEM, ANE, GBS, transverse myelitis, and flaccid paralysis have all been associated with these viral respiratory infections. Having an understanding of the direct neurotropism of such viruses is imperative to understanding pathogenesis, clinical presentation and potential treatment paradigms aimed at improving morbidity and mortality.

Dr. Robinson has received honoraria for serving as an expert legal witness for traumatic brain injury. Dr. Busl disclosed that she does not have potential conflicts of interest.

For information regarding this article, E-mail: christopher.robinson@neurology.ufl.edu

REFERENCES

- Shi T, Arnott A, Semogas I, et al: The etiological role of common respiratory viruses in acute respiratory infections in older adults: A systematic review and meta-analysis. *J Infect Dis* 2019; pii:ijy662
- Akins PT, Belko J, Uyeki TM, et al: H1N1 encephalitis with malignant edema and review of neurologic complications from influenza. *Neurocrit Care* 2010; 13:396–406
- Algahtani H, Subahi A, Shirah B: Neurological complications of middle east respiratory syndrome coronavirus: A report of two cases and review of the literature. *Case Rep Neurol Med* 2016; 2016:3502683
- Antonucci R, Chiappe S, Porcella A, et al: Bronchiolitis-associated encephalopathy in critically-ill infants: An underestimated complication? *J Matern Fetal Neonatal Med* 2010; 23:431–436
- Arbour N, Day R, Newcombe J, et al: Neuroinvasion by human respiratory coronaviruses. *J Virol* 2000; 74:8913–8921
- Arnold JC, Singh KK, Milder E, et al: Human metapneumovirus associated with central nervous system infection in children. *Pediatr Infect Dis J* 2009; 28:1057–1060
- Mariotti P, Iorio R, Frisullo G, et al: Acute necrotizing encephalopathy during novel influenza A (H1N1) virus infection. *Ann Neurol* 2010; 68:111–114
- Bohmwald K, Gálvez NMS, Ríos M, et al: Neurologic alterations due to respiratory virus infections. *Front Cell Neurosci* 2018; 12:386
- Liebner S, Dijkhuizen RM, Reiss Y, et al: Functional morphology of the blood-brain barrier in health and disease. *Acta Neuropathol* 2018; 135:311–336
- Dahm T, Rudolph H, Schwerk C, et al: Neuroinvasion and inflammation in viral central nervous system infections. *Mediators Inflamm* 2016; 2016:8562805
- Swanson PA II, McGavern DB: Portals of viral entry into the central nervous system. In: *The Blood-Brain Barrier in Health and Disease*, Volume Two: Pathophysiology and Pathology. Dorovini-Zis K (Ed). CRC Press, Cleveland, OH, 2015, p 23

12. Suen WW, Prow NA, Hall RA, et al: Mechanism of West Nile virus neuroinvasion: A critical appraisal *Viruses* 2014; 6:2796–2825
13. Tyrrell DA, Bynoe ML: Cultivation of a novel type of common-cold virus in organ cultures. *Br Med J* 1965; 1:1467–1470
14. Vabret A, Dina J, Brison E, et al: [Human coronaviruses]. *Pathol Biol (Paris)* 2009; 57:149–160
15. Guan Y, Zheng BJ, He YQ, et al: Isolation and characterization of viruses related to the SARS coronavirus from animals in southern China. *Science* 2003; 302:276–278
16. Chan-Yeung M, Xu RH: SARS: Epidemiology. *Respirology* 2003; 8(Suppl):S9–S14
17. Gu J, Gong E, Zhang B, et al: Multiple organ infection and the pathogenesis of SARS. *J Exp Med* 2005; 202:415–424
18. Nassar MS, Bakhrebah MA, Meo SA, et al: Middle East Respiratory Syndrome Coronavirus (MERS-CoV) infection: Epidemiology, pathogenesis and clinical characteristics. *Eur Rev Med Pharmacol Sci* 2018; 22:4956–4961
19. Li YC, Bai WZ, Hashikawa T: The neuroinvasive potential of SARS-CoV2 may be at least partially responsible for the respiratory failure of COVID-19 patients. *J Med Virol* 2020 Feb 27. [online ahead of print]
20. Huang C, Wang Y, Li X, et al: Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; 395:497–506
21. Xu J, Zhong S, Liu J, et al: Detection of severe acute respiratory syndrome coronavirus in the brain: Potential role of the chemokine mig in pathogenesis. *Clin Infect Dis* 2005; 41:1089–1096
22. Ding Y, He L, Zhang Q, et al: Organ distribution of severe acute respiratory syndrome (SARS) associated coronavirus (SARS-CoV) in SARS patients: Implications for pathogenesis and virus transmission pathways. *J Pathol* 2004; 203:622–630
23. Li YC, Bai WZ, Hirano N, et al: Coronavirus infection of rat dorsal root ganglia: Ultrastructural characterization of viral replication, transfer, and the early response of satellite cells. *Virus Res* 2012; 163:628–635
24. Morfopoulou S, Brown JR, Davies EG, et al: Human coronavirus OC43 associated with fatal encephalitis. *N Engl J Med* 2016; 375:497–498
25. Yeh EA, Collins A, Cohen ME, et al: Detection of coronavirus in the central nervous system of a child with acute disseminated encephalomyelitis. *Pediatrics* 2004; 113:e73–e76
26. Turgay C, Emine T, Ozlem K, et al: A rare cause of acute flaccid paralysis: Human coronaviruses. *J Pediatr Neurosci* 2015; 10:280–281
27. Sharma K, Tengsupakul S, Sanchez O, et al: Guillain-Barré syndrome with unilateral peripheral facial and bulbar palsy in a child: A case report. *SAGE Open Med Case Rep* 2019; 7:2050313X19838750
28. Lau KK, Yu WC, Chu CM, : Possible central nervous system infection by SARS coronavirus. *Emerg Infect Dis* 2004; 10:342–344
29. Arbour N, Ekandé S, Côté G, et al: Persistent infection of human oligodendrocytic and neuroglial cell lines by human coronavirus 229E. *J Virol* 1999; 73:3326–3337
30. Fazzini E, Fleming J, Fahn S: Cerebrospinal fluid antibodies to coronavirus in patients with Parkinson's disease. *Mov Disord* 1992; 7:153–158
31. Tsai LK, Hsieh ST, Chao CC, et al: Neuromuscular disorders in severe acute respiratory syndrome. *Arch Neurol* 2004; 61:1669–1673
32. Arabi YM, Harthi A, Hussein J, et al: Severe neurologic syndrome associated with Middle East respiratory syndrome corona virus (MERS-CoV). *Infection* 2015; 43:495–501
33. Mao LWM, Chen S, He Q, et al: Neurological Manifestations of Hospitalized Patients with COVID-19 in Wuhan, China: A retrospective case series study. *medRxiv* 2020 Feb 25. [online ahead of print]
34. Fan HH, Wang LQ, Liu WL, et al: Repurposing of clinically approved drugs for treatment of coronavirus disease 2019 in a 2019-novel coronavirus (2019-nCoV) related coronavirus model. *Chin Med J (Engl)* 2020 Mar 6. [online ahead of print]
35. Zhou YH, Qin YY, Lu YQ, et al: Effectiveness of glucocorticoid therapy in patients with severe novel coronavirus pneumonia: protocol of a randomized controlled trial. *Chin Med J (Engl)* 2020 Mar 5. [online ahead of print]
36. Gao J, Tian Z, Yang X: Breakthrough: Chloroquine phosphate has shown apparent efficacy in treatment of COVID-19 associated pneumonia in clinical studies. *Biosci Trends* 2020; 14:72–73
37. Dolin R: Influenza. In: *Harrison's Principles of Internal Medicine*. Sixteenth Edition. Kasper DLBE, Fauci AS, Hauser SL, et al (Eds). McGraw Hill, New York, NY, 2005, pp 1066–1070
38. Uyeki TM, Bernstein HH, Bradley JS, et al: Clinical practice guidelines by the infectious diseases Society of America: 2018 update on diagnosis, treatment, chemoprophylaxis, and institutional outbreak management of seasonal influenza. *Clin Infect Dis* 2019; 68:e1–e47
39. Iuliano AD, Roguski KM, Chang HH, et al: Global Seasonal Influenza-associated Mortality Collaborator Network: Estimates of global seasonal influenza-associated respiratory mortality: A modelling study. *Lancet* 2018; 391:1285–1300
40. Jang H, Boltz D, Sturm-Ramirez K, et al: Highly pathogenic H5N1 influenza virus can enter the central nervous system and induce neuroinflammation and neurodegeneration. *Proc Natl Acad Sci U S A* 2009; 106:14063–14068
41. Fineberg HV: Pandemic preparedness and response—lessons from the H1N1 influenza of 2009. *N Engl J Med* 2014; 370:1335–1342
42. Sellers SA, Hagan RS, Hayden FG, et al: The hidden burden of influenza: A review of the extra-pulmonary complications of influenza infection. *Influenza Other Respir Viruses* 2017; 11:372–393
43. Surana P, Tang S, McDougall M, et al: Neurological complications of pandemic influenza A H1N1 2009 infection: European case series and review. *Eur J Pediatr* 2011; 170:1007–1015
44. Toovey S: Influenza-associated central nervous system dysfunction: A literature review. *Travel Med Infect Dis* 2008; 6:114–124
45. Sachedina N, Donaldson LJ: Paediatric mortality related to pandemic influenza A H1N1 infection in England: An observational population-based study. *Lancet* 2010; 376:1846–1852
46. Mizuguchi M: Acute necrotizing encephalopathy of childhood: A novel form of acute encephalopathy prevalent in Japan and Taiwan. *Brain Dev* 1997; 19:81–92
47. Abdelrahman HS, Safwat AM, Alsagheir MM: Acute necrotizing encephalopathy in an adult as a complication of H1N1 infection. *BJR Case Rep* 2019; 5:20190028
48. Ochi N, Takahashi K, Yamane H, et al: Acute necrotizing encephalopathy in an adult with influenza A infection. *Ther Clin Risk Manag* 2018; 14:753–756
49. Mizuguchi M, Abe J, Mikkaichi K, et al: Acute necrotizing encephalopathy of childhood: A new syndrome presenting with multifocal, symmetric brain lesions. *J Neurol Neurosurg Psychiatry* 1995; 58:555–561
50. Lee YJ, Smith DS, Rao VA, et al: Fatal H1N1-related acute necrotizing encephalopathy in an adult. *Case Rep Crit Care* 2011; 2011:562516
51. Okumura A, Mizuguchi M, Kidokoro H, et al: Outcome of acute necrotizing encephalopathy in relation to treatment with corticosteroids and gammaglobulin. *Brain Dev* 2009; 31:221–227
52. Togashi T, Matsuzono Y, Narita M: Epidemiology of influenza-associated encephalitis-encephalopathy in Hokkaido, the northernmost island of Japan. *Pediatr Int* 2000; 42:192–196
53. Martin A, Reade EP: Acute necrotizing encephalopathy progressing to brain death in a pediatric patient with novel influenza A (H1N1) infection. *Clin Infect Dis* 2010; 50:e50–e52
54. Morishima T, Togashi T, Yokota S, et al: Collaborative Study Group on Influenza-Associated Encephalopathy in Japan: Encephalitis and encephalopathy associated with an influenza epidemic in Japan. *Clin Infect Dis* 2002; 35:512–517
55. Tam CC, O'Brien SJ, Rodrigues LC: Influenza, campylobacter and mycoplasma infections, and hospital admissions for Guillain-Barre syndrome, England. *Emerg Infect Dis* 2006; 12:1880–1887
56. Tam CC, O'Brien SJ, Petersen I, et al: Guillain-Barré syndrome and preceding infection with campylobacter, influenza and Epstein-Barr virus in the general practice research database. *PLoS One* 2007; 2:e344
57. Grave C, Boucheron P, Rudant J, et al: Seasonal influenza vaccine and Guillain-Barre syndrome: A self-controlled case series study. *Neurology* 2020 Feb 25. [online ahead of print]
58. Hurwitz ES, Nelson DB, Davis C, et al: National surveillance for Reye syndrome: A five-year review. *Pediatrics* 1982; 70:895–900

59. Nakai Y, Itoh M, Mizuguchi M, et al: Apoptosis and microglial activation in influenza encephalopathy. *Acta Neuropathol* 2003; 105:233–239
60. Taubenberger JK: The origin and virulence of the 1918 “Spanish” influenza virus. *Proc Am Philos Soc* 2006; 150:86–112
61. Gamboa ET, Wolf A, Yahr MD, et al: Influenza virus antigen in postencephalitic parkinsonism brain. Detection by immunofluorescence. *Arch Neurol* 1974; 31:228–232
62. McCall S, Henry JM, Reid AH, et al: Influenza RNA not detected in archival brain tissues from acute encephalitis lethargica cases or in postencephalitic Parkinson cases. *J Neuropathol Exp Neurol* 2001; 60:696–704
63. Treanor JJ, Hayden FG, Vrooman PS, et al: Efficacy and safety of the oral neuraminidase inhibitor oseltamivir in treating acute influenza: A randomized controlled trial. US Oral Neuraminidase Study Group. *JAMA* 2000; 283:1016–1024
64. Jefferson T, Jones MA, Doshi P, et al: Neuraminidase inhibitors for preventing and treating influenza in healthy adults and children. *Cochrane Database Syst Rev* 2014; (4):CD008965
65. Del Mar C, Doshi P, Hama R, et al: Neuraminidase inhibitors for influenza complications. *Lancet* 2014; 384:1260–1261
66. King AMQ, Lefkowitz EJ, Mushegian AR, et al: Changes to taxonomy and the international code of virus classification and nomenclature ratified by the international committee on taxonomy of viruses (2018). *Arch Virol* 2018; 163:2601–2631
67. Pilar Orive FJ, Casado Flores J, García Teresa MA, et al: [Acute respiratory infections in pediatric intensive care units. A multicenter prospective study]. *An Esp Pediatr* 1998; 48:138–142
68. Binder W, Thorsen J, Borczuk P: RSV in adult ED patients: Do emergency providers consider RSV as an admission diagnosis? *Am J Emerg Med* 2017; 35:1162–1165
69. Eisenhut M: Extrapulmonary manifestations of severe respiratory syncytial virus infection—a systematic review. *Crit Care* 2006; 10:R107
70. Kho N, Kerrigan JF, Tong T, et al: Respiratory syncytial virus infection and neurologic abnormalities: retrospective cohort study. *J Child Neurol* 2004; 19:859–864
71. Sweetman LL, Ng YT, Butler JJ, et al: Neurologic complications associated with respiratory syncytial virus. *Pediatr Neurol* 2005; 32:307–310
72. Kneyber MC, Brandenburg AH, de Groot R, et al: Risk factors for respiratory syncytial virus associated apnoea. *Eur J Pediatr* 1998; 157:331–335
73. Lindgren C, Grögaard J. Reflex apnoea response and inflammatory mediators in infants with respiratory tract infection. *Acta Paediatr* 1996; 85:798–803.
74. Edwards KM, Zhu Y, Griffin MR, et al: New Vaccine Surveillance Network: Burden of human metapneumovirus infection in young children. *N Engl J Med* 2013; 368:633–643
75. Vehapoglu A, Turel O, Uygur Sahin T, et al: Clinical significance of human metapneumovirus in refractory status epilepticus and encephalitis: Case report and review of the literature. *Case Rep Neurol Med* 2015; 2015:131780
76. Mergeay M, Coeckelbergh E, De Cauwer H, et al: An adult case of metapneumovirus-induced acute encephalitis. *Acta Neurol Belg* 2019; 119:645–648
77. Jeannot N, van den Hoogen BG, Schefold JC, et al: Cerebrospinal fluid findings in an adult with human metapneumovirus-associated encephalitis. *Emerg Infect Dis* 2017; 23:370
78. Schildgen O, Glatzel T, Geikowski T, et al: Human metapneumovirus RNA in encephalitis patient. *Emerg Infect Dis* 2005; 11:467–470
79. Schieble JH, Fox VL, Lennette EH: A probable new human picornavirus associated with respiratory diseases. *Am J Epidemiol* 1967; 85:297–310
80. Schmidt NJ, Lennette EH, Ho HH: An apparently new enterovirus isolated from patients with disease of the central nervous system. *J Infect Dis* 1974; 129:304–309
81. Gear JH: Nonpolio causes of polio-like paralytic syndromes. *Rev Infect Dis* 1984; 6(Suppl 2):S379–S384
82. Messacar K, Asturias EJ, Hixon AM, et al: Enterovirus D68 and acute flaccid myelitis—evaluating the evidence for causality. *Lancet Infect Dis* 2018; 18:e239–e247
83. Berlin LE, Rorabaugh ML, Heldrich F, et al: Aseptic meningitis in infants < 2 years of age: Diagnosis and etiology. *J Infect Dis* 1993; 168:888–892
84. Wang Y, Zou G, Xia A, et al: Enterovirus 71 infection in children with hand, foot, and mouth disease in Shanghai, China: Epidemiology, clinical feature and diagnosis. *Virol J* 2015; 12:83
85. Messacar K, Schreiner TL, Van Haren K, et al: Acute flaccid myelitis: A clinical review of US cases 2012–2015. *Ann Neurol* 2016; 80:326–338
86. Greninger AL, Naccache SN, Messacar K, et al: A novel outbreak enterovirus D68 strain associated with acute flaccid myelitis cases in the USA (2012–14): A retrospective cohort study. *Lancet Infect Dis* 2015; 15:671–682
87. Chen CY, Chang YC, Huang CC, et al: Acute flaccid paralysis in infants and young children with enterovirus 71 infection: MR imaging findings and clinical correlates. *AJNR Am J Neuroradiol* 2001; 22:200–205
88. Maloney JA, Mirsky DM, Messacar K, et al: MRI findings in children with acute flaccid paralysis and cranial nerve dysfunction occurring during the 2014 enterovirus D68 outbreak. *AJNR Am J Neuroradiol* 2015; 36:245–250
89. Schubert RD, Hawes IA, Ramachandran PS, et al: Pan-viral serology implicates enteroviruses in acute flaccid myelitis. *Nat Med* 2019; 25:1748–1752
90. Bove R: Social media in the age of the “New Polio”. *N Engl J Med* 2019; 380:1195–1197
91. Lema CL, Cisterna DM, Freire MC: [Neurologic disease due to adenovirus infection]. *Medicina (B Aires)* 2005; 65:196–200
92. Soeur M, Wouters A, de Saint-Georges A, et al: Meningoencephalitis and meningitis due to an adenovirus type 5 in two immunocompetent adults. *Acta Neurol Belg* 1991; 91:141–150
93. Landry ML, Hsiung GD: Adenovirus-associated meningoencephalitis in a healthy adult. *Ann Neurol* 1988; 23:627–628
94. Lee KS, Lee BL, Heo YJ: Acute encephalopathy with biphasic seizures and late reduced diffusion associated with adenoviral pneumonia. *Child Neurol Open* 2019; 6:2329048X19826288
95. Huang YC, Huang SL, Chen SP, et al: Adenovirus infection associated with central nervous system dysfunction in children. *J Clin Virol* 2013; 57:300–304
96. Wright WF, Pinto CN, Palisoc K, et al: Viral (aseptic) meningitis: A review. *J Neurol Sci* 2019; 398:176–183
97. Dawood N, Desjobert E, Lumley J, et al: Confirmed viral meningitis with normal CSF findings. *BMJ Case Rep* 2014; 2014:bcr2014203733
98. McCarthy VP, Zimmerman AW, Miller CA: Central nervous system manifestations of parainfluenza virus type 3 infections in childhood. *Pediatr Neurol* 1990; 6:197–201