

SCIENTIFIC REPORTS



OPEN

Poly-L-ornithine promotes preferred differentiation of neural stem/progenitor cells via ERK signalling pathway

Received: 22 May 2015
Accepted: 28 September 2015
Published: 27 October 2015

Hongfei Ge, Liang Tan, Pengfei Wu, Yi Yin, Xin Liu, Hui Meng, Gaoyu Cui, Nan Wu, Jiangkai Lin, Rong Hu & Hua Feng

Neural stem/progenitor cells (NSPCs) replacement therapies are the most attractive strategies to restore an injured brain. Key challenges of such therapies are enriching NSPCs and directing them differentiation into specific neural cell types. Here, three biomaterial substrates Poly-L-ornithine (PO), Poly-L-lysine (PLL) and fibronectin (FN) were investigated for their effects on proliferation and differentiation of rat NSPCs, and the underlying mechanisms were also explored. The results showed PO significantly increased NSPCs proliferation and induced preferred differentiation, compared with PLL and FN. Checking protein markers of several neural cell subtypes, it is showed PO significantly induced NSPCs expressing Doublecortin (DCX) and Olig2, one for neuroblasts and young neurons and the other for young oligodendrocytes. It is suggested the ERK signaling pathway was involving in this process because an ERK antagonist U0126 could inhibit PO's effects mentioned above, as well as an ERK pathway agonist Ceramide C6 could enhance them. Given that both neurons and oligodendrocytes are the most vulnerable cells in many neurological diseases, PO-induced preferred differentiation into neurons and oligodendrocytes is a potential paradigm for NSPCs-based therapies.

Neural stem/progenitor cells (NSPCs), known as their characteristics of self-renewal, undifferentiated features and capacity to give birth to all neural lineage cell types^{1–3}, especially neuronal subtypes^{4–11}, have aroused much attention because of their potential to ameliorate various neurological diseases and injuries, meanwhile, provide a suitable cell source for transplantation therapies¹². Currently, there are two culture systems to obtain NSPCs known as neurosphere culture system^{13,14} and adherent culture system coated with different substrates^{15–18}. Especially, different substrates employed in adherent culture system, which could mimic the physiological microenvironment for culturing and expanding stem cells, have attracted great interests to date.

Although many substrates have been tried to fulfill above purpose including poly-L-ornithine (PO)¹⁹, Poly-L-Lysine (PLL)²⁰, Laminin^{21–23}, and Fibronectin (FN)¹⁹, their effects on proliferation and differentiation of NSPCs still remain elusive and which one is better for enriching NSPCs and directing their differentiation into preferred neural subtypes needs to be fully explored. PLL alone or combined with FN or laminin is widely used for the attachment of NSPCs to explore their morphology, proliferation, migration and differentiation²⁴. While, some studies suggest that PLL could enhance the likelihood of host inflammatory responses²⁵. Recent studies have shown that PO is less immunogenic than PLL and possesses some other advantages^{26,27}. FN, which is an extracellular glycoprotein that binds both cell integrins and other ECM molecules, plays a pivotal role in cell adhesion, survival and differentiation²⁸. However,

Department of Neurosurgery and Key Laboratory of Neurotrauma, Southwest Hospital, Third Military Medical University, Chongqing 400038, China. Correspondence and requests for materials should be addressed to R.H. (email: huchrong@aliyun.com) or H.F. (email: fenghua8888@vip.163.com)




Figure 1. Cell morphology and molecular marker expressions of rat NSPCs. (A) The suspended growth of neurospheres was notably observed. (B) The immunostaining of the Nestin expression (red) of the NSPCs in neurosphere before seeded on substrates. (C) The immunostaining of the Nestin expression (red) of the NSPCs after seeded on PO. (D) The immunostaining of the Sox2 expression (green) of the NSPCs after seeded on PO. Cell nuclei was stained with DAPI in blue. Scale bar: 20 μm .

Sun, T. *et al.* report that NSPCs on FN-coated dish, lose their proliferation potential after P6²⁹. As a result, it is necessary to explore the outcomes caused by different substrates on the biological behaviors of NSPCs.

In this present study, three substrates (PO, PLL and FN) were tested for their ability to promote proliferation and preferred differentiation of NSPCs. Meanwhile, the likely underlying mechanism(s) was explored. The aim of this study is to investigate their different effects on the proliferation and differentiation of NSPCs and therefore look for a more suitable biomaterial candidate to mimic the physiological microenvironment for expanding NSPCs and directing their preferred differentiation. At the same time, try to elucidate the possible underlying mechanism(s), which might provide a favorable cell replacement strategy for basic and clinical research associated with various neurological diseases and injuries.

Results

Culture and immunofluorescence identification for NSPCs. For preparation of NSPCs, fresh tissues were dissected from neocortical tissues from E14.5 Sprague–Dawley rats. The suspended growth of neurospheres was notably observed after 3 days cultured in the neurosphere culture system (Fig. 1A). Meanwhile, cells expressing Nestin, a marker for NSPCs, reached to 50–60% of total cells in a neurosphere, which was consistent with the previous study¹⁵ (Fig. 1B). In the adherent culture system, NSPCs also expressed Nestin (Fig. 1C, red) and Sox2 (Fig. 1D, green), another marker for NSPCs, after seeded on PO, PLL or FN (data not shown).

PO, PLL and PN showed no different effects on survival of NSPCs. First, we evaluated the effects of PO, PLL and FN on the death/survival of NSPCs by using cell death/survival assays through flow cytometry. As shown in Fig. 2, though there was a tendency that there were more cell death with




Figure 2. PO, PLL and FN showed no different effects on death/survival of NSPCs. (A) Typical illustrations showing death of NSPCs growing on PLL, PO and FN respectively, through flow cytometry on day 14 ($n = 5$). (B) Summarized graph showing the quantitative results from flow cytometry assay on day 3, 7 and 14 ($n = 5$). (C) Cell death of NSPCs determined by LDH assay on day 3, 7 and 14 growing on PLL, PO and FN respectively ($n = 6$). N.S. indicates no significant difference among PLL, PO and FN.

the prolonged culture time, no considerable difference was observed among PO, PLL and FN on day 3, 7 and 14 post cultured. The data suggested that these substrates have no different influence on the death/survival of NSPCs.

PO significantly increased proliferation of NSPCs. Next, we investigated the effects of PO, PLL and FN on the proliferation of NSPCs. Here, NSPCs were cultured in the enrichment medium with 20 ng/ml bFGF and 20 ng/ml EGF according to the well-known standard method. First, CCK8 assay illustrated that the absorbance at 450nm was higher in NSPCs growing on PO on day 7 and 14 in comparison with PLL or FN (Fig. 3A), which implied that NSPCs growing on PO proliferated more vigorously than those on PLL or FN. Second, Ki-67 assay showed that the co-labeling percentage of Ki-67 and Nestin in NSPCs cultured on PO was significantly higher than that on PLL or FN on day 7 and 14 (Fig. 3B, C). Third, western blotting data showed that the expression level of Nestin of NSPCs growing on PO, which was preferentially expressed in NSPCs, was markedly higher than that on PLL or FN on day 7 and 14 (Fig. 3D, E). Together, these data indicated that PO has a better ability to promote the proliferation of NSPCs comparing with PLL or FN.

PO induced preferred neuronal and oligodendrocytic differentiation of NSPCs. To investigate their effects on the differentiation of NSPCs, western blotting assays were performed. During this process, NSPCs were cultured in differentiation medium without 20 ng/ml bFGF and 20 ng/ml EGF. As shown in Fig. 4A, B, the expression of Doublecortin (DCX), which is preferentially expressed in neuroblasts and as the marker of young neurons, obviously increased in NSPCs growing on PO at all checking time points compared with PLL or FN. Their effects on glial cell fate, such as astrocyte or oligodendrocyte, were also examined. The expression level of glial fibrillary acidic protein (GFAP), which is widely used to identify astroglia in the brain, had no significant difference at all time points among different substrates (Fig. 4A, D). However, the expression of Olig2, expressed by young oligodendrocytes, significantly increased in NSPCs




Figure 3. PO significantly increased proliferation of NSPCs. (A) Bar graph showing the proliferation of NSPCs growing on different substrates by using CCK8 assay on day 3, 7 and 14 (n=6). *P < 0.05, #P < 0.01, One-Way ANOVA followed by Tukey's post hoc test. N.S. indicates no significant difference. (B) Bar graph showing the proliferation ability of NSPCs growing on different substrates by using Ki-67 assay on day 3, 7 and 14 (n=3). *P < 0.05, #P < 0.01, One-Way ANOVA followed by Tukey's post hoc test. N.S. indicates no significant difference. (C) Immunostaining images showing the percentage of Ki-67 positive cells in NSPCs growing on PO was higher than that on PLL or FN. The NSPCs were stained with antibody for Ki-67 (red) and antibody for Nestin (green), cell nuclei was stained with DAPI in blue. Co-labeling of Ki-67 and Nestin suggests actively proliferating NSPCs. Scale bar: 20 μm. (D) Western blotting showing the level of Nestin expressed by NSPCs growing on different substrates on day 14. (E) Summarized graph showing the expression level of Nestin of NSPCs growing on PO was markedly higher than that on PLL or FN on day 7 and 14. Blotting bands were analyzed using the Image Lab™ software for relative density and normalized to β-actin controls. *P < 0.05, #P < 0.01, One-Way ANOVA followed by Tukey's post hoc test. N.S. indicates no significant difference.




Figure 4. PO induced preferred neuronal and oligodendrocytic differentiation of NSPCs. (A) Western blotting bands showing three neural lineage differentiation of NSPCs by checking specific protein markers (DCX for young neurons, GFAP for astrocytes and Olig2 for oligodendrocytes) on day 3, 7 and 14. (B–D) Bar graphs summarizing quantitative results from (A) on day 3, 7 and 14, respectively. Bands were analyzed using the Image Lab™ software for relative density and normalized to Tubulin or GAPDH controls. * $P < 0.05$, # $P < 0.01$, One-Way ANOVA followed by Tukey's post hoc test. N.S. indicates no significant difference.




Figure 5. Activation of ERK was increased in NSPCs growing on PO. (A) Blotting bands showing p-ERK expression in NSPCs growing on PLL, PO and FN on day 3, 7 and 14. (B) Bar graphs summarizing quantitative results from (A). Bands were analyzed using the Image Lab™ software for relative density and normalized to total ERK controls. # $P < 0.01$, One-Way ANOVA followed by Tukey's post hoc test.

cultured on PO on day 7 and 14, compared to PLL or FN (Fig. 4A,C). These data indicated that NSPCs cultured on PO were more likely to give birth to young neurons and oligodendrocytes, representing a better candidate directing neuronal or oligodendrocytic differentiation of NSPCs.

ERK activation involves in PO-induced preferred neuronal differentiation from NSPCs. ERK has been reported to be implicated in the ECM-stimulated (including cell culture substrate) cell biological behaviors such as cell growth, proliferation and differentiation³⁰. To unravel the potential mechanisms for preferential fate choice of NSPCs induced by PO, and whether ERK signaling pathway was involved in this process, the following tests were conducted. The activation of ERK was evaluated via measuring




Figure 6. The crucial role of ERK in NSPCs differentiation into neuronal lineage. (A) Illustration showing the experimental procedure. (B) Immunostaining images displaying the influence of ERK signalling on the differentiation of NSPCs growing on PO, PLL and FN into DCX⁺, GFAP⁺ and Olig2⁺ cells using ERK antagonist or agonist on day 14. Scale bar: 20 μ m. (C–E) Bar graphs summarizing quantitative results from (B). #P < 0.01, One-Way ANOVA followed by Tukey's post hoc test. N.S. indicates no significance.

ERK phosphorylation (p-ERK) in western blot assay. As shown in Fig 5A, B, the expression of p-ERK in NSPCs cultured on PO was remarkably higher than that on PLL or FN at all checking time points. To provide direct evidence for the involvement of ERK in PO-induced preferred neuronal/oligodendrocytic differentiation, NSPCs cultured on PO were treated with a specific ERK antagonist U0126 (10 μ M) or agonist Ceramide C6 (10 μ M). Immunostaining data showed that antagonist significantly reduced the number of DCX⁺ cells (Fig. 6B, C) compared with the vehicle control. However, it had no obvious effect on number of GFAP⁺ cells (Fig. 6B, D) or Olig2⁺ cells (Fig. 6B, E). Instead, with the addition of 10 μ M Ceramide C6 could enhance more differentiation of NSPCs growing on PO into young neurons obviously (Fig. 6B, C). However, there was no effect on the astroglial and oligodendrocytic differentiating capacity (Fig. 6B, D, E). Taken together, these data indicated that the activation of ERK signaling pathway might contribute to PO-induced preferred differentiation of NSPCs into neurons, but not oligodendrocytes.

Discussion

The utility of neural stem/progenitor cells (NSPCs), with the ability of self-renewal and differentiation into functional neural cells^{4–11,31}, holds promise for cell therapy-based treatment of many neurological disorders such as Parkinson's disease³², traumatic brain injury³³ and ischemic attack^{34–36}. How to expand NSPCs and induce preferred differentiation into specific neural cell lineage is an attractive field to date. The adherent culture system with substrate(s) could mimic the physiological microenvironment for culturing and highly efficient expansion of NSPCs. Among the widely used substrates for culturing NSPCs such as PO, PLL and FN, this study, to our limited knowledge, for the first time demonstrates that PO promotes proliferation of NSPCs and preferential differentiation into neuronal and oligodendrocytic cell types, compared with PLL or FN. Given that both neurons and oligodendrocytes are the most vulnerable cells in many neurological diseases and injuries, PO-induced preferred differentiation into neurons and oligodendrocytes is a favorable strategy for NSPCs-based therapies.

During last decades, PO, PLL and FN are the most widely used culturing substrates for investigating NSPCs²⁴. They can provide a good physical support for the attachment of NSPCs, therefore benefit survival of NSPCs in culture conditions. Few studies compared their effects on the proliferation and preferred differentiation of NSPCs. However, recent findings make comparative study necessary. For example, some evidences indicate that PLL is likely to enhance inflammatory responses²⁵. Furthermore, it's reported that adherent NSPCs on FN would lose their potentiality of proliferation after P6²⁹. These

evidence suggest that different biomaterials might have distinct effects on characteristics of NSPCs. In this study, we found PLL, FN, PO had no different effects on the survival of NSPCs, as reported previously^{37,38}. Furthermore, we indeed found PO promoted the proliferation of NSPCs and preferential differentiation into neuronal and oligodendrocytic cells, compared to PLL or FN. In addition, previous studies have found that PO coating significantly increases the mechanical strength of the alginate microcapsule, which could provide a physical support for the attachment of NSPCs during cell-based therapy²⁴ and results in lower shear stress to increase biocompatibility³⁹. While, Flanagan LA *et al.* revealed that human NSPCs grown on PO for 5 days were less well spread¹⁸. The reason of this discrepancy between the two results might be the difference in coating procedure, cell density and difference in species. However, it provides a clue for the combination of PO and laminin for the culture of NSPCs and even for the improvement of biomaterials. Together, these data suggest PO hold the potential serving as a better biomaterial candidate than PLL and FN for enriching neurons and/or oligodendrocytes from NSPCs to treating neurological disorders in the future.

Although these conventional biomaterials are widely employed, the underlying mechanisms whereby NSPCs binding to substrates and undergoing proliferation and differentiation remain open. Usually, substrates in ECM offer outside-in signals for cells mediated by an array of integrins on the cell surface. Especially, β_1 -integrin receptor (such as $\alpha_5\beta_1$, $\alpha_3\beta_1$ and $\alpha_6\beta_1$ integrin complexes^{40–42}) has been shown to be crucial for NSPCs proliferation and neurogenesis^{43,44}. However, the underlying mechanisms for how integrins transmit the signals across the cell membrane to cytoplasm remain elusive. Data in this research have shown the activation of ERK1/2 is required for PO-induced preferred neuronal differentiation from NSPCs. It has been reported that ERK1/2 is implicated in neuronal differentiation^{30,45,46}. Therefore, it is needed to be further investigated how ERK1/2 acts with integrins to mediate PO signaling preferred neuronal differentiation. However, the PO-induced preferential differentiation into oligodendrocyte from NSPCs is not dependent on ERK1/2 activation, because the number of Olig2⁺ cells differentiated from NSPCs is not affected by the addition of ERK1/2 antagonist or agonist. This implies the mechanism whereby PO induced preferred oligodendrocytic differentiation is different from neuronal one. The mechanism(s) underlying PO-induced oligodendrocytic differentiation is not clear and needs to be determined in the future research.

In summary, our study has revealed pivotal insights into the activity of PO on NSPCs proliferation and differentiation, and might open new avenues for NSPCs-based therapies specifically promoting neuronal and oligodendrocytic differentiation in brain disorders.

Methods

Animal. This study was performed in accordance with the China's animal welfare legislation for the care and use of animals and approved by the Third Military Medical University Chongqing, China. We did our best to minimize the number of animals and decrease their suffering. E14.5 Sprague-Dawley rats were sacrificed after anesthetized with pentobarbital (60 mg/kg intraperitoneally).

Primary Neural Stem/Progenitor Cells Culture. Neocortical tissues obtained from E14.5 Sprague-Dawley rats were dipped in 0.25% Trypsin Protease (HyClone™, SV30031.01) for 30 minutes at 37°C. Soybean trypsin inhibitor (2.8 mg/mL, Roche Diagnosis, Mannheim, Germany) was in addition to inhibit trypsin activity. After washing twice with DMEM/F12 medium (HyClone™, SH30023.01), the tissue samples were triturated using a fire-polished Pasteur pipette and passed through a 100 μ m Nylon cell strainer (BD Falcon, 352360) to harvest dissociated cell suspensions. Then, they were seeded at an initial cell density of 1×10^5 cells/mL in DMEM/F12 medium supplemented with 20 ng/ml recombinant murine FGF-basic (PeproTech, 450–33), 20 ng/ml recombinant murine EGF (PeproTech, 315–09), and 2% B-27 supplement (Life Technologies, 17504–044) for 24 hours at 37°C under humidified 5% CO₂ conditions. NSPCs suspension were plated onto 35-mm different pre-coated dishes (18 hrs) after washing twice with sterile cell culture grade water according to the manufacturer's instructions respectively. The pre-coated substrates were as follows: 100 μ g/ml Poly-L-lysine (sigma, P4832), 10 μ g/ml poly-L-ornithine (Sigma, P4957) and 20 μ g/ml Fibronectin (sigma, F2006). Half volume of culture medium was exchanged every 3 days. U0126 (MCE, HY-12031) and Ceramide C6 (Santa cruz, sc-3527) were dissolved in DMSO and dissolved in medium without mitogens at a concentration of 10 μ M.

Immunohistochemistry. For fluorescence immunocytochemistry, NSPCs adhered to different pre-coated coverslips were fixed with 4% paraformaldehyde in 0.01 M phosphate-buffered saline (pH 7.4) for 2 hours at room temperature and blocked with 5% v/v fetal bovine serum or with 0.5% v/v Triton-X 100 (Sigma-Aldrich, $\times 100$) in PBS. NSPCs were incubated with rabbit polyclonal to Ki67 (Abcam, 1:200), rabbit polyclonal to Doublecortin (Abcam, 1:200), rabbit polyclonal to glial fibrillary acidic protein (GFAP) (Proteintech Group, Inc, 1:100), rabbit polyclonal to Olig2 (Proteintech Group, Inc, 1:100), rabbit polyclonal to Sox2 (Abcam, 1:200), or mouse monoclonal to Nestin (Abcam, 1:250) overnight at 4°C and then goat anti-rabbit Cy3 or anti-mouse FITC (Proteintech Group, Inc, 1:200), or donkey anti-rabbit FITC or anti-mouse Cy3 (Beyotime, 1:250) for 2 hours at room temperature. All cultures were counterstained with the DNA-binding dye 4'-6-Diamidino-2-phenylindole (DAPI, 2 mg/ml in PBS) for 10 minutes at room temperature. Then, Coverslips were mounted onto glass slides and the images were

obtained from a Zeiss UV 780 Meta confocal microscope (Carl Zeiss), and examined by AxioVision 4 (Carl Zeiss AG) or Adobe Photoshop CS5 software.

Western Blotting. NSPCs adhered to different precoated dishes were homogenized with lysis buffer containing 0.025 M Tris-HCl, pH 8.0, 0.15 M NaCl, 0.001 M EDTA, 1% Nonidet P-40, 10% glycerol; pH 7.4, and a protease inhibitor mixture (Complete, Roche). The protein concentration was measured by enhanced BCA Protein Assay Kit (Beyotime, P0010S). Proteins (15 µg/lane) were separated by 10% SDS-PAGE under reducing conditions and electroblotted to polyvinylidene difluoride membranes (Roche, 3010040001). Membranes were incubated in blocking with 5% skimmed milk or 3% bovine serum albumin (BSA) in TBST at room temperature for 2 hours. Membranes were then incubated with rabbit polyclonal to Doublecortin (Abcam, 1:2000), rabbit polyclonal to glial fibrillary acidic protein (GFAP) (Proteintech Group, Inc, 1:1000), rabbit polyclonal to Olig2 (Proteintech Group, Inc, 1:1000), mouse monoclonal to Nestin (Abcam, 1:2500), rabbit polyclonal to p44/42 MAPK(Erk1/2) (CST, 1:1000), rabbit polyclonal to Phospho-p44/42 MAPK (Erk1/2) (CST, 1:1000), or mouse monoclonal to GAPDH (Zsgb-bio, 1:1000) overnight at 4 °C. After incubation with peroxidase-conjugated (HRP)-conjugated secondary IgG (Zsgb-bio, 1:5000) for 2 hours at room temperature. All membranes were detected by ChemiDoc™ XRS+ imaging system using the Pierce Fast Western Blot Kits (Thermo Scientific, 35065). Densitometric measurement of each membrane was performed using Image Lab™ image acquisition and analyzed by Image Lab™ software. GAPDH, an internal control, was used to normalize the expression level of each protein.

Cell Proliferation Assay. Cell proliferation was assessed by Cell Counting Kit-8, that determines the cell viability in cell proliferation through WST reduction assay (Dojindo, CK04-05), which detects dehydrogenase activities of viable cells. 100 µl of cell suspension (~10000 cells/well) was dispensed in a 96-well cell culture cluster with different pre-coated substrates, and they were incubated with 10% (v/v) WST solution for 2.5 h at 37 °C. Then, the absorbance of the culture medium at a test wavelength of 450 nm was determined using a microplate reader and a reference wavelength of 630 nm as well.

NSPCs adhered to different precoated dishes were dissociated using StemPro® Accutase® Cell Dissociation Reagent (Life Technologies, A1110501), and the dissociated cell suspensions were seeded in 24-well cell culture plates ($2-5 \times 10^5$ cells/ml) for fluorescence to visualise the percentage of Ki-67⁺ cells as described above.

In vitro cytotoxicity assay. Lactate dehydrogenase (LDH) releasing levels were measured to evaluate cytotoxicity. LDH, released from injured cells in the medium, indicated the integrity of the cell membrane. Neurosphere culture supernatants were collected on day 3, 7 and 14. Cells were lysed with 2% Triton X-100 (Sigma, ×100) for 15 min to release all LDH from the cytoplasm after collecting the last supernatants. Cell lysis, served as positive controls, determined the maximal LDH release. LDH release was detected using the absorbance of the culture medium at a test wavelength of 450 nm with a microplate reader (Nanjing Jiancheng bioengineering institute, A020-2) according to the manufacturer's instructions and it was shown as a rate of LDH released in the medium to total cellular ($n = 4$).

Annexin-V Assay. Flow cytometry was used to detect Annexin-V staining via an Annexin V-FITC Apoptosis Detection Kit (Beyotime). Adherent cells dissociated using StemPro® Accutase® Cell Dissociation Reagent (Life Technologies, A1110501) were collected and processed in accordance with the manufacturer recommendation. After 10 minutes of incubation with Annexin V-FITC at room temperature, cells were resuspended and incubated in binding buffer. Then Propidium Iodide was added into cell suspension for 5 minutes on ice. Cells were analyzed using a FACScan flow cytometer (Becton Dickinson).

Statistical methods. All data were showed as mean ± SEM and statistical analyses were carried out with SPSS v17.0 (SPSS Inc, Chicago, IL). Unpaired Student's t test or one-way ANOVA followed by Tukey's post hoc test was used to define statistical significance, and $p < 0.05$ was considered statistically significant.

References

1. Keller, G. M. *In vitro* differentiation of embryonic stem cells. *Curr Opin Cell Biol* **7**, 862–9 (1995).
2. Smith, A. G. Embryo-derived stem cells: of mice and men. *Annu Rev Cell Dev Biol* **17**, 435–62 (2001).
3. Gadue, P., Huber, T. L., Nostro, M. C., Kattman, S. & Keller, G. M. Germ layer induction from embryonic stem cells. *Exp Hematol* **33**, 955–64 (2005).
4. Lee, S. H., Lumelsky, N., Studer, L., Auerbach, J. M. & McKay, R. D. Efficient generation of midbrain and hindbrain neurons from mouse embryonic stem cells. *Nat Biotechnol* **18**, 675–9 (2000).
5. Carpenter, M. K. *et al.* Enrichment of neurons and neural precursors from human embryonic stem cells. *Exp Neurol* **172**, 383–97 (2001).
6. Tropepe, V. *et al.* Direct neural fate specification from embryonic stem cells: a primitive mammalian neural stem cell stage acquired through a default mechanism. *Neuron* **30**, 65–78 (2001).
7. Pachernik, J., Esner, M., Bryja, V., Dvorak, P. & Hampl, A. Neural differentiation of mouse embryonic stem cells grown in monolayer. *Reprod Nutr Dev* **42**, 317–26 (2002).

8. Wichterle, H., Lieberam, I., Porter, J. A. & Jessell, T. M. Directed differentiation of embryonic stem cells into motor neurons. *Cell* **110**, 385–97 (2002).
9. Ying, Q. L., Stavridis, M., Griffiths, D., Li, M. & Smith, A. Conversion of embryonic stem cells into neuroectodermal precursors in adherent monoculture. *Nat Biotechnol* **21**, 183–6 (2003).
10. Watanabe, K. *et al.* Directed differentiation of telencephalic precursors from embryonic stem cells. *Nat Neurosci* **8**, 288–96 (2005).
11. Fico, A. *et al.* High-throughput screening-compatible single-step protocol to differentiate embryonic stem cells in neurons. *Stem Cells Dev* **17**, 573–84 (2008).
12. Polak, J. M. & Bishop, A. E. Stem cells and tissue engineering: past, present, and future. *Ann N Y Acad Sci* **1068**, 352–66 (2006).
13. Reynolds, B. A., Tetzlaff, W. & Weiss, S. A multipotent EGF-responsive striatal embryonic progenitor cell produces neurons and astrocytes. *J Neurosci* **12**, 4565–74 (1992).
14. Deleyrolle, L. P. & Reynolds, B. A. Isolation, expansion, and differentiation of adult Mammalian neural stem and progenitor cells using the neurosphere assay. *Methods Mol Biol* **549**, 91–101 (2009).
15. Nakaji-Hirabayashi, T., Kato, K., Arima, Y. & Iwata, H. Oriented immobilization of epidermal growth factor onto culture substrates for the selective expansion of neural stem cells. *Biomaterials* **28**, 3517–29 (2007).
16. Nakaji-Hirabayashi, T., Kato, K. & Iwata, H. Surface-anchoring of spontaneously dimerized epidermal growth factor for highly selective expansion of neural stem cells. *Bioconjug Chem* **20**, 102–10 (2009).
17. Chetty, S. *et al.* Stress and glucocorticoids promote oligodendrogenesis in the adult hippocampus. *Mol Psychiatry* **19**, 1275–83 (2014).
18. Flanagan, L. A., Rebaza, L. M., Derzic, S., Schwartz, P. H. & Monuki, E. S. Regulation of human neural precursor cells by laminin and integrins. *J Neurosci Res* **83**, 845–56 (2006).
19. Hazel, T. & Muller, T. Culture of neuroepithelial stem cells. *Curr Protoc Neurosci* **Chapter 3**, Unit 3 1 (2001).
20. Purcell, E. K., Singh, A. & Kipke, D. R. Alginate composition effects on a neural stem cell-seeded scaffold. *Tissue Eng Part C Methods* **15**, 541–50 (2009).
21. Raghavan, S. & Bitar, K. N. The influence of extracellular matrix composition on the differentiation of neuronal subtypes in tissue engineered innervated intestinal smooth muscle sheets. *Biomaterials* **35**, 7429–40 (2014).
22. Wang, L. & Marquardt, T. Direct live monitoring of heterotypic axon-axon interactions *in vitro*. *Nat Protoc* **7**, 351–63 (2012).
23. Ho, M., Yu, D., Davidson, M. C. & Silva, G. A. Comparison of standard surface chemistries for culturing mesenchymal stem cells prior to neural differentiation. *Biomaterials* **27**, 4333–9 (2006).
24. Darrabie, M. D., Kendall, W. F. Jr. & Opara, E. C. Characteristics of Poly-L-Ornithine-coated alginate microcapsules. *Biomaterials* **26**, 6846–52 (2005).
25. Strand, B. L., Gaserod, O., Kulseng, B., Espevik, T. & Skjak-Baek, G. Alginate-polylysine-alginate microcapsules: effect of size reduction on capsule properties. *J Microencapsul* **19**, 615–30 (2002).
26. Brunetti, P. *et al.* Immunoprotection of pancreatic islet grafts within artificial microcapsules. *Int J Artif Organs* **14**, 789–91 (1991).
27. Calafiore, R. *et al.* Effects of alginate/polyaminoacidic coherent microcapsule transplantation in adult pigs. *Transplant Proc* **29**, 2126–7 (1997).
28. Schwarzbauer, J. E. & DeSimone, D. W. Fibronectins, their fibrillogenesis, and *in vivo* functions. *Cold Spring Harb Perspect Biol* **3** (2011), doi: 10.1101/cshperspect.a005041.
29. Sun, T. *et al.* A comparison of proliferative capacity and passaging potential between neural stem and progenitor cells in adherent and neurosphere cultures. *Int J Dev Neurosci* **29**, 723–31 (2011).
30. Chan, W. S. *et al.* Differential regulation of proliferation and neuronal differentiation in adult rat spinal cord neural stem/progenitors by ERK1/2, Akt, and PLCgamma. *Front Mol Neurosci* **6**, 23 (2013).
31. Tornero, D. *et al.* Human induced pluripotent stem cell-derived cortical neurons integrate in stroke-injured cortex and improve functional recovery. *Brain* **136**, 3561–77 (2013).
32. Kim, J. H. *et al.* Dopamine neurons derived from embryonic stem cells function in an animal model of Parkinson's disease. *Nature* **418**, 50–6 (2002).
33. Alvarez, Z. *et al.* Neurogenesis and vascularization of the damaged brain using a lactate-releasing biomimetic scaffold. *Biomaterials* **35**, 4769–81 (2014).
34. Park, K. J., Park, E., Liu, E. & Baker, A. J. Bone marrow-derived endothelial progenitor cells protect postischemic axons after traumatic brain injury. *J Cereb Blood Flow Metab* **34**, 357–66 (2014).
35. Karelina, K., Alzate-Correa, D. & Obrietan, K. Ribosomal S6 kinase regulates ischemia-induced progenitor cell proliferation in the adult mouse hippocampus. *Exp Neurol* **253**, 72–81 (2014).
36. Tang, Y. *et al.* Neural stem cell protects aged rat brain from ischemia-reperfusion injury through neurogenesis and angiogenesis. *J Cereb Blood Flow Metab* **34**, 1138–47 (2014).
37. Goetz, A. K. *et al.* Temporally restricted substrate interactions direct fate and specification of neural precursors derived from embryonic stem cells. *Proc Natl Acad Sci USA* **103**, 11063–8 (2006).
38. Harper, M. M., Ye, E. A., Blong, C. C., Jacobson, M. L. & Sakaguchi, D. S. Integrins contribute to initial morphological development and process outgrowth in rat adult hippocampal progenitor cells. *J Mol Neurosci* **40**, 269–83 (2010).
39. Thu, B. *et al.* Alginate polycation microcapsules. II. Some functional properties. *Biomaterials* **17**, 1069–79 (1996).
40. Wojcik-Stanaszek, L., Gregor, A. & Zalewska, T. Regulation of neurogenesis by extracellular matrix and integrins. *Acta Neurobiol Exp (Wars)* **71**, 103–12 (2011).
41. Nakaji-Hirabayashi, T., Kato, K. & Iwata, H. Improvement of neural stem cell survival in collagen hydrogels by incorporating laminin-derived cell adhesive polypeptides. *Bioconjug Chem* **23**, 212–21 (2012).
42. Lathia, J. D. *et al.* Patterns of laminins and integrins in the embryonic ventricular zone of the CNS. *J Comp Neurol* **505**, 630–43 (2007).
43. Belvindrah, R., Hankel, S., Walker, J., Patton, B. L. & Muller, U. Beta1 integrins control the formation of cell chains in the adult rostral migratory stream. *J Neurosci* **27**, 2704–17 (2007).
44. Loulier, K. *et al.* beta1 integrin maintains integrity of the embryonic neocortical stem cell niche. *PLoS Biol* **7**, e1000176 (2009).
45. Li, Z., Theus, M. H. & Wei, L. Role of ERK 1/2 signaling in neuronal differentiation of cultured embryonic stem cells. *Dev Growth Differ* **48**, 513–23 (2006).
46. Barnabe-Heider, F. & Miller, F. D. Endogenously produced neurotrophins regulate survival and differentiation of cortical progenitors via distinct signaling pathways. *J Neurosci* **23**, 5149–60 (2003).

Acknowledgements

This work was supported by grants from National Natural Science Foundation of China (No. 81371340), National Key Basic Research Program of China (2014CB541603), and the Key Project of Natural Science Foundation of Chongqing (CSTC2013jjB0125).

Author Contributions

H.F. and R.H. designed the study and wrote the manuscript; H.F.G., L.T. and P.F.W. performed the study, data analysis and interpretation; Y.Y. and X.L. performed the data analysis; H.M. and G.Y.C. performed western blotting, data analysis and interpretation; N.W. and J.K.L. provided illustrations; H.F., H.F.G., R.H. and X.L. contributed in writing the manuscript. All authors reviewed the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Ge, H. *et al.* Poly-L-ornithine promotes preferred differentiation of neural stem/progenitor cells via ERK signalling pathway. *Sci. Rep.* **5**, 15535; doi: 10.1038/srep15535 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>