

RESEARCH ARTICLE

Physical inactivity in healthy, obese, and diabetic adults in Germany: An analysis of related socio-demographic variables

Stephanie Linder^{1*}, Karim Abu-Omar¹, Wolfgang Geidl², Sven Messing², Mustafa Sarshar³, Anne K. Reimers¹, Heiko Ziemainz¹

1 Division of Physical Activity and Public Health, Department of Sport Science and Sport, Friedrich-Alexander University Erlangen-Nuremberg, Erlangen, Bavaria, Germany, **2** Division of Exercise and Health, Department of Sport Science and Sport, Friedrich-Alexander University Erlangen-Nuremberg, Erlangen, Bavaria, Germany, **3** Division of Health and Physical Activity, Department of Sport Science, Otto-von-Guericke University, Magdeburg, Germany

* stephanie.linder@fau.de



OPEN ACCESS

Citation: Linder S, Abu-Omar K, Geidl W, Messing S, Sarshar M, Reimers AK, et al. (2021) Physical inactivity in healthy, obese, and diabetic adults in Germany: An analysis of related socio-demographic variables. PLoS ONE 16(2): e0246634. <https://doi.org/10.1371/journal.pone.0246634>

Editor: Rasheed Ahmad, Dasman Diabetes Institute, KUWAIT

Received: August 8, 2020

Accepted: January 25, 2021

Published: February 9, 2021

Copyright: © 2021 Linder et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data cannot be shared publicly because this study is a secondary data analysis of public use files (except the NBS data set) and there are legal restrictions on sharing these data sets (the license agreement on the data sets explicitly state that disclosure to third parties is excluded). However, the public use file nature of the data ensures public access to the data. All data sets (with exception of the NBS) are available for public use from the respective organisations. The scientific use files GEDA (2009/2010/2012/2014),

Abstract

Background

Adults with diabetes or obesity are more likely to be physically inactive than healthy adults. Physical activity is essential in the management of both diseases, necessitating targeted interventions in these groups. This study analysed physical inactivity (defined as not taking part in leisure-time physical activity) in over 100,000 adults in Germany considering their body mass index and the presence of diabetes. Furthermore, the relationship between specific socio-demographic factors with physical inactivity was investigated, particularly focusing diabetic and obese people, to refine the identification of risk-groups for targeted interventions on physical activity promotion.

Methods

Data from 13 population-based health surveys conducted in Germany from 1997 to 2018 were used. The relevant variables extracted from these datasets were merged and employed in the analyses. We included data from 129,886 individuals in the BMI analyses and 58,311 individuals in the diabetes analyses. Logistic regression analyses were performed to identify the importance of six socio-demographic variables (age, sex/gender, education, income, employment, and migration) for the risk of physical inactivity.

Results

Obese and diabetic people reported a higher prevalence of physical inactivity than those who were not affected. Logistic regression analyses revealed advanced age, low education level, and low household income as risk factors for physical inactivity in all groups. A two-sided migration background and unemployment also indicated a higher probability of physical inactivity.

DEGS1, BGS98, GSTel03 can be requested from the Research Data Centre (FDZ) of the Robert-Koch-Institute (https://www.rki.de/DE/Content/Forsch/FDZ/FDZ_node.html). Allbus data (1998, 2004, 2014) and GESIS Panel are available via the GESIS data stock catalogue (DBS, <http://search.gesis.org>). Share Wave Data (Wave 5 doi: [10.6103/SHARE.w5.710](https://doi.org/10.6103/SHARE.w5.710)) are distributed to registered users through the SHARE Research Data Center (<http://www.share-project.org/data-access.html>). For NBS see doi: [10.1055/s-2007-984440](https://doi.org/10.1055/s-2007-984440).

Funding: The SHARE data collection has been primarily funded by the European Commission through FP5 (QLK6-CT- 2001-00360), FP6 (SHARE-I3: RII-CT-2006-062193, COMPARE: CIT5-CT-2005-028857, SHARELIFE: CIT4-CT-2006-028812) and FP7 (SHARE-PREP: N°211909, SHARE-LEAP: N°227822, SHARE M4: N°261982). Additional funding from the German Ministry of Education and Research, the U.S. National Institute on Aging (U01_AG09740-13S2, P01_AG005842, P01_AG08291, P30_AG12815, R21_AG025169, Y1-AG-4553-01, IAG_BSR06-11, OGHA_04-064) and from various national funding sources is gratefully acknowledged (see www.share-project.org).

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: BMI, Body Mass Index; CI, Confidence Interval; OR, Odds Ratio; PA, Physical Activity.

Conclusion

Similar socio-demographic barriers appear to be important determinants of physical inactivity, regardless of BMI status or the presence of diabetes. However, physical activity promoting interventions in obese and diabetic adults should consider the specific disease-related characteristics of these groups. A special need for target group specific physical activity programmes in adults from ethnic minorities or of advanced age was further identified.

Introduction

Diabetes and obesity are critical challenges to public health [1]. In Germany, nearly five million adults have diabetes [2, 3]. A similar pattern is present for obesity (body mass index (BMI) ≥ 30 kg/m²), with almost a quarter of the German population affected [4]. In addition, being obese [5] is a crucial risk factor for the development of type 2 diabetes [3, 6].

Studies have highlighted physical inactivity as a major modifiable lifestyle factor in the prevention and management of both conditions [7, 8] since increasing physical activity (PA) precipitates significant health and weight loss benefits [9]. Nevertheless, adults with diabetes or obesity continue to be particularly inactive and are more likely to fail to meet PA recommendations than healthy individuals [10, 11]. Public health professionals have therefore identified the need for programs to promote PA in these groups [12]. Many studies have attempted to better understand the different propensities of physical inactivity, consequently, they have identified variables that influence people's PA behaviour. In addition to age and sex/gender [13], socio-demographic factors, particularly income, education, employment status, and migration background, have been repeatedly proven to influence PA [14–18]. An analysis of factors that influence PA behaviour is essential to develop appropriate PA interventions [19, 20]. However, due to the generally small number of study participants, socio-demographic factors associated with physical inactivity have not yet been extensively examined among people with obesity or diabetes. Mapping and characterising such factors and potential social gradients for different sub-groups of obese people could refine the identification of target groups to enhance the tailoring of interventions designed to promote PA. Additionally, such an investigation is necessary to assess whether socio-demographic factors for physical inactivity may differ across vulnerable groups.

While there is growing evidence, that in particular PA in the context of leisure time has promoting health benefits [21, 22], accurate data on leisure-time PA among obese and diabetic individuals in Germany remains scarce. Therefore, the present study investigates physical inactivity defined as not taking part in any leisure-time PA among individuals with normal weight, overweight or obesity and with or without a diagnosis of diabetes using a large sample of adults in Germany. To uncover a detailed picture of physically inactive individuals, we further stratified results by age, educational level, and household income. Since the evidence indicates several socio-demographic factors which influence physical inactivity, a further goal was to analyse the association between different socio-demographic variables for physical inactivity stratified by BMI and the presence or absence of diabetes. Our analysis intends to provide new insights into the propensity of physical inactivity at leisure time and its socio-demographic predictors among obese and diabetic individuals to support the future adoption and tailoring of PA promotion programs for these risk groups.

Table 1. Data sets included in the analysis.

Year(s) data were collected	Survey	Institution collecting the data	Age range of respondents	Sample size
1997–1999	BGS 98	Robert Koch Institute	18–79	7,124
1998	Allbus	Leibniz Institute for the Social Sciences	18+	3,234
2003	GSTel03	Robert Koch Institute	18+	8,318
2004	Allbus	Leibniz Institute for the Social Sciences	18+	2,946
2006	NBS	GFK Nürnberg	10+	10,554
2008–2011	DEGS1	Robert Koch Institute	18–79	7,987
2009	GEDA 2009	Robert Koch Institute	18+	21,262
2010	GEDA 2010	Robert Koch Institute	18+	22,050
2012	GEDA 2012	Robert Koch Institute	18+	19,294
2013	SHARE-WAVE 5	Max Planck Institute for Social Law and Social Policy	50+	5,752
2013–2018	GESIS Panel	Leibniz Institute for the Social Sciences	18+	7,599
2014	Allbus	Leibniz Institute for the Social Sciences	18+	3,471
2014	GEDA 2014	Robert Koch Institute	18+	24,016

<https://doi.org/10.1371/journal.pone.0246634.t001>

Method

Participants and data collection

This analysis was based on pooled data from 13 population-based national health surveys (conducted between 1997 and 2018) involving 143,709 participants. Seven surveys were part of a nationwide Health Monitoring System administered by the Robert Koch-Institute [23–29] four surveys were conducted by the Leibniz Institute for the Social Sciences [30–33] and one each by the Max Planck Institute for Social Law and Social Policy [34–36] and the GFK Nürnberg [37]. All surveys were self-reported and comprised several major sections related to information about general socio-demographic and socio-economic characteristics, health status, and PA behaviour. Since we conducted a secondary data analysis of the given data sets, ethical approval is not applicable. An overview of the included surveys is presented in Table 1.

Study design

Two sub-samples were formed by stratifying the total population based on BMI and diabetes. The ‘BMI sample’ included data from adults aged 18 years and older. The final sample consisted of 129,886 individuals. The ‘diabetes sample’ exclusively included people aged 50 years and over because in more than 50% of the cases, diabetes is diagnosed at or after the age of 50 years [38]. The final sample consisted of 58,311 individuals. Individuals with BMI < 18.5 kg/m² were excluded from the analysis.

Specificities of the sub-samples

BMI sample. Based on the self-reported height and weight data from the surveys, BMI was calculated as weight in kilograms divided by the square of height in metres. The participants were divided into five categories based on the classification of the World Health Organisation [5]: normal weight (18.5 to <25 kg/m²), overweight (25 to <30 kg/m²), obesity grade I (30 to <35 kg/m²), obesity grade II (35 to <40 kg/m²) and obesity grade III (\geq 40 kg/m²).

Diabetes sample. Different questions were used to identify diabetes in the surveys: participants were asked if they suffer from diabetes, had ever received a medical diagnosis, or received a diagnosis within the last 12 months. Participants who answered ‘No’ to these

questions were included in the non-diabetic group, and those who answered 'Yes' were included in the diabetic group.

Measures

Physical inactivity. PA was assessed related to the duration of leisure-time PA per week within the last three months (e.g., GEDA 2009) or generally (e.g., GESIS Panel, Allbus). Regarding the information on weekly duration, the responses were grouped in hours as follows: no PA, ≤ 2 h, 2–4 h, and >4 h. Most datasets included questions on leisure-time PA only (BGS 1998, GEDA 2009, 2010, 2012, SHARE 5, DEGS1). Adults were identified as being physically inactive and thus assigned to the group 'no PA' if they reported to never engage in leisure-time physical activities (BGS 1998, NBS 2006). Respondents to SHARE 5 were coded as being physically inactive if they reported almost never or never engaging in leisure-time PA. Respondents to GEDA 2009, GEDA 2010, GEDA 2012 and DEGS1 were assigned to the group 'no PA' if they engaged in no leisure-time PA in the past week. Adults who indicated that they were physically active were allocated to the groups ' ≤ 2 h' to ' >4 h' according to their stated duration of leisure-time PA.

Socio-demographic factors. Since we pooled secondary data sets the analysis of socio-demographic variables was restricted to those that are routinely collected as part of behavioural risk factor surveillance. Participants answered basic questions about their socio-demographic background. Six socio-demographic variables were included in the analysis. For each variable separate categories were formed: (1) sex/gender (male, female); (2) age (for BMI sample: 18–<30 years, 30–<60 years, 60–<80 years, and 80+ years; for diabetes sample: 50–<60 years, 60–<70 years, 70–<80 years, and 80+ years); (3) education level (low (9–10 years schooling), medium (10–11 years schooling), and high (12–13 years schooling)); (4) household income (bottom 20%, 20–40%, 40–60%, 60–80%, and top 20%); (5) migration background (none, one-sided, and two-sided) and (6) employment (unemployed, employed).

Statistical analysis

All analyses were conducted separately for the BMI sample and the diabetes sample. The results of the analysis of the diabetes sample are presented in the appendix only.

Descriptive statistics (frequencies) were used to determine PA levels of all groups and to illustrate the distribution of physical inactivity (corresponds to the 'no PA' group) for the variables age, education level and household income. Binary logistic regression analyses were conducted to investigate the associations between the six socio-demographic variables with physical inactivity as being the dependent variable. The odds ratio (OR) was estimated for each BMI subgroup separately. This allows to investigate the socio-demographic correlates of not taking part in leisure-time PA for the different BMI subgroups. The significance level of all tests was based on $p < .05$. All analyses were performed using the IBM SPSS statistical software package, version 21.

Results

Sample characteristics

Table 2 presents the detailed characteristics of the whole sample. In all, 129,886 adults (44.8% male) were included; 47% of them were normal weight, 36.4% were overweight, and 16.6% were obese (grade I: 12.2%, grade II: 3.1%, grade III: 1.3%). Most adults were in the age group of 30–59 years (55.3%), had no migration background (86.8%), and were employed (79.6%).

Table 2. Sociodemographic characteristics by BMI subgroups in n (% cumulated).

Variables	BMI subgroups (in kg/m ²) (n = 129,886)				
	18,5 -< 25 (n = 61,029)	25 -< 30 (n = 47,253)	30 -< 35 (n = 15,864)	35 -< 40 (n = 4,045)	≥ 40 (n = 1,695)
Sex/Gender (%Male)	23,479 (38.5)	26,930 (57.0)	8,057 (50.8)	1,681 (41.6)	612 (36.1)
Employment (% unemployed)	6,512 (16.1)	5,476 (18.0)	2,122 (20.9)	597 (22.9)	277 (24.1)
Age (in years)					
18 -< 30	14,166 (23.4)	4,260 (9.0)	1,010 (6.4)	237 (5.9)	138 (8.2)
30 -< 60	32,892 (54.2)	25,472 (54.0)	8,378 (52.9)	2,251 (55.7)	1,008 (59.7)
60 -< 80	11,950 (19.7)	15,885 (33.7)	5,975 (37.7)	1,468 (36.3)	499 (29.5)
80+	1,630 (2.7)	1,571 (3.3)	478 (3.0)	83 (2.1)	44 (2.6)
Education level					
Low	14,194 (24.2)	17,538 (38.6)	7,373 (48.4)	1,994 (51.3)	859 (52.2)
Middle	16,539 (28.2)	11,758 (25.9)	3,764 (24.7)	971 (25.0)	409 (24.8)
High	27,854 (47.5)	16,138 (35.5)	4,103 (26.9)	924 (23.8)	379 (23.0)
Household income					
Bottom 20%	10,687 (18.6)	7,252 (16.5)	2,997 (20.5)	918 (24.7)	481 (30.6)
20–40%	11,201 (19.4)	9,015 (20.5)	3,365 (23.1)	898 (24.2)	359 (22.8)
40–60%	12,527 (21.8)	10,272 (23.4)	3,392 (23.3)	825 (22.2)	305 (19.4)
60–80%	10,673 (18.5)	8,375 (19.1)	2,547 (17.5)	582 (15.7)	227 (14.4)
Top 20%	12,506 (21.7)	9,021 (20.5)	2,284 (15.7)	495 (13.3)	201 (12.8)
Migration background					
Non	28,025 (86.0)	20,907 (87.6)	6,514 (86.6)	1,626 (86.6)	670 (87.1)
One-sided	1,445 (4.4)	977 (4.1)	344 (4.6)	90 (4.8)	41 (5.3)
Two-sided	3,108 (9.5)	1,989 (8.3)	660 (8.8)	161 (8.6)	58 (7.5)
Diabetes (% diabetic)	1,431 (2.6)	3,061 (7.2)	2,074 (14.5)	867 (23.7)	407 (27.0)

<https://doi.org/10.1371/journal.pone.0246634.t002>

PA levels

The prevalence of physical inactivity was lowest among normal weight individuals (26.2%), followed by overweight individuals (34%) and was highest among people with obesity grade III (56.7%). Only small group differences were found for '≤2 h', with percentages ranging between 21.6% (obesity grade III) and 26.4% (normal weight) and '2–4 h', with percentages ranging between 15.9% (obesity grade I) and 23.8% (normal weight). Furthermore, 23.6% of the normal-weight individuals engaged in PA for more than four hours per week; lower percentages were found for overweight people (18.7%) and the lowest for individuals with obesity grade II (9.6%) and III (8.5%). An overview of the percentages of all groups is presented in [Fig 1](#).

Socio-demographic characteristics among inactive individuals by age, education and household income

The prevalence of physical inactivity was lower among individuals with a high education level across all BMI groups. The highest difference was found in the normal-weight individuals (22.3%) and the lowest difference was found for individuals with obesity grade III (14.9%). With an increase in age, the proportion of people not exercising was higher. The percentage of physical inactivity in the age group of 18–29 years was significantly higher among individuals with obesity grade III (44.2%) compared to normal weight individuals (17.5%). This difference disappeared in the age group of ≥ 80 years (normal weight: 59.5%; obesity grade III: 68.2%).




Fig 1. Percentage distribution of PA categories (frequencies) for the BMI sample.

<https://doi.org/10.1371/journal.pone.0246634.g001>

The percentage of physical inactivity was lower among those with higher household income. Fig 2 demonstrates the percentage distribution of each variable across the BMI groups.

Relationship between socio-demographic variables and the risk of physical inactivity

Table 3 presents the results of the logistic regression analyses examining the relationship between the socio-demographic variables and the risk of physical inactivity. Being male was found to be associated with a slightly lower risk of physical inactivity compared to females, significant results were revealed for overweight (OR: 0.90, 95% CI: 0.82, 0.98), obesity grade I (OR: 0.82, 95% CI: 0.71, 0.95) and grade II (OR: 0.75, 95% CI: 0.56, 0.99). Compared with the reference age group of 18–29 years, the odds of physical inactivity were higher in the age groups of 30–59 years and ≥60 years. The highest risks were found for adults with obesity




Fig 2. Physical inactivity (in %) for education, household income and age across the BMI groups.

<https://doi.org/10.1371/journal.pone.0246634.g002>

Table 3. Binary logistic regression on dependent variable physical inactivity for the BMI sample^a.

	18,5 -< 25 (kg/m ²)	25 -< 30 (kg/m ²)	30 -< 35 (kg/m ²)	35 -< 40 (kg/m ²)	≥ 40 (kg/m ²)
Variable category	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
Male	0.96 (0.89–1.04)	0.90 (0.82–0.98)*	0.82 (0.71–0.95)*	0.75 (0.56–0.99)*	0.67 (0.44–1.04)
18 -< 30 years (Ref.)					
30 -< 60 years	2.04 (1.84–2.28)	1.76 (1.50–2.07)**	2.04 (1.48–2.80)**	1.26 (0.69–2.29)	1.08 (0.52–2.28)
60 -< 80 years	1.98 (1.72–2.28)	1.76 (1.47–2.10)**	2.12 (1.51–2.97)**	1.52 (0.80–2.91)	1.08 (0.46–2.52)
Low education (Ref.)					
Middle education	0.56 (0.51–0.62)*	0.64 (0.58–0.71)**	0.69 (0.58–0.83)**	0.84 (0.60–0.97)*	0.81 (0.49–2.52)
High education	0.42 (0.38–0.46)	0.47 (0.42–0.52)**	0.51 (0.42–0.60)**	0.65 (0.45–0.94)*	0.57 (0.32–0.99)*
Income level bottom 20% (Ref.)					
Income level 20–40%	0.93 (0.83–1.05)	0.93 (0.81–1.07)	0.74 (0.59–0.92)*	0.64 (0.43–0.97)	1.06 (0.59–1.89)
Income level 40–60%	0.76 (0.67–0.85)	0.76 (0.67–0.87)**	0.76 (0.61–0.94)*	0.62 (0.41–0.95)*	0.84 (0.44–1.60)
Income level 60–80%	0.67 (0.59–0.77)	0.68 (0.59–0.79)**	0.70 (0.55–0.89)*	0.40 (0.25–0.63)*	0.96 (0.46–2.03)
Income level top 20%	0.57 (0.50–0.66)	0.56 (0.48–0.65)**	0.64 (0.50–0.83)*	0.70 (0.42–1.16)**	0.93 (0.44–1.99)
No migration (Ref.)					
One-sided migration	0.90 (0.74–1.10)	0.91 (0.73–1.13)	1.26 (0.91–1.74)	1.32 (0.73–2.39)	0.93 (0.34–2.50)
Two-sided migration	1.53 (1.34–1.74)*	1.13 (0.96–1.34)	1.03 (0.77–1.37)	0.85 (0.51–1.44)	1.39 (0.56–3.46)
Unemployed	1.69 (1.44–1.98)**	1.48 (1.24–1.76)**	1.17 (0.89–1.52)	1.11 (0.74–1.68)	1.05 (0.57–1.95)
Diabetes	1.73 (1.34–2.22)**	1.33 (1.11–1.58)*	1.70 (1.38–2.10)**	1.34 (0.96–1.87)	0.99 (0.60–1.63)

Note: OR > 1 indicating a higher risk of reporting not to engage in PA.

* = p-value < 0.05,

** = p-value < 0.001, OR = Odds Ratio, 95% CI = 95% confidence interval of the odds ratio.

^a Model adjusted for sex, age, education, household income, migration background, employment, diabetes.

<https://doi.org/10.1371/journal.pone.0246634.t003>

grade I (OR_{30-<60}: 2.04, 95% CI 1.48, 2.80; OR_{60-<80}: 2.12, 95% CI 1.51, 2.97) and normal weight (OR_{30-<60}: 2.04, 95% CI 1.84, 2.28; OR_{60-<80}: 1.98, 95% CI 1.72, 2.28). In terms of household income, the odds for the top 20% group were between 7% (OR_{obesityIII}: 0.93, 95% CI: 0.44, 1.99) and 44% (OR_{overweight}: 0.56, 95% CI: 0.48, 0.65) lower than that for the bottom 20% group. The odds among people with a high education level were between 35% (OR_{obesityII}: 0.65, 95% CI: 0.45, 0.94) and 58% (OR_{normalweight}: 0.42, 95% CI: 0.38, 0.46) lower than in people with a low education level. Among ethnic minorities, the risk of inactivity was higher for two-sided migration (both parents belong to ethnic minorities) for normal weight, overweight and obesity grades I and III. For obesity grades I and II, higher odds were found for a one-sided migration (one parent belonging to an ethnic minority) compared to the reference group. Being unemployed was found to be associated with a significantly higher risk for normal weight (OR: 1.69, 95% CI: 1.44, 1.98) and overweight (OR: 1.48, 95% CI: 1.24, 1.76). Having diabetes was associated with a significantly higher risk of inactivity in normal weight (OR: 1.73, 95% CI: 1.34, 2.22), overweight (OR: 1.33, 95% CI: 1.11, 1.58) and obesity grade I (OR: 1.70, 95% CI: 1.38, 2.10).

Discussion

Summary of results

The current study aimed to examine patterns of physical inactivity (defined as not taking part in leisure-time PA) and to identify socio-demographic predictors of physical inactivity among different BMI groups and people with diabetes to support the development of targeted interventions promoting PA among individuals with obesity and diabetes. Overall, our results

revealed that in obese as well as in individuals with diabetes, socio-demographic predictors of physical inactivity were similar compared to those among normal weight and non-diabetic adults: a higher risk of inactivity was found in people of older age and those with low levels of education or low household income. The prevalence of physical inactivity was higher among people with obesity and diabetes than among normal weight and non-diabetic individuals, respectively. Considerable differences were also found between the obesity grades, with an increase of 11.7% from grade I to grade III. Nevertheless, every eighth obese and third diabetic person reported being physically active during their leisure-time for more than two hours per week, which is in accordance with the German recommendations for physical activity [39].

Implications for PA interventions

The calculated extent of PA corresponds to earlier findings from Germany [40] and other countries [20, 41] confirming the well-established relationship between physical inactivity and obesity. The lack of PA among people with obesity and diabetes is problematic given the importance of adequate PA for the treatment and control of both diseases [42]. Those affected may have additional barriers to PA such as more serious physical disabilities [43], perceived discomfort during training [44], and feeling inferior and inadequate [45, 46] which influence their willingness to be physically active. Nevertheless, some individuals with obesity and diabetes engaged in leisure-time PA, even for more than four hours weekly. This indicates that barriers to PA among this population can be overcome. In this regard, knowledge regarding the specific challenges obese and diabetic adults face when it comes to PA is important for planning interventions that target them [47].

In addition to disease-related barriers, this study demonstrates that socio-demographic factors may be fundamental in influencing the PA behaviour of adults who are obese or diabetic. The key role of socio-demographic factors was observed across all subgroups independent of body weight and the presence of diabetes. In accordance with previous studies, our results indicated a lower risk of physical inactivity among men [48, 49]. In the present study, markers of low socioeconomic status such as low educational levels and less household income were associated with a higher risk of physical inactivity, consistent with earlier studies [20, 50–52]. Individuals with lower socioeconomic status generally appear to have greater barriers to exercise [53]. This implies the need for policy makers to consider social gradients when designing PA programs particularly if they target diabetic or obese adults.

Our findings support previous research demonstrating obese and diabetic older people to be particularly physically inactive [54–57]. Given the increasing proportion of these groups in the general population, health care providers should actively approach affected older people for PA interventions motivating them to engage in regular PA. The high level of physical inactivity among the elderly population can result from physical limitations due to age-related loss of physical function [58] or specific motivation barriers [41, 59]. However, obese, and diabetic people face additional limitations. Diabetes is associated with a risk of mental disability, which may further lead to lower participation in PA [56]. In addition, the functional effects of obesity may exacerbate the age-related decline in physical functions, and it has been found that the self-reported mobility of obese people is reduced compared to lean older adults [60]. When developing PA interventions for older people, specific disease-related conditions of obesity and diabetes which influence PA behaviour should also be considered [56].

Earlier studies are consistent with our results, indicating a higher risk of physical inactivity among people with an migration background [18, 61, 62]. Migrant populations in high-income countries are particularly affected by type 2 diabetes and obesity, underlining the need for PA interventions for these groups [63]. When planning such interventions, the existence of

migration-related barriers on multiple levels should also be considered, including cultural and religious beliefs, environmental barriers, and (inter-)personal factors [64–67]. Programmes to improve access to sports clubs for migrants, such as the programme ‘Integration through Sport’ of the German Olympic Sports Federation [68], are promising. However, there is a need for interventions that specifically target older migrants as well as those with obesity and diabetes to positively influence their PA behaviour.

Strengths and limitations

We pooled data from several public-health surveys to create a large and nationwide sample. For countries such as Germany, this is the only means to yield comprehensive information on specific target groups with different socio-demographic and health-related backgrounds. We were therefore able to analyse the outcomes stratified by three grades of obesity, which has rarely been examined previously in this context [69]. The selected variables are robust and have withstood the test of time, which justifies the pooling of surveys from different years. Consideration must be given to potential limitations related to the pooling procedure. The cross-sectional design precludes the assessment of causal relationships. Moreover, the pooling and self-report assessment design prevented the gathering of precise information about PA behaviour (such as intensity and duration) or type of diabetes. The interpretation of the ‘no PA’ group as an indicator of physical inactivity must be assessed with caution. In several data sets people were coded as physically inactive, when they reported a lack of PA in the past week, while in others they were asked whether they were active in a typical week. Future studies should also include PA in other domains to provide a more complete picture of the ‘real’ overall levels of PA considering that these could also promote health. In particular, the assessment of PA in the domains of work or for transport would have been quite relevant in this regard. Self-reported data has further restrictions as the accuracy of self-reports is often limited due to recall errors, social desirability bias and misinterpretation of questions (i.e., differences in the interpretation of the term PA [70, 71]). Moreover, studies generally report a poor concordance between self-report and objective measures of PA [72]. However, the possibility of recall and measurement errors tend to be greater for light and moderate intensity activities than for vigorous ones [73]. High correlations between self-reported and measured height and weight have been reported [74]. Nevertheless, overweight and obese people tend to underestimate their body weight [75] and at the same time to overestimate their PA to an even higher degree than normal weight subjects [76]. In addition, although BMI is a typical method of measuring obesity, it is a poor diagnostic tool for identifying obesity in the general population [77].

Conclusion

In conclusion, the current study confirms that adults in Germany with diabetes or obesity are less physically active at leisure-time than those without diabetes and normal BMI, respectively. Identical socio-demographic factors for engaging in PA appears to be crucial in all groups. This will require intervention approaches to increase PA that reach the identified vulnerable target groups, particularly the older population and adults with a migration background who are also affected by obesity or diabetes.

Supporting information

S1 Fig. Percentage distribution of PA (categories (frequencies) for the diabetes sample. (TIF)

S2 Fig. Physical inactivity (in %) for education, household income and age across the diabetes groups.

(TIF)

S1 Table. Binary logistic regression on dependent variable physical inactivity for the diabetes sample. ^a Note: OR > 1 indicating a higher risk of reporting not to engage in PA. * = p-value <0.05, ** = p-value <0.001, OR = Odds Ratio, 95% CI = 95% confidence interval of the odds ratio. ^a Model adjusted for sex, age, education, household income, migration background, employment, BMI.

(DOCX)

Author Contributions**Conceptualization:** Stephanie Linder, Karim Abu-Omar, Anne K. Reimers, Heiko Ziemainz.**Data curation:** Sven Messing.**Formal analysis:** Stephanie Linder, Sven Messing, Mustafa Sarshar.**Funding acquisition:** Karim Abu-Omar.**Methodology:** Sven Messing, Mustafa Sarshar.**Project administration:** Heiko Ziemainz.**Resources:** Karim Abu-Omar, Sven Messing.**Supervision:** Anne K. Reimers, Heiko Ziemainz.**Writing – original draft:** Stephanie Linder.**Writing – review & editing:** Karim Abu-Omar, Wolfgang Geidl, Mustafa Sarshar, Anne K. Reimers, Heiko Ziemainz.**References**

1. World Health Organization. Global status report on noncommunicable diseases 2014. Vol. (No. WHO/N. 2014.
2. Heidemann C, Kuhnert R, Born S, Scheidt-Nave C. 12-Month prevalence of known diabetes mellitus in Germany. *J Heal Monit [Internet]*. 2017; 2(1):43–50. Available from: <http://edoc.rki.de/oa/articles/rei427ccep96/PDF/27H3uMDZmc4wQ.pdf>
3. Dagenais GR, Auger P, Bogaty P, Gerstein H, Lonn E, Yi Q, et al. Increased occurrence of diabetes in people with ischemic cardiovascular disease and general and abdominal obesity. *Can J Cardiol*. 2003 Nov; 19(12):1387–1391. PMID: [14631473](https://pubmed.ncbi.nlm.nih.gov/14631473/)
4. Mensink GBM, Schienkiewitz A, Haftenberger M, Lampert T, Ziese T, Scheidt-Nave C. Übergewicht und Adipositas in Deutschland: Ergebnisse der Studie zur Gesundheit Erwachsener in Deutschland (DEGS1). *Bundesgesundheitsblatt—Gesundheitsforsch—Gesundheitsschutz*. 2013; 56(5–6):786–94.
5. World Health Organization. Obesity: Preventing and managing the global epidemic. World Health Organization—Technical Report Series. 2000.
6. Hu F., Mansons J.E., Stampfer M.J., Colditz G., Liu S., Solomon C.G., & Willett WC. Diet, lifestyle, and the risk of type 2 diabetes mellitus in women. *N Engl J Med*. 2001; 345(11):790–7. <https://doi.org/10.1056/NEJMoa010492> PMID: [11556298](https://pubmed.ncbi.nlm.nih.gov/11556298/)
7. Church, Timothy S, Cheng Y.J., Earnest C.P., Barlow C.E., Gibbons L.W., Priest E.L., & Blair SN. Exercise Capacity and Body Composition as Predictors of Mortality Among Men. *Diabetes Care*. 2004; 27(1):83–8. <https://doi.org/10.2337/diacare.27.1.83> PMID: [14693971](https://pubmed.ncbi.nlm.nih.gov/14693971/)
8. Sigal RJ, Kenny GP, Wasserman DH, Castaneda-Sceppa C. Physical Activity/Exercise and Type 2 Diabetes [Internet]. *Am Diabetes Assoc*. 2004 [cited 2020 Jul 29]. Available from: <https://care.diabetesjournals.org/content/27/10/2518.short> <https://doi.org/10.2337/diacare.27.10.2518> PMID: [15451933](https://pubmed.ncbi.nlm.nih.gov/15451933/)

9. Saris WHM, Blair SN, Van Baak MA, Eaton SB, Davies PSW, Di Pietro L, et al. How much physical activity is enough to prevent unhealthy weight gain? Outcome of the IASO 1st stock conference and consensus statement. *Obes Rev*. 2003; 4(2):101–14. <https://doi.org/10.1046/j.1467-789x.2003.00101.x> PMID: 12760445
10. Vanden Bosch ML, Robbins LB, Anderson K. Correlates of Physical Activity in Middle-Aged Women With and Without Diabetes. *West J Nurs Res* [Internet]. 2015 Dec 1 [cited 2020 Jul 29]; 37(12):1581–603. Available from: https://journals.sagepub.com/doi/abs/10.1177/0193945914541333?casa_token=BIPWGI5XjJwAAAAA:kfoJYromVquEUbYyq-9FG3ROw953C3PNQcO_-dFBnfhKkdB_lqjTxLn43qH8U7dinbJ9oawwb2cmMg <https://doi.org/10.1177/0193945914541333> PMID: 25004922
11. Petersen L, Schnohr P, Sørensen TIA. Longitudinal study of the long-term relation between physical activity and obesity in adults. *Int J Obes* [Internet]. 2004 [cited 2020 Jul 29]; 28(1):105–12. Available from: <https://www.nature.com/articles/0802548> <https://doi.org/10.1038/sj.ijo.0802548> PMID: 14647181
12. Lee IM, Paffenbarger RS. Associations of light, moderate, and vigorous intensity physical activity with longevity: The Harvard Alumni Health Study. *Am J Epidemiol*. 2000; 151(3):293–9. <https://doi.org/10.1093/oxfordjournals.aje.a010205> PMID: 10670554
13. Trost SG, Owen N, Bauman AE, Sallis JF, Brown W. Correlates of adults' participation in physical activity: Review and update. *Med Sci Sports Exerc* [Internet]. 2002 [cited 2020 Aug 8]; 34(12):1996–2001. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/12471307> <https://doi.org/10.1097/00005768-200212000-00020> PMID: 12471307
14. O'Donoghue G, Kennedy A, Puggina A, Aleksovska K, Buck C, Burns C, et al. Socio-economic determinants of physical activity across the life course: A “DEterminants of DIet and Physical ACTivity” (DEDI-PAC) umbrella literature review. Vol. 13, *PLoS ONE*. Public Library of Science; 2018.
15. Gidlow C, Johnston LH, Crone D, Ellis N, James D. A systematic review of the relationship between socio-economic position and physical activity. *Health Educ J* [Internet]. 2006 Dec [cited 2020 Aug 8]; 65(4):338–67. Available from: www.sagepub.com
16. Demarest S, Van Oyen H, Roskam AJ, Cox B, Regidor E, Mackenbach JP, et al. Educational inequalities in leisure-time physical activity in 15 European countries. *Eur J Public Health* [Internet]. 2014 [cited 2020 Aug 8]; 24(2):199–204. Available from: <https://academic.oup.com/eurpub/article-abstract/24/2/199/447727> <https://doi.org/10.1093/eurpub/ckt061> PMID: 23748597
17. Marques A, Sarmento H, Martins J, Saboga Nunes L. Prevalence of physical activity in European adults—Compliance with the World Health Organization's physical activity guidelines. *Prev Med (Baltim)* [Internet]. 2015 [cited 2020 Aug 8]; 81:333–8. Available from: <https://www.sciencedirect.com/science/article/pii/S0091743515003035> <https://doi.org/10.1016/j.yjmed.2015.09.018> PMID: 26449407
18. Rommel A, Saß AC, Born S, Ellert U. Die gesundheitliche Lage von Menschen mit Migrationshintergrund und die Bedeutung des sozioökonomischen Status: Erste Ergebnisse der Studie zur Gesundheit Erwachsener in Deutschland (DEGS1). *Bundesgesundheitsblatt—Gesundheitsforsch—Gesundheitsschutz* [Internet]. 2015 [cited 2020 Aug 8]; 58(6):543–52. Available from: <https://europepmc.org/article/med/25824135>
19. Ae B, Rs R, Jf S, Jc W, Rjf L, Bw M, et al. Correlates of physical activity: why are some people physically active and others not—Supplementary appendix. *Lancet* [Internet]. 2012 [cited 2020 Aug 8]; 6736(12). Available from: <https://www.sciencedirect.com/science/article/pii/S0140673612607351>
20. Marques A, Martins J, Peralta M, Catunda R, Nunes LS. European adults' physical activity socio-demographic correlates: A cross-sectional study from the European Social Survey. *PeerJ* [Internet]. 2016 [cited 2020 Aug 8]; 2016(6). Available from: <https://peerj.com/articles/2066/> <https://doi.org/10.7717/peerj.2066> PMID: 27280072
21. Lahti J, Holstila A, Lahelma E, Rahkonen O. Leisure-time physical activity and all-cause mortality. *PLoS One*. 2014 Jul 2; 9(7). <https://doi.org/10.1371/journal.pone.0101548> PMID: 24988295
22. Holtermann A, Hansen J, . . . HB-B journal of sports, 2012 undefined. The health paradox of occupational and leisure-time physical activity. *bjsm.bmj.com* [Internet]. [cited 2021 Jan 20]; <http://bjsm.bmj.com/>
23. Robert Koch Institute Department of Epidemiology and Health Monitoring. German National Health Interview and Examination Survey 1998. Scientific Use File 4th version. 2016.
24. Robert Koch Institute Department of Epidemiology and Health Monitoring. German Telephone Health Survey (GSTel03). Scientific Use File second version. 2014.
25. Robert Koch Institute Department of Epidemiology and Health Monitoring. German Health Interview and Examination Survey for Adults (DEGS1). Scientific Use File first version. 2015;
26. Robert Koch Institute Department of Epidemiology and Health Monitoring. German Health Update 2009 (GEDA 2009). Scientific Use File second version. 2014.
27. Robert Koch Institute Department of Epidemiology and Health Monitoring. German Health Update 2010 (GEDA 2010). Scientific Use File third version. 2014.

28. Robert Koch Institute Department of Epidemiology and Health Monitoring. German Health Update 2012 (GEDA 2012). Scientific Use File first version. 2014.
29. Robert Koch Institute Department of Epidemiology and Health Monitoring. German Health Update 2014/2015-EHIS (GEDA 2014/2015-EHIS). Scientific Use File first version. 2018.
30. GESIS. ALLBUS/GGSS 1998 (Allgemeine Bevölkerungsumfrage der Sozialwissenschaften/German General Social Survey 1998). ZA3000 Data file Version 2.0.0. 2012.
31. GESIS. ALLBUS/GGSS 2004 (Allgemeine Bevölkerungsumfrage der Sozialwissenschaften/German General Social Survey 2004). ZA3762 Data file Version 2.0.0. 2011.
32. GESIS. GESIS Panel—Extended Edition. ZA 5664 Data file Version 24.0.0. 2018.
33. GESIS. ALLBUS/GGSS 2014 (Allgemeine Bevölkerungsumfrage der Sozialwissenschaften/German General Social Survey 2014). ZA5240 Data file Version 2.2.0. 2018.
34. Börsch-Supan A, Brandt M, Hunkler C, Kneip T, Korbmayer J, Malter F, et al. Data resource profile: The survey of health, ageing and retirement in Europe (share). *Int J Epidemiol* [Internet]. 2013 [cited 2020 Aug 8]; 42(4):992–1001. Available from: <https://academic.oup.com/ije/article-abstract/42/4/992/657275> <https://doi.org/10.1093/ije/dyt088> PMID: 23778574
35. Malter F, Börsch-Supan A, editors. SHARE Wave 5: Innovations & Methodology. Munich: MEA, Max Planck Institute for Social Law and Social Policy; 2015.
36. Börsch-Supan A. Survey of Health, Ageing and Retirement in Europe (SHARE) Wave 5. Release version: 1.0.0. SHARE-ERIC. Data set. 2015.
37. Rütten A, Abu-Omar K, Adlwarth W, Meierjürgen R. Bewegungssarme lebensstile—Zur klassifizierung unterschiedlicher zielgruppen für eine gesundheitsförderliche körperliche aktivierung. *Gesundheitswesen* [Internet]. 2007 Jul [cited 2020 Aug 8]; 69(7):393–400. Available from: <http://www.thieme-connect.de/DOI/DOI?10.1055/s-2007-984440>
38. Jacobs E, Rathmann W, Tönnies T, Arendt D, Marchowicz M, Veith L, et al. Age at diagnosis of Type 2 diabetes in Germany: a nationwide analysis based on claims data from 69 million people. *Diabet Med* [Internet]. 2019 Aug 27 [cited 2020 Aug 8]; dme.14100. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/dme.14100> PMID: 31390484
39. Pfeifer K, Rütten A. National Recommendations for Physical Activity and Physical Activity Promotion [Internet]. Vol. 79, Gesundheitswesen, Supplement. 2017 [cited 2020 Aug 8]. https://opus4.kobv.de/opus4-fau/files/7827/Bewegungsempfehlung_OPUS.pdf
40. Luzak A, Heier M, Thorand B, Laxy M, Nowak D, Peters A, et al. Physical activity levels, duration pattern and adherence to WHO recommendations in German adults. *PLoS One*. 2017 Feb 1; 12(2). <https://doi.org/10.1371/journal.pone.0172503> PMID: 28245253
41. Hays LM, Clark DO. Correlates of physical activity in a sample of older adults with type 2 diabetes. *Diabetes Care* [Internet]. 1999 [cited 2020 Aug 8]; 22(5):706–12. Available from: <https://care.diabetesjournals.org/content/22/5/706.short> <https://doi.org/10.2337/diacare.22.5.706> PMID: 10332670
42. Li G, Zhang P, Wang J, Gregg EW, Yang W, Gong Q, et al. The long-term effect of lifestyle interventions to prevent diabetes in the China Da Qing Diabetes Prevention Study: a 20-year follow-up study. *Lancet* [Internet]. 2008 [cited 2020 Aug 8]; 371(9626):1783–9. Available from: <https://www.researchgate.net/publication/5348531> [https://doi.org/10.1016/S0140-6736\(08\)60766-7](https://doi.org/10.1016/S0140-6736(08)60766-7) PMID: 18502303
43. Gregg EW, Beckles GLA, Williamson DF, Leveille SG, Langlois JA, Engelgau MM, et al. Diabetes and physical disability among older U.S. adults. *Diabetes Care* [Internet]. 2000 [cited 2020 Aug 8]; 23(9):1272–7. Available from: <https://care.diabetesjournals.org/content/23/9/1272.short> <https://doi.org/10.2337/diacare.23.9.1272> PMID: 10977018
44. Thomas N, Alder E, Leese GP. Barriers to physical activity in patients with diabetes. *Postgrad Med J* [Internet]. 2004 [cited 2020 Aug 8]; 80(943):287–91. Available from: <http://pmj.bmj.com/> <https://doi.org/10.1136/pgmj.2003.010553> PMID: 15138320
45. Bombak AE. Obese persons' physical activity experiences and motivations across weight changes: A qualitative exploratory study Energy balance-related behaviours. *BMC Public Health*. 2015 Nov 14; 15(1). <https://doi.org/10.1186/s12889-015-2459-x>
46. Toft BS, Uhrenfeldt L. The lived experiences of being physically active when morbidly obese: A qualitative systematic review. *Int J Qual Stud Health Well-being* [Internet]. 2015 Sep 22 [cited 2020 Aug 8]; 10. Available from: <http://dx.doi.org/10.3402/qhw.v10.28577> PMID: 26400462
47. Catania JA, Dolcini MM, Sallis JF, Owen N, Fotheringham MJ. A commentary on Sallis, Owen, and Fotheringham's perspective on "behavioral epidemiology: A systematic framework to classify phases of research on health promotion and disease prevention" [Internet]. Vol. 24, *Annals of Behavioral Medicine*. 2002 [cited 2020 Aug 8]. Available from: https://idp.springer.com/authorize/casa?redirect_uri=https://link.springer.com/article/10.1007%25252FBF02895665%3Fref%3Dvidupdatez.com/

- image&casa_token=HJN8Cvsx0HIAAAAA:wK1T6hc9ue6eHvW3cyRdlgjxWE0CfvVkpMRIKehRdfyt4BTcj5FICmH1CHNn6HzBoxUg2XSMBpxOduYdw
48. Pitsavos C, Panagiotakos DB, Lentzas Y, Stefanadis C. Epidemiology of leisure-time physical activity in socio-demographic, lifestyle and psychological characteristics of men and women in Greece: The ATTICA Study. *BMC Public Health*. 2005 Apr 18; 5. <https://doi.org/10.1186/1471-2458-5-37> PMID: 15836794
 49. Bauman A, Bull F, Chey T, Craig CL, Ainsworth BE, Sallis JF, et al. The international prevalence study on physical activity: Results from 20 countries. *Int J Behav Nutr Phys Act*. 2009 Mar 31; 6. <https://doi.org/10.1186/1479-5868-6-21> PMID: 19335883
 50. Varo J, . . . MM-G-I journal, 2003 undefined. Distribution and determinants of sedentary lifestyles in the European Union. *academic.oup.com* [Internet]. [cited 2021 Jan 18]; <https://academic.oup.com/ije/article-abstract/32/1/138/642829>
 51. Morrato EH, Hill JO, Wyatt HR, Ghushchyan V, Sullivan PW. Physical activity in U.S. adults with diabetes and at risk for developing diabetes, 2003. *Diabetes Care* [Internet]. 2007 [cited 2020 Aug 8]; 30(2):203–9. Available from: <https://care.diabetesjournals.org/content/30/2/203.short> <https://doi.org/10.2337/dc06-1128> PMID: 17259482
 52. Sliemers SM, Schuit AJ, Chinapaw MJM, Seidell JC, van Mechelen W. Disagreement in physical activity assessed by accelerometer and self-report in subgroups of age, gender, education and weight status. *Int J Behav Nutr Phys Act*. 2009 Mar 25; 6.
 53. Chinn DJ, White M, Harland J, Drinkwater C, Raybould S. Barriers to physical activity and socioeconomic position: Implications for health promotion. *J Epidemiol Community Health*. 1999; 53(3):191–2. <https://doi.org/10.1136/jech.53.3.191> PMID: 10396499
 54. Haley C, Andel R. Correlates of physical activity participation in community-dwelling older adults. *J Aging Phys Act* [Internet]. 2010 [cited 2020 Aug 8]; 18(4):375–89. Available from: <https://journals.humankinetics.com/view/journals/japa/18/4/article-p375.xml> <https://doi.org/10.1123/japa.18.4.375> PMID: 20956840
 55. Moschny A, Platen P, Klaßen-Mielke R, Trampisch U, Hinrichs T. Physical activity patterns in older men and women in Germany: A cross-sectional study. *BMC Public Health*. 2011; 11. <https://doi.org/10.1186/1471-2458-11-559> PMID: 21752288
 56. Zhao G, Ford ES, Li C, Balluz LS. Physical activity in U.S. older adults with diabetes mellitus: Prevalence and correlates of meeting physical activity recommendations. *J Am Geriatr Soc*. 2011 Jan; 59(1):132–7. <https://doi.org/10.1111/j.1532-5415.2010.03236.x> PMID: 21226683
 57. Riebe D, Blissmer BJ, Greaney ML, Ewing Garber C, Lees FD, Clark PG. The relationship between obesity, physical activity, and physical function in older adults. *J Aging Health* [Internet]. 2009 Dec [cited 2020 Aug 8]; 21(8):1159–78. Available from: <http://jah.sagepub.com> <https://doi.org/10.1177/0898264309350076> PMID: 19897781
 58. Horne M, Skelton DA, Speed S, Todd C. Perceived barriers to initiating and maintaining physical activity among South Asian and White British adults in their 60s living in the United Kingdom: A qualitative study. *Ethn Heal*. 2013 Dec 1; 18(6):626–45. <https://doi.org/10.1080/13557858.2013.814762> PMID: 23834070
 59. Franco MR, Tong A, Howard K, Sherrington C, Ferreira PH, Pinto RZ, et al. Older people's perspectives on participation in physical activity: A systematic review and thematic synthesis of qualitative literature. *Br J Sports Med* [Internet]. 2015 [cited 2020 Aug 8]; 49(19):1268–76. Available from: <http://dx.doi.org/10.1136/bjsports-2014-094015> PMID: 25586911
 60. Jenkins KR. Obesity's Effects on the Onset of Functional Impairment among Older Adults. *Gerontologist* [Internet]. 2004 [cited 2020 Aug 8]; 44(2):206–16. Available from: <https://academic.oup.com/gerontologist/article-abstract/44/2/206/876633> <https://doi.org/10.1093/geront/44.2.206> PMID: 15075417
 61. Kobel S, Kettner S, Hermeling L, Dreyhaupt J, Steinacker JM. Objectively assessed physical activity and weight status of primary school children in Germany with and without migration backgrounds. *Public Health* [Internet]. 2019 [cited 2020 Aug 8]; 173:75–82. Available from: <https://www.sciencedirect.com/science/article/pii/S0033350619301544> <https://doi.org/10.1016/j.puhe.2019.05.009> PMID: 31255961
 62. Lämmle L, Worth A, Bös K. Socio-demographic correlates of physical activity and physical fitness in German children and adolescents. *Eur J Public Health* [Internet]. 2012 [cited 2020 Aug 8]; 22(6):880–4. Available from: <https://academic.oup.com/eurpub/article-abstract/22/6/880/546109> <https://doi.org/10.1093/eurpub/ckr191> PMID: 22219503
 63. Meeks K, Freitas-Da-Silva D, Adeyemo A, Beune E, Modesti P, Stronks K, et al. Disparities in type 2 diabetes among ethnic minority groups resident in Europe—a meta-analysis. *Eur J Public Health* [Internet]. 2015 [cited 2020 Aug 8]; 25(suppl_3). Available from: <https://academic.oup.com/eurpub/article->

[abstract/25/suppl_3/ckv174.014/2484261](https://doi.org/10.1007/s11739-015-1302-9) <https://doi.org/10.1007/s11739-015-1302-9> PMID: 26370238

64. Langøien LJ, Terragni L, Rugseth G, Nicolaou M, Holdsworth M, Stronks K, et al. Systematic mapping review of the factors influencing physical activity and sedentary behaviour in ethnic minority groups in Europe: A DEDIPAC study. Vol. 14, *International Journal of Behavioral Nutrition and Physical Activity*. BioMed Central Ltd.; 2017.
65. O'Driscoll T, Banting LK, Borkoles E, Eime R, Polman R. A systematic literature review of sport and physical activity participation in culturally and linguistically diverse (CALD) migrant populations. *J Immigr Minor Heal* [Internet]. 2014 [cited 2020 Aug 8]; 16(3):515–30. Available from: [https://doi.org/10.1007/s10903-013-9857-x](https://doi.org/10.1007/s10903-013-9857-x&casa_token=EFa3JyfTznEAAAAA:IWkakXuB5cmwzaDKOe3xMgT5P8UrpMs2n6VlqLA2hpBaU6G3vPRbWbtpdU28aW6S6QU4gUjghf70j4FQ) PMID: 23771744
66. Caperchione CM, Kolt GS, Mummery WK. Physical activity in culturally and linguistically diverse migrant groups to Western society: A review of barriers, enablers and experiences. Vol. 39, *Sports Medicine*. 2009. p. 167–77. <https://doi.org/10.2165/00007256-200939030-00001> PMID: 19290674
67. Waxenegger A, Mayerl H, Stolz E, Rásky É, Freidl W. Impact of migration background on health of children and adolescents in Austria results of the Austrian health survey 2014. *Cent Eur J Public Health* [Internet]. 2018 [cited 2020 Aug 8]; 26(2):132–6. Available from: <https://europepmc.org/article/med/28701003> <https://doi.org/10.21101/cejph.a4917> PMID: 30102502
68. Sportbund DS-F am MDO, 2014 undefined. Integration Durch Sport—Programmkonzeption.
69. Slagter SN, Van Vliet-Ostapchouk J V., Van Beek AP, Keers JC, Lutgers HL, Van Der Klauw MM, et al. Health-Related quality of life in relation to obesity grade, type 2 diabetes, metabolic syndrome and inflammation. *PLoS One*. 2015 Oct 16; 10(10). <https://doi.org/10.1371/journal.pone.0140599> PMID: 26474291
70. Ashe MC, Miller WC, Eng JJ, Noreau L. Older adults, chronic disease and leisure-time physical activity. *Gerontology* [Internet]. 2009 [cited 2020 Aug 8]; 55(1):64–72. Available from: <https://www.karger.com/article/Abstract/141518> <https://doi.org/10.1159/000141518> PMID: 18566534
71. Krems C, Lührmann PM, Neuhäuser-Berthold M. Physical activity in young and elderly subjects. *J Sports Med Phys Fitness* [Internet]. 2004 [cited 2020 Aug 8]; 44(1):71–6. Available from: <http://search.proquest.com/openview/e2e3c054b788959696a7ddef9ebe6d52/1?pq-origsite=scholar&cbl=4718> PMID: 15181393
72. Steene-Johannessen J, . . . SA-M and science, 2016 undefined. Are self-report measures able to define individuals as physically active or inactive? *ncbi.nlm.nih.gov* [Internet]. [cited 2021 Jan 18]; <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6235100/>
73. Livingstone M, Robson P, McCarthy S, Kiely M, Harrington K, Browne P, et al. Physical activity patterns in a nationally representative sample of adults in Ireland. *Public Health Nutr*. 2001; 4(5a):1107–16. <https://doi.org/10.1079/phn2001192> PMID: 11820924
74. Kuczmarski M, Kuczmarski R, Dietetic MN-J of the A, 2001 undefined. Effects of age on validity of self-reported height, weight, and body mass index: findings from the Third National Health and Nutrition Examination Survey, 1988–1994. Elsevier [Internet]. [cited 2021 Jan 18]; <https://www.sciencedirect.com/science/article/pii/S0002822301000086>
75. Morrissey SL, Whetstone LM, Cummings DM, Owen LJ. Comparison of self-reported and measured height and weight in eighth-grade students. *J Sch Health* [Internet]. 2006 [cited 2020 Aug 8]; 76(10):512–5. Available from: <https://academic.oup.com/aje/article-abstract/115/2/223/77664> <https://doi.org/10.1111/j.1746-1561.2006.00150.x> PMID: 17096824
76. Nelson DE, Holtzman D, Bolen J, Stanwyck CA, Mack KA. Reliability and validity of measures from the Behavioral Risk Factor Surveillance System (BRFSS). *Soz Präventivmed* [Internet]. 2001 [cited 2020 Aug 8]; 46(SUPPL. 1). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/11851091>.
77. Lavie CJ, McAuley PA, Church TS, Milani RV., Blair SN. Obesity and cardiovascular diseases: Implications regarding fitness, fatness, and severity in the obesity paradox. Vol. 63, *Journal of the American College of Cardiology*. Elsevier USA; 2014. p. 1345–54.