



# A systematic review of infant feeding food allergy prevention guidelines - can we AGREE?

Sandra L. Vale<sup>a\*</sup>, Monique Lobb<sup>b</sup>, Merryn J. Netting<sup>c</sup>, Kevin Murray<sup>d</sup>, Rhonda Clifford<sup>e</sup>,  
Dianne E. Campbell<sup>f</sup> and Sandra M. Salter<sup>e</sup>

## ABSTRACT

Food allergy is a significant issue worldwide, particularly in Westernised countries. There is no clear explanation why food allergy appears to have increased so rapidly in recent years, particularly in young children, hence ongoing research to identify effective primary prevention strategies. Food allergy prevention guidelines for health professionals have been developed based on existing clinical trial evidence for effective translation and implementation. As these guidelines underpin clinical practice, it is important to ensure robust processes of development. We conducted a systematic review to identify food allergy prevention guidelines for health professional use; to compare the recommendations made by the identified guideline documents; and to assess the quality of the identified guideline documents.

We searched Medline, EMBASE, CINAHL, Scopus, Global Health and Guidelines International Network for the period 1990 to 13 August 2019, to identify articles referring to English-language food allergy prevention guidelines or the guidelines themselves. A grey literature search of Google Scholar and reference checking was also undertaken. The guidelines were compared for recommendation similarities and differences. An Appraisal Guidelines for Research and Evaluation (AGREE II) appraisal was undertaken to assess guideline quality.

The electronic database search yielded 1121 publications and reference checking identified an additional 16 publications. After title, abstract and full text screening, data extraction was undertaken on 156 publications and with additional reference checking, 28 food allergy prevention guidelines and advice documents were identified. Comparison of the recommendations within the guidelines and advice documents indicated the greatest variation in recommendations related to exclusive breastfeeding and timing of solid food introduction. Eight of the 10 guidelines and none of the 18 advice documents met the quality threshold set by the reviewers. Overall, documents specifically termed "guidelines" scored better than advice documents when assessed using the AGREE II tool.

Variation in recommendations may create confusion for health professionals and result in inconsistent advice being provided to parents, and less translation of the evidence into actual food allergy reduction in the population. Appraisal using the AGREE II tool identified that there is considerable room for improvement in the development of guidelines and advice documents for food allergy prevention. The AGREE II appraisal identified common areas of poorer quality development and/or documentation of processes to inform future guideline development. Based

<sup>a</sup>School of Allied Health, Faculty of Health and Medical Sciences, The University of Western Australia, 35 Stirling Highway, Perth, WA, 6009, Australia

\*Corresponding author. E-mail: [sandravale@iinet.net.au](mailto:sandravale@iinet.net.au)

Full list of author information is available at the end of the article.

<http://doi.org/10.1016/j.waojou.2021.100550>

Received 5 November 2020; Received in revised form 14 April 2021; Accepted 26 April 2021

Online publication date xxx

1939-4551/© 2021 The Author(s). Published by Elsevier Inc. on behalf of World Allergy Organization. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

on this study, we recommend the use of validated guideline development tools, to direct food allergy prevention guideline review or development. Use of the AGREE II tool, to direct the review and development of guidelines, is very likely to improve guideline quality.

**Keywords:** Allergy prevention, Guidelines, Food allergy, Infant feeding

## INTRODUCTION

Food allergy is a significant issue worldwide, particularly in Westernised countries.<sup>1-4</sup> While good epidemiological data are lacking, it is estimated that worldwide, more than 220 million people have a food allergy.<sup>5-8</sup> In developed countries, food allergy is more common in children,<sup>9</sup> with verified food allergy prevalence ranging from 6% to 10% in infants and 2% and 5% in adults.<sup>10-13</sup> There is also evidence of a high prevalence of food allergy in developing countries, with a 2.5% incidence of challenge-proven food allergy observed in South Africa in 2015<sup>14</sup> and reported prevalence in China increasing from 3.5% to 7.7% between 1999-2009.<sup>14</sup>

Food allergy impacts greatly on the quality of life of children and their caregivers,<sup>15,16</sup> and contributes significant direct health costs for the healthcare system and even larger costs for families with a food allergic child.<sup>17</sup> Preventing food allergy is, therefore, a logical step in minimising the mortality, significant morbidity, and related costs associated with this condition.

Food allergy prevention strategies based on delaying introduction of common food allergens in high risk individuals have been largely ineffective,<sup>18,19</sup> and, consequently, the search for effective primary prevention strategies has shifted to interventions including: timeframe for exclusive breastfeeding, breastmilk substitutes, early introduction of foods including common food allergen introduction, vitamin D and omega-3 fatty acid supplementation, and modification of the maternal and infant microbiome.<sup>20-24</sup>

In 2015, several randomised controlled trials (RCTs) and a meta-analysis of these trials examining the effect of early introduction of food allergens on the development of food allergy were published.<sup>24-26</sup> Results from these trials have led

to changes in infant feeding advice for food allergy prevention.<sup>24-26</sup> Several large RCTs have examined the effect of early introduction of egg into the diet compared with delayed introduction and have shown some evidence that, depending on the baseline risk status of the treatment group, prevention of IgE-egg sensitisation or egg allergy may be associated with earlier introduction of egg.<sup>25,27,28</sup> However, the Enquiring about Tolerance (EAT) study which examined the effect of introduction of 6 foods (cow's milk, egg, peanut, wheat, fish, sesame) to the diet of exclusively breastfed infants from 3 months of age,<sup>26</sup> and the Prevention of Egg allergy with Tiny amount of InTake (PETIT) study in infants with well controlled eczema (high risk)<sup>28</sup> showed the greatest benefits with cooked egg.

The Learning Early About Peanut (LEAP) study, which randomised 640 high risk infants aged 4-11 months to consume or avoid peanut until 60 months of age, was a pivotal peanut allergy prevention study.<sup>24</sup> This study demonstrated an 86.1% relative reduction in peanut allergy prevalence in the consumption group compared to the control group.<sup>24</sup> These studies provide a foundation for evidence-based food allergy prevention guidelines and advice documents.

Clinical practice guidelines are designed for health professionals and are important to ensure the best health outcomes for patients.<sup>29,30</sup> Food allergy prevention guidelines for health professionals should be developed based on existing clinical trial evidence for effective translation and implementation. As guidelines are intended to underpin clinical practice, it is important that high quality evidence is integrated, and the development and reported process is robust.

The Appraisal Guidelines for Research and Evaluation (AGREE II) tool<sup>31,32</sup> is a validated

instrument used to assess guideline quality.<sup>33,34</sup>

The AGREE II tool assesses guidelines across 6 domains: scope/purpose; stakeholder involvement; rigor of development; presentation; applicability; and editorial independence. Assessing the quality of guidelines (including advice documents) is important to determine (1) if adequate guideline development processes were used; and (2) where the guideline documents differ in quality based on their nominated title (eg, guideline vs. consensus statement) and processes used in development.

This systematic review aimed to identify food allergy prevention guidelines for health professional use; to compare the recommendations made by the identified guideline documents; and to assess the quality of the identified guideline documents.

## METHODOLOGY

In this review, the word "guideline" is defined as any document termed an evidence-based guideline, expert recommendation, consensus statement, joint statement, position paper, or clinical report/guidance document, in accordance with the World Health Organisation (WHO) definition of "any document that contains recommendation for clinical health practice or public health policy".<sup>35</sup> The AGREE II consortium acknowledges that documents specifically titled as "guidelines" generally score higher when appraised using the AGREE II tool than non-official guideline documents.<sup>36</sup> Hence, for this AGREE II assessment, a comparison has been made between documents specifically named "guidelines" by their authors in their title and all other documents which have been grouped together as "advice documents".

### Search strategy

The literature was systematically searched to identify guidelines and advice documents developed for health professional use for the primary prevention of food allergy. A two-phase search was employed with the initial phase identifying both guideline documents and publications that referred to a guideline document. The second phase involved sourcing all the guideline documents identified in phase 1.

The following databases were searched for the period 1990 to 13 August 2019: Medline, EMBASE, CINAHL, Scopus, Global Health and Guidelines International Network. A grey literature search of Google Scholar was also undertaken. In addition, a search of the reference lists from publications included in the full-text screen and references from identified guideline documents was undertaken. English language restrictions were applied. Guideline documents and studies reporting guideline documents between January 1990 and the date of the search (13 August 2019) were sought.

The following search terms were used: ("health professionals" OR "general practitioners" OR "nurses" OR "dietitians" OR "dieticians") AND ("food allergy" OR "food hypersensitivity" OR "allergy") AND ("guidelines" OR "guideline" OR "policies" OR "policy" OR "strategy" OR "recommendation" OR "statement" OR "protocol" OR "consensus" OR "clinical practice") AND ("prevention" OR "primary prevention").

### Article selection

Publications identified from the search were exported to Endnote reference management software, version 8 (Clarivate Analytics, Philadelphia, PA), duplicates removed, then uploaded to Covidence systematic review software (Veritas Health Innovation, Melbourne, Australia) for screening. Two reviewers independently screened the titles and abstracts of all publications identified in the search. All publications meeting the inclusion criteria were retained. Where the titles and abstracts provided insufficient details, full-text publications were retrieved and screened again by both reviewers against the inclusion criteria. All disagreements were resolved by discussion between the reviewers without the need for moderation.

Two reviewers independently extracted data of interest using standardised data extraction forms developed for this review. The following information, where available, was extracted for each publication: authors; article title; name of guideline document; year of guideline document; name of organisation; country. Once identified through data extraction, the guideline documents were retrieved.

## Selection criteria

For phase 1 of the search, the following were included: guideline documents whose stated purpose was the primary prevention of food allergy (including the original guideline document and articles referring to such guideline documents); guideline documents and articles in English; and guideline documents intended specifically for health professional use. Guideline documents for stated purposes other than primary prevention of food allergy were included only if they provided detailed, specific recommendations regarding food allergy prevention within their scope.

For phase 2 of the search, English language guideline documents whose stated purpose was the primary prevention of food allergy for health professionals were included; and guideline documents for stated purposes other than primary prevention of food allergy were included only if they provided detailed, specific recommendations regarding food allergy prevention within their scope.

If one professional organisation or government published more than one guideline document, all versions meeting the selection criteria were included in the review.

## Guideline comparison

The recommendations contained within the guideline documents relating to maternal diet during pregnancy and lactation; breastfeeding substitutes; solid food timing; advice regarding introduction of common food allergens; specific advice regarding egg and peanut introduction; and spacing of introduction of new foods, were retrieved as these are key factors in relation to food allergy prevention. The guidelines were compared for their recommendations relating to these factors. Other interventions such as Vitamin D, omega-3 fatty acid supplementation, and modification of the maternal diet and the infant microbiome, were not included in this review.

## Quality appraisal of guideline documents

Quality assessment of all identified guideline documents was undertaken independently by 2 reviewers who reviewed and scored each guideline document using the AGREE II tool.<sup>31,32</sup> The AGREE II

tool assesses guidelines using the following domains: scope/purpose (objectives, question, population); stakeholder involvement (group membership, target population, target users); rigor of development (search methods, evidence criteria, evidence strengths and limitations, recommendations, benefits and harms considerations, recommendations and evidence link, external review, and updating procedures); presentation clarity (specific, unambiguous recommendations, management options, and identifiable key recommendations); applicability (application facilitators and barriers, implementation of advice/tools, resource implications, and monitor/audit criteria); and editorial independence (funding body, competing interests).

The reviewers referred to the AGREE II tool with the user's manual<sup>34</sup> when assessing the guideline documents and were masked to scores assigned by the other reviewer. Each domain has a different number of quality assessment questions, each requiring a score between 1 and 7 (7 being the highest score). The quality scores were synthesised and domain scores for each guideline document calculated according to the AGREE II manual protocol.<sup>34</sup> Domain scores are calculated by subtracting the minimum possible score for the domain from the obtained score for the domain; this is then divided by the maximum possible score for the domain minus the minimum possible score for the domain; this score is multiplied by 100 to achieve a percentage.<sup>34</sup>

## Interpretation of domain scores

The AGREE II tool does not specify cut-off scores equating to guideline quality.<sup>34</sup> For this review, consistent with other reviews,<sup>37,38</sup> the quality threshold for guideline acceptability was defined as guideline documents achieving at least 50% for Domain 3 (rigor of development) and at least 50% for at least 2 other domains.

## Statistical analysis

Data were analysed in SPSS 26.0 (SPSS Inc, Chicago, IL). Descriptive statistics for the AGREE II assessment were obtained, and comparison of the means of the guidelines compared to the advice

Organisation	Name of document	Author specified type of document	Region	Year
American Academy of Paediatrics	Effects of early nutritional interventions on the development of atopic disease in infants and children: The role of maternal dietary restriction, breastfeeding, timing of introduction of complementary foods and hydrolysed formulas <sup>39</sup>	Clinical report/ Guidance	United States	2008
American Academy of Paediatrics	Consensus communication on early peanut introduction and the prevention of peanut allergy in high-risk infants <sup>40</sup>	Consensus communication	United States	2015
American Academy of Paediatrics	The effects of early nutritional interventions on the development of atopic disease in infants and children: The role of maternal dietary restriction, breastfeeding, hydrolysed formulas and timing of introduction of allergenic complementary foods <sup>41</sup>	Clinical report/ Guidance	United States	2019
American College of Allergy, Asthma and Immunology (ACAAI)	Food allergy and introduction of solid foods to infants: a consensus document <sup>42</sup>	Consensus document	United States	2006
Asia Pacific Association of Paediatric Allergy, Respiriology & Immunology (APAPARI)	Early introduction of allergenic foods for the prevention of food allergy from an Asian perspective - An APAPARI consensus statement <sup>43</sup>	Consensus statement	Asia	2017
Australasian Society of Clinical Immunology and Allergy (ASCIA)	Australasian Society of Clinical Immunology and Allergy position statement: summary of allergy prevention in children <sup>44</sup>	Position statement	Australia & New Zealand	2005
Australasian Society of Clinical Immunology and Allergy (ASCIA)	Infant feeding advice <sup>45</sup>	Advice	Australia & New Zealand	2008
Australasian Society of Clinical Immunology and Allergy (ASCIA)	ASCIA Guidelines for infant feeding and allergy prevention <sup>46</sup>	Guideline	Australia & New Zealand	2016
	ASCIA Guidelines: Infant feeding and allergy prevention <sup>47</sup>			

(continued)

Organisation	Name of document	Author specified type of document	Region	Year
British Society for Allergy & Clinical Immunology (BSACI)	Preventing food allergy in higher risk infants: guidance for healthcare professionals <sup>48</sup>	Guidance	United Kingdom	2018
	Implementing primary prevention of food allergy in infants: New BSACI guidance published <sup>49</sup>			
Canadian Paediatric Society (CPS) and Canadian Society of Allergy and Clinical Immunology (CSACI)	Dietary exposures and allergy prevention in high-risk infants <sup>50</sup>	Joint statement	Canada	2013
Canadian Paediatric Society (CPS) and Canadian Society of Allergy and Clinical Immunology (CSACI)	Timing of introduction of allergenic solids for infants at high risk <sup>51</sup>	Practice point	Canada	2019
European Academy of Allergy and Clinical Immunology (EAACI)	Dietary prevention of allergic diseases in infants and small children <sup>52</sup>	Recommendations	Europe	2004
European Academy of Allergy and Clinical Immunology (EAACI)	EAACI Food Allergy and Anaphylaxis Guidelines. Primary prevention of food allergy <sup>53</sup>	Guideline	Europe	2014
European Society for Paediatric Allergology and Clinical Immunology (ESPACI) and European Society for Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN)	Dietary products used in infants for treatment and prevention of food allergy <sup>54</sup>	Joint statement	Europe	1999
European Society for Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN)	Complementary feeding: A commentary by the ESPGHAN Committee on Nutrition <sup>55</sup>	Position paper	Europe	2007
European Society for Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN)	Complementary feeding: A position paper by the European Society for Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN) Committee on Nutrition <sup>56</sup>	Position paper	Europe	2017
Finnish Allergy Programme	Allergy in children: practical recommendations of the Finish Allergy Programme 2008-2018 for prevention, diagnosis and treatment <sup>57</sup>	Recommendations	Finland	2012

(continued)



Organisation	Name of document	Author specified type of document	Region	Year
German Society for Allergology and Clinical Immunology (DGAKI) and the German Society for Paediatric and Adolescent Medicine (DGKJ)	Allergy Prevention <sup>58</sup>	Clinical practice guideline	Germany	2009
German Society for Allergology and Clinical Immunology (DGAKI) and the German Society for Paediatric and Adolescent Medicine (DGKJ)	S3-Guideline on allergy prevention: 2014 update <sup>59</sup>	Guideline	Germany	2014
Hong Kong Institute of Allergy (HKIA)	Guidelines for allergy prevention in Hong Kong <sup>60</sup>	Guideline	Hong Kong	2015
	Guidelines for Allergy Prevention in Hong Kong <sup>61</sup>			
Hong Kong Institute of Allergy (HKIA)	HKIA position paper on prevention of peanut allergy in high risk infants <sup>62</sup>	Position paper	Hong Kong	2016
Italian Society of Preventative and Social Paediatrics (ISPSP), the Italian Society of Paediatric Allergy and Immunology (ISPAI) and the Italian Society of Pediatrics (ISP)	Prevention of food and airway allergy: consensus of the Italian Society of Preventative and Social Paediatrics, the Italian Society of Paediatric Allergy and Immunology, and Italian Society of Pediatrics <sup>63</sup>	Consensus statement	Italy	2016
Japanese Society of Paediatric Allergy and Clinical Immunology (JSPACI)	Japanese guidelines for food allergy 2017 <sup>64</sup>	Guideline	Japan	2017
National Institute of Allergy and Infectious Diseases (NIAID)	Guidelines for the Diagnosis and Management of Food Allergy in the United States: Report of the NIAID-Sponsored Expert Panel <sup>65</sup>	Guideline	United States	2010
National Institute of Allergy and Infectious Diseases (NIAID)	NIAID Addendum guidelines for prevention of peanut allergy in the United States <sup>66</sup>	Guideline	United States	2017
Philippine Society of Allergy, Asthma and Immunology (PSAAI) and the Philippine Society for Paediatric Gastroenterology, Hepatology and Nutrition (PSPGHN)	Dietary prevention of allergic diseases in children: the Philippine guidelines <sup>67</sup>	Guideline	Philippines	2017

(continued)

Organisation	Name of document	Author specified type of document	Region	Year
Scientific Advisory Committee on Nutrition (SACN) and Committee on Toxicity of Chemicals in food Consumer products and the Environment (COT)	Assessing health benefits and risks of the introduction of peanut and hen's egg into the infant diet before six months of age in the UK <sup>68</sup>	Joint statement	United Kingdom	2018
Academy of Medicine, Singapore Ministry of Health (AMS-MOH)	Management of food allergy <sup>69</sup>	Clinical practice guideline	Singapore	2010
	Academy of medicine, Singapore-Ministry of Health clinical practice guidelines: management of food allergy <sup>70</sup>			

**Table 1. (Continued)** Summary of included guideline documents

documents was undertaken using an independent t-test.

## RESULTS

The electronic database search yielded 1121 publications and reference checking identified an additional 18 publications. After removal of duplicates (n = 477); title and abstract screening was undertaken on 660 publications; 239 publications underwent full-text screening. Data extraction was undertaken on 156 publications to identify guideline documents including: the name of the guideline, creator/owner, country, and date. Two additional guideline documents were identified through reference searching. This yielded 28 guideline documents (from 17 organisations) over a period of 21 years ([Supplemental Figure 1](#)). The AGREE II appraisal was undertaken on all 28 guideline documents identified.

### Identified guideline documents

A summary of included guideline documents is provided in [Table 1](#).<sup>39-70</sup> Of the 28 food allergy prevention guideline documents, 10 were specifically titled as "guidelines",<sup>46,51,55-57,60,62-64,66</sup> and 18 were titled as consensus statements, position statements, joint statements or recommendations and were grouped together as "advice documents".<sup>39-45,48,50-52,54-57,62,63,68</sup> Where an organisation had more than 1 version of their guideline document, all versions were

included. A timeline of the 28 food allergy prevention guidelines is provided in [Fig. 1](#).

### Guideline comparison

The recommendations within the 28 guidelines document in relation to maternal diet, exclusive breastfeeding, breastmilk substitutes, timing of solid food introduction, and any recommendations for specific food introduction are summarised in [Table 2](#).


#### Maternal diet

Of the 28 guideline documents, 17<sup>39,41,44-47,50,52,53,57-61,63-65,67,69,70</sup> included recommendations regarding maternal diet during pregnancy and lactation. Eleven guideline documents did not include maternal dietary recommendations as they were guidelines specific to infant feeding. Of the 17 with maternal diet recommendations, all stipulated "no dietary restrictions".<sup>39,41,44-47,50,52,53,57-61,63-65,67,69,70</sup> In addition to the "no dietary restrictions" recommendation, 4 guideline documents also recommended a healthy balanced maternal diet;<sup>46,47,53,58,59</sup> 3 guideline documents recommended the inclusion of fish;<sup>46,47,58,59</sup> and 1 document stipulated that fish oil supplements were not recommended.<sup>63</sup>

#### Exclusive breastfeeding

Twenty-five guideline documents made recommendations regarding exclusive breastfeeding.<sup>39,41-61,63-65,67-69</sup> The remaining 3 made





2019	<b>AAP 2019:</b> cow's milk formula; solids from 4-6 months; don't delay common allergens	<b>CPS/CSACI 2019:</b> HRI – solids around 6 months not before 4 months; don't delay common allergens			
2018	<b>BSACI 2018:</b> cow's milk formula; solids around 6 months not before 4 months; HRI – solids from 4 months; don't delay common allergens	<b>SACN/COT 2018:</b> solids around 6 months; introduce egg and peanut around 6 months			
2017	<b>APAPARI 2017:</b> HRI – allergy testing; solids from 6 months; don't delay common allergens	<b>ESPGHAN 2017:</b> solids not before 17 weeks; don't delay common allergens	<b>JSPACI 2017:</b> solids from 5-6 months; don't delay common allergens	<b>NIAID 2017:</b> solids from 4-6 months; don't delay common allergens	<b>PSAAI/PSPGHN 2017:</b> HRI – pHF or eHF for at least 6 months; solids from 6 months; cooked egg 4-6 months; wheat before 6 months; fish 6-9 months; peanut 4-11 months
2016	<b>ASCIA 2016:</b> cow's milk formula; solids from around 6 months not before 4 months; don't delay common allergens; peanut and egg before 12 months	<b>HKIA 2016:</b> HRI – skin prick test before peanut introduction; don't delay common allergens	<b>ISPSP/ISP/ISP 2016:</b> cow's milk formula; solids after 4th month and if possible after 6th month; don't delay common allergens		
2015	<b>AAP 2015:</b> HRI - introduce peanut between 4-11 months of age	<b>HKIA 2015:</b> HRI – consider hydrolysed formula; solids from 4-6 months; neither withhold nor encourage common allergens			
2014 - 2012	<b>FAP 2012:</b> standard cow's milk formula; solids from 4-6 months; feed wheat and oats by 6 months; don't delay common allergens	<b>CPS/CSACI 2013:</b> hydrolysed formula; solids from 6 months; don't delay common allergens	<b>DGAKI/DGKJ 2014:</b> hydrolysed formula until 4 months; solids from over 4 months; don't delay common allergens; fish before 12 months	<b>EAACI 2014:</b> HRI - hydrolysed formula until 6 months; solids from 4-6 months; neither withhold nor encourage common allergens	
2010 - 2009	<b>DGAKI/DGKJ 2009:</b> pHF or eHF until 4 months; solids from end of 4th month; don't delay common allergens; fish before 12 months	<b>NIAID 2010:</b> HRI – consider hydrolysed formula; solids from 4-6 months; don't delay common allergens	<b>AMS-MOH 2010:</b> HRI-hydrolysed formula; avoid cow's milk first 5 days of life; solids from 4-6 months		
2008	<b>AAP 2008:</b> solids from 4-6 months; don't delay common allergens	<b>ASCIA 2008:</b> HRI- pHF; solids from around 4-6 months; don't delay common allergens			
2007 - 2006	<b>ACAAI 2006:</b> cow's milk formula; solids from 6 months; don't delay common allergens	<b>ESPGHAN 2007:</b> solids not before 17 weeks; don't delay common allergens			
2005 - 1999	<b>ESPACI/ESPGHAN 1999:</b> HRI – hydrolysed formula; solids from 5 months	<b>EAACI 2004:</b> eHF until 4-6 months; pHF possible benefit; solids from 6 months but at least 4 months; don't delay common allergens	<b>ASCIA 2005:</b> HRI- pHF or eHF; solids from 4-6 months; avoidance of peanuts, nuts and shellfish may be recommended		

Fig. 1 Guideline document timeline

Guideline document	Maternal diet (pregnancy and breastfeeding)	Breastfeeding	Breastmilk substitutes	Solid food timing	Peanut and egg
AAP 2019	No restrictions - not applicable for 2015 document - no change from 2008 document	Exclusive BF for at least 4 months - not applicable for 2015 document - no change from 2008 document	Hydrolysed formula not recommended - not applicable for 2015 document - 2008 document 'soy formula not recommended'	4-6 months of age: Do not delay common allergens - not applicable for 2015 document - no change from 2008 document	HRI - earliest age of peanut introduction is 4-6 months and consider evaluation of HRI by allergist before peanut introduction; Infants with mild-moderate eczema - earliest age of peanut introduction is around 6 months; Infants with no eczema or any food allergy - earliest age of peanut introduction is age appropriate and based on family meals and culture 2015 document 'For HRI - introduce peanut between 4 and 11 months of age; Consider evaluation of HRI by allergist before peanut introduction'
ACAAI 2006	Not applicable	Exclusive BF for 6 months	Standard cow's milk formula	6 months of age; Delay introduction of common allergens; Cooked, homogenised foods should be preferred over fresh counterparts if reduced allergenicity (e.g. beef and kiwifruit); egg, peanut, tree nuts fish and seafood introduction requires caution	Peanut and egg introduction requires caution

APAPARI 2017	Not applicable	Continue BF alongside solid food introduction up to 2 years if possible, according to cultural practice	Not specified	HRI - recommend allergy testing to egg and peanut; At risk infants - no delay in introduction of allergenic foods; Healthy infants - 6 months of age	Allergy testing for HRI prior to introduction of egg and peanut
ASCIA 2016	No restrictions; Healthy balanced diet; Up to 3 serves oily fish/week during pregnancy - 2005 and 2008 documents only stipulated 'no restrictions'	At least 6 months (where possible) and for as long as mother and infant wish to continue - No change from 2008 document - 2005 document 'exclusive BF for 4-6 months'	All infants - standard cow's milk formula - 2008 document 'pHF for HRI; soy milk is not recommended' - 2005 document 'pHF or eHF for HRI; soy milk is not recommended'	When infant is ready, around 6 months, not before 4 months; Introduce all common allergens; egg should be cooked; Continue to regularly include in infant diet once introduced - 2008 document 'From around 4-6 months; Introduce all common allergens' - 2005 document 'From 4 to 6 months; Introduce; peanut, nuts and shellfish for the first 2-4 years of life may be recommended. However, peanut, nut and shellfish avoidance may be recommended'	Introduce cooked egg and peanut before 12 months of age; Procedure for high risk infants - 2008 document 'Do not delay' - 2005 document 'Peanut avoidance for first 2-4 years may be recommended'
BSACI 2018	Not applicable	Exclusive BF for around 6 months; Continue to breastfeed while introducing solids if possible	Standard cow's milk formula	From around 6 months, but not before 4 months, when infant is ready; HRI - parents may wish to introduce solids from 4 months, cooked egg then peanut should be	HRI - may benefit from introduction of peanut and egg from 4 months alongside other foods

(continued)

Guideline document	Maternal diet (pregnancy and breastfeeding)	Breastfeeding	Breastmilk substitutes	Solid food timing	Peanut and egg
				given, then other allergenic foods; egg should be cooked; Introduce before 12 months of age; Continue to regularly include in infant diet once introduced	
CPS and CSACI 2019	Not applicable - 2013 document 'no restrictions'	Breastfeed for up to 2 years and beyond - 2013 document 'Exclusive BF for first 6 months'	Not applicable - 2013 document 'Hydrolysed formula; Soy formula not recommended'	HRI - around 6 months of age but not before 4 months; All other infants - around 6 months; Introduce all common allergens; Continue to regularly include in infant diet once introduced - 2013 document 'From 6 months of age; introduce all common allergens'	Do not delay - No change from 2013 document
DGAKI and DGKJ 2014	Balanced and varied diet; No restrictions; Fish should form part of the maternal diet - No change from 2009 document	Predominantly breastfed up to 4 months of age - 2009 document 'Exclusive BF up to 4 months of age'	Hydrolysed infant formula until 4 months of age; Soy based formula is not recommended for allergy prevention - 2009 specified pHF or eHF, otherwise no change	From over the age of 4 months; Common allergens should not be delayed; Fish should be introduced by 12 months of age - No change from 2009 document	Not specified - No change from 2009 document
EAACI 2014	No restrictions; No supplements while breastfeeding - 2004 only stipulated 'no restrictions'	Exclusive BF for 4-6 months - 2004 document 'Exclusive BF for at least 4 months'	HRI - hydrolysed formula until 4 months of age then standard cow's milk formula; All other infants - standard cow's milk formula; Soy and hydrolysed	From 4 to 6 months of age, when infant is ready; Neither withhold nor encourage exposure of common food allergens	Not specified - No change from 2004 document

			<p>formulas not recommended</p> <ul style="list-style-type: none"> <li>- 2004 document 'eHF until 4-6 months; pHF may have an effect; Soy formulas not recommended'</li> </ul>	<ul style="list-style-type: none"> <li>- 2004 document 'Preferably 6 months but at least 4 months of age; No evidence for restrictive diets beyond 6 months for common food allergens'</li> </ul>	
ESPGHAN 2017	<p>Not applicable</p> <ul style="list-style-type: none"> <li>- No change from 1999 to 2017 documents</li> </ul>	<p>Continue BF while introducing solid foods</p> <ul style="list-style-type: none"> <li>- 2007 document 'Exclusive BF for around 6 months</li> <li>- 1999 document 'Exclusive BF for 4-6 months'</li> </ul>	<p>Not specified</p> <ul style="list-style-type: none"> <li>- No change from 2007 document</li> <li>- 1999 document 'HRI - reduced allergenicity formula; All other infants - standard cow's milk formula'</li> </ul>	<p>Not before 17 weeks; Do not delay common food allergens</p> <ul style="list-style-type: none"> <li>- No change from 2007 document</li> <li>- 1999 document 'From 5 months; no information specified regarding common food allergens'</li> </ul>	<p>High risk infants - introduce peanut between 4 and 11 months</p> <ul style="list-style-type: none"> <li>- 2007 document 'Do not delay'</li> <li>- Not specified in 1999 document</li> </ul>
Finish Allergy Program 2012	<p>No restrictions</p>	<p>Exclusive BF for 4-6 months</p>	<p>Standard cow's milk formula</p>	<p>From 4 to 6 months while continuing BF; Introduce wheat and oats by 6 months of age</p>	<p>Do not delay</p>
HKIA 2016	<p>Not applicable</p> <ul style="list-style-type: none"> <li>- 2015 document 'Healthy diet during pregnancy; No restrictions'</li> </ul>	<p>Not applicable</p> <ul style="list-style-type: none"> <li>- 2015 document 'At least 4-6 months'</li> </ul>	<p>Not applicable</p> <ul style="list-style-type: none"> <li>- 2015 document 'HRI - consider hydrolysed formula if exclusive breastfeeding is not possible'</li> </ul>	<p>HRI - SPT before introduction encouraged; Low risk infants - introduce peanut upon introduction of foods; Do not delay common food allergens</p> <ul style="list-style-type: none"> <li>- 2015 document 'From 4 to 6 months of age when developmentally ready; Neither withhold nor encourage exposure to common food allergens'</li> </ul>	<p>HRI - SPT; negative and mild positive SPT - 6g peanut protein/wk 3 times/wk until 5 years of age; Positive SPT - oral peanut challenge, include peanut if negative challenge an avoid peanut if positive challenge</p> <ul style="list-style-type: none"> <li>- Not specified in 2015 document</li> </ul>

(continued)

Guideline document	Maternal diet (pregnancy and breastfeeding)	Breastfeeding	Breastmilk substitutes	Solid food timing	Peanut and egg
ISPSP and ISPAI and ISP 2016	Fish oil supplementation not recommended	Exclusive BF for at least 4 months (possibly 6 months)	Standard cow's milk formula	After the 4th month and if possible after the 6th month; Introduce common food allergens in the same way as for children without allergic risk	Not specified
JSPACI 2017	No restrictions	Insufficient evidence to indicate superiority of BF in the prevention of allergic disease	Insufficient evidence to support the use of hydrolysed formula	From 5 to 6 months of age when developmentally ready; Do not delay common food allergens	Introduce peanuts sooner rather than later after weaning
NIAID 2017	Not applicable - 2010 document 'no restrictions'	Not applicable - 2010 document 'Exclusive BF until 4-6 months of age, unless BF is contraindicated for medical reasons'	Not applicable - 2010 document 'HRI - hydrolysed formulas may be considered; Soy formula not recommended'	4-6 months of age; Introduce common food allergens from 4 to 6 months of age - No change from 2010 document	HRI - earliest age of peanut introduction is 4-6 months and consider evaluation of HRI by allergist before peanut introduction; Infants with mild-moderate eczema - earliest age of peanut introduction is around 6 months; Infants with no eczema or any food allergy - earliest age of peanut introduction is age appropriate and based on family meals and culture - Not specified in 2010 document
PSAAI and PSPGHN 2017	No increased intake of certain foods recommended; No restrictions	Exclusive BF for at least 3-6 months	HRI - pHF or eHF recommended for at least 6 months; Soy milk not recommended	From 6 months of age; Cooked egg at 4-6 months; wheat before 6 months; fish at 6-9 months; peanut at 4-11 months	Cooked egg at 4-6 months; peanut at 4-11 months



SACN and COT 2018	Not applicable	Exclusive BF for around 6 months	Not specified	Around 6 months of age; No information regarding common food allergens	Introduce peanut and egg around 6 months of age; If history of eczema or suspected food allergy, medical advice before peanut introduction may be sought; once introduced, peanut and egg should continue to be consumed as part of the usual infant diet
AMS-MOH 2010	No restrictions	Exclusive BF for at least 4-6 months	HRI - hydrolysed formula recommended; Avoid cow's milk formula in the first 5 days of life	4-6 months of age for all infants; No information regarding common food allergens	Not specified

**Table 2. (Continued)** Summary of comparison of recommendations. Abbreviations: BF = breastfeeding; HRI = High risk infants; SPT = Skin prick test; AAP = American Academy of Pediatrics; ACAAI = American College of Allergy, Asthma and Immunology; APAPARI = Asia Pacific Association of Paediatric Allergy, Respiriology & Immunology; ASCIA = Australasian Society of Clinical Immunology and Allergy; BSACI = British Society for Allergy & Clinical Immunology; CPS = Canadian Paediatric Society; CSACI = Canadian Society of Allergy and Clinical Immunology; DGAKI = German Society for Allergology and Clinical Immunology; DGKJ = German Society for Paediatric and Adolescent Medicine; EAACI = European Academy for Allergy and Clinical Immunology; ESPACI = European Society for Paediatric Allergology and Clinical Immunology; ESPGHAN = European Society for Paediatric Gastroenterology, Hepatology and Nutrition; HKIA = Hong Kong Institute of Allergy; ISPSP = Italian Society of Preventative and Social Paediatrics; ISPAI = Italian Society of Paediatric Allergy and Immunology; ISP = Italian Society of Pediatrics; JSPACI = Japanese Society of Paediatric Allergy and Clinical Immunology; NIAID = National Institute of Allergy and Infectious Diseases; PSAAI = Philippine Society of Allergy, Asthma and Immunology; PSPGHN = Philippine Society for Paediatric Gastroenterology, Hepatology and Nutrition; SACN = Scientific Advisory Committee on Nutrition; COT = Committee on Toxicity of Chemicals in food, consumer products and the environment; AMS-MOH = Academy of Medicine, Singapore Ministry of Health

	Type of document for comparison	Domain 1 (%)	Domain 2 (%)	Domain 3 (%)	Domain 4 (%)	Domain 5 (%)	Domain 6 (%)	Meets quality threshold
ASCIA 2016	Guideline	94.4	52.7	60.4	83.3	29.2	79.2	Yes
DGAKI/DGKJ 2009	Guideline	88.9	47.2	66.7	72.2	6.2	70.8	Yes
DGAKI/DGKJ 2014	Guideline	97.2	4.4	78.1	75.0	47.9	83.8	Yes
EAACI 2014	Guideline	94.4	91.7	89.6	91.7	56.2	54.2	Yes
HKIA 2015	Guideline	50.0	2.7	19.8	61.6	4.2	45.8	No
JSPACI 2017	Guideline	58.3	41.7	22.9	61.1	12.5	62.5	No
NIAID 2010	Guideline	100.0	91.7	82.3	86.1	27.1	62.5	Yes
NIAID 2017	Guideline	97.2	91.7	71.9	88.9	20.8	50.0	Yes
PSAAI/PSPGHN 2017	Guideline	94.4	91.7	74.0	86.1	8.3	70.8	Yes
AMS-MOH 2010	Guideline	88.9	88.9	52.1	91.7	29.2	12.5	Yes
AAP 2008	Advice document	88.9	13.9	19.8	47.2	2.1	4.2	No
AAP 2015	Advice document	75.0	22.2	21.9	41.7	6.2	4.2	No
AAP 2019	Advice document	83.3	25.0	28.1	52.8	2.1	62.5	No
ACAAI 2006	Advice document	88.9	36.1	33.3	58.3	12.5	91.7	No
APAPARI 2017	Advice document	75.0	8.3	20.8	36.1	16.7	4.2	No
ASCIA 2008	Advice document	88.9	36.1	20.1	50.0	14.6	62.5	No
ASCIA 2005	Advice document	72.2	30.6	24.0	52.8	4.2	12.5	No
BSACI guidance	Advice document	66.7	58.3	13.5	66.7	16.7	45.8	No
CPS/CSACI 2013	Advice document	91.7	50.0	25.0	66.7	12.5	4.2	No
CPS/CSACI 2019	Advice document	97.2	16.7	12.5	50.0	4.2	4.2	No
EAACI 2004	Advice document	61.1	25.0	46.9	44.4	8.3	8.3	No
ESPACI/ESPGHAN 1999	Advice document	91.7	52.8	12.5	61.1	2.1	4.2	No
ESPGHAN 2007	Advice document	94.4	36.1	17.7	58.3	8.3	12.5	No

ESPGHAN 2017	Advice document	100.0	50.0	28.1	50.0	16.7	4.2	No
Finnish AP 2012	Advice document	69.4	36.1	12.5	41.7	10.4	8.3	No
HKIA 2016	Advice document	44.4	19.4	7.3	33.3	8.3	0.0	No
ISPSP/ISPAI/ISP 2016	Advice document	83.3	33.3	44.8	61.1	6.2	33.3	No
SACN/COT 2018	Advice document	94.4	38.9	47.9	63.9	6.2	4.2	No

Table 3. Domain Scores - Guidelines and advice documents


no exclusive feeding recommendations, as these guideline documents were specifically related to peanut introduction.<sup>40,62,66</sup> Of those that did make breastfeeding recommendations, 6 recommended "exclusive/predominantly exclusive breastfeeding for at least 4 months".<sup>39,41,52,58,59,63</sup> "Exclusive breastfeeding for 4-6 months" was recommended by 7 guideline documents;<sup>44,53,54,57,60,61,65,69,70</sup> 7 recommended "exclusive breastfeeding for 6 months or around 6 months or at least 6 months".<sup>42,45-50,55,68</sup> Two guideline documents provided no recommendation regarding exclusive breastfeeding but stipulated that breastfeeding "should continue up to 2 years or longer".<sup>43,52</sup>

### Breastmilk substitutes

Twenty of the 28 guideline documents provided recommendations for breastmilk substitutes for primary prevention of allergy;<sup>39,41,42,44-50,52-54,57-61,63-65,67,69,70</sup> 4 of the remaining documents focussed on solid food introduction, and, therefore, were not intended to include breastfeeding information,<sup>40,51,62,66</sup> and 4 made no recommendation.<sup>43,55,56,68</sup> Of these, soy formula was not recommended by 10 documents.<sup>39,44,45,50,52,53,58,59,65,67</sup> Specific recommendations for high risk infants were made in 8 documents,<sup>44,45,53,54,60,61,65,67,69,70</sup> with partially hydrolysed formula (pHF) recommended in all 8 documents<sup>44,45,53,54,60,61,65,67,69,70</sup> and extensively hydrolysed formula (eHF) recommended in 7 of the 8 documents.<sup>44,53,54,60,61,65,67,69,70</sup> Where documented, the majority of documents that recommended the use of hydrolysed formulas were based on evidence suggesting potential reduction in allergic disease generally;<sup>44,45,50</sup> atopic eczema;<sup>52-54,60,61,67,69,70</sup> asthma and allergic rhinitis;<sup>60,61,67</sup> food intolerance;<sup>44,50,52,53</sup> or food allergy.<sup>52,53,65,69,70</sup>

### Timing of solid food introduction into the infant diet

Twenty-seven documents made recommendations regarding the timing of solid food introduction.<sup>39,41-70</sup> The remaining guideline document was specific to peanut introduction, hence made no general solid food timing recommendations.<sup>40</sup> There was variability in the wording for timing of the introduction of solid foods; however, all were



**Fig. 2** Domain scores comparison for guidelines versus advice documents

within the range of 4 to 6 months. The most common wording (used in 10 documents) stipulated "from 4 to 6 months";<sup>39,41,44,45,53,57,60,61,65,66,69,70</sup> 4 documents stipulated "from 6 months";<sup>42,50,67,68</sup> 3 documents stipulated "around 6 months but not before 4 months".<sup>46-49,51</sup> When comparing timing of solid food introduction in guideline documents developed between 2015 and 2019 with such recommendations,<sup>40,41,43,46-49,51,56,60,61,63,64,66-68</sup> there was still similar variability in the wording of recommendations regarding timing of solid food introduction.

Specific recommendations for high risk infants were included in 4 documents.<sup>43,48,49,51,62</sup> Of these, one document recommended introduction of solid foods "at 4 months" (compared to "6 months" for infants not at risk),<sup>48,49</sup> and 1 document recommended introduction of solids "around 6 months but not before 4 months";<sup>51</sup> while 2 documents recommended allergy testing before introduction of solid foods.<sup>43,62</sup>

### Common food allergens

Twenty-four documents made recommendations regarding timing of common food allergen introduction.<sup>39,41-53,55-67</sup> Sixteen documents recommended that the common food allergen introduction should not be delayed;<sup>39,41-43,45-47,50-52,55,56,58,59,62,64-66</sup> 1 document recommended delaying common food allergen introduction;<sup>42</sup> and 1 document recommended delaying peanut, nuts and shellfish.<sup>44</sup> Three documents recommended neither withholding nor encouraging common food allergen

introduction.<sup>53,60,61,63</sup> Five documents made recommendations about specific foods;<sup>57-59,62,67</sup> 1 recommended wheat and oat introduction by 6 months of age;<sup>57</sup> 2 recommending fish introduction by 12 months of age;<sup>58,59</sup> and 1 recommended egg, wheat, fish, and peanut introduction at specified ages.<sup>67</sup> Of those guideline documents produced between 2016 to 2019,<sup>41,43,46-49,51,56,62-64,66-68</sup> all 12 stipulated that common food allergen introduction should not be delayed.

### Recommendations specific to peanut and egg introduction

Nineteen documents included recommendations regarding peanut and egg.<sup>40-51,55-57,63,64,66-68</sup> Of these, 6 indicated peanut and egg should not be delayed,<sup>39,45,50,51,55,57</sup> and 1 stipulated introduction by 12 months of age.<sup>46,47</sup> One document recommended caution regarding introduction,<sup>42</sup> and another recommended avoidance.<sup>44</sup> Special advice for high risk infants was provided in 8 documents,<sup>40,41,43,48,49,56,62,66,68</sup> with 6 of these documents recommending testing/evaluation by an allergist.<sup>40,41,43,62,66,68</sup> Six documents were updated after publication of the LEAP study and include peanut allergy specific information.<sup>41,46,47,51,56,62,66</sup>

### Spacing of introduction of new foods

Only 5 documents made recommendations regarding the spacing (time between introduction of each new food) of introducing new foods, with

all 5 documents recommended introducing 1 new food at a time.<sup>42,44,46,47,51,55</sup>

### Quality appraisal using AGREE II

An overview of the domain scores for each of the 28 guideline documents is presented in Table 3. Eight<sup>46,47,53,58,59,65-67,69,70</sup> of the 10 guidelines<sup>46,47,53,58-61,64-67,69,70</sup> examined achieved the quality threshold, and of these, 1 guideline<sup>53</sup> scored equal to or greater than 50% across all domains. None of the advice documents<sup>39-45,48-52,54-57,62,63,68</sup> met the quality threshold. Domain 3 (rigor of development) was considered integral to developing a quality guideline: only the 8 guidelines<sup>46,47,53,58,59,65-67,69,70</sup> that met the quality threshold achieved at least 50% for this domain.

The highest scores were achieved for Domain 1 (scope and purpose) and Domain 4 (clarity of presentation) with mean scores of 83.2% and 61.9%, respectively. Domains 5 (applicability) and 6 (editorial independence) achieved the lowest mean scores of 14.3% and 34.3%, respectively.

Domains 2 (stakeholder engagement) and 6 (editorial independence) had the greatest range in scores, both having a range of 91.7% despite having different minimum scores (2.7% and 0.0% respectively).

A comparison of mean scores for the guidelines and advice documents identified a significant difference at the 5% level for all domains except Domain 1 (scope and purpose) for which both guidelines and advice documents achieved the highest mean scores of 86.4 and 81.5, respectively (Supplemental Table 1). Fig. 2 shows a graphical comparison of the domain scores for guidelines versus advice documents.

## DISCUSSION

To our knowledge, this systematic review provides the only review comparing food allergy prevention guidelines (created over a 21-year timeframe, 1999-2019) and the first comprehensive appraisal of the quality of these guidelines. In searching for primary prevention guidelines for food allergy, we identified 28 food allergy prevention documents that meet the WHO definition

of a guideline,<sup>35</sup> even though they varied in their titles (eg, guidelines, consensus statement, joint statement) and their processes of development. Since undertaking this study, European Academy of Allergy and Clinical Immunology (EAACI) updated their 2014 guidelines which were in press at the time of finalising this article for publication.<sup>71</sup> The updated EAACI guidelines were not included in the AGREE II appraisal nor the comparison of guideline recommendations; however, some references have been made to EAACI guidelines recommendations in the discussion.

### Comparison and consistency of recommendations

Children may be sensitised *in utero* or while breastfeeding, and as such, maternal diet during pregnancy and breastfeeding has become an increasing focus for food allergy prevention.<sup>72,73</sup> As such, key factors which should be included in guidelines, in relation to food allergy prevention, include maternal diet during pregnancy and breastfeeding; timeframe for exclusive breastfeeding; breastmilk substitutes; timing of introduction of solid foods; and recommendations regarding common food allergen introduction.<sup>21,74</sup> Other interventions such as vitamin D and omega 3- fatty acid supplementation and modification of the maternal and infant microbiome, remain more controversial<sup>20</sup> and were considered beyond the scope of this review.

Some guideline documents were clearly designed to contain comprehensive recommendations relating to all aspects of infant feeding and allergy prevention;<sup>39,41,44-50,52,53,57-61,63-65,67,69,70</sup> whereas others were clearly deliberately targeted at single recommendations, such as those specifically developed or updated in response to the LEAP study results<sup>24</sup> to provide advice specifically related to the prevention of peanut allergy.<sup>40,51,62,66,68</sup> As such, comparisons regarding the comprehensiveness of all guidelines are not necessarily overly meaningful.

Overall, all 28 guideline documents were consistent in recommending no maternal dietary restrictions during pregnancy and breastfeeding for allergy prevention. However, 4 documents provided additional recommendations regarding

maternal diet, relating to a balanced diet<sup>46,47,58-61</sup> and include fish.<sup>58,59</sup> In the case of the most recent Australasian Society of Clinical Immunology and Allergy (ASCI) guideline<sup>46,47</sup> the recommendation regarding a balanced maternal diet was included to be consistent with national dietary recommendations,<sup>75</sup> and factors such as this may be the reason for inclusion in the other guideline documents.

Less consistency was observed in the recommendations relating to duration of exclusive breastfeeding. The challenge with recommendations regarding exclusive breastfeeding is balancing the WHO guidelines,<sup>76</sup> for exclusive breastfeeding until 6 months (primarily to ensure adequate nutrition in all infants in developing as well as developed regions) against primary prevention of allergic disease. This may also be the case for infant feeding guidelines for the general population within different countries. In Australia, for example, general population infant feeding guidelines<sup>77</sup> aim to be consistent with the WHO guidelines, stipulating "at around 6 months" with no mention of 4 months; whereas the evidence for food allergy prevention supports introduction of common food allergens from around 6 months, but not before 4 months, indicating that if the infant is ready at 4 months, solids can be introduced.<sup>24-28</sup> Therefore, food allergy prevention guidelines are likely to provide different advice to general infant feeding recommendations which may cause some confusion for healthcare providers and parents. Further to this, despite recent studies being unable to make conclusions regarding the role of breastfeeding in the prevention of food allergy development,<sup>78</sup> guidelines continue to make recommendations regarding exclusive breastfeeding. However, interestingly, the updated EAACI guidelines make no recommendations regarding timing of exclusive breastfeeding, nor the timing of introduction of solids.<sup>71</sup>

While breastfeeding is promoted for its many benefits,<sup>79</sup> recommendations regarding infant formulas are important for mothers who cannot breastfeed or who choose not to breastfeed. Overall, there was consistency in not recommending soy formula for allergy

prevention<sup>39,44,45,50,52,53,58,65,67</sup> and some consistency regarding the use of hydrolysed infant formulas in high risk infants.<sup>44,45,50,52-54,58-61,65,67,69,70</sup> However, in the last 5 years (2015-2019), in response to a recent systematic review and meta-analysis,<sup>80</sup> a trend to recommend standard cow's milk formula for all infants including those at high risk of allergy was observed in the newer guidelines<sup>41,46-49,63</sup> compared to older guidelines which tended to recommend hydrolysed infant formula.<sup>60,61,67</sup>

While the introduction of solid foods is an important milestone,<sup>21</sup> it is also important for the child's immediate nutritional status and long-term health, including immune programming.<sup>81</sup> In our review, recommendations regarding the timing of solid food introduction varied in wording usually related to their recommendations regarding duration of exclusive breastfeeding. The most common recommendation for solid food introduction was "4-6 months".<sup>39,41,44,45,53,57,58,65,66,69,70</sup> When considering only the 14 guideline documents published between 2015 and 2019, greater variation in recommendations regarding solid food introduction was observed than in the previous 5-year period. In the Australian context, this may be explained by the need to align with current infant feeding advice for the general population, as timing of solid food introduction is dependent on the recommendations relating to exclusive breastfeeding.<sup>21</sup>

There was more consistency with regard to recommendations regarding common allergenic food introduction, with recommendations that it should not be delayed. In the last 5 years, (2015-2019), 10 out of 14 documents recommended that common food allergen introduction should not be delayed, consistent with recent studies to support "early" introduction of common food allergens.<sup>41,43,46-49,51,56,62-64,66</sup>

With the publishing of the LEAP study,<sup>24</sup> we would expect to see guideline documents including recommendations specific to peanut and possibly other specific foods. From 2016 to 2019, recommendations regarding specific food allergens, most commonly egg and peanut, were included in four guideline documents.<sup>46,47,66-68</sup>



### Overall quality of the guideline documents

Of the 28 food allergy prevention guideline documents identified, only 8 met the AGREE II quality threshold and of these all eight were titled "guidelines" by their authors. Further, only 1 guideline achieved a domain score of at least 50% across all domains. Therefore, most documents either did not have robust processes of development in place, or they did not fully report their processes of development. Rigour of development (Domain 3) was considered a critical component of guideline quality for this AGREE II assessment, as it relates to the evidence base used to underpin the recommendations as well as the process of formulating the recommendations. It is important to communicate development processes so that health professionals can easily identify whether guidelines are evidence-based. Ensuring that high quality evidence underpins recommendations for food allergy prevention is critical, particularly as the evidence has changed over time and the implications of poor advice from a low quality guideline can have a lifelong impact on the individual. For example, the evidence regarding food allergen introduction has changed from delaying common food allergen introduction to introducing common food allergens within the first year of life.<sup>24-28</sup> If guidelines with recommendations about common food allergen introduction are not up to date with current evidence, the "window of opportunity" to introduce the common food allergens and hence potentially prevent food allergy, could be missed. The updated EAACI guidelines document has used the AGREE II framework as the basis for the guideline development.<sup>71</sup>

As food allergy is common, parents concerned about the risk of their baby developing food allergies seek guidance from health professionals, particularly in relation to introducing common food allergens into their baby's diet. Clinical immunology/allergy specialists, general practitioners, paediatricians, child health nurses, and dietitians all rely on food allergy prevention guidelines to provide evidence-based advice to parents. Health professionals need to consider the robustness of the evidence underpinning the recommendations contained within each guideline document, as clinical practice should be governed by best available evidence.<sup>29,30</sup> In this review, we

found that some guideline documents lacked adequate reporting of the relevant stakeholders involved in the development process (domain 2), particularly consumer representation. We also found a lack of detailed information regarding funding sources and clarification regarding author conflicts of interest (Domain 6). These two domains (2 and 6) had the greatest variation in scores compared to other domain score ranges. This indicates that documentation of the stakeholders involved in the development process (domain 2) and the editorial independence (domain 6), ranged from poorly described or lacking to well documented within each domain. The main reason for the lower scores pertaining to stakeholder involvement, was how clearly consumer engagement was documented, if at all. Given that improving patient outcomes is the focus of clinical practice guidelines,<sup>82</sup> the consumer perspective is important. Timely and appropriate consumer engagement can provide insight into the relevance, practicality, and achievability of proposed recommendations, and can also contribute to development of appropriate implementation strategies after guidelines have been developed.<sup>82</sup>

Most documents reviewed clearly detailed the scope and purpose of the document (Domain 1) and recommendations were clearly presented (Domain 4), thereby meeting the 50% quality threshold. This allows health professionals to easily identify the aim of the document, the target audience, and the recommendations with regards to food allergy prevention. Having clearly documented food allergy prevention recommendations assists busy health professionals to easily identify the advice they should be providing to parents.

There are often barriers to implementing recommendations as evidenced by low scores in Domain 5 (applicability). Guidelines should consider potential outcomes and implementation factors such as how to successfully carry out the intervention, cost effectiveness, and the workforce required to implement the guideline recommendations.<sup>82</sup> This is important for food allergy prevention, because if recommendations are not achievable, the potential to reduce food allergy is lost. For example, if screening for peanut allergy by skin prick testing is recommended

prior to introducing peanut, access to skin prick testing must occur before the recommended timeframe for introduction to peanut (ie, before 1 year of age or earlier in some settings). These issues will be specific to the region for which the guidelines are being developed. In the United States where allergy specialists are accessible, screening prior to peanut introduction is recommended.<sup>66</sup> However, in Australia, allergy testing prior to peanut introduction is not feasible, as access to allergy specialists to conduct skin prick testing to screen for peanut sensitisation is unlikely before the child is 12 months of age, and the opportunity to introduce peanut to prevent peanut allergy would be missed. As a result, the ASCIA guidelines do not recommend screening for peanut sensitisation prior to peanut introduction.<sup>46,47</sup> Tools for both health professionals and parents to support implementation of food allergen prevention guidelines are also important to help allay fears and encourage recommendation adoption. As the updated EAACI guideline has used the AGREE II tool in developing the guideline, the barriers and facilitators to implementation as well as audit criteria and resource implications have been considered and clearly communicated.<sup>71</sup>

A question to consider is whether food allergy prevention guideline documents which score poorly using the AGREE II tool translate to having poor quality recommendations. The documents titled guidelines by their authors were more likely to meet the quality threshold than the advice documents, of which none met the quality threshold. Does this mean that the advice document recommendations are substandard? While the AGREE II tool appraises the methodology of guideline development and reporting, it does not appraise the clinical appropriateness of the recommendations themselves, or alignment with evidence.<sup>32</sup> Domain 3 (rigor of development) evaluates the process of identifying and grading the evidence that underpins the recommendations, to the extent it relies on this information being communicated within the document or in supporting material that is publicly available. Clearly reporting the process of guideline development may increase health professional confidence to adhere to and implement the guidelines, an important factor

with food allergy prevention. If health professionals fail to implement food allergy prevention guidelines, the "window of opportunity" to introduce common food allergens in the first year of life could be missed, resulting in an increased risk of food allergy. As some food allergies will be lifelong, this poses a significant impact on quality of life<sup>15,16</sup> and, in some cases, poses a threat to life.

## CONCLUSION

This review identified 28 food allergy prevention guideline documents published over 21 years. While we have compared the recommendations within the identified guideline documents for similarities, we must acknowledge that some of the variation in the recommendations may relate to country specific issues such as the general-population-based infant feeding recommendations. Overall, the greatest variation in recommendations was related to duration of exclusive breastfeeding and timing of solid food introduction. This may create confusion for health professionals and result in inconsistent advice being provided to parents, and less translation of the evidence into actual food allergy reduction in the population at large.

Documents specifically termed guidelines by the authors in their title scored higher using the AGREE II appraisal, principally because they included more information about their development process. Assessment using the AGREE II tool identified areas of improvement for future guideline development. The 2 key areas for improvement in food allergy prevention guidelines identified by this study include: documentation of stakeholder involvement, particularly consumer engagement, and clear documentation of editorial independence. Based on this study, we would recommend use of validated guideline development tools to direct development and review of food allergy prevention guidelines and ensure robust development and reporting processes.

### Funding

Not applicable.

### Consent for publication

All authors consent to this work being published.

**Author contributions**

All authors have contributed substantially to the writing of this publication.

**Availability of data and materials**

Data is available upon request.

**Potential competing interests**

Ms. Vale reports I am employed by the Australasian Society of Clinical Immunology and Allergy.

Ms Lobb has nothing to disclose.

Dr. Netting has nothing to disclose.

Dr. Murray has nothing to disclose.

Dr. Clifford has nothing to disclose.

Prof. Campbell reports other from DBV Technologies, other from DBV Technologies, other from Allergenix, other from Westmead Fertility Centre, and Research Grant (P.I., collaborator or consultant; pending and received grants) - National Health and Medical Research Council of Australia (NHMRC), Allergy and Immunology Foundation of Australasia, Nestle Health Sciences outside the submitted work.

Dr. Salter has nothing to disclose.

**Ethics approval**

Not applicable.

**Appendix A Supplementary data**

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.waojou.2021.100550>.

**Author details**

<sup>a</sup>School of Allied Health, Faculty of Health and Medical Sciences, The University of Western Australia, 35 Stirling Highway, Perth, WA, 6009, Australia. <sup>b</sup>The University of Western Australia, 35 Stirling Highway, Perth, WA, 6009, Australia. <sup>c</sup>The University of Adelaide, Walkerville, SA, 5081, Australia. <sup>d</sup>School of Population and Global Health, The University of Western Australia, Perth, WA, 6009, Australia. <sup>e</sup>School of Allied Health, The University of Western Australia, 35 Stirling Highway, Perth, WA, 6009, Australia. <sup>f</sup>The University of Sydney, Westmead, NSW, 2145, Australia.

**REFERENCES**

- Jackson KD, Howie LD, Akinbami OJ. *Trends in Allergic Conditions Among Children: United States, 1997-2011*; NCHS Data Brief, No. 121. Hyattsville, MD, USA: National Center for Health Statistics; 2013.
- Sicherer SH, Muñoz-Furlong A, Godbold JH, Sampson HA. US prevalence of self-reported peanut, tree nut, and sesame allergy: 11-year follow-up. *J Allergy Clin Immunol*. 2010;125:1322-1326.
- McKean M, Caughey AB, Leong RE, Wong A, Cabana MD. The timing of infant food introduction in families with a history of atopy. *Clin Pediatr*. 2015;54:745-751.
- Osborne NJ, Koplin JJ, Martin PE, et al. Prevalence of challenge-proven IgE-mediated food allergy using population-based sampling and predetermined challenge criteria in infants. *J Allergy Clin Immunol*. 2011;127(3):668-676.
- Sicherer SH, Sampson HA. Food allergy: a review and update on epidemiology, pathogenesis, diagnosis, prevention, and management. *J Allergy Clin Immunol*. 2018;141:41-58.
- Dunlop JH, Keet CA. Epidemiology of food allergy. *Immunol Allergy Clin Immunol*. 2018;38:13-25.
- Sicherer SH. Epidemiology of food allergy. *J Allergy Clin Immunol*. 2011;127:594-602.
- Jones SM, Burks AW. Food allergy. *N Engl J Med*. 2017;377:1168-1176.
- Berin MC, Sampson HA. Food allergy: an enigmatic epidemic. *Trends Immunol*. 2013;34:390-397.
- Schussler E, Sobel J, Hsu J, et al. Workgroup report by the joint task force involving American Academy of allergy, asthma & immunology (AAAAI); food allergy, anaphylaxis, dermatology and drug allergy (FADDA) Adverse reactions to foods committee and adverse reactions to drugs, biologicals, and latex committee; and the centers for disease control and prevention botulism clinical treatment guidelines work group - allergic reactions to botulinum antitoxin: a systematic review. *Clin Infect Dis*. 2017;66:S65-S72.
- Sasaki M, Koplin JJ, Dharmage SC, et al. Prevalence of clinic-defined food allergy in early adolescence: the School Nuts study. *J Allergy Clin Immunol*. 2017;141(1):391-398. <https://doi.org/10.1016/j.jaci.2017.05.041>.
- Australian Institute of Health and Welfare. *Chronic Diseases and Associated Risk Factors in Australia*; 2001. AIHW cat no. PHE 33. Available from: <https://www.aihw.gov.au/reports/chronic-disease/associated-risk-factors-australia-2001/contents/table-of-contents>.
- De Martinis M, Sirufo MM, Suppa M, Ginaldi L. New perspectives in food allergy. *Int J Mol Sci*. 2020;21:1474. <https://doi.org/10.3390/ijms21041474>.
- Agnes SY, Leung GW, Wong K, Tang MLK. Food allergy in the developing world. *J Allergy Clin Immunol*. 2018;141(1):76-78 E1. <https://doi.org/10.1016/j.jaci.2017.11.008>.
- Avery NJ, King RM, Knight S, Hourihane JOB. Assessment of quality of life in children with peanut allergy. *Pediatr Allergy Immunol*. 2003;4:378-382. <https://doi.org/10.1034/j.1399-3038.2003.00072.x>.
- Sicherer SH, Noone SA, Muñoz-Furlong A. The impact of childhood food allergy on quality of life. *Ann Allergy Asthma Immunol*. 2001;87(2):461-464.
- Gupta R, Holdford D, Bilaver L, Dyer A, Holl JL, Meltzer D. The economic impact of childhood food allergy in the United States. *JAMA Pediatr*. 2013;167(11):1026-1031.
- Chin B, Chan ES, Goldman RD. Early exposure to food and food allergy in children. *Can Fam Physician*. 2014;60(4):338-339.
- Chan ES, Abrams EM, Hildebrand KJ, Watson W. Early introduction of foods to prevent food allergy. *Allergy Asthma Clin Immunol*. 2018;14(Suppl 2):57. <https://doi.org/10.1186/s13223-018-0286-1>.
- du Toit George, Tsakok Teresa, Lack Simon, Lack Gideon. Prevention of food allergy. *J Allergy Clin Immunol*.

- 2016;137(4):998-1010. <https://doi.org/10.1016/j.jaci.2016.02.005>.
21. Netting MJ, Allen KJ. Advice about infant feeding for allergy prevention: a confusing picture for Australian consumers? *JPCH*. 2017;53:870-875.
22. Grimshaw KE, Allen K, Edwards CA, et al. Infant feeding and allergy prevention: a review of current knowledge and recommendations. A EuroPrevall state of the art paper. *Allergy*. 2009;64:1407-1416.
23. Koplin JJ, Allen KJ. Optimal timing for solids introduction - why are the guidelines always changing? *Clin Exp Allergy*. 2013;43:826-834.
24. Du Toit G, Roberts G, Sayre PH, et al. Randomized trial of peanut consumption in infants at risk for peanut allergy. *N Engl J Med*. 2015;372(9):803-813, 26.
25. Perkin MR, Logan K, Tseng A, et al. Randomized trial of introduction of allergenic foods in breast-fed infants. *N Engl J Med*. 2016;374(18):1733-1743.
26. Perkin MR, Logan K, Marris T, et al. Enquiring about Tolerance (EAT) study: feasibility of an early allergenic food introduction regimen. *J Allergy Clin Immunol*. 2016;137(5):1477-1486. e8.
27. Lavery WJ, Assa'ad A. How to prevent food allergy during infancy: what has changed since 2013? *Curr Opin Allergy Clin Immunol*. 2018;18:265-270.
28. Natsme O, Kabashima S, Nakazato J, et al. Two-step egg introduction for prevention of egg allergy in high-risk infants with eczema (PETIT): a randomized, double-blind, placebo-controlled trial. *Lancet*. 2017;389:276-286.
29. Shekelle Paul G, Woolf Steven H, Eccles Martin, Grimshaw Jeremy. Developing guidelines. *BMJ*. 1999;318:593. <https://doi.org/10.1136/bmj.318.7183.593>.
30. Consensus report, Institute of Medicine. *Clinical Practice Guidelines We Can Trust*; 2011. March 23 <http://www.iom.edu/Reports/2011/Clinical-Practice-Guidelines-We-Can-Trust.aspx>. Accessed January 13, 2012.
31. Brouwers MC, Kerkvliet K, Spithoff K. The AGREE Reporting Checklist: a tool to improve reporting of clinical practice guidelines. *BMJ*. 2016;352:i1152. <https://doi.org/10.1136/bmj.i1152>.
32. Brouwers MC, Kho ME, Browman GP, Burgers JS, Cluzeau F, Feder G. AGREE Next Steps Consortium. AGREE II: advancing guideline development, reporting and evaluation in health care. *CMAJ (Can Med Assoc J)*. 2010;182(18):E839-E842. <https://doi.org/10.1503/cmaj.090449>.
33. The AGREE Reporting Checklist: a tool to improve reporting of clinical practice guidelines. *BMJ*. 2016;352:i1152. <https://doi.org/10.1136/bmj.i1152>.
34. AGREE Next Steps Consortium. *The AGREE II Instrument*; 2017. Retrieved April 25th, 2020 from <http://www.agreetrust.org>.
35. World Health Organisation. *WHO Handbook for Guideline Development*. second ed. WHO Library Cataloguing-in-Publication Data; 2014. ISBN 978 92 4 154896 0.
36. AGREE II FAQ. <https://www.agreetrust.org/resource-centre/agree-ii/faq-agree-ii-2/Last>. Accessed June 6, 2020.
37. Zhang Z, Guo J, Su G, Li J, Wu H, Xie X. Evaluation of the quality of guidelines for myasthenia gravis with the AGREE II instrument. *PLoS One*. 2014;9(11), e111796. <https://doi.org/10.1371/journal.pone.0111796>.
38. Gao Y, Wang J, Luo X, et al. Quality appraisal of clinical practice guidelines for diabetes mellitus published in China between 2007 and 2017 using the AGREE II instrument. *BMJ Open*. 2019;9, e022392. <https://doi.org/10.1136/bmjopen-2018-022392>.
39. Greer FR, Sicherer SH, Burks AW. Effects of early nutritional interventions on the development of atopic disease in infants and children: the role of maternal dietary restriction, breastfeeding, timing of introduction of complementary foods, and hydrolyzed formulas. *Pediatrics*. 2008;121(1):183-191. <https://doi.org/10.1542/peds.2007-3022>.
40. Fleischer DM, Sicherer S, Greenhawt M, et al. Consensus communication on early peanut introduction and the prevention of peanut allergy in high-risk infants. *Ann Allergy Asthma Immunol*. 2015 Aug;115(2):87-90. <https://doi.org/10.1016/j.anai.2015.06.001>. Epub 2015 Jun 27. PMID: 26122934.
41. Greer FR, Sicherer SH, Burks AW. Committee on nutrition; section on allergy and immunology. The effects of early nutritional interventions on the development of atopic disease in infants and children: the role of maternal dietary restriction, breastfeeding, hydrolyzed formulas, and timing of introduction of allergenic complementary foods. *Pediatrics*. 2019 Apr;143(4), e20190281. <https://doi.org/10.1542/peds.2019-0281>. Epub 2019 Mar 18. PMID: 30886111.
42. Fiocchi A, Assa'ad A, Bahna S. Adverse reactions to foods committee; American College of allergy, asthma and immunology. Food allergy and the introduction of solid foods to infants: a consensus document. Adverse reactions to foods committee, American College of allergy, asthma and immunology. *Ann Allergy Asthma Immunol*. 2006 Jul;97(1):10-20. [https://doi.org/10.1016/s1081-1206\(10\)61364-6](https://doi.org/10.1016/s1081-1206(10)61364-6). quiz 21, 77, PMID: 16892776.
43. Tham EH, Shek LP, Van Bever HP, et al. Asia Pacific association of pediatric allergy, Respiriology & immunology (APAPARI). Early introduction of allergenic foods for the prevention of food allergy from an Asian perspective-an Asia Pacific association of pediatric allergy, Respiriology & immunology (APAPARI) consensus statement. *Pediatr Allergy Immunol*. 2018 Feb;29(1):18-27. <https://doi.org/10.1111/pai.12820>. Epub 2017 Nov 22. PMID: 29068090.
44. Prescott SL, Tang MLK. The Australasian Society of Clinical Immunology and Allergy position statement: summary of allergy prevention in children. *Med J Aust*. 2005;182(9):464-467. <https://doi.org/10.5694/j.1326-5377.2005.tb06787.x>.
45. Australasian Society of Clinical Immunology and Allergy. *ASCIa Infant Feeding Advice*; 2010. [https://www.allergy.org.au/images/stories/hp/info/ASCIa\\_Infant\\_Feeding\\_Advice\\_2010.pdf](https://www.allergy.org.au/images/stories/hp/info/ASCIa_Infant_Feeding_Advice_2010.pdf). Accessed January 27, 2020.
46. Joshi PA, Smith J, Vale S, Campbell DE. The Australasian Society of Clinical Immunology and Allergy infant feeding for allergy prevention guidelines. *MJA*. 2019;210(2):89-93. <https://doi.org/10.5694/mja2.12102>.
47. Australasian Society of Clinical Immunology and Allergy (ASCIa). *ASCIa Guidelines: Infant Feeding and Allergy Prevention*; 2016. <https://allergy.org.au/hp/papers/infant-feeding-and-allergy-prevention>. Accessed January 27, 2020.



48. Stiefel G, Anagnostou K, Boyle RJ, et al. BSACI guideline for the diagnosis and management of peanut and tree nut allergy. *Clin Exp Allergy*. 2017;47:719-739. <https://doi.org/10.1111/cea.12957>.
49. Turner PJ, Feeney M, Meyer R, Perkin MR, Fox AT. Implementing primary prevention of food allergy in infants: new BSACI guidance published. *Clin Exp Allergy*. 2018;48:912-915. <https://doi.org/10.1111/cea.13218>.
50. Chan ES, Cummings C. Canadian paediatric society, community Paediatrics committee and allergy section. Dietary exposures and allergy prevention in high-risk infants: a joint statement with the Canadian society of allergy and clinical immunology. *Paediatr Child Health*. 2013;18(10):545-554. <https://doi.org/10.1093/pch/18.10.545>.
51. Abrams EM, Hildebrand K, Blair B, Chan ES. Timing of introduction of allergenic solids for infants at high risk. *Paediatr Child Health*. 2019 Feb;24(1):56-57. <https://doi.org/10.1093/pch/pxy195>. Epub 2019 Feb 15. PMID: 30833823; PMCID: PMC6376285.
52. Muraro A, Dreborg S, Halken S, et al. Dietary prevention of allergic diseases in infants and small children. Part III: critical review of published peer-reviewed observational and interventional studies and final recommendations. *Pediatr Allergy Immunol*. 2004 Aug;15(4):291-307. <https://doi.org/10.1111/j.1399-3038.2004.00127.x>. PMID: 15305938.
53. Muraro A\*, Halken S\*, Arshad SH, et al. EAACI food allergy and anaphylaxis guidelines. Primary prevention of food allergy. *Allergy*. 2014;69:590-601. <https://doi.org/10.1111/all.12398>.
54. Høst A, Koletzko B, Dreborg S, et al. Dietary products used in infants for treatment and prevention of food allergy. Joint statement of the European society for paediatric Allergology and clinical immunology (ESPACI) committee on hypoallergenic formulas and the European society for paediatric Gastroenterology, Hepatology and nutrition (ESPGHAN) committee on nutrition. *Arch Dis Child*. 1999 Jul;81(1):80-84. <https://doi.org/10.1136/adc.81.1.80>. PMID: 10373144; PMCID: PMC1717972.
55. Agostoni C, Decsi T, Fewtrell M, et al, ESPGHAN Committee on Nutrition. Complementary feeding: a commentary by the ESPGHAN committee on nutrition. *J Pediatr Gastroenterol Nutr*. 2008 Jan;46(1):99-110. <https://doi.org/10.1097/01.mpg.0000304464.60788.bd>. PMID: 18162844.
56. Fewtrell M, Bronsky J, Campoy C, et al. Complementary feeding: a position paper by the European society for paediatric Gastroenterology, Hepatology, and nutrition (ESPGHAN) committee on nutrition. *J Pediatr Gastroenterol Nutr*. 2017 Jan;64(1):119-132. <https://doi.org/10.1097/MPG.0000000000001454>. PMID: 28027215.
57. Pelkonen AS, Kuitunen M, Dunder T, et al. Allergy in children: practical recommendations of the Finnish Allergy Programme 2008-2018 for prevention, diagnosis, and treatment. *Pediatr Allergy Immunol*. 2012 Mar;23(2):103-116. <https://doi.org/10.1111/j.1399-3038.2012.01298.x>. PMID: 22432881.
58. Mucbe-Borowski C, Kopp M, Reese I, Sitter H, Werfel T, Schäfer T. Allergy prevention. *Dtsch Arztebl Int*. 2009;106(39):625-631. <https://doi.org/10.3238/arztebl.2009.0625>.
59. Schäfer T, Bauer CP, Beyer K, et al. S3-Guideline on allergy prevention: 2014 update: guideline of the German society for Allergology and clinical immunology (DGAKI) and the German society for pediatric and adolescent medicine (DGKJ). *Allergo J Int*. 2014;23(6):186-199. <https://doi.org/10.1007/s40629-014-0022-4>.
60. Chan AWM, Chan JKC, Lee TH, Leung TF, Tam AYC. *Guidelines for Allergy Prevention in Hong Kong*; 2015. [http://www.allergy.org.hk/HKIA%20-%20Guidelines%20for%20Allergy%20Prevention%20in%20Hong%20Kong%20\(Final\).pdf](http://www.allergy.org.hk/HKIA%20-%20Guidelines%20for%20Allergy%20Prevention%20in%20Hong%20Kong%20(Final).pdf). Accessed April 13, 2020.
61. Chan AWM, Chan JKC, Tam AYC, Lee TH. Guidelines for allergy prevention in Hong Kong. *Medical Practice*. 2016;22(3):279-285.
62. Chan JKC, Chan AWM, Ho MHK, Lee TH. *HKIA Position Paper on Prevention of Peanut Allergy in High Risk Infants*; 2016. [http://www.allergy.org.hk/HKIA%20-%20Guidelines%20for%20Prevention%20of%20Peanut%20Allergy%20\(Final\).pdf](http://www.allergy.org.hk/HKIA%20-%20Guidelines%20for%20Prevention%20of%20Peanut%20Allergy%20(Final).pdf). Accessed April 13, 2020.
63. di Mauro G, Bernardini R, Barberi S, et al. Prevention of food and airway allergy: consensus of the Italian society of preventive and social Paediatrics, the Italian society of paediatric allergy and immunology, and Italian society of Pediatrics. *World Allergy Organ J*. 2016;9:28. <https://doi.org/10.1186/s40413-016-0111-6>. Published 2016 Aug 18.
64. Ebisawa M, Ito K, Fujisawa T. Committee for Japanese pediatric guideline for food allergy, the Japanese society of pediatric allergy and clinical immunology, the Japanese society of Allergology. Japanese guidelines for food allergy 2017. *Allergol Int*. 2017 Apr;66(2):248-264. <https://doi.org/10.1016/j.alit.2017.02.001>. Epub 2017 Mar 10. PMID: 28285847.
65. Boyce JA, Assa'ad A, Burks AW, et al, NIAID-Sponsored Expert Panel. Guidelines for the diagnosis and management of food allergy in the United States: report of the NIAID-sponsored expert panel. *J Allergy Clin Immunol*. 2010;126(6 Suppl):S1-S58. <https://doi.org/10.1016/j.jaci.2010.10.007>.
66. Togias A, Cooper SF, Acebal ML, et al. Addendum guidelines for the prevention of peanut allergy in the United States: report of the National Institute of Allergy and Infectious Diseases-sponsored expert panel. *World Allergy Organ J*. 2017;10(1):1. <https://doi.org/10.1186/s40413-016-0137-9>. Published 2017 Jan 6.
67. Recto MST, Genuino MLG, Castor MAR, et al. Dietary primary prevention of allergic diseases in children: the Philippine guidelines. *Asia Pac Allergy*. 2017;7(2):102-114. <https://doi.org/10.5415/apallergy.2017.7.2.102>.
68. *Joint Statement from the Scientific Advisory Committee on Nutrition and the Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment. Assessing the Health Benefits and Risks of the Introduction of Peanut and Hen's Egg into the Infant Diet before Six Months of Age in the UKA*; 2018. <https://cot.food.gov.uk/sites/default/files/jointsacncotalergystatementfinal2.pdf>. Accessed March 17, 2020.
69. Ministry of Health Singapore. *Management of Food Allergy: AMS-MOH Clinical Practice Guidelines 2*; 2010. [https://www.moh.gov.sg/docs/librariesprovider4/guidelines/cpg\\_management-of-food-allergy.pdf](https://www.moh.gov.sg/docs/librariesprovider4/guidelines/cpg_management-of-food-allergy.pdf). Accessed March 9, 2020.
70. Lee BW, Aw MM, Chiang WC, et al. Academy of medicine, Singapore-Ministry of Health clinical practice guidelines: management of food allergy. *Singapore Med J*. 2010 Jul;51(7):599-607. Erratum in: *Singapore Med J*. 2013 Aug;54(8):474. Aw, M [corrected to Aw, M M]; Shek, L [corrected to Shek, L P]. PMID: 20730402.

71. Halken Susanne, et al. EAACI guideline: preventing the development of food allergy in infants and young children (2020 update). *Pediatr Allergy Immunol*. 2020. <https://doi.org/10.1111/pai.13496>.
72. Jeurink PV, Knipping K, Wiens F, et al. Importance of maternal diet in the training of the infant's immune system during gestation and lactation. *Crit Rev Food Sci Nutr*. 2019;59(8): 1311-1319.
73. Wang Y, Allen KJ, Koplin JJ. Dietary intervention for preventing food allergy in children. *Curr Opin Pediatr*. 2017;29(6):704-710.
74. Grimshaw KE, Allen K, Edwards CA, et al. Infant feeding and allergy prevention: a review of current knowledge and recommendations. A EuroPrevall state of the art paper. *Allergy*. 2009;64:1407-1416.
75. National Health and Medical Research Council. *The Australian Dietary Guidelines: Summary*; 2013. [https://www.eatforhealth.gov.au/sites/default/files/files/the\\_guidelines/n55a\\_australian\\_dietary\\_guidelines\\_summary\\_book.pdf](https://www.eatforhealth.gov.au/sites/default/files/files/the_guidelines/n55a_australian_dietary_guidelines_summary_book.pdf). Accessed June 6, 2020.
76. World Health Organization, UNICEF. *Global Strategy for Infant and Young Child Feeding*. Geneva, Switzerland: World Health Organization; 2003.
77. National Health and Medical Research Council. *Infant Feeding Guidelines*. Canberra: The Council; 2012.
78. Greer FR, Sicherer SH, Burks AW, et al. The effects of early nutritional interventions on the development of atopic disease in infants and children: the role of maternal dietary restriction, breastfeeding, hydrolyzed formulas, and timing of introduction of allergenic complementary foods. *Pediatrics*. 2019;143.
79. Faucher MA. An updated scientific review of the benefits of breastfeeding with additional resources for use in everyday practice. *J Midwifery Wom Health*. 2012;57:422-423.
80. Boyle RJ, Ierodiakonou D, Khan T, et al. Hydrolysed formula and risk of allergic or autoimmune disease: systematic review and meta-analysis. *BMJ*. 2016 Mar 8;352:i974. <https://doi.org/10.1136/bmj.i974>. PMID: 26956579; PMCID: PMC4783517.
81. Prescott SL, Smith P, Tang M, et al. The importance of early complementary feeding in the development of oral tolerance: concerns and controversies. *Pediatr Allergy Immunol*. 2008;19: 375-380.
82. Shekelle P, Aronson MD, Givens J. Overview of clinical practice guidelines. *UpToDate*. Apr. 2020;8. [https://www.uptodate-com.ezproxy.library.uwa.edu.au/contents/overview-of-clinical-practice-guidelines/print?search=Overview%20of%20clinical%20practice%20guidelines&source=search\\_result&selectedTitle=1~150&usage\\_type=default&display\\_rank=1](https://www.uptodate-com.ezproxy.library.uwa.edu.au/contents/overview-of-clinical-practice-guidelines/print?search=Overview%20of%20clinical%20practice%20guidelines&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1). Accessed July 26, 2020.