



OPEN

A comprehensive list of the *Bunyavirales* replication promoters reveals a unique promoter structure in *Nairoviridae* differing from other virus families

Yutaro Neriya¹, Shohei Kojima², Arata Sakiyama³, Mai Kishimoto⁴, Takao Iketani⁵, Tadashi Watanabe⁶, Yuichi Abe⁷, Hiroshi Shimoda⁸, Keisuke Nakagawa⁹, Takaaki Koma¹⁰ & Yusuke Matsumoto¹¹✉

Members of the order *Bunyavirales* infect a wide variety of host species, including plants, animals and humans, and pose a threat to public health. Major families in this order have tri-segmented negative-sense RNA genomes, the 5' and 3' ends of which form complementary strands that serve as a replication promoter. Elucidation of the mechanisms by which viral polymerases recognize the promoter to initiate RNA synthesis is important for understanding viral replication and pathogenesis, and developing antivirals. A list of replication promoter configuration patterns may provide details on the differences in the replication mechanisms among bunyaviruses. By using public sequence data of all known bunyavirus species, we constructed a comprehensive list of the replication promoters comprising 40 nucleotides in both the 5' and 3' ends of the genome that form a specific complementary strand. Among tri-segmented bunyaviruses, members of the family *Nairoviridae*, including the highly pathogenic Crimean-Congo hemorrhagic fever virus, have evolved a GC-rich promoter structure differing from that of other families. The unique promoter structure might be related to the large genome size of the family *Nairoviridae* among bunyaviruses, and the large genome architecture might confer pathogenic advantages. The promoter list provided in this report is useful for predicting the virus family-specific replication mechanisms of bunyaviruses.

Bunyavirales is a new order proposed by the International Committee on Taxonomy of Viruses (ICTV) in 2016. It consists of 12 families of divergent viruses: *Arenaviridae*, *Cruliviridae*, *Fimoviridae*, *Hantaviridae*, *Leishbuviridae*, *Mypoviridae*, *Nairoviridae*, *Peribunyaviridae*, *Phasmaviridae*, *Phenuiviridae*, *Tospoviridae*, and *Wupedeviridae*. Members of *Bunyavirales* have a segmented single-stranded negative-sense or ambisense RNA genome. The

¹School of Agriculture, Utsunomiya University, 350 Mine-machi, Utsunomiya, Tochigi 321-8505, Japan. ²Genome Immunobiology RIKEN Hakubi Research Team, RIKEN Center for Integrative Medical Sciences, 1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama, Kanagawa 230-0045, Japan. ³Department of Bacteriology, Graduate School of Medicine, Osaka City University, 1-4-3, Asahi-machi, Abeno-ku, Osaka 545-8585, Japan. ⁴Division of Molecular Pathobiology, International Institute for Zoonosis Control, Hokkaido University, North 20, West 10 Kita-ku, Sapporo 001-0020, Japan. ⁵Laboratory of Pathogenesis of Viral Infectious Diseases, Tokyo University of Agriculture and Technology, 3-8-1 Saiwai-chou, Fuchu, Tokyo 183-8509, Japan. ⁶Department of Virology, Graduate School of Medicine, University of the Ryukyus, Uehara 207, Nishihara-cho, Nakagami-gun, Okinawa 903-0215, Japan. ⁷Division of Molecular Diagnostics, Aichi Cancer Center, 1-1, Kanokoden, Chikusa-ku, Nagoya 464-8681, Japan. ⁸Laboratory of Veterinary Microbiology, Joint Faculty of Veterinary Medicine, Yamaguchi University, 1677-1, Yoshida, Yamaguchi 753-8515, Japan. ⁹Laboratory of Veterinary Microbiology, Joint Department of Veterinary Medicine, Faculty of Applied Biological Sciences, Gifu University, 1-1 Yanagido, Gifu 501-1193, Japan. ¹⁰Department of Microbiology, Tokushima University Graduate School of Biomedical Sciences, 3-18-15 Kuramoto, Tokushima 770-8503, Japan. ¹¹Department of Microbiology and Cell Biology, Tokyo Metropolitan Institute of Medical Science, 2-1-6 Kamikitazawa, Setagaya-ku, Tokyo 156-8506, Japan. ✉email: matsumoto-ys@igakuken.or.jp

families *Arenaviridae*, *Hantaviridae*, *Nairoviridae*, *Peribunyaviridae*, and *Phenuiviridae* include several important pathogens that can cause severe diseases in animals, including humans, while the families *Fimoviridae*, *Phasmaviridae*, *Phenuiviridae*, and *Tospoviridae* include pathogens associated with plant diseases.

Major groups of bunyaviruses possess tri-segmented negative-sense RNA genomes, and share the same genetic organization consisting of three segments, *i.e.*, the small (S), medium (M), and large (L) segments, based on their relative sizes. Each segment acts as a template for the replication of a positive-sense antigenome, and for the transcription of mRNA. The S segment encodes the nucleocapsid protein (NP), the M segment encodes a glycosylated polyprotein precursor (GPC) that is cleaved into envelope spike proteins Gn and Gc, and the L segment encodes the L protein, an RNA-dependent RNA polymerase (RdRp) responsible for the transcription and replication of the three RNA segments. The RNA synthesis activities of the three RNA segments are regulated by nucleotide (nt) sequences within the 3' and 5' untranslated regions (UTRs), which flank the S, M, and L open reading frames. The terminal nts of 3' and 5' UTRs exhibit complementarity, and such sequences have been shown to bind to, and influence the activity of, viral RdRp, promoting transcription to yield a 5'-capped mRNA by using cleaved host mRNA as a primer, and replication that results in the synthesis of a full-length copy of the genome template.

Bunyavirus promoters are composed of two promoter elements, *i.e.*, promoter element 1 (PE1), the genomic extreme complement region, and PE2, the complement region located behind PE1, which was first described in Bunyamwera virus (BUNV) of the family *Peribunyaviridae*^{1,2}. PE1 comprises approximately 10–15 nts located at the extreme termini of the genome that are strictly conserved among all three segments. These nts have been shown to interact with L protein at separate sites in La Crosse virus (LACV) of the family *Peribunyaviridae*^{3,4}. PE2 comprises segment-specific nts at subsequent positions that are required to form canonical Watson–Crick base-pairing with corresponding nts at the opposite end of the template^{1,2,5}. These RdRp–RNA and RNA–RNA interactions are thought to account for the pseudocircular form of viral ribonucleoprotein complexes^{6,7}. In BUNV, sequence changes within PE1 have a significant effect on promoter function, but adjacent nts within PE2 are highly resistant to sequence changes, provided that their interterminal Watson–Crick base-pairing potential is maintained¹. For the family *Nairoviridae*, little is known on the roles of the 3'- and 5'-terminal UTRs in regulating RNA synthesis. As with other *Bunyavirales* members, the UTRs of all nairoviruses comprise highly conserved terminal proximal nts (PE1) shared by all three segments, followed by less conserved regions that are segment-specific nts (PE2). The importance of these segment-specific nts in RNA synthesis has been partially examined using a minigenome reporter assay in non-pathogenic Hazara virus (HAZV), which is closely related to Crimean-Congo hemorrhagic fever virus (CCHFV)⁸. PE1 and PE2 were found to be separated by a spacer region, which exhibited a critical requirement to be short in length and lack base-pairing ability. Taken together, the accumulated data indicate that the promoter structure of bunyaviruses differs among virus families. Understanding these properties is essential for developing antivirals targeting viral RNA synthesis and processing.

To characterize the promoter structure of diverse bunyaviruses, we constructed a list of all viral promoters that exhibit complementarity and each of the nt counts within the first 40 nts of both the 5' and 3' (anti)genomic ends. We found that the promoters of the family *Nairoviridae* differ from those of other virus families, and have characteristics unique to the family, which has a large genome size.

Results

Construction of the list of promoters in the order *Bunyavirales*. This study aimed to characterize the promoter structure of all virus species in the order *Bunyavirales*, including the families *Arenaviridae*, *Cruliviridae*, *Fimoviridae*, *Hantaviridae*, *Leishbuviridae*, *Myoviridae*, *Nairoviridae*, *Peribunyaviridae*, *Phasmaviridae*, *Phenuiviridae*, *Tospoviridae*, and *Wupedeviridae*, which are tri-segmented or multi-segmented viruses, as summarized in Fig. 1. The complementarity of the 5' and 3' extreme 40 nts of the genomic ends of virus genomes registered in the ICTV list (<https://ictv.global/taxonomy>) was analyzed. Because there are incomplete genome sequences in the National Center for Biotechnology Information (NCBI) database that do not precisely cover the genome extremes, we selected viral sequences with complete complementarity in the terminal +1 to +3 nts (some exceptions with non-complementary +1 nt are included). The complement structure of 5' and 3' genomic ends (positive-sense form) as well as the genome length, counts of G:C/A:U complementarity, and counts of each nt (A, U, G and C) in the promoter region were calculated by using an automatic calculating system based on an Excel file (Supplementary Table 1), and the results are tabulated in Supplementary Table 2. A dataset was generated for each virus species that had complete data for all segments, including: tri-segmented bunyaviruses *Arenaviridae* (2 species), *Cruliviridae* (3 species), *Hantaviridae* (24 species), *Myoviridae* (1 species), *Nairoviridae* (20 species), *Peribunyaviridae* (56 species), *Phasmaviridae* (2 species), *Phenuiviridae* (47 species), *Tospoviridae* (19 species) and *Wupedeviridae* (1 species); multi-segmented bunyaviruses *Fimoviridae* (17 species) and *Phenuiviridae* (14 species); and di-segmented bunyavirus *Arenaviridae* (23 species; characterized in Supplementary Fig. 1).

Characteristics of the replication promoters of five major virus families. For five major tri-segmented virus families, *i.e.*, *Peribunyaviridae*, *Phenuiviridae*, *Tospoviridae*, *Hantaviridae*, and *Nairoviridae*, we examined the conservation of the nts in the 40 nts at the promoter region using the sequence generator WebLogo⁹ (Fig. 2A, which shows representative M segments, and Supplementary Figs. 2 and 3). The promoters differed among viruses: the initial nt was adenosine (A) in *Peribunyaviridae*, *Phenuiviridae* and *Tospoviridae*, and was uridine (U) in *Hantaviridae* and *Nairoviridae*. These promoters were further categorized into those starting with a tri-nt repeat (5'-AGUAGU and 5'-UAGUAG) and those starting with a di-nt repeat (5'-ACAC, 5'-AGAG and 5'-UCUC) (Fig. 2A,B). We next examined the percentages of G:C and A:U complementarity at every nt position in the promoter region among the virus species in each virus family (Fig. 2A). The complementarity conformation was remarkably different among virus families. G:C complementarity was relatively

ICTV list: 590 virus species in the *Bunyavirales* order**→ 5'- and 3'-complementarity (*) analysis of 40 nts (Supplementary Table 1)**

```

5' -AGUAGUGUACUCCUACAUUAGAAAAUUUAAAAUUAUAC
    ***** ***** ** * * * * *
3' -UCAUCACACGAGGAUGUAUUCUUUUAACAUGAAAAACUU

```

Promoter list of *Bunyavirales* (Supplementary Table 2)

Genome length / Promoter structure / G:C and A:U complementarity counts / Each nts counts

Tri-segment	Multi-segment
<i>Arenaviridae</i> (2)	<i>Fimoviridae</i> (17)
<i>Cruliviridae</i> (3)	<i>Phenuiviridae</i> (14)
<i>Hantaviridae</i> (24)	<i>Arenaviridae</i> (23)
<i>Myxoviridae</i> (1)	
<i>Nairoviridae</i> (20)	
<i>Peribunyaviridae</i> (56)	
<i>Phasmaviridae</i> (2)	
<i>Phenuiviridae</i> (47)	
<i>Tospoviridae</i> (19)	
<i>Wupedeviridae</i> (1)	

Figure 1. Construction of the promoter list of bunyaviruses. A schematic diagram of the promoter list construction. The promoter structure of 590 species of bunyaviruses registered in the ICTV list was analyzed. The genome length, promoter structure, G:C and A:U complementarity counts, and each nt (A, U, G, and C) count in the 40 nts of the promoters in the genomic ends are listed in Supplementary Table 2. The calculations were performed in an Excel file using automatic calculations for the promoter region as described in Supplementary Table 1.

higher in the virus genomes with a promoter starting with U than in those with a promoter starting with A. The genomes of the families *Hantaviridae* and *Nairoviridae* contain high G:C complementarity at the 13–16 nt and 17–21 nt positions, respectively. In some viruses belonging to the family *Phenuiviridae*, a shift of 1 nt at the 10th position from the 5' extreme appeared to increase the complementarity of the subsequent 5' and 3' ends¹⁰, but it did not increase the total G:C complementarity frequency in the promoter region of *Phenuiviridae* (data not shown). It has been reported that HAZV has a promoter composed of two complementary regions of PE1 and PE2 separated by a spacer region formed by non-complementary sequences at the 13–16 nt position⁸. We found the same feature in most virus species of the *Nairoviridae* family (Fig. 2 and Supplementary Fig. 3).

To analyze the promoter structure in more depth, the G:C and A:U complementarity in the 40-nt promoter region was determined in three segments of the five virus families. The average complementarity count of virus species in each virus family is shown in Fig. 3A. The A:U complementarity counts were higher than the G:C complementarity counts in all segments for all virus families. However, G:C complementarity was particularly higher in the promoters of the family *Nairoviridae* than in those of other virus families (Fig. 3A). Each nt (A, U, G, and C) in the promoter region was counted, and the average value within each virus family is shown in Fig. 3B. In *Phenuiviridae*, *Tospoviridae*, and *Hantaviridae*, A in the 5' end and U in the 3' end were frequent in all segments. In *Peribunyaviridae*, both A and U were abundant at the 5' and 3' ends. In contrast, in the family *Nairoviridae*, C and G were more frequent at the 5' and 3' ends, respectively, than in other virus families. We demonstrated that the family *Nairoviridae* had more G:C complementarity as well as higher G/C counts in the 40 nts of the promoter region than other virus families, which is suggestive of stronger affinity for base pairing at both genomic ends.

Genome length of virus families in the order *Bunyavirales*. The promoter structure of *Nairoviridae* differed from that of other tri-segmented virus families in that it had high G:C complementarity and a non-complementary nt spacer region. To investigate the relationship between these features and the characteristics of the viral genomes, the genome lengths of all viral species of the five virus families were studied. Figure 4A shows the average total genome length (combined length of the L, M, and S segments) of all virus species in the five virus families. The full genome lengths of the families *Peribunyaviridae*, *Phenuiviridae*, and *Hantaviridae* were comparable, while the genome length of the family *Tospoviridae* was larger, and that of the family *Nairoviridae* was the largest. The length of each segment was also examined in all virus species, and the average lengths within virus families are shown in Fig. 4B. The family *Tospoviridae* had relatively large L, M, and S segments. The L segment of *Nairoviridae* was the largest among all segments of all viruses.

Genome length of viruses in the family *Nairoviridae*. Among tri-segmented viruses belonging to the order *Bunyavirales*, only the family *Nairoviridae* includes highly pathogenic viruses categorized as biosafety level (BSL)-4 pathogens that cause hemorrhagic fever in humans, such as CCHFV¹¹. We hypothesized that the high pathogenicity of this virus family in mammals may be related to its large genome size. We examined the length




Figure 2. Characteristics of bunyavirus replication promoters in the M segment. (A) Sequence conservation and nt complementarity counts in the promoters of five major virus families. Sequence conservation was analyzed using the sequence logo generator WebLogo. The percentages of G:C and A:U complementarity in the promoter region (1–40 nts) among virus species in each family are shown as a bar graph. Virus species in the following families were examined: *Peribunyaviridae* (n = 56), *Phenuiviridae* (n = 47), *Tospoviridae* (n = 19), *Hantaviridae* (n = 24), and *Nairoviridae* (n = 20). (B) Characteristics of promoters. Virus promoters were divided into those that start with A and those that start with U, and further characterized according to the presence of a 3-nt- or 2-nt repeat. *Phenuiviridae* and *Tospoviridae* possess an ambisense genome.

of the available sequences annotated to “*Nairoviridae*” in the NCBI database. We first selected virus genome sequences possessing 5'-UCUC—GAGA-3' ends, which are the most conserved genomic end sequences in nai-




Figure 3. Counts of G:C and A:U complementarity, and each nt (A, U, G, and C) in the promoters. (A and B) The counts of G:C and A:U complementarity (A) and each nt (B) in the 40 nts of the promoters. Bars represent the means and standard deviations. Virus species in the following families were examined: *Peribunyaviridae* (n = 56), *Phenuiviridae* (n = 47), *Tospoviridae* (n = 19), *Hantaviridae* (n = 24), and *Nairoviridae* (n = 20). ** $p < 0.01$, * $p < 0.05$, two-tailed unpaired Student's *t*-test (G:C vs. A:U).

roviruses, in the L, M and S segments. The lengths of these sequences are shown in Fig. 5A. The family *Nairoviridae* contains two highly pathogenic viruses in mammals, i.e., CCHFV and Nairobi sheep disease virus (NSDV), which have a mortality rate of 30% and 90% in humans and small ruminants, respectively^{12,13}. The length of the sequences of CCHFV and NSDV are shown as red and yellow bars, respectively, in Fig. 5A. The lengths of the M segments of CCHFV and NSDV were found to be the longest among virus genome sequences possessing 5'-UCUC—GAGA-3' ends.




Figure 4. Genome length of bunyaviruses. (A) The total genome length (L segment + M segment + S segment) of each virus family. (B) The lengths of the L, M, and S segments in each virus family. Bars represent the means and standard deviations. Virus species in the following families were examined: *Peribunyaviridae* (n = 56), *Phenuiviridae* (n = 47), *Tospoviridae* (n = 19), *Hantaviridae* (n = 24), and *Nairoviridae* (n = 20). ** $p < 0.01$, one-way analysis of variance followed by Tukey's test, *** $p < 0.01$, in comparison to the other four families.

Discussion

In this study, we tabulated the replication promoter structures of all known virus species in the order *Bunyavirales*. Our analysis focused on five major tri-segmented virus families, and the results indicated that the genomes can be divided into two categories: those with a promoter starting with A (families *Peribunyaviridae*, *Phenuiviridae*, and *Tospoviridae*) and those with a promoter starting with U (families *Hantaviridae* and *Nairoviridae*; Fig. 2). It has been shown that viral RNA polymerases have the ability to initiate RNA synthesis with a purine (G or A), but not with a pyrimidine (U or C)¹⁴. Therefore, the 5'-U genomic end of *Hantaviridae* and *Nairoviridae* is unconventional. The genomes of LACV and Rift Valley fever virus (RVFV) of the families *Peribunyaviridae* and *Phenuiviridae*, respectively, contain a 5'-triphosphate end that starts with A (5'-pppA)^{15,16}. The 5'-pppA is generated by viral RdRp that recognizes the opposite U as the template. As seen in several segmented and non-segmented RNA viral polymerases^{17–19}, bunyaviral RdRp synthesizes RNA from an internal nt, and not from the terminus of the template. In LACV, RNA synthesis is initiated with A using the U at the +4 position of the antigenome (3'-UCAUCA) as the template during genome replication⁴. The elongated product, 5'-pppAGU, is realigned to the +1 to +3 position of the antigenome template (3'-UCAUCA), and is further elongated to generate 5'-pppAGUAGU. Accordingly, the position of U responsible for RNA synthesis initiation is presumably the +3 position in the *Phenuiviridae* (3'-UGUG) and *Tospoviridae* (3'-UCUC) antigenomes. This indicates that the 5'-pppAC and 5'-pppAG products realign to the 3'-UGUG and 3'-UCUC of the antigenomes, respectively, and are further elongated to generate 5'-pppACAC and 5'-pppAGAG, respectively, which are precise complementary chains of the antigenome templates. In contrast, the genomes of Hantaan virus (HTNV) in the family *Hantaviridae*, and CCHFV in the family *Nairoviridae* contain a 5'-monophosphate end^{15,20}, suggesting an unconventional RNA processing event during replication. In HTNV, RNA synthesis is initiated with an internal G at the +3 position by using a C of the 3'-AUC \overline{A} UC of the antigenome as the template²⁰. Subsequently, the elongated 5'-pppGUA product realigns to the 3'-AUC \overline{A} UC to further produce 5'-pppGUAGUA. Then, the extreme 5'-pppG is removed by an endoribonuclease activity of viral RdRp to produce 5'-pUAGUA (5'-monophosphate end)²⁰. The endoribonuclease activity of RdRp is responsible for the cap snatching that cleaves the 5' end of the host mRNA for use as a transcription primer²¹. The -1 position of viral mRNA of HTNV is G, indicating that viral RdRp can cleave host mRNA after the G nt (cleave GpN to produce G/pN) during transcription. In *Nairoviridae*, the -1 position of viral mRNA is C^{22,23}, and it is also generated via the cap-snatching mechanism. Similar to the RNA synthesis in *Hantaviridae*, it is supposed that nairoviral RNA synthesis is internally initiated with 5'-pppC at the +2 position by using the G of 3'-AGA of the antigenome as the template. Subsequently, the 5'-pppCU product would realign to the 3'-AGA, and be further elongated to generate 5'-pppCUCU. The 5'-pppC would then be removed, resulting in the production of a 5'-monophosphate end. Therefore, although the hantaviral RdRp is a conventional enzyme that initiates RNA synthesis with a purine (G), the nairoviral RdRp is considered to be an unconventional enzyme that can initiate with a pyrimidine (C). Such a difference may be important for the targeting of novel antivirals specific for nairoviral diseases.

Our analysis additionally confirmed that most bunyaviral genomes begin with a di- or tri-nt repeat (Fig. 2), which has been suggested previously^{20,24}. The repeats can determine the initiation site for RdRp (e.g., +2 in *Nairoviridae*, +3 in *Hantaviridae*, *Phenuiviridae* and *Tospoviridae*, and +4 in *Peribunyaviridae*), which is important for the prime-realign RNA synthesis mechanism. The biological significance of the internal position of RNA synthesis initiation is unclear. It is likely that the di-nt repeat is restricted to virus families possessing an




Figure 5. Genome length of virus species in the family *Nairoviridae*. (A) Genome length of each segment of nairovirus sequences (5'-UCUC—GAGA-3') registered in the NCBI Refseq database. Bars represent the precise sequence length of each virus. CCHFV and NSDV are shown in red and yellow. (B) Schematic diagram of GPC encoded in the M segment of nairoviruses. DUGV: Dugbe virus, TFLV: Tofla virus, ERVEV: Erve virus.

ambisense genome, such as *Phenuiviridae* and *Tospoviridae*, as well as *Nairoviridae* (for which only CCHFV has been reported)²⁵. This suggests that the ambisense coding property may be related to the di-nt repetition in

the genomic ends. If this is true, analysis of genomic end repetition patterns may enable the elucidation of new transcripts in various bunyaviral genomes.

The *Nairoviridae* promoter appears to have high G:C complementarity in the 17 to 21-nts region (Fig. 2A and Supplementary Fig. 2), and this likely reflects the high G:C complementarity rate at the promoter region (Fig. 3A). Interestingly, this GC-rich dsRNA region is located after a spacer region composed of non-complementary bases around the 14th position in all three segments, as has been reported previously in HAZV and CCHFV^{8,26}. We have previously suggested the possibility that the HAZV polymerase can recognize this GC-rich dsRNA as a promoter element essential for RNA synthesis initiation via an unidentified domain of the L protein⁸. This kind of specific protein-RNA interaction has been proposed to be a suitable target for antivirals against CCHFV, which is closely related to HAZV. Our comprehensive analysis of the promoter list also suggested that this kind of strategy may be applicable for all viruses belonging to the family *Nairoviridae*.

In bunyaviruses, genome replication in each segment is regulated by the segment-specific promoter strength, but the variations in nts (A, U, G, and C) in each promoter region do not differ significantly among the L, M, and S segments in all virus families, except for *Nairoviridae* (Fig. 3B). It is possible that the promoter strength among segments is determined by slight differences in the promoter structure that do not affect the total complementarity counts or nt variations. Viruses in the family *Phenuiviridae* and CCHFV of the family *Nairoviridae* have an ambisense S segment, but there is no nt variation pattern in the promoter that is unique to the S segment (Fig. 2B). This suggests that the nt variation in the promoter was not affected by the presence of the ambisense segment during the viral evolution process. On the other hand, the nt variation in the promoter of the nairoviral L segment was different from that of the M and S segments, *i.e.*, it was observed to have less G and C at the 5' and 3' ends, respectively (Fig. 2B). The nairoviral L segment is remarkably long when compared to other nairoviral segments and the genomes of other virus families (Fig. 4B). This large genome size may be associated with the promoter structure.

It remains unclear why the genome of the family *Nairoviridae* is so large. *Nairoviridae* is the only tri-segmented virus family that includes hemorrhagic fever viruses classified as BSL-4 pathogens, such as CCHFV. We hypothesized that the large genome size of the family *Nairoviridae* may be related to its high pathogenesis in mammals. Although the length of the L segment in *Nairoviridae* is the longest among all bunyaviruses, it is not particularly long among the highly pathogenic viruses in this family (Fig. 5A). Rather, our analysis confirmed that among viruses in the family *Nairoviridae*, the M segment is the largest segment in two highly pathogenic viruses in mammals, CCHFV and NSDV, suggesting that the M segment contains factors involved in viral pathogenesis. The M segment encodes GPC that is first translated as a polyprotein from mRNA, and further cleaved into Gn, Gc, and other accessory or uncharacterized proteins. A schematic diagram of several representative nairovirus GPCs is shown in Fig. 5B. GPC contains an N-terminal signal peptide and multiple membrane-spanning domains, and is processed by signal peptidases to generate an N-terminal pre-Gn protein, C-terminal pre-Gc protein, and a double-membrane-spanning NSm protein. The pre-Gn and pre-Gc are subsequently processed by furin-like or subtilisin kexin isozyme-1 proteases to generate a mucin-like protein containing a large number of O-glycosylation sites, a protein designated as GP38 (-like), virion envelope glycoprotein Gn, and virion envelope glycoprotein Gc²⁷. We showed that although the sizes of Gn and Gc are similar among virus species, those of the O-glycosylation sites and GP38-like protein are different; in particular, they are larger in CCHFV and NSDV (Fig. 5B). This suggests that these regions may be determinants of the pathogenicity of *Nairoviridae*. It has been proposed that GP38 is involved in CCHFV particle formation and viral infectivity²⁷. Analysis of convalescent patient sera showed high titers of CCHFV GP38 antibodies, which indicated the immunogenicity of this protein in humans during natural CCHFV infection. In a mouse model, an antibody against GP38 could protect the animals from a heterologous CCHFV challenge, indicating an association between GP38 and the high pathogenesis of CCHFV²⁸. Our present analysis indicates that there is an association between the N-terminal GPC region and viral pathogenesis not only in CCHFV, but also in other highly pathogenic nairoviruses, including NSDV.

In conclusion, we constructed a comprehensive list of the promoters in *Bunyavirales* that included all virus families in this order. Studies on the RNA synthesis mechanism of *Bunyavirales* have been limited to only a few virus species. Analysis of the conservation in all promoter structures is useful for the prediction of RNA synthesis mechanisms in uncharacterized and newly identified bunyaviruses. The automatic promoter-characterizing system (Supplementary Table 1) is applicable for all bunyaviruses for which the precise genomic end sequences are known.

Methods

List of bunyavirus promoters. In total, 590 bunyavirus species were registered in the ICTV list (<https://talk.ictvonline.org/>) on December 7th, 2021. The complete sequences of the L, M and S segments of bunyaviruses available on the NCBI associated with the GenBank accession numbers listed in Supplementary Table 2 were used for the analysis. After obtaining the full-length genome sequences, the sequences were input in the "Sequence" column in Supplementary Table 1, and the extreme 40 nts of each of the 5' and 3' ends, the complementarity between the 5' and 3' ends of the sequences, and the counts of G:C and A:U complementarity and each of the nts (A, U, G, and C) in the promoter region were calculated automatically. In Supplementary Table 1, the results of the L segments were input as representative data. Conservation of the nts in the promoter was analyzed by using the sequence logo generator WebLogo (<https://weblogo.berkeley.edu/logo.cgi>).

Analysis of the genome length of nairoviruses. Sequences annotated as "*Nairoviridae*" were downloaded from the NCBI refseq database on January 16th, 2022. There were 5272 *Nairoviridae* sequences in the database. We first checked for the presence of the extreme promoter sequence 5'-UCUCA in the 8-nt ends of the sequences. The promoter sequence was present in both ends of 368 *Nairoviridae* sequences (Supplementary

Table 3), and the lengths of these sequences were calculated using a custom Python script. The codes used for this analysis are available on GitHub (https://github.com/shohei-kojima/Arenaviridae_overhang_analysis_2022).

Amino acid sequence map of theairovirus glycoprotein. The structural characteristics of theairovirus glycoprotein were predicted using TMHMM-2.0 for the transmembrane protein²⁹, SignalP-6.0 for the signal cleavage site³⁰, and NetOGlyc-4.0 for the O-linked glycosylation sites³¹. Data on the glycoprotein sequences were collected from UniProt (<https://www.ebi.ac.uk/uniprot/index>). The UniProt accession numbers were: CCHFV, Q8JSZ3; NSDV, A0A0A7H811; Dugbe virus, Q02004; Tofla virus, A0A0U5AG15; HAZV, A6XIP3; and Erve virus, J3S7E1. GP38-like regions were found using the Protein Basic Local Alignment Search Tool (BLASTp) based on the amino acid sequences of the CCHFV and Dugbe virus GPC.

Statistical analysis. Statistical analyses were performed with Prism software (version 9.1.2; GraphPad, San Diego, CA, USA). Statistical significance was assigned when p values were < 0.05. Inferential statistical analysis was performed by a two-tailed unpaired Student's *t*-test or one-way analysis of variance followed by Tukey's test, as appropriate.

Data availability

All databases used in this study are available from DDBJ/ENA/GenBank (<https://www.ddbj.nig.ac.jp/about/insdc-e.html>). The accession numbers of viral sequences used in this study are listed in Supplementary Tables 1, 2 and 3.

Received: 17 May 2022; Accepted: 30 July 2022

Published online: 09 August 2022

References

- Barr, J. N. & Wertz, G. W. Bunyamwera bunyavirus RNA synthesis requires cooperation of 3'- and 5'-terminal sequences. *J. Virol.* **78**, 1129–1138 (2004).
- Kohl, A., Dunn, E. F., Lowen, A. C. & Elliott, R. M. Complementarity, sequence and structural elements within the 3' and 5' non-coding regions of the Bunyamwera orthobunyavirus S segment determine promoter strength. *J. Gen. Virol.* **85**, 3269–3278 (2004).
- Gerlach, P., Malet, H., Cusack, S. & Reguera, J. Structural insights into bunyavirus replication and its regulation by the vRNA promoter. *Cell* **161**, 1267–1279 (2015).
- Arragain, B. *et al.* Structural snapshots of La Crosse virus polymerase reveal the mechanisms underlying *Peribunyaviridae* replication and transcription. *Nat. Commun.* **13**, 902. <https://doi.org/10.1038/s41467-022-28428-z> (2022).
- Barr, J. N., Rodgers, J. W. & Wertz, G. W. The Bunyamwera virus mRNA transcription signal resides within both the 3' and the 5' terminal regions and allows ambisense transcription from a model RNA segment. *J. Virol.* **79**, 12602–12607 (2005).
- Ariza, A. *et al.* Nucleocapsid protein structures from orthobunyaviruses reveal insight into ribonucleoprotein architecture and RNA polymerization. *Nucleic Acids Res.* **41**, 5912–5926 (2013).
- Reguera, J., Malet, H., Weber, F. & Cusack, S. Structural basis for encapsidation of genomic RNA by La Crosse Orthobunyavirus nucleoprotein. *Proc. Natl. Acad. Sci.* **110**, 7246–7251 (2013).
- Matsumoto, Y., Ohta, K., Kolakofsky, D. & Nishio, M. A minigenome study of Hazaraairovirus genomic promoters. *J. Virol.* **93**, e02118–e02118 (2019).
- Crooks, G. E., Hon, G., Chandonia, J. M. & Brenner, S. E. WebLogo: A sequence logo generator. *Genome Res.* **14**, 1188–1190 (2004).
- Ren, F., Zhou, M., Deng, F., Wang, H. & Ning, Y. J. Combinatorial minigenome systems for emerging Banyangviruses reveal viral reassortment potential and importance of a protruding nucleotide in genome “panhandle” for promoter activity and reassortment. *Front. Microbiol.* **11**, 599. <https://doi.org/10.3389/fmicb.2020.00599> (2020).
- Weidmann, M. *et al.* Biosafety standards for working with Crimean-Congo hemorrhagic fever virus. *J. Gen. Virol.* **97**, 2799–2808 (2016).
- Dai, S., Deng, F., Wang, H. & Ning, Y. Crimean-Congo hemorrhagic fever virus: Current advances and future prospects of antiviral strategies. *Viruses* **13**, 1195. <https://doi.org/10.3390/v13071195> (2021).
- Krasteva, S., Jara, M., Frias-De-Diego, A. & Machado, G. Nairobi sheep disease virus: A historical and epidemiological perspective. *Front. Vet. Sci.* **7**, 419. <https://doi.org/10.3389/fvets.2020.00419> (2020).
- Banerjee, A. K. 5'-terminal cap structure in eucaryotic messenger ribonucleic acids. *Microbiol. Rev.* **44**, 175–205 (1980).
- Habjan, M. *et al.* Processing of genome 5' termini as a strategy of negative-strand RNA viruses to avoid RIG-I-dependent interferon induction. *PLoS ONE* **3**, e2032. <https://doi.org/10.1371/journal.pone.0002032> (2008).
- Obijeski, J. F., Bishop, D. H., Palmer, E. L. & Murphy, F. A. Segmented genome and nucleocapsid of La Crosse virus. *J. Virol.* **20**, 664–675 (1976).
- Garcin, D. & Kolakofsky, D. Tacaribe arenavirus RNA synthesis in vitro is primer dependent and suggests an unusual model for the initiation of genome replication. *J. Virol.* **66**, 1370–1376 (1992).
- Deflubé, L. R. *et al.* Ebolavirus polymerase uses an unconventional genome replication mechanism. *Proc. Natl. Acad. Sci.* **116**, 8535–8543 (2019).
- Shareef, A. M., Ludeke, B., Jordan, P., Deval, J. & Fearn, R. Comparison of RNA synthesis initiation properties of non-segmented negative strand RNA virus polymerases. *PLOS Pathog.* **17**, e1010151. <https://doi.org/10.1371/journal.ppat.1010151> (2021).
- Garcin, D. *et al.* The 5' ends of Hantaan virus (*Bunyaviridae*) RNAs suggest a prime-and-realign mechanism for the initiation of RNA synthesis. *J. Virol.* **69**, 5754–5762 (1995).
- Patterson, J. L., Holloway, B. & Kolakofsky, D. L. Crosse virions contain a primer-stimulated RNA polymerase and a methylated cap-dependent endonuclease. *J. Virol.* **52**, 215–222 (1984).
- Jeeva, S., Cheng, E., Ganaie, S. S. & Mir, M. A. Crimean-Congo hemorrhagic fever virus nucleocapsid protein augments mRNA translation. *J. Virol.* **91**, e00636–e00636 (2017).
- Jin, H. & Elliott, R. M. Non-viral sequences at the 5' ends of Dugbeairovirus S mRNAs. *J. Gen. Virol.* **74**, 2293–2297 (1993).
- Ferron, F., Weber, F., de la Torre, J. C. & Reguera, J. Transcription and replication mechanisms of *Bunyaviridae* and *Arenaviridae* L proteins. *Virus Res.* **234**, 118–134 (2017).
- Barnwal, B., Karlberg, H., Mirazimi, A. & Tan, Y. J. The non-structural protein of Crimean-Congo hemorrhagic fever virus disrupts the mitochondrial membrane potential and induces apoptosis. *J. Biol. Chem.* **291**, 582–592 (2016).
- Mega, D. F. *et al.* Mutagenic analysis of Hazaraairovirus nontranslated regions during single- and multistep growth identifies both attenuating and functionally critical sequences for virus replication. *J. Virol.* **94**, e00357–e00357 (2020).

27. Freitas, N. *et al.* The interplays between Crimean-Congo hemorrhagic fever virus (CCHFV) M segment-encoded accessory proteins and structural proteins promote virus assembly and infectivity. *PLoS Pathog.* **16**, e1008850. <https://doi.org/10.1371/journal.ppat.1008850> (2020).
28. Mishra, A. K. *et al.* Structure and characterization of Crimean-Congo hemorrhagic fever virus GP38. *J. Virol.* **94**, e02005-e2019 (2020).
29. Krogh, A., Larsson, B., von Heijne, G. & Sonnhammer, E. L. L. Predicting transmembrane protein topology with a hidden markov model: Application to complete genomes¹ Edited by F. Cohen. *J. Mol. Biol.* **305**, 567–580 (2001).
30. Teufel, F. *et al.* SignalP 6.0 predicts all five types of signal peptides using protein language models. *Nat. Biotechnol.* **40**, 1023–1025 (2022).
31. Steintoft, C. *et al.* Precision mapping of the human O-GalNAc glycoproteome through simplecell technology. *EMBO J.* **32**, 1478–1488 (2013).

Author contributions

Y.N., S.K., A.S., M.K., T.I., T.W., Y.A., H.S., K.N., T.K. and Y.M. conceived and designed the study, performed the analyses, and analyzed the data. Y.N. generated the automatic promotor calculator (Supplementary Table 1). S.K. performed the genome length analysis of *Nairoviridae*. A.S. designed the schematic diagram of the nairovirus GPC. Y.M. wrote the manuscript. All authors have read and agreed to the content of the manuscript.

Funding

This work was supported by Takeda Science Foundation, and Tokyo Biochemical Foundation, Japan (to Y.M.).

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-022-17758-z>.

Correspondence and requests for materials should be addressed to Y.M.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022