



MEETING ABSTRACT

Open Access

# PW01-024 – Phenotypic analysis of a *MEFV* negative FMF cohort

A Soriano<sup>1</sup>, D Rigante<sup>2</sup>, L Cerrito<sup>1</sup>, C Fonnesu<sup>1</sup>, L Sicignano<sup>1</sup>, A Gallegos<sup>3</sup>, R Manna<sup>1\*</sup>

From 7th Congress of International Society of Systemic Auto-Inflammatory Diseases (ISSAID) Lausanne, Switzerland. 22-26 May 2013

## Introduction

Familial Mediterranean fever (FMF) is an inherited autosomal recessive disorder, ethnically restricted and commonly found among individuals of Mediterranean descent, caused by Mediterranean Fever gene (*MEFV*) mutations on the chromosome 16. It is the most frequent periodic febrile syndrome among autoinflammatory syndromes. Eighty % of patients with FMF have *MEFV* mutations, while around 20% do not have mutations.

## Objectives

We analysed epidemiological and clinical characteristics, as well as treatment schedules of a large cohort of FMF patients without any *MEFV* mutations, who responded to colchicine, in order to identify further clinical features of this specific subgroup.

## Methods

Epidemiological and clinical details of 344 patients attending the Periodic Fevers Research Centre in a period of 15 years were analysed. We selected patients without *MEFV* mutations, in whom diagnosis was established by the Tel-Hashomer criteria. We finally compared the clinical findings of *MEFV*-negative population with the *MEFV*-positive one.

## Results

Genetic testing by *MEFV* analysis was performed in all patients (n = 344); 41 patients (14%, 20 males and 21 females) negative for *MEFV* mutations were selected and studied. Similarly with *MEFV* positive patients, in our case-series, most *MEFV*-negative ones came from Southern and Central Italy. The mean age of FMF onset was 21.8 years, differently from what observed in

*MEFV*-positive population, in which the mean age was 15. The frequency of attacks went from less than 1 attack/month (in 26%) to 1-2 attacks/month (in 54%) and more than 2 attacks/months (in 19.5%). The mean duration of each attack was  $83.9 \pm 8.91$  hours. The typical clinical signs of FMF attacks were: fever (T max  $39.4^{\circ}\text{C} \pm 0.12$ , present in 100% of patients), articular pain (76%), abdominal pain (63.4%), oral aphthosis (44%), and chest pain (37%). Thirty-one out of 41 patients had joint involvement in terms of arthritis (21.5%), arthralgias (25%), arthromyalgias (32%), and myalgias (21.5%). Attacks were controlled with a mean dose of colchicine of 1.5 mg/day in all patients (vs a mean dose of 1.3 mg/die in the *MEFV*-positive population). No statistically significant difference was detected in terms of frequency and duration of attacks, as well as in symptoms distribution and colchicine dosage between *MEFV*-negative and positive populations.

## Conclusion

Analysis of our *MEFV*-negative series of Italian patients revealed a higher prevalence of late-onset FMF, whereas the percentage distribution of symptoms was similar to *MEFV*-positive patients. These results support the hypothesis of involvement of other low-penetrance genetic systems in the FMF clinical expression.

## Disclosure of interest

None declared.

## Authors' details

<sup>1</sup>Periodic Fevers Research Centre, Catholic University of Sacred Heart, Rome, Italy. <sup>2</sup>Department of Pediatrics, Catholic University of Sacred Heart, Rome, Italy. <sup>3</sup>Nuestra Señora del Prado Hospital, Talavera de la Reina, Spain.

<sup>1</sup>Periodic Fevers Research Centre, Catholic University of Sacred Heart, Rome, Italy

Full list of author information is available at the end of the article

Published: 8 November 2013

#### References

1. Soriano A, Manna R: **Familial Mediterranean fever: new phenotypes.** *Autoimmun Rev* 2012, **12**:31-7.
2. Rigante D: **The fresco of autoinflammatory diseases from the pediatric perspective.** *Autoimmun Rev* 2012, **11**:348-56.
3. Manna R, Cerquaglia C, Curigliano V, Fonnesu C, Giovinale M, Verrecchia E, Montalto M, De Socio G, Soriano A, La Regina M, Gasbarrini G: **Clinical features of familial Mediterranean fever: an Italian overview.** *Eur Rev Med Pharmacol Sci* 2009, **13**(Suppl 1):51-3.

doi:10.1186/1546-0096-11-S1-A77

**Cite this article as:** Soriano *et al.*: PW01-024 – Phenotypic analysis of a MEFV negative FMF cohort. *Pediatric Rheumatology* 2013 **11**(Suppl 1):A77.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

