

RESEARCH

Open Access

Involvement of NT3 and P75^{NTR} in photoreceptor degeneration following selective Müller cell ablation

Weiyong Shen*, Ling Zhu, So-Ra Lee, Sook H Chung and Mark C Gillies

Abstract

Background: Neurotrophins can regulate opposing functions that result in cell survival or apoptosis, depending on which form of the protein is secreted and which receptor and signaling pathway is activated. We have recently developed a transgenic model in which inducible and patchy Müller cell ablation leads to photoreceptor degeneration. This study aimed to examine the roles of mature neurotrophin-3 (NT3), pro-NT3 and p75 neurotrophin receptor (P75^{NTR}) in photoreceptor degeneration in this model.

Methods: Transgenic mice received tamoxifen to induce Müller cell ablation. Changes in the status of Müller and microglia cells as well as expression of mature NT3, pro-NT3 and P75^{NTR} were examined by immunohistochemistry and Western blot analysis. Recombinant mature NT3 and an antibody neutralizing P75^{NTR} were injected intravitreally 3 and 6 days after Müller cell ablation to examine their effects on photoreceptor degeneration and microglial activation.

Results: We found that patchy loss of Müller cells was associated with activation of surviving Müller cells and microglial cells, concurrently with reduced expression of mature NT3 and upregulation of pro-NT3 and P75^{NTR}. Intravitreal injection of mature NT3 and a neutralizing antibody to P75^{NTR}, either alone or in combination, attenuated photoreceptor degeneration and the beneficial effect was associated with inhibition of microglial activation.

Conclusions: Our data suggest that Müller cell ablation alters the balance between the protective and deleterious effects of mature NT3 and pro-NT3. Modulation of the neuroprotective action of mature NT3 and pro-apoptotic pro-NT3/P75^{NTR} signaling may represent a novel pharmacological strategy for photoreceptor protection in retinal disease.

Keywords: Müller cell, Microglia, Neurotrophins, p75^{NTR}, NT3, Retinal degeneration

Background

Neurotrophins are a family of neurotrophic factors including at least four neurotrophins: nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), and neurotrophins (NT)-3 and-4 [1]. They are synthesized as precursors that can be either cleaved intracellularly or secreted uncleaved as pro-neurotrophins. Neurotrophins bind to two distinct types of receptors, the Trk receptor tyrosine kinase family and the p75 neurotrophin receptor (p75^{NTR}) [1,2]. In the retina, NGF preferentially binds TrkA, which is almost exclusively expressed by

ganglion cells. BDNF and NT4 preferentially bind TrkB, which is widely expressed by ganglion, horizontal, dopaminergic amacrine and Müller cells. NT3 preferentially binds TrkC that is expressed by photoreceptors and Müller cells. Trk receptors preferentially respond to mature neurotrophins promoting neuronal survival, differentiation and synaptic function [1,2]. The expression of p75^{NTR} is almost exclusively confined to Müller cells in the retina [2-4]. In contrast to mature neurotrophins, pro-neurotrophins exert their apoptotic effect *via* a receptor complex containing p75^{NTR} and sortilin [2,5]. Thus, neurotrophins can regulate opposing cellular functions that result in cell survival or apoptosis, depending

* Correspondence: weiyong.shen@sydney.edu.au

Save Sight Institute, the University of Sydney, 8 Macquarie Street, Sydney 2000, Australia

on which form of the protein is secreted and which receptor and signaling pathway is activated.

Many different types of insult can potently induce pro-neurotrophins and p75^{NTR}. Accumulation of pro-NGF and upregulation of p75^{NTR} have been found to be positively correlated with accelerated retinal neurodegeneration in diabetes [6-8]. Upregulation of p75^{NTR} has been observed during light-induced photoreceptor degeneration [2], ocular hypertension [9], ischemic injury [10] and optic nerve axotomy [3,11]. Genetic ablation of p75^{NTR} or biochemical blockage of p75^{NTR} activation attenuates neuronal death induced by pro-neurotrophins [2,5]. Binding of pro-NGF to p75^{NTR} has been reported to induce robust expression of neurotoxic factors, suggesting that ligand activation of p75^{NTR} in Müller cells may activate neurotoxic pathways through a paracrine mechanism that negates the protective effect of mature neurotrophins [3,12]. Notably, previous studies indicate that NGF and BDNF can be secreted as pro-forms in the retina under pathological conditions [2,4,13,14]. However, the involvement of pathological pro-NT3/P75^{NTR} signaling in photoreceptor degeneration remains to be elucidated.

Progressive dysfunction and death of photoreceptors is the major cause of loss of vision in most retinal diseases. There is increasing evidence that Müller cells are important for photoreceptor health [15,16]. We recently generated a transgenic model using a portion of the regulatory region of the retinaldehyde binding protein 1 (Rlbp1) gene as a cell-specific promoter along with a CreER/Lox-P approach for inducible Müller cell-specific gene targeting [17]. These Rlbp1-CreER transgenic mice were crossed with Rosa-DTA176 mice, a transgenic line carrying an attenuated form of the diphtheria toxin fragment A (DTA176) gene, for Müller cell ablation following tamoxifen induction [17]. Selective Müller cell ablation in adult mice led to photoreceptor degeneration, blood-retinal barrier breakdown and deep retinal neovascularisation [17]. These changes are common, critical features of many retinal diseases such as macular telangiectasia [18-20], age-related macular degeneration [21,22], diabetic retinopathy [23,24] and ischemic retinopathy [25]. In this study, we have utilized this unique transgenic model to examine the roles of abnormal expression of mature NT3, pro-NT3 and P75^{NTR} in the photoreceptor degeneration after selective Müller cell ablation.

Methods

Conditional Müller cell ablation in transgenic mice

Animal studies were performed in accordance with the Association for Research in Vision and Ophthalmology statement and were approved by The University of Sydney Animal Ethics Committee. Rlbp1-CreER mice were crossed with Rosa-DTA176 mice to produce Rlbp-CreER-

DTA176 transgenic mice, which were used for conditional, selective Müller cell ablation as we have previously described [17]. Animals were screened by PCR to identify those carrying both Rlbp1 and DTA176 genes. Selective Müller cell ablation in transgenic mice was induced by daily intraperitoneal injection of tamoxifen (TMX, 3 mg in 0.2 ml sunflower oil) for 4 consecutive days at 6–8 weeks of age [17]. Mice not carrying the Rlbp1 Müller cell-specific promoter but carrying the DTA176 gene were used as controls in this study.

Cryosection and flat-mount immunohistochemistry (IHC)

Eyes were briefly fixed in 4% paraformaldehyde for 5 min, and then anterior segments were removed. After post-fixation in 4% paraformaldehyde for 1 h, eye cups were either transferred to PBS containing 30% sucrose and then embedded in optimal cutting temperature compound for cryosection IHC or placed in PBS for retinal flat-mount IHC. For cryosection IHC, frozen sections were blocked with 5% normal goat serum and incubated with an antibody (Ab) to glutamine synthetase (GS, mouse monoclonal, 1:100; Millipore no. MAB302), glial fibrillary acidic protein (GFAP, rabbit polyclonal, 1:250; Dako no. Z0334), P75^{NTR} (rabbit polyclonal, 1:250; a gift from Dr. Moses V. Chao, New York University, School of Medicine; no. 9651) and ionized calcium binding adaptor molecule 1 (Iba-1, rabbit polyclonal, 1:500, Wako no. 019–19741). Bound antibodies were detected with Alexa Fluor 488 or 594-conjugated goat or donkey secondary antibodies (1:1,000; Invitrogen).

For flat-mount staining, dissected eye cups were fixed in 4% paraformaldehyde for 1 h and then placed in PBS at +4°C overnight. On the next day, retinas were isolated, rinsed in PBS and permeabilized with 1% Triton-X-100 containing 5% normal goat serum blocking solution for 2 h. Retinas were incubated in 100 µl of solution containing peanut-agglutinin (PNA) conjugated with Alexa Fluor 488 or 594 (10 µg/ml, Invitrogen, no. L-21409 and L-32459) to label cone photoreceptor outer segments and an Ab against Iba1 (1:500, Wako no. 019–19741) for retinal microglia in 0.1 M PBS with 1% BSA and 0.5% Triton X-100 overnight at +4°C. Retinal whole-mounts were counterstained with Hoechst for 5 min before mounting onto slides for confocal laser scanning microscopy. Images were processed and analyzed using computer-based image analysis software to determine the percentage of PNA- or Iba-1 stained area per field of view as described previously [17,26]. In brief, a gradient detection algorithm was applied to the original digital image and binary thresholding performed on the gradient image by selecting its mean gray value as the threshold. This procedure allowed sufficient identification of the subject profiles to calculate the percentages of PNA- or Iba-1-stained area per field of view.

Intravitreal injection of mature NT3 and an antibody against P75^{NTR}

Intravitreal injection was performed using a 32-gauge needle attached to a Hamilton syringe as we have previously described [17]. Intravitreal injections were performed in transgenic mice 3 and 6 days after TMX-induced Müller cell ablation, with one eye receiving 2 μ l of testing reagent and the other eye receiving 2 μ l of BSS as a control in each mouse. Doses injected were: (1) NT3, 0.4 μ g (R&D Systems, catalog no. 267N3/CF); (2) P75^{NTR} rabbit polyclonal Ab (no. 9651, 1:1 dilution, a gift from Dr. Moses V. Chao); (3) 0.4 μ g NT3 + 1:1 dilution of P75 Ab. Injected eyes were enucleated 10 days after TMX-induced Müller cell ablation for retinal whole-mount staining to examine changes in cone photoreceptor outer segments and microglial activation as described above.

Western blot

For Western blot, proteins were extracted from retinas and their concentrations determined by DC protein assay. Equal amounts of protein were subjected to SDS-polyacrylamide gel electrophoresis then transferred to a PVDF membrane for Western blot. Membranes were probed with antibodies to GS (mouse monoclonal, 1:1,000; Millipore, no. MAB302), GFAP (mouse monoclonal, 1:5,000; Neomarker, no. MS-280-P), P75^{NTR} (rabbit polyclonal, a gift from Dr. Moses V. Chao, no. 9651), NT3 (rabbit polyclonal, 1:1,000, Alomone Laboratory, no. ANT-003), pro-NT3 (rabbit polyclonal, 1:500, Alomone Laboratory, no. ANT-012) and rhodopsin (mouse monoclonal, 1:500, Millipore no. MAB5356), guanine nucleotide-binding protein subunit alpha-1 (GNAT1, rabbit polyclonal, 1:500, Santa Cruz no. sc-389) and G α protein transducin (G α t, mouse monoclonal, 1:2,000, BD Transduction Laboratories no. 610589). After incubation with secondary antibodies conjugated with horseradish peroxidase, protein bands were visualized using the G:Box BioImaging systems and quantified using the GeneTools image scanning and analysis package. Protein expression was normalized to α -/ β -tubulin (rabbit polyclonal, 1:2,000; Cell Signaling no. 2148), which serves as a loading control.

Statistics

Results are expressed as mean \pm SEM. Data were analyzed using paired or un-paired *t*-test with a *p* value <0.05 accepted as statistically significant.

Results

Patchy loss of Müller cells was accompanied by activation of surviving Müller cells

We have previously shown patchy loss of Müller cells in Rlbp1-CreER-DTA176 transgenic mice, which can be observed as early as 1 day and becomes stable 14 days after induction with TMX [17]. We performed double-


label IHC for GS and GFAP to examine the status of surviving Müller cells after selective Müller cell ablation (Figure 1A-D). In the control retina, strong GS immunoreactivity was observed across the neuroretina from the inner limiting membrane to the outer limiting membrane, with cell bodies localized to the inner nuclear layer, while GFAP expression was only observed in cells around the inner limiting membrane and in the outer plexiform layer (Figure 1A). In transgenic mice, reactive activation of surviving Müller cells was observed as early as 1 day after TMX treatment (Figure 1B), became more profound from 7 days (Figure 1C) and lasted for at least 3 months (Figure 1D) after TMX-induced Müller cell ablation. Western blot analysis demonstrated significant reduction in GS expression and upregulation of GFAP 7d after Müller cell ablation (Figure 1E and F). Thus, patchy loss of Müller cells induces reactive activation of surviving Müller cells in this model.

Upregulation of P75^{NTR} in transgenic retinas after selective Müller cell ablation

As P75^{NTR} is predominantly expressed in Müller cells in the retina [2-4], we next examined whether these activated Müller cells overexpress P75^{NTR}. Müller cells in the control retina expressed P75^{NTR} weakly (Figure 2A-C). By contrast, patchy loss of Müller cells was accompanied by strong immunostaining for P75^{NTR} in surviving Müller cells 7 and 14 days after Müller cell ablation (Figure 2D-I). Western blot results showed significant upregulation of P75^{NTR} and GFAP in transgenic mice 7 and 14 days after TMX treatment (Figure 2J and K). These results indicate that activated Müller cells overexpress P75^{NTR}.

Differential expression of mature NT3, pro-NT3, rod and cone phototransduction proteins

We next examined changes in mature NT3, pro-NT3, rod and cone phototransduction proteins after Müller cell ablation (Figure 3). Western blots were performed on retinal lysates using antibodies to detect mature NT3, pro-NT3, rhodopsin, guanine nucleotide-binding protein subunit alpha-1 (GNAT1) and G α protein transducin (G α t) 7 days after Müller cell ablation (Figure 3A and F). GNAT1 and G α t are proteins essential for rod and cone phototransduction [27-29]. Quantitative analysis of protein densitometry revealed significant reduction in mature NT3 and upregulation of pro-NT3, which resulted in a decreased ratio of NT3:pro-NT3 (Figure 3A-D). The differential expression of mature NT3 and pro-NT3 was accompanied by significant reduction in rhodopsin (Figure 3E), GNAT1 (Figure 3G) and G α t (Figure 3H) in TG mice, indicating that selective Müller cell ablation causes damage to both rod and cone photoreceptors.




Activation of retinal microglial cells after selective Müller cell ablation

Microglia and Müller cells are prominent participants in retinal responses to injury and diseases. There is evidence that the interaction between activated microglia and Müller cells can initiate a program of bidirectional microglia-Müller cell signaling that augments initial inflammatory responses to retinal injury [30]. We performed cryosection IHC using an Ab against Iba1 to examine microglial activation after Müller cell ablation (Figure 4). In the control retina, resting microglia cells were observed in the ganglion cell layer, inner plexiform layer and, occasionally, in the outer plexiform layer (Figure 4A). These resting microglia cells showed small somas with thin and ramified cells processes. They were

predominantly confined to the inner retina but were found in neither the outer nuclear layer nor in the sub-retinal space (Figure 4A). Activation of microglia cells was observed in transgenic mice as early as 1 day after TMX-induced Müller cell ablation, as evidenced by expansion of their soma size and thickening of their cell processes (Figure 4B-F, arrows). Activated microglia cells were observed in the subretinal space from 1 day and frequently observed from 7 days after induced Müller cell ablation (Figure 4B-F, arrowheads).


We further conducted double-label immunostaining on retinal whole mounts to map microglial activation with cone photoreceptor damage. Iba1 Ab was used to identify microglial cells, and lectin PNA was used to label cone photoreceptor outer segments (Figure 5). The



control retina showed dense staining for PNA but lack of microglial infiltration at the level of photoreceptor outer segments (Figure 5A-C). In TG mice, most microglia retained slender cell processes, and PNA-staining showed limited loss of cone photoreceptor outer segments with only a few photoreceptor nuclei protruding into the subretinal space 7 days after induced Müller cell ablation (Figure 5D-F). Pronounced activation of microglial cells was evidenced by expansion of their soma size, which was clearly observed 11 days after Müller cell ablation (Figure 5G-I). Most microglia 11 days after Müller cell ablation (Figure 5H) had larger soma than at day 7 (Figure 5E), which is reflected by the increased area of microglial staining at day 11 compared with day 7 (Figure 5K) when the number of microglia seemed similar at both time points (Figure 5E and H). This was associated with more loss of photoreceptor outer segments at day 11 than at day 7 (Figure 5J).

Photoreceptor protection after intravitreal injection of mature NT3 and P75^{NTR} Ab

Since selective Müller cell ablation resulted in reduced expression of mature NT3 and upregulation of pro-NT3 and P75^{NTR}, we reasoned that intravitreal supplementation of exogenous mature NT3 and receptor blocking of P75^{NTR} might protect photoreceptors. We tested this hypothesis by performing intravitreal injections of recombinant mature NT3 and a neutralizing Ab to P75^{NTR} 3 and 6 days after TMX-induced Müller cell ablation. Changes in cone photoreceptor outer segments were analyzed 4 days after the last intravitreal injection (Figure 6). In control mice, intravitreal injection of NT3 in combination with P75^{NTR} Ab neither induced photoreceptor outer segment loss nor caused microglial infiltration in the subretinal space (Figure 6A and B). In TG mice, however, treatments with NT3 and P75^{NTR} Ab, either alone or in combination, reduced cone photoreceptor outer



segment loss and photoreceptor nuclei protrusion when compared with eyes receiving BSS (Figure 6C-F). Quantitative analysis of changes in PNA-stained cone photoreceptor outer segments confirmed the beneficial effect of NT3 and P75^{NTR} Ab treatments on photoreceptor protection (Figure 6G).

Inhibition of microglia after intravitreal injection of mature NT3 and P75^{NTR} Ab

We next examined the extent of microglial activation after intravitreal injection of mature NT3 and the P75^{NTR} neutralizing Ab. Consistent with our earlier findings that severe photoreceptor damage was accompanied by massive activation of microglia cells in this transgenic model (Figures 4 and 5), marked activation of microglia cells and loss of cone photoreceptor outer segments were

observed in eyes receiving BSS (Figure 7A-C). In contrast, inhibition of microglial activation was observed in eyes receiving NT3, P75 Ab or both (Figure 7D, G and J). The extent of inhibition of microglial activation corresponded well with the extent of photoreceptor protection in eyes receiving of NT3, P75 Ab or the combined treatment (Figure 7E, I and L). Quantitative analysis of Iba-1 staining retinal whole mounts showed that intravitreal injection of NT3 and P75^{NTR} Ab, either alone or in combination, significantly inhibited microglial activation (Figure 7M).

Discussion

We have demonstrated here that patchy loss of Müller cells induced photoreceptor degeneration accompanied by activation of surviving Müller cells and microglia in this model. Photoreceptor degeneration was concomitant with




Figure 4 Microglial activation after selective Müller cell ablation. Immunohistochemistry was performed using an Ab against ionized calcium-binding adaptor molecule 1 (*Iba-1*, green) for microglia and nuclei were counterstained with Hoechst (red) to reveal retinal structure. **(A)** Control (Ct) mice showing “resting” microglia cells in the ganglion cell layer (GCL), inner plexiform layer (IPL) and occasionally the outer plexiform layer (OPL), but not in the outer nuclear layer (ONL) or the subretinal space (SP). **(B-F)** Activation of microglia cells was observed in transgenic (TG) mice as early as 1 day (1d, **B**) after TMX-induced Müller cell ablation, as reflected by increased soma size and thickened cell processes in the inner retina (**B-F**, arrows). Microglia cells were observed in the subretinal space from 1d and frequently seen from 7d after TMX treatment (**B-F**, arrowheads). Note: protrusion of photoreceptor nuclei into the subretinal space in (**D-F**) and obvious thinning ONL in (**F**) 3 months (3 m) after TMX-induced Müller cell ablation. Scale bars: **A-F**, 50 μ m.

reduced expression of mature NT3 and upregulation of pro-NT3 and P75^{NTR}. We then demonstrated that intravitreal injection of mature NT3 and a neutralizing Ab to P75^{NTR}, either alone or in combination, attenuated photoreceptor degeneration concurrently with inhibition of microglial activation. Our results suggest that aberrant expression of NT3, pro-NT3 and P75^{NTR} is associated with photoreceptor degeneration and that treatment with mature NT3 together with inhibition of pro-NT3/P75^{NTR} signaling may represent a novel pharmacological strategy for protecting photoreceptors in retinal disease.

The primary role of Müller cells in the healthy retina is to maintain retinal homeostasis. The importance of Müller cells for photoreceptor health is indicated by the photoreceptor degeneration that developed after selective Müller cell ablation in our transgenic model [17]. We demonstrated that while TMX induction of DTA expression in Müller cells induced patches of Müller cell loss, surviving Müller cells were reactively activated. Reactive activation of Müller cells has been observed in retinal damage caused by every conceivable insult, including ischemia, trauma, degeneration, inflammation or neuronal excitotoxicity [31-35]. Müller cell activation is believed to be important for protection or repair of neurons in the early stages of neuronal damage [32-35]. However, activated Müller cells may overreact to the

primary events and mediate secondary injury through generation of neurotoxic substances such as inflammatory mediators, free radicals, lipid peroxidation, increased levels of extracellular calcium and release of excitatory amino acids [3,12,15,36,37].

The role of p75^{NTR} as an apoptotic receptor is implicated in promoting apoptosis of neurons in the central nervous system following various injuries [2,38-40]. The involvement of p75^{NTR} in photoreceptor cell death has been previously reported in various animal models of retinal degenerations [2,4,41,42]. In the present study, we have particularly examined changes in P75^{NTR} expression 7 and 14 days after TMX treatment because these are the times at which the rate of photoreceptor apoptosis peaks in this model [17]. Double-label IHC showed strong immunoreactivity for P75^{NTR} in surviving Müller cells, and Western blot analysis showed significant upregulation of P75^{NTR} and GFAP at these two time points. These findings suggest that activated Müller cells overexpress P75^{NTR}. There is evidence that activation of the pro-NGF/P75^{NTR} signaling contributes to retinal neuronal injury through a paracrine mechanism in the retina [3,12]. It has been reported that activation of the pro-NGF/P75^{NTR} signaling induced robust expression of neurotoxic substances such as tumor necrosis factor alpha and α 2 macroglobulin [3,12]. Whether




Figure 5 Mapping cone photoreceptor outer segment loss with infiltration of activated microglial cells after Müller cell ablation. (A-I) Double labeling using peanut-agglutinin (PNA)-conjugated with Alexa Fluor-594 for cone photoreceptor outer segments (A, D and G, red) and an Ab against ionized calcium binding adaptor molecule 1 (IBA-1) for microglia (B, E and H, green) on retinal whole mounts. (A-C) Images from a control (Ctl) retina. (D-I) Images from transgenic (TG) mice 7 and 11 days after tamoxifen-induced Müller cell ablation. (C, F and I) Merged images. Arrows in (F and I) point to areas of photoreceptor degeneration where degenerate cell bodies with nuclei stained blue protrude into the subretinal space. (J and K) Quantitative analysis of changes in PNA-labeled cone photoreceptor outer segments (J) and infiltration of activated microglial cells (K) 7 and 11 days after Müller cell ablation. * $P < 0.01$, TG vs. Ctl and $^{\dagger}P < 0.01$, 11d vs. 7d; unpaired t -test; $n = 12-14$ at each time point. Scale bars in A-F: 100 μ m.

this also occurs in our transgenic model warrants further investigation.

Photoreceptor cell injury in animal models of retinal degenerations is often accompanied by microglial activation.

We found activated microglial cells in the subretinal space as early as 1 and 3 days after TMX-induced Müller cell ablation. Marked activation of microglial cells was observed as photoreceptors were degenerating, and our





Figure 6 Intravitreal injection of recombinant NT3 and blockage of P75^{NTR} protected photoreceptor degeneration.

Intravitreal injection was performed in transgenic mice 3d and 6d after tamoxifen induced Müller cell ablation, with one eye receiving the testing reagent and the contralateral eye receiving balanced salt solution (BSS). Doses of injection: (1) NT3, 0.4 µg; (2) P75^{NTR} rabbit polyclonal Ab (1:1 dilution) and (3) 0.4 µg NT3 + 1:1 dilution of P75^{NTR} Ab. Eyes were enucleated 10d after TMX treatment for flat-mount staining using fluorescence-conjugated peanut-agglutinin (PNA).

(A and B) PNA-labeled cone photoreceptor outer segments (green) and nuclear counterstaining in a control (Ctl) retina. (C-F) Images from transgenic (TG) mice showing changes in cone photoreceptor outer segments and protrusion of photoreceptor nuclei into the subretinal space (red, indicated by asterisks) in eyes receiving BSS (C), NT3 (D), P75^{NTR} Ab (E) and a combination of NT3 and P75^{NTR} Ab (F).

(G) Quantitative analysis of PNA staining showed that intravitreal injection of NT3 and P75 Ab protected the loss of cone photoreceptor outer segments, with a combined treatment more effective than either alone. N = 9 – 11/group. Scale bars in A-F: 100 µm.

further analysis revealed a close association between microglial infiltration in the subretinal space and loss of photoreceptor outer segments. This close spatial relationship between the activated microglial cells and photoreceptor damage in this study suggests that microglial cells might be involved in photoreceptor degeneration. Microglial cells are the resident tissue macrophages and primary immune cells of the central nervous system and retina. Under basal conditions, “resting” microglia demonstrate ramified morphologies and extend fine processes through nearby neural parenchyma. Resting microglial cells play key roles in housekeeping functions such as pruning excess or dysfunctional synapses [43,44], distributing supportive growth factors to active neurons [45] and regulating synaptic function [46,47]. However, activated microglial cells phagocytose cell debris and aggregated proteins while concurrently secreting toxic compounds during injury in the central nervous system [48]. Activated microglial cells are reported to secrete cytotoxic factors including free oxygen intermediates, proteases and excitatory amino acids, which may induce neuronal degeneration [48-50]. Microglial cells also secrete tumor necrosis factor-α [51,52], interleukin-1 [30,53,54] and the pro-forms of NGF, BDNF, NT3 and NT4 [4]. It has been reported that microglia-derived pro-NGF promotes photoreceptor cell death via activation of pro-NGF/P75^{NTR} signaling [4]. Recent studies indicate that activated microglia and Müller cells can influence each other to initiate a program of bidirectional microglia-Müller cell signaling, thereby contributing to the neuroinflammatory response and photoreceptor death [30,55]. Therefore, it is conceivable that activated Müller cells and microglia may contribute to photoreceptor degeneration via release of neurotoxic compounds in our transgenic mice. Our observations are



consistent with a previous study that reported that accumulation of activated microglia in the outer retina occurred concurrently with the wave of photoreceptor degeneration in a mouse model of subretinal hemorrhage [56].

The photoreceptor degeneration after Müller cell ablation could be due to either reduced NT3/TrkC signaling or enhanced pro-neurotrophin/p75^{NTR} signaling or both [2,13,57]. It has been reported that Müller cells, RPE cells and ganglion cells express mature NT3 [58-61] and activated microglial cells express pro-NT3 [4]. As Muller cells are the major source of neurotrophic factors such as NT3 [60], basic fibroblast growth factor and ciliary neurotrophic factor [62-64], loss of neurotrophic support from Muller cells would affect the survival of photoreceptors. Our previous study showed that intravitreal supplementation of ciliary neurotrophic factor attenuates photoreceptor injury in this transgenic model [17]. In the present study, we found that activation of microglial and surviving Müller cells was concomitant with significant overexpression of pro-NT3 and P75^{NTR} and reduction in mature NT3 after Müller cell ablation. We hypothesized that supplementation with exogenous mature NT3 and inhibition of pro-neurotrophins/P75^{NTR} signaling would protect photoreceptors. We tested this hypothesis by intravitreal injections of recombinant mature NT3 and a neutralizing Ab against P75^{NTR}. Quantitative analysis of PNA-stained cone photoreceptor outer segments showed intravitreal injection of NT3 and inhibition of P75^{NTR}, either alone or in combination, attenuated photoreceptor degeneration. These results further confirm a potentially critical role of abnormal expression of NT3, pro-NT3 and P75^{NTR} in photoreceptor degeneration. Previous studies have shown that activation of pro-NGF/P75^{NTR} signaling contributes to retinal neuronal damage in animal models of glaucoma, retinal degenerations and diabetes [2-4,6,8,9,65]. A recent study reported that exogenous delivery of the NT-3 gene can be neuroprotective in an animal model of focal cerebral injury [66]. There is evidence that mature NT3 stimulates Müller cells to produce bFGF, and this effect can be blocked by a NT3 neutralizing Ab [2]. Dorrell et al. have reported that targeted delivery of NT4 to activated Müller cells protects retinas from neuronal degeneration in *Vldlr*^{-/-} mice [67]. To our knowledge, our study is the first to show direct evidence that overexpressions of pro-NT3 and P75^{NTR} as well as downregulation of mature NT3 are associated with photoreceptor degeneration and that regulation of pro-NT3/P75^{NTR} signaling is effective in photoreceptor protection.

We also found that intravitreal injections of NT3 and inhibition of P75^{NTR} resulted in reduced microglial activation when they were given individually or together. In eyes receiving BSS injection, massive migration of activated microglial cells was accompanied by loss of

photoreceptor outer segments. A recent study indicated that migration of activated microglial cells into the sub-retinal space transforms the environment of the outer retina from an immune-privileged zone into a highly proinflammatory region, which, in turn, potentiates cellular apoptosis and photoreceptor degeneration [56]. We found that the degree of microglial inhibition appeared to map well with the extent of photoreceptor protection after intravitreal injection of mature NT3 and the P75^{NTR} Ab. Since microglial cells can be activated by cell debris and aggregated proteins released from damaged tissue, the inhibition of microglia that we observed after treatment with mature NT3 might be due to a protective effect of NT3 on photoreceptors that would have limited the extent of damage. For eyes receiving the P75^{NTR} neutralizing Ab, it is possible that blocking P75^{NTR} reduced the production of inflammatory cytokines, chemotactic cytokines and adhesion molecules from activated Müller cells, thus preventing microglial activation and protecting photoreceptors [3,12,30,56,68]. The combined treatment by NT3 and P75^{NTR} Ab is likely to result in dual effects on neuroprotection and microglial inhibition.

Conclusions

We have shown that patchy loss of Müller cells induces photoreceptor degeneration, which is accompanied by activation of surviving Müller cells and microglia. These changes were associated with reduced expression of mature NT3 and upregulation of pro-NT3 and P75^{NTR}. The ability of intravitreal treatment with NT3 and the P75^{NTR} neutralizing Ab to protect photoreceptors and inhibit microglial activation indicates that manipulation of neuronal-Müller cell-microglial interactions may be a novel strategy for neuroprotection in retinal disease.

Abbreviations

BDNF: Brain-derived neurotrophic factor; DTA: Diphtheria toxin fragment A; Gat: Gα protein transducing; GFAP: Glial fibrillary acidic protein; GNAT1: Guanine nucleotide-binding protein subunit alpha-1; GS: Glutamine synthetase; Iba-1: Ionized calcium binding adaptor molecule 1; IHC: Immunohistochemistry; NGF: Nerve growth factor; NT-3: Neurotrophin-3; P75^{NTR}: p75 neurotrophin receptor; PNA: Peanut-agglutinin; Rlbp1: Retinaldehyde binding protein 1; TMX: Tamoxifen.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

WS was involved in the study design, performed the experiments, analyzed the data and images, and wrote the manuscript. LZ contributed to genotyping and Western blot analysis. SL contributed to genotyping, IHC, Western blot and data analysis. SC assisted with in vivo experiments. MG was involved in the study design and writing the manuscript. All authors have read and approved the final version of the manuscript.

Acknowledgements

This study was supported by grants from Lowy Medical Research Institute, the National Health and Medical Research Council (APP1028393, APP1050373) and Ophthalmic Research Institute of Australia. Mark Gillies is a

fellow of Sydney Medical School Foundation and supported by a National Health and Medical Research Council (Australia) Practitioner Fellowship.

Received: 12 August 2013 Accepted: 1 November 2013

Published: 14 November 2013

References

1. Friedman WJ: Proneurotrophins, seizures, and neuronal apoptosis. *Neuroscientist* 2010, **16**:244–252.
2. Harada T, Harada C, Nakayama N, Okuyama S, Yoshida K, Kohsaka S, Matsuda H, Wada K: Modification of glial-neuronal cell interactions prevents photoreceptor apoptosis during light-induced retinal degeneration. *Neuron* 2000, **26**:533–541.
3. Bai Y, Dergham P, Nedev H, Xu J, Galan A, Rivera JC, ZhiHua S, Mehta HM, Woo SB, Sarunic MV, Neet KE, Saragovi HU: Chronic and acute models of retinal neurodegeneration TrkA activity are neuroprotective whereas p75NTR activity is neurotoxic through a paracrine mechanism. *J Biol Chem* 2010, **285**:39392–39400.
4. Srinivasan B, Roque CH, Hempstead BL, Al-Ubaidi MR, Roque RS: Microglia-derived pro-nerve growth factor promotes photoreceptor cell death via p75 neurotrophin receptor. *J Biol Chem* 2004, **279**:41839–41845.
5. Jansen P, Giehl K, Nyengaard JR, Teng K, Lioubinski O, Sjoegaard SS, Breiderhoff T, Gotthardt M, Lin F, Eilers A, Petersen CM, Lewin GR, Hempstead BL, Willnow TE, Nykjaer A: Roles for the pro-neurotrophin receptor sortilin in neuronal development, aging and brain injury. *Nat Neurosci* 2007, **10**:1449–1457.
6. Al-Gayyar MM, Matragoon S, Pillai BA, Ali TK, Abdelsaid MA, El-Remessy AB: Epicatechin blocks pro-nerve growth factor (proNGF)-mediated retinal neurodegeneration via inhibition of p75 neurotrophin receptor expression in a rat model of diabetes [corrected]. *Diabetologia* 2011, **54**:669–680.
7. Ali TK, Al-Gayyar MM, Matragoon S, Pillai BA, Abdelsaid MA, Nussbaum JJ, El-Remessy AB: Diabetes-induced peroxynitrite impairs the balance of pro-nerve growth factor and nerve growth factor, and causes neurovascular injury. *Diabetologia* 2011, **54**:657–668.
8. Al-Gayyar MM, Mysona BA, Matragoon S, Abdelsaid MA, El-Azab MF, Shanab AY, Ha Y, Smith SB, Bollinger KE, El-Remessy AB: Diabetes and overexpression of proNGF cause retinal neurodegeneration via activation of RhoA pathway. *PLoS One* 2013, **8**:e54692.
9. Rudzinski M, Wong TP, Saragovi HU: Changes in retinal expression of neurotrophins and neurotrophin receptors induced by ocular hypertension. *J Neurobiol* 2004, **58**:341–354.
10. Lonngrén U, Napankangas U, Lafuente M, Mayor S, Lindqvist N, Vidal-Sanz M, Hallböök F: The growth factor response in ischemic rat retina and superior colliculus after brimonidine pre-treatment. *Brain Res Bull* 2006, **71**:208–218.
11. Hu B, Yip HK, So KF: Localization of p75 neurotrophin receptor in the retina of the adult SD rat: an immunocytochemical study at light and electron microscopic levels. *Glia* 1998, **24**:187–197.
12. Lebrun-Julien F, Bertrand MJ, De Backer O, Stellwagen D, Morales CR, Di Polo A, Barker PA: ProNGF induces TNFα-dependent death of retinal ganglion cells through a p75NTR non-cell-autonomous signaling pathway. *Proc Natl Acad Sci USA* 2010, **107**:3817–3822.
13. Ibanez CF, Simi A: p75 neurotrophin receptor signaling in nervous system injury and degeneration: paradox and opportunity. *Trends Neurosci* 2012, **35**:431–440.
14. Harada T, Harada C, Kohsaka S, Wada E, Yoshida K, Ohno S, Mamada H, Tanaka K, Parada LF, Wada K: Microglia-Müller glia cell interactions control neurotrophic factor production during light-induced retinal degeneration. *J Neurosci* 2002, **22**:9228–9236.
15. Bringmann A, Pannicke T, Grosche J, Francke M, Wiedemann P, Skatchkov SN, Osborne NN, Reichenbach A: Müller cells in the healthy and diseased retina. *Prog Retin Eye Res* 2006, **25**:397–424.
16. Reichenbach A, Bringmann A: New functions of Müller cells. *Glia* 2013, **61**:651–678.
17. Shen W, Fruttiger M, Zhu L, Chung SH, Barnett NL, Kirk JK, Lee S, Coorey NJ, Killingsworth M, Sherman LS, Gillies MC: Conditional Müller cell ablation causes independent neuronal and vascular pathologies in a novel transgenic model. *J Neurosci* 2012, **32**:15715–15727.
18. Powner MB, Gillies MC, Tretiach M, Scott A, Guymer RH, Hageman GS, Fruttiger M: Perifoveal Müller cell depletion in a case of macular telangiectasia type 2. *Ophthalmology* 2010, **117**:2407–2416.

19. Powner MB, Gillies MC, Zhu M, Vevis K, Hunyor AP, Fruttiger M: **Loss of Muller's cells and photoreceptors in macular telangiectasia type 2.** *Ophthalmology* 2013, **120**:2344–2352.
20. Issa PC, Gillies MC, Chew EY, Bird AC, Heeren TF, Peto T, Holz FG, Scholl HP: **Macular telangiectasia type 2.** *Prog Retin Eye Res* 2013, **34**:49–77.
21. Adler R, Curcio C, Hicks D, Price D, Wong F: **Cell death in age-related macular degeneration.** *Mol Vis* 1999, **5**:31.
22. Ambati J, Fowler BJ: **Mechanisms of age-related macular degeneration.** *Neuron* 2012, **75**:26–39.
23. Stitt AW, Lois N, Medina RJ, Adamson P, Curtis TM: **Advances in our understanding of diabetic retinopathy.** *Clin Sci (Lond)* 2013, **125**:1–17.
24. Fletcher EL, Phipps JA, Ward MM, Puthussery T, Wilkinson-Berka JL: **Neuronal and glial cell abnormality as predictors of progression of diabetic retinopathy.** *Curr Pharm Des* 2007, **13**:2699–2712.
25. Fletcher EL, Downie LE, Hatzopoulos K, Vessey KA, Ward MM, Chow CL, Pianta MJ, Vingrys AJ, Kalloniatis M, Wilkinson-Berka JL: **The significance of neuronal and glial cell changes in the rat retina during oxygen-induced retinopathy.** *Doc Ophthalmol* 2010, **120**:67–86.
26. Shen WY, Lai CM, Graham CE, Binz N, Lai YK, Eade J, Guidolin D, Ribatti D, Dunlop SA, Rakoczy PE: **Long-term global retinal microvascular changes in a transgenic vascular endothelial growth factor mouse model.** *Diabetologia* 2006, **49**:1690–1701.
27. Pearson RA, Barber AC, Rizzi M, Hippert C, Xue T, West EL, Duran Y, Smith AJ, Chuang JZ, Azam SA, Luhmann UF, Benucci A, Sung CH, Bainbridge JW, Carandini M, Yau KW, Sowden JC, Ali RR: **Restoration of vision after transplantation of photoreceptors.** *Nature* 2012, **485**:99–103.
28. Gopalakrishna KN, Boyd KK, Artemyev NO: **Comparative analysis of cone and rod transducins using chimeric Galpha subunits.** *Biochemistry* 2012, **51**:1617–1624.
29. Chen F, Ng PS, Faul KF, Lee RH: **Cone photoreceptor betagamma-transducin: posttranslational modification and interaction with phosducin.** *Invest Ophthalmol Vis Sci* 2003, **44**:4622–4629.
30. Wang M, Ma W, Zhao L, Fariss RN, Wong WT: **Adaptive Muller cell responses to microglial activation mediate neuroprotection and coordinate inflammation in the retina.** *J Neuroinflammation* 2011, **8**:173.
31. Mori T, Tan J, Arendash GW, Koyama N, Nojima Y, Town T: **Overexpression of human S100B exacerbates brain damage and perinfarct gliosis after permanent focal ischemia.** *Stroke* 2008, **39**:2114–2121.
32. Ridet JL, Malhotra SK, Privat A, Gage FH: **Reactive astrocytes: cellular and molecular cues to biological function.** *Trends Neurosci* 1997, **20**:570–577.
33. Streit WJ, Walter SA, Pennell NA: **Reactive microgliosis.** *Prog Neurobiol* 1999, **57**:563–581.
34. Unger JW: **Glial reaction in aging and Alzheimer's disease.** *Microsc Res Tech* 1998, **43**:24–28.
35. Wilson JX: **Antioxidant defense of the brain: a role for astrocytes.** *Can J Physiol Pharmacol* 1997, **75**:1149–1163.
36. Bringmann A, Iandiev I, Pannicke T, Wurm A, Hollborn M, Wiedemann P, Osborne NN, Reichenbach A: **Cellular signaling and factors involved in Muller cell gliosis: neuroprotective and detrimental effects.** *Prog Retin Eye Res* 2009, **28**:423–451.
37. Coorey NJ, Shen W, Chung SH, Zhu L, Gillies MC: **The role of glia in retinal vascular disease.** *Clin Exp Optom* 2012, **95**:266–281.
38. Beattie MS, Harrington AW, Lee R, Kim JY, Boyce SL, Longo FM, Bresnahan JC, Hempstead BL, Yoon SO: **ProNGF induces p75-mediated death of oligodendrocytes following spinal cord injury.** *Neuron* 2002, **36**:375–386.
39. Friedman WJ: **Neurotrophins induce death of hippocampal neurons via the p75 receptor.** *J Neurosci* 2000, **20**:6340–6346.
40. Harrington AW, Leiner B, Blechschmitt C, Arevalo JC, Lee R, Morl K, Meyer M, Hempstead BL, Yoon SO, Giehl KM: **Secreted proNGF is a pathophysiological death-inducing ligand after adult CNS injury.** *Proc Natl Acad Sci USA* 2004, **101**:6226–6230.
41. Santos AM, Lopez-Sanchez N, Martin-Oliva D, de la Villa P, Cuadros MA, Frade JM: **Sortilin participates in light-dependent photoreceptor degeneration in vivo.** *PLoS One* 2012, **7**:e36243.
42. Srinivasan B, Wang Z, Brun-Zinkernagel AM, Collier RJ, Black RA, Frank SJ, Barker PA, Roque RS: **Photic injury promotes cleavage of p75NTR by TACE and nuclear trafficking of the p75 intracellular domain.** *Mol Cell Neurosci* 2007, **36**:449–461.
43. Trapp BD, Wujek JR, Criste GA, Jalabi W, Yin X, Kidd GJ, Stohman S, Ransohoff R: **Evidence for synaptic stripping by cortical microglia.** *Glia* 2007, **55**:360–368.
44. Wake H, Moorhouse AJ, Jinno S, Kohsaka S, Nabekura J: **Resting microglia directly monitor the functional state of synapses in vivo and determine the fate of ischemic terminals.** *J Neurosci* 2009, **29**:3974–3980.
45. Walton NM, Sutter BM, Laywell ED, Levkoff LH, Kearns SM, Marshall GP II, Scheffler B, Steindler DA: **Microglia instruct subventricular zone neurogenesis.** *Glia* 2006, **54**:815–825.
46. Bessis A, Bechade C, Bernard D, Roumier A: **Microglial control of neuronal death and synaptic properties.** *Glia* 2007, **55**:233–238.
47. Roumier A, Pascual O, Bechade C, Wakselman S, Poncer JC, Real E, Triller A, Bessis A: **Prenatal activation of microglia induces delayed impairment of glutamatergic synaptic function.** *PLoS One* 2008, **3**:e2595.
48. Schwartz M, Shechter R: **Systemic inflammatory cells fight off neurodegenerative disease.** *Nat Rev Neurol* 2010, **6**:405–410.
49. Banati RB, Gehrmann J, Schubert P, Kreutzberg GW: **Cytotoxicity of microglia.** *Glia* 1993, **7**:111–118.
50. Boje KM, Arora PK: **Microglial-produced nitric oxide and reactive nitrogen oxides mediate neuronal cell death.** *Brain Res* 1992, **587**:250–256.
51. Sawada M, Kondo N, Suzumura A, Marunouchi T: **Production of tumor necrosis factor-alpha by microglia and astrocytes in culture.** *Brain Res* 1989, **491**:394–397.
52. Kaur C, Rathnasamy G, Ling EA: **Roles of activated microglia in hypoxia induced neuroinflammation in the developing brain and the retina.** *J Neuroimmune Pharmacol* 2013, **8**:66–78.
53. Rivera JC, Sitaras N, Noueihed B, Hamel D, Madaan A, Zhou T, Honore JC, Quiniou C, Joyal JS, Hardy P, Sennlaub F, Lubell W, Chemtob S: **Microglia and interleukin-1beta in ischemic retinopathy elicit microvascular degeneration through neuronal semaphorin-3A.** *Arterioscler Thromb Vasc Biol* 2013, **33**:1881–1891.
54. Yoshida N, Ikeda Y, Notomi S, Ishikawa K, Murakami Y, Hisatomi T, Enaida H, Ishibashi T: **Laboratory evidence of sustained chronic inflammatory reaction in retinitis pigmentosa.** *Ophthalmology* 2013, **120**:e5–e12.
55. Rutar M, Natoli R, Provis JM: **Small interfering RNA-mediated suppression of Ccl2 in Muller cells attenuates microglial recruitment and photoreceptor death following retinal degeneration.** *J Neuroinflammation* 2012, **9**:221.
56. Zhao L, Ma W, Fariss RN, Wong WT: **Minocycline attenuates photoreceptor degeneration in a mouse model of subretinal hemorrhage microglial: inhibition as a potential therapeutic strategy.** *Am J Pathol* 2011, **179**:1265–1277.
57. Harrington AW, Kim JY, Yoon SO: **Activation of Rac GTPase by p75 is necessary for c-jun N-terminal kinase-mediated apoptosis.** *J Neurosci* 2002, **22**:156–166.
58. Agarwal N, Agarwal R, Kumar DM, Ondricek A, Clark AF, Wordinger RJ, Pang IH: **Comparison of expression profile of neurotrophins and their receptors in primary and transformed rat retinal ganglion cells.** *Mol Vis* 2007, **13**:1311–1318.
59. Li R, Wen R, Banzon T, Maminishkis A, Miller SS: **CNTF mediates neurotrophic factor secretion and fluid absorption in human retinal pigment epithelium.** *PLoS One* 2011, **6**:e23148.
60. Taylor S, Srinivasan B, Wordinger RJ, Roque RS: **Glutamate stimulates neurotrophin expression in cultured Muller cells.** *Brain Res Mol Brain Res* 2003, **111**:189–197.
61. Tauris J, Gustafsen C, Christensen EI, Jansen P, Nykjaer A, Nyengaard JR, Teng KK, Schwarz E, Ovesen T, Madsen P, Petersen CM: **Proneurotrophin-3 may induce Sortilin-dependent death in inner ear neurons.** *Eur J Neurosci* 2011, **33**:622–631.
62. LaVail MM, Unoki K, Yasumura D, Matthes MT, Yancopoulos GD, Steinberg RH: **Multiple growth factors, cytokines, and neurotrophins rescue photoreceptors from the damaging effects of constant light.** *Proc Natl Acad Sci USA* 1992, **89**:11249–11253.
63. Rhee KD, Ruiz A, Duncan JL, Hauswirth WW, Lavail MM, Bok D, Yang XJ: **Molecular and cellular alterations induced by sustained expression of ciliary neurotrophic factor in a mouse model of retinitis pigmentosa.** *Invest Ophthalmol Vis Sci* 2007, **48**:1389–1400.
64. Walsh N, Valter K, Stone J: **Cellular and subcellular patterns of expression of bFGF and CNTF in the normal and light stressed adult rat retina.** *Exp Eye Res* 2001, **72**:495–501.
65. Matragoon S, Al-Gayyar MM, Mysona BA, Abdelsaid MA, Pillai BA, Neet KE, Fagan SC, El-Remessy AB: **Electroporation-mediated gene delivery of cleavage-resistant pro-nerve growth factor causes retinal neuro- and vascular degeneration.** *Mol Vis* 2012, **18**:2993–3003.

66. Zhang J, Shi Q, Yang P, Xu X, Chen X, Qi C, Lu H, Zhao B, Zheng P, Zhang P, Liu Y: **Neuroprotection of neurotrophin-3 against focal cerebral ischemia/reperfusion injury is regulated by hypoxia-responsive element in rats.** *Neuroscience* 2012, **222**:1–9.
67. Dorrell MI, Aguilar E, Jacobson R, Yanes O, Gariano R, Heckenlively J, Banin E, Ramirez GA, Gasmi M, Bird A, Siuzdak G, Friedlander M: **Antioxidant or neurotrophic factor treatment preserves function in a mouse model of neovascularization-associated oxidative stress.** *J Clin Invest* 2009, **119**:611–623.
68. Lebrun-Julien F, Morquette B, Douillette A, Saragovi HU, Di Polo A: **Inhibition of p75(NTR) in glia potentiates TrkA-mediated survival of injured retinal ganglion cells.** *Mol Cell Neurosci* 2009, **40**:410–420.

doi:10.1186/1742-2094-10-137

Cite this article as: Shen *et al.*: Involvement of NT3 and P75^{NTR} in photoreceptor degeneration following selective Müller cell ablation. *Journal of Neuroinflammation* 2013 **10**:137.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

