



Published in final edited form as:

*Mucosal Immunol.* 2014 May ; 7(3): 602–614. doi:10.1038/mi.2013.79.

## TNF $\alpha$ -dependent development of lymphoid tissue in the absence of ROR $\gamma$ t<sup>+</sup> Lymphoid Tissue Inducer cells

Glauca C. Furtado<sup>1</sup>, Michelle E. Pacer<sup>1</sup>, Gerold Bongers<sup>1</sup>, Cecile Benezech<sup>2</sup>, Zhengxiang He<sup>1</sup>, Lili Chen<sup>1</sup>, M Cecilia Berin<sup>1</sup>, George Kollias<sup>3</sup>, Jorge H. Caamaño<sup>2</sup>, and Sergio A. Lira<sup>1</sup>

<sup>1</sup>Immunology Institute, Icahn School of Medicine at Mount Sinai, 1425 Madison Ave, Box 1630, New York, NY 10029-6574

<sup>2</sup>IBR-MRC Centre for Immune Regulation, College of Medical and Dental Sciences, University of Birmingham, Birmingham B15 2TT, UK

<sup>3</sup>Biomedical Sciences Research Center Alexander Fleming, Vari, Greece

### Abstract

Lymphoid tissue often forms within sites of chronic inflammation. Here we report that expression of the proinflammatory cytokine TNF $\alpha$  drives development of lymphoid tissue in the intestine. Formation of this ectopic lymphoid tissue was not dependent on the presence of canonical ROR $\gamma$ t<sup>+</sup> lymphoid tissue inducer (LTi) cells, because animals expressing increased levels of TNF $\alpha$  but lacking ROR $\gamma$ t<sup>+</sup> LTi cells (*TNF/Rorc(gt)*<sup>-/-</sup> mice) developed lymphoid tissue in inflamed areas. Unexpectedly, such animals developed several lymph nodes that were structurally and functionally similar to those of wild type animals. TNF $\alpha$  production by F4/80<sup>+</sup> myeloid cells present within the anlagen was important for activation of stromal cells during the late stages of embryogenesis and for the activation of an organogenic program that allowed development of lymph nodes. Our results show that lymphoid tissue organogenesis can occur in the absence of LTi cells and suggest that interactions between TNF $\alpha$ -expressing myeloid cells and stromal cells have an important role in secondary lymphoid organ formation.

### Introduction

Lymphoid organs are critical for generation of adaptive immune response. Secondary lymphoid organs (SLO) are formed at predefined areas during embryogenesis whereas tertiary lymphoid organs (TLO) are formed after birth in tissues with ongoing inflammatory processes<sup>1,2</sup>. Both secondary and tertiary lymphoid organs have lymphocytes that are topologically segregated, and diverse sets of myeloid and stromal cells. In addition, they have specialized vasculature such as high endothelial venules (HEV), and a lymphatic network.

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:[http://www.nature.com/authors/editorial\\_policies/license.html#terms](http://www.nature.com/authors/editorial_policies/license.html#terms)

Address correspondence to: Glauca C. Furtado or Sergio A. Lira, Immunology Institute, Icahn School of Medicine at Mount Sinai, 1425 Madison Ave, Box 1630, New York, NY 10029-6574, Phone: 212-659-8043, FAX: 212-849-2525.

#### Competing financial interests

The authors declare no financial interests

The two major cell types involved in lymph node organogenesis are the hematopoietic lymphoid tissue inducer (LTi) cells and non-hematopoietic lymphoid tissue stromal “organizer cells” (LTo)<sup>1</sup>. Clustering of LTi and LTo cells is an essential step in lymph node development<sup>3</sup>. Animals that are deficient in the nuclear retinoid orphan receptor (ROR) $\gamma$ , encoded by the *Rorc* gene, or the negative regulator of basic helix-loop-helix protein signaling Id2, lack LTi cells and therefore fail to form lymph nodes and Peyer’s patches<sup>4-6</sup>. The current model for development of lymphoid organs posits that LTi cells originate in the fetal liver from common lymphoid progenitors and that they migrate to the sites where the lymph nodes are formed (lymph node anlagen)<sup>1, 7</sup>. At these sites, binding of the TNF $\alpha$  family ligand Receptor Activator of NF-kB (RANKL) to its receptor RANK induces the differentiation and survival of LTi cells and trigger expression of LT $\alpha$ 1 $\beta$ 2 on their surface<sup>3, 8-11</sup>. A key step in the development of LNs is the engagement of Lymphotoxin  $\alpha$ 1 $\beta$ 2 (LT $\alpha$ 1 $\beta$ 2) expressed by LTi cells to its receptor LT $\beta$ R on mesenchymal organizer cells<sup>12, 13</sup>. This interaction promotes upregulation of intracellular adhesion molecule (ICAM-1), vascular cell adhesion molecule (VCAM-1) and mucosal addressin cell adhesion molecule (MAdCAM-1) on the surface of LTo cells<sup>14, 15</sup> and the expression of the chemokines CCL19, CCL21, and CXCL13<sup>7</sup>.

Animals genetically deficient in LT- $\alpha$  and LT $\beta$ R do not form lymph nodes or Peyer’s patches<sup>10, 12, 16</sup>. Furthermore, genetic deletion of molecules in the LT $\beta$ R signaling pathway (NF-kappa B non canonical pathway) such as NF-kappa B-inducible kinase (NIK)<sup>17</sup> and RelB<sup>18</sup> precludes LN formation. While the role of LT $\alpha$ 1 $\beta$ 2/LT $\beta$ R is firmly established in the process of lymphoid organogenesis, the role of other members of the TNF $\alpha$  superfamily is unclear.

Female mice injected in utero with LT $\beta$ R-Ig fusion protein retain cervical and mesenteric lymph nodes (mLN) but fail to form other lymph nodes<sup>19, 20</sup>. However, simultaneous treatment LT $\beta$ R-Ig fusion protein and anti-TNFR1 antibody, or LT $\beta$ R-Ig plus anti-TNF $\alpha$  antibodies, prevents development of all lymph nodes<sup>21</sup>, which suggests that TNF $\alpha$  has a role in mLN organogenesis. However, TNF $\alpha$  or TNF-R1-deficient mice have all lymph nodes, including mLN, but they fail to form B cell follicles. These results suggest that TNF $\alpha$  activity in lymphoid organogenesis may be secondary to other TNF $\alpha$  members such as LT. However, simultaneous deficiency of TNFR1 and RelA abrogates the development of all lymph nodes, despite the presence of a normal complement of LT $\alpha$ 1 $\beta$ 2<sup>+</sup> LTi cells<sup>22</sup>. Thus, the role of TNF $\alpha$  in lymphoid organogenesis remains poorly defined.

Here we used *TNF*<sup>ARE/+</sup> mice, a well-established model of human inflammatory disease, to study the role of TNF $\alpha$  in lymphoid organogenesis. These animals express increased levels of TNF $\alpha$  under basal conditions, due to mutation in the 3’ region of the *Tnfa* gene that causes higher stability of its mRNA and, consequently, increased levels of TNF $\alpha$  protein<sup>23</sup>. Intercross of *TNF*<sup>ARE/+</sup> mice with *Rorc*( $\gamma$ )<sup>-/-</sup> mice led to the generation of *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice. Surprisingly, *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice developed TLO and several SLO (mesenteric, axillary and cervical LN and others) despite the lack of the classical ROR $\gamma$ <sup>+</sup> LTi cells. Development of lymph nodes was mechanistically linked to activation of stromal cells by TNF $\alpha$  produced by myeloid cells present in the anlagen, and expression of

molecules involved in lymphoid organogenesis. These results establish that lymphoid organogenesis can occur in the absence of *Rorc* if there is increased TNF $\alpha$  signaling.

## Results

### Increased expression of TNF $\alpha$ promotes development of TLO in the absence of LTi cells

Two types of lymphoid aggregates can be identified in the intestine of adult mice: isolated lymphoid follicles (ILF) and tertiary lymphoid organs (TLO). ILFs are genetically programmed clusters of B cells present at the base of the villi, that require ROR $\gamma$ <sup>+</sup>LTi cells and LT $\beta$ R signaling for their formation<sup>5, 24–26</sup>. TLO are composed by large clusters of B220<sup>+</sup> cells that contain CD3<sup>+</sup> lymphocytes, and are formed in response to infection or inflammation<sup>27, 28</sup>. To further define the role of LTi cells and TNF $\alpha$  in the formation of lymphoid aggregates in the intestine we examined the presence of these structures in the ileum of *TNF*<sup>ARE/+</sup> mice. The inflammatory infiltrates in the ileum are composed of neutrophils, macrophages, and T cells that are distributed throughout the submucosa and muscular layers and sometimes reach the serosa. Large mononuclear aggregates rich in B cells, or TLO, are also found in the terminal ileum of the *TNF*<sup>ARE/+</sup> mice<sup>29</sup>. To determine whether the formation of these aggregates is dependent on ROR $\gamma$ <sup>+</sup>LTi cells we crossed *Rorc*( $\gamma$ )<sup>-/-</sup> mice with *TNF*<sup>ARE/+</sup> mice to generate *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice. Histological analysis of the terminal ileum of age-matched wild type (WT), *Rorc*( $\gamma$ )<sup>-/-</sup>, *TNF/Rorc*( $\gamma$ )<sup>+/+</sup> and *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice at 16–20 weeks of age showed that *TNF/Rorc*( $\gamma$ )<sup>+/+</sup> and *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice, but not WT or *Rorc*( $\gamma$ )<sup>-/-</sup> mice, had marked submucosal inflammation, vilus blunting, patchy transmural inflammation, and lymphoid aggregates (Figure 1a). The lymphoid aggregates in *TNF/Rorc*( $\gamma$ )<sup>+/+</sup> and *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice contained large clusters of B220<sup>+</sup> B cells and few CD3<sup>+</sup> T cells (Figure 1b and 1c), which were absent in *Rorc*( $\gamma$ )<sup>-/-</sup> mice. These results indicate that TLO can be formed in the ileum in the absence of ROR $\gamma$ <sup>+</sup>LTi cells.

### *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice develop secondary lymphoid organs

*Rorc* is essential for development of secondary lymphoid organs<sup>5</sup>. As expected, no lymph nodes were found in the *Rorc*( $\gamma$ )<sup>-/-</sup> mice examined (Figure 2a). However, we were surprised to find that 100% of the *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice had fully developed mesenteric LN (mLN) that were grossly indistinguishable from those found in WT mice. Axillary (Figure 2b), cervical (Figure 2c), brachial, inguinal, para-aortic, and peripancreatic LN were also present at lower frequency (Figure 2d). Mediastinal and popliteal LN, as well as Peyer's patches, were not observed in these animals.

To further characterize the structure of the LNs present in *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice we performed immunostaining. LNs of WT and *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice had segregated T and B cell areas, PNA<sup>d+</sup> high endothelial venules, an extensive lymphatic network, ER-TR7<sup>+</sup> lymph node stroma and CD35<sup>bright</sup> follicular dendritic cells (Figure 2e). To determine if these LN were functional we immunized *TNF/Rorc*( $\gamma$ )<sup>+/+</sup> and *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice orally with OVA and cholera toxin 7 times at 1-wk intervals and assessed OVA-specific antibody serum titers by ELISA (Figures 2f). The serum levels of OVA-specific IgA and IgG were similar between *TNF/Rorc*( $\gamma$ )<sup>+/+</sup> and *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice, indicating that both strains

responded to oral immunization with OVA. We next examined if cells from the mesenteric lymph nodes could produce cytokines after immunization. mLNs were collected and cultured with media alone or with 50µg/ml of OVA. Supernatants were harvested 72 hours later and IFN $\gamma$  and IL-17 were measured by ELISA. As shown in Figure 2g, similar levels of IFN $\gamma$  were produced by mesenteric LN cells of *TNF/Rorc( $\gamma$ )<sup>+/+</sup>* and *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice. As expected, IL-17 was not detected in *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* cells since ROR $\gamma$ t is required for IL-17 production (Figure 2h)<sup>30</sup>. Collectively these results indicate that increased expression of TNF $\alpha$  can drive the formation of secondary lymphoid organs in the absence of ROR $\gamma$ t<sup>+</sup> LTi cells.

### **F4/80<sup>+</sup>CD11b<sup>+</sup> cells are the source of TNF $\alpha$ in the mLN anlagen**

Our results suggested that a ROR $\gamma$ t-independent cell type could play a role in the formation of SLO in *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice. To start addressing this hypothesis we first examined the cellular composition of the mLN of *Rorc( $\gamma$ )<sup>-/-</sup>* and *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice at P1. Very few lymphocytes were present to the mLN anlagen of *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice at this stage (Figure 3a). F4/80<sup>+</sup>, NK1.1<sup>+</sup> and CD11c<sup>+</sup> cells were the most abundant CD45<sup>+</sup> leukocytes present in the mLN anlagen of *Rorc( $\gamma$ )<sup>-/-</sup>* and *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice, but their relative proportions were comparable. CD11c<sup>+</sup> cells in the mLN of *Rorc( $\gamma$ )<sup>-/-</sup>* and *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice did not express c-Kit (Figure 3b) and thus were distinct from the c-Kit<sup>+</sup>CD11c<sup>+</sup> lymphoid tissue initiator cells shown to be important in the formation of Peyer's patches (PP)<sup>31</sup>. Further flow cytometric analyses showed that the F4/80<sup>+</sup> cells comprised two populations: F480<sup>hi</sup>/CD11b<sup>low</sup>/MHC II<sup>neg</sup>/CD11c<sup>-</sup> and F4/80<sup>low</sup>/CD11b<sup>hi</sup>/MHC II<sup>pos</sup>/CD11c<sup>+</sup> cells (Fig. 3c). These results indicate that there were no marked differences in the type and relative number of leukocytes in the mLN anlagen of *Rorc( $\gamma$ )<sup>-/-</sup>* and *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice at P0.5-P1

We asked next if TNF $\alpha$  was expressed in the mLN during development. TNF $\alpha$  expression was detected in the mLN anlagen of WT mice at steady state during embryogenesis (Supplemental Figure 1b). Next we compared the levels of TNF $\alpha$  mRNA in mLN anlagen of *Rorc( $\gamma$ )<sup>-/-</sup>* and *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice, and found it to be upregulated in the latter at all embryonic and postnatal stages examined (Supplemental Figure 1a). We then used flow cytometry to determine the cellular source of TNF $\alpha$  in the mLN anlagen. In WT mice, TNF $\alpha$  was detected in F4/80<sup>+</sup> myeloid cells as early as E15.5 in WT mLN while TNFR1 was expressed in both myeloid and CD45<sup>-</sup> stromal cells (Supplemental Figure 1c). Analysis of the mLN anlagen of P0.5-P1 *Rorc( $\gamma$ )<sup>-/-</sup>* and *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice showed that TNF $\alpha$  was mainly produced by CD45<sup>+</sup>F4/80<sup>+</sup> cells and not by other CD45<sup>+</sup> or stromal (CD45<sup>-</sup>) cells (Figure 4a). A two fold increase in the production of TNF $\alpha$  by F4/80<sup>+</sup> cells was observed in the mLN anlagen of *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice. Further flow cytometric analyses showed that TNF $\alpha$  was expressed by both F4/80<sup>hi</sup>/CD11b<sup>low</sup> and F4/80<sup>low</sup>/CD11b<sup>hi</sup> cells (Figure 4b). Together these results indicate that: 1) TNF $\alpha$  is physiologically expressed by F4/80<sup>+</sup> cells in the mLN anlagen of WT mice during embryogenesis, 2) that TNF $\alpha$  is expressed by F4/80<sup>+</sup> cells in both *Rorc( $\gamma$ )<sup>-/-</sup>* mice *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice, and, 3) that TNF $\alpha$  expression is increased during embryogenesis and early postnatal life in the mLN of *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice compared to *Rorc( $\gamma$ )<sup>-/-</sup>* mice.

### TNF $\alpha$ does not bypass the requirement of ID2 for lymphoid organogenesis

Id2-deficient mice lack LTi cells<sup>6</sup>, NK cells<sup>6</sup> and fetal CD11b<sup>+</sup> myeloid cells<sup>5</sup> in the lymph node anlagen, and do not develop SLO. We had shown above that TNF $\alpha$  overexpression bypasses the requirement for *Rorc*( $\gamma$ )<sup>+</sup> cells in SLO formation, thus we investigated next if TNF $\alpha$  would bypass the requirement for *Id2* in lymph node organogenesis. To do so, we intercrossed TNF<sup>ARE/+</sup> mice with *Id2*<sup>-/-</sup> mice to generate *TNF/Id2*<sup>-/-</sup> mice (Figure 5). None of the *TNF/Id2*<sup>-/-</sup> mice examined at birth (n=7) had mLN. We also examined the presence of F4/80<sup>+</sup> myeloid cells, and found them to be present in the mLN anlagen of WT and *TNF/Id2*<sup>+/-</sup> mice but absent in *TNF/Id2*<sup>-/-</sup> mice (Figure 5, dashed lines), and in *Id2*<sup>-/-</sup> mice, in agreement with previous reports<sup>5</sup>. Myeloid cell migration to the mLN anlagen of *TNF/Id2*<sup>-/-</sup> mice was impaired and, strikingly, no LNs were formed in these mice. These results indicate that TNF $\alpha$  does not bypass the requirement for *Id2* in lymphoid organogenesis and suggest that TNF $\alpha$ -producing F4/80<sup>+</sup>CD11b<sup>+</sup> cells or NK cells are important for development of lymph nodes in *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice.

### NK cells are not critical for development of mLN in *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice

Because Id2-deficient mice have defective NK cell development<sup>6</sup> it remained possible that NK cells played a role in the formation of SLO. To test this hypothesis, we first examined if NK cells were present in the anlagen. As shown in Figure 3, NK cells were present in the mLN anlagen of *Rorc*( $\gamma$ )<sup>-/-</sup> and *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> at P1. To determine if they played a role in SLO development we depleted them from *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice. To do so, we injected pregnant mothers at E15 and E18 with 200 $\mu$ g of isotype control or with the anti-NK monoclonal antibody PK136, which depletes NK cells in vivo. *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> offspring received additional injection of 100 $\mu$ g of control or PK136 on days 0, 3, 6 and 9. On day 15 the mLN were collected and the number of NK cells and the formation of mLN were examined (Supplementary Figure 2a and 2b). Treatment of *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice with anti-PK136 caused a complete reduction in the number of NK cells in the mLN (Supplementary Fig. 2a), but did not prevent normal development of mLN (Supplementary Fig. 2b). These results indicate that NK cells do not contribute significantly to SLO formation in *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice and suggest that the F4/80<sup>+</sup> cells are the important cells in the process, as they are the sole source of TNF $\alpha$  in the *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> lymph node anlagen.

### TNF $\alpha$ triggers expression of several genes involved in lymphoid organogenesis

To determine how TNF $\alpha$  expression by myeloid cells could contribute to lymph node organogenesis, we compared the transcriptomes of the mLN anlagen of *Rorc*( $\gamma$ )<sup>-/-</sup> and *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice at postnatal day 1 (P1), using the Illumina gene arrays (Figure 6a and 6b). Consistent with a TNF $\alpha$ -driven signature, the highest expressed genes in the mLN of *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice were acute-phase response genes (*Saa3* and *Serpina-3g*). Expression of several genes involved in lymphoid organogenesis such as *Cxcl13*, *Lymphotoxin beta* (*Ltb*), *Relb*, *Ccl19* and *Madcam-1*, was increased in the mLN of *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice. Expression of macrophage related genes (*Lyz2*, *Lyz1*, *csfr1*, *Mmp9*), MHC molecules (H2-M2) and chemokines (*Ccl5*, *Cxcl10* and *Cxcl16*) were also increased in the mLN of *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice. To validate and extend these findings we performed qPCR analysis (Figure 6c). Expression of *Cxcl13*, RANKL, *Ltb*, was confirmed to be upregulated in the

mLN of *TNF/Rorc(γ)<sup>-/-</sup>* animals at different stages of postnatal development. Interestingly, transcripts for *LTα* were significantly upregulated in the mLN of *TNF/Rorc(γ)<sup>-/-</sup>* mice after P1, but not at earlier time points. These results indicate that increased expression of TNFα promotes expression of genes involved in lymphoid tissue organogenesis during embryogenesis.

### TNFα induces stromal cell maturation

Maturation of mesenchymal stromal cells into organizer cell is a key step in lymphoid organogenesis<sup>1</sup>. The existing evidence suggests that activation of the stromal cells is mediated by interaction of LTα1β2 present on the RORγ<sup>+</sup> LTi cells with LTβR expressed on stromal cells. This interaction leads to upregulation of ICAM-1, VCAM-1 and MAdCAM-1 expression on the surface of the stromal cells<sup>14, 32</sup>. To examine whether stromal maturation to “organizer” cells could occur in the absence of LTi cells, we analyzed the presence of ICAM-1<sup>hi</sup>VCAM-1<sup>hi</sup> cells in the mLN region of *Rorc(γ)<sup>-/-</sup>* and *TNF/Rorc(γ)<sup>-/-</sup>* (Figure 7a) and WT mice (Figure 7b) mice by flow cytometry. Cells were gated in the CD45<sup>-</sup> stromal cell population. ICAM-1<sup>hi</sup>VCAM-1<sup>hi</sup> cells were present at a significantly higher frequency in the mLN of *TNF/Rorc(γ)<sup>-/-</sup>* mice at E18.5 onwards when compared with the same region of *Rorc(γ)<sup>-/-</sup>* mice<sup>33</sup> (Figure 7a and 7c). One day after birth the frequency of ICAM-1<sup>hi</sup>VCAM-1<sup>hi</sup> cells in mLN stroma of *TNF/Rorc(γ)<sup>-/-</sup>* was higher than that of *Rorc(γ)<sup>-/-</sup>* mice, but comparable to that of WT mice (Figure 7b). Another parameter of stromal cell activation is the production of chemokines. To examine if the stromal cells from *TNF/Rorc(γ)<sup>-/-</sup>* mice produced chemokines, we sorted CD45<sup>-</sup> cells from the mLN anlagen and performed QPCR analyses. Sorted stromal cells (CD45<sup>-</sup>) from the mLN anlagen of *TNF/Rorc(γ)<sup>-/-</sup>* mice at P1 expressed increased levels of Cxcl13, Ccl19 and Ccl21 mRNA when compared to sorted stromal cells of *Rorc(γ)<sup>-/-</sup>* mice (Figure 7d). Taken together these results indicate that increased levels of TNFα are sufficient to induce lymph node stromal cell maturation in the absence of LTi cells.

### TNFα overexpression does not bypass the requirement for LTβR signaling in lymphoid organogenesis

The high expression levels of LTβR ligands in the mLN of *TNF/Rorc(γ)<sup>-/-</sup>* mice after birth suggested a role for Lymphotoxin-LTβR signaling in the development of LN in this model. To determine if this was indeed the case, we crossed *TNF<sup>ARE/+</sup>* mice with *LTβR*-deficient animals (referred to as *TNF/LTβR<sup>-/-</sup>* mice) (Figure 8). With exception of mLN, no lymph nodes, Peyer's Patches and TLO were found in any of the *TNF/LTβR<sup>-/-</sup>* mice animals examined (age 3–36 wk, n = 19) (Figures 8a, 8b and Supplementary Figure 3a). The mLN of *TNF/LTβR<sup>-/-</sup>* mice were markedly abnormal as shown by the absence of B cell follicles and T cell areas and the impaired recruitment of lymphocytes to these organs (Figure 8c). In addition, HEVs appear to be absent and the number of lymphatic vessels is also reduced. Finally, they lacked CD35<sup>bright</sup> FDC, and had an aberrant ER-TR7<sup>+</sup> stroma.

*LTβR*-deficient mice have severe splenic defects that include loss of T/B cell segregation and an abnormal stroma<sup>12</sup>. Interestingly, the spleen of *TNF/LTβR<sup>-/-</sup>* mice displayed normal T/B cell distribution, had MAdCAM-1<sup>+</sup> cells and a normal ER-TR7<sup>+</sup> cell network (Supplementary Figure 3b). Thus, TNFα overexpression can compensate for the absence of



LT $\beta$ R signaling and promote development of a normal spleen. Together the results indicate that TNF $\alpha$  overexpression corrects the splenic defects, but not the lack of SLO associated with abrogation of LT $\beta$ R signaling.

### Influx of hematopoietic cells into the anlagen of *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice

To gain further insights into the mechanisms of SLO formation, we compared the kinetics of hematopoietic cell recruitment to the mLN anlagen of WT, *Rorc( $\gamma$ )<sup>-/-</sup>* and *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice. A significant number of CD45<sup>+</sup> cells were present in mLN of WT mice at P0 while very few cells were detected in the mesenteric area of both *Rorc( $\gamma$ )<sup>-/-</sup>* and *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice (Figure 9). In contrast, by P5 the mLN anlagen of *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice appeared to be populated by CD45<sup>+</sup> cells while almost no bone marrow derived cells were present in a similar area in *Rorc( $\gamma$ )<sup>-/-</sup>* mice. Consistent with our flow cytometric data, F4/80<sup>+</sup> cells were present within the anlagen of WT and *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mLN, and their frequency was proportionally increased in the latter strain.

### Grafting of *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* anlagen under the kidney capsule induces lymphoid neogenesis

Grafting of re-aggregates of embryonic and neonatal ROR $\gamma$ <sup>+</sup>LTi cells and LTo cells under the kidney capsule of adult mice promotes the formation of structures that resemble lymph nodes that recruit and organize host T and B cells<sup>34</sup>. At P0 the mLN of WT mice contain mostly LTi cells and myeloid cells and are not organized. Kidney grafts of wild type mLN result in organized tissues populated by host lymphocytes only 2–3 weeks after grafting. To test whether the anlagen of *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice could promote development of LN-like structures in adult mice, we grafted them under the kidney capsule of *Rorc( $\gamma$ )<sup>-/-</sup>* recipient mice (Figure 10). Three weeks later the kidneys were removed and processed for histological analysis. All (100%) animals transplanted with WT mLN anlagen (n = 3) and 71% of those transplanted with *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* anlagen (n = 14) developed lymphoid aggregates under the kidney capsule. These aggregates, contained segregated T and B cell areas, HEV and lymphatic vessels, and were similar to host LN. We conclude that mLN anlagen of *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice can promote the formation of lymphoid organs at a non-predestined site in adult mice in the absence of ROR $\gamma$ <sup>+</sup>LTi cells, and that this response is not dependent on systemically increased levels of TNF $\alpha$ .

## Discussion

ROR $\gamma$  is a transcription factor encoded by the *Rorc* gene whose expression is critical for development of embryonic LTi cells and other types of group 3 innate lymphoid cells<sup>35</sup>. Our results show that lymphoid organogenesis can occur in the absence of *Rorc*, provided that there is increased expression of TNF. Formation of most lymph nodes under these circumstances is dependent on LT $\beta$ R signaling.

A body of work supports the notion that SLO and TLO development share common mechanisms. However, evidence first derived from analysis of CCL21-driven transgenic models<sup>36</sup> and other models<sup>37</sup>, suggested that their development differs, as canonical LTi cells, critical for SLO development, were shown not to be absolutely required for TLO

development. Here we show that the generation of TLO in the ileum of *TNF ARE/+* animals is also independent on ROR $\gamma$ <sup>+</sup> LTi cells. Our results complement those of Eberl and colleagues that showed that *Rorc*( $\gamma$ )-deficient mice that received an inflammatory insult such as DSS-induced colitis, develop TLO in the colon<sup>27</sup>. Together these studies demonstrate that inflammatory stimuli promote development of TLO in different areas of the intestine (ileum and colon) in the absence of ROR $\gamma$ <sup>+</sup> LTi cells, and implicate TNF $\alpha$  as an important factor in their generation, since its expression is elevated in the ileum of *TNF ARE/+* mice and in the colon of DSS-treated animals<sup>29, 38</sup>. Importantly, the development of TLO in the intestine of *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice also demonstrates that the formation of these organs is independent of Th17<sup>+</sup> cells and other *Rorc*( $\gamma$ )-dependent members of the growing family of innate lymphoid cells.

While recent experimental evidence supports the concept that TLO formation can occur in the absence of canonical LTi cells, the bulk of the literature suggests that they are critical for development of SLO. The current model for SLO formation suggests that Id2<sup>+</sup>ROR $\gamma$ <sup>+</sup>CD3<sup>-</sup>CD4<sup>+</sup>IL-7R $\alpha$ <sup>+</sup>LT $\alpha$ 1b2<sup>+</sup>RANK<sup>+</sup>RANKL<sup>+</sup>LTi cells are key drivers of lymphoid organogenesis based on the fact that *Id2*-, *Rorc*-, *IL-7Ra*- and *LTa*-deficient mice lack SLO<sup>1, 39</sup>. Exceptions to this rule include nasal associated lymphoid tissue (NALT), whose formation takes place after birth and is not dependent on LT $\alpha$  and *Rorc*( $\gamma$ )<sup>40</sup> and milky spots of the omentum and fat associated lymphoid clusters is also independent of LTi cells<sup>41</sup>. Here we show that many SLO can form in the absence of ROR $\gamma$ <sup>+</sup>LTi cells, provided that the basal levels of TNF $\alpha$  are increased. Mesenteric, axillary and cervical lymph nodes were found in 60–100% of *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice. Other lymph nodes such as brachial, inguinal, para aortic and peripancreatic were found in more than 10% of the mice, whereas popliteal and mediastinal lymph nodes were not detected. The lymph nodes detected in *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> mice were positioned in the same region as WT nodes, had similar architecture and cellularity, and could mount an efficient immune response after immunization.

How could TNF $\alpha$  promote organogenesis in the absence of LTi cells? Here we show that TNF $\alpha$  is produced at higher levels during embryogenesis in the *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> than in *Rorc*( $\gamma$ )<sup>-/-</sup> mice, and that the TNFR1 receptor, is expressed by stromal cells. Stromal cell activation by LTi cells is critical for the generation of lymph nodes. LT $\alpha$ 1 $\beta$ 2 - expressed by LTi cells binds to LT $\beta$ R expressed by stromal cells, which activates both canonical and non-canonical NF- $\kappa$ B signaling pathways to promote the latter cells to become mature stromal “organizer” cells that express increased levels of ICAM-1, VCAM-1 and MAdCAM-1 and the B and T cell chemoattractants CXCL13, CCL19 and CCL21<sup>1, 39</sup>. We suggest that increased levels of TNF $\alpha$  functionally compensated for the lack of LT $\beta$ R signaling during embryogenesis and contributed to the maintenance of a functional anlagen. This hypothesis is supported by our observations that stromal cells in the anlagen of *TNF/Rorc*( $\gamma$ )<sup>-/-</sup>, but not in the *Rorc*( $\gamma$ )<sup>-/-</sup> mice, are activated. They express higher levels of ICAM-1 and VCAM-1 during embryogenesis and immediately after birth. Furthermore, cells in the *TNF/Rorc*( $\gamma$ )<sup>-/-</sup> anlagen expressed increased levels of the TNF $\alpha$ -inducible chemokines CCL2 and CXCL10. These chemokines, acting in concert with TNF $\alpha$ , could promote recruitment of additional hematopoietic cells. At birth, influx of hematopoietic cells could further contribute to



organogenesis. Interestingly, we noted that the expression of LT ligands increased after birth. This could reflect either increased expression of LT ligands by resident non-LTi cells or reflect increased influx of hematopoietic cells that express LT ligands. We favor the second hypothesis because we have observed increased influx of hematopoietic cells in the *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* anlagen during the first 5 days of life. The increased expression of LT ligands is absolutely critical for normal lymph node development as shown by the analysis of the TNF/LT $\beta$ R mutants. In the newborn *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* anlagen, LT ligands could potentially synergize with TNF $\alpha$  to activate the transcription of several molecules related to lymph node organogenesis, macrophage function, and inflammation. A recent report has shown the synergistic effect of TNF $\alpha$  signaling together with the alternative NF- $\kappa$ B pathway to drive high expression levels of Ccl21, Cxcl13, Vcam-1, Icam-1 and Madcam-1 in spleens<sup>42</sup>. Of note, we have detected expression of Cxcl13, Ccl19 and Ccl21, by stromal cells located in the mLN of *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice at P1. These chemokines induce lymphoid organogenesis when expressed in vivo<sup>43-45</sup>, and their expression by stromal cells in the anlagen could account for the formation of LN in *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice.

Myeloid cells are essential for the maintenance<sup>46</sup>, organization<sup>47</sup>, and vascularization<sup>48</sup> of TLO. Results presented here suggest that they may also contribute to development of SLO, working in part as LTi-like cells. We show here that TNF $\alpha$  is expressed by myeloid cells present in the anlagen of WT mLN as early as E15.5. Clusters of fetal CD11b<sup>+</sup> cells and LTi cells are observed at early stages of WT LN development<sup>5</sup>. The generation of the CD11b<sup>+</sup> cells does not require ROR $\gamma$ t because these myeloid cells are still present in the LN anlagen of *Rorc( $\gamma$ )<sup>-/-</sup>* mice, as shown here and in <sup>5</sup>. The increased number of myeloid cells that express stable *TNF $\alpha$*  mRNA contribute to high levels of this protein in the lymph node anlagen in *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice. Consistent with a role for myeloid cells contributing to lymph node development in the *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mouse model, is the observation that absence of these cells (and thus the source of TNF $\alpha$  necessary to activate the stromal cells) in lymph node anlagen of *TNF/Id2<sup>-/-</sup>* mice resulted in the failure to rescue the formation of these organs.

Our results indicate that the development of lymph nodes in *TNF/Rorc( $\gamma$ )<sup>-/-</sup>* mice appears to be delayed when compared to their WT counterparts. In addition, it is not clear why Peyer's patches and some lymph nodes do not develop in this mouse model. The development of these structures may be dependent on the local numbers or phenotype of myeloid cells in those locations. Additional research will be necessary to uncover the origin of these myeloid cells and the factors mediating their influx into different lymph node anlagen.

TNF $\alpha$  receptor ligation activates the NF- $\kappa$ B classical pathway, which involves the I $\kappa$ B kinase and results in the activation of RelA. LT $\beta$ R ligation activates both the NF- $\kappa$ B classical and alternative pathways<sup>18</sup>. The alternative NF- $\kappa$ B pathway is mediated by the NF- $\kappa$ B-inducing kinase (NIK) and results in the activation of NF- $\kappa$ B2/Relb. Because animals genetically deficient in *Ltbr*, *Nik*, and *Relb* do not form lymph nodes<sup>12, 17, 18</sup> it was concluded that the alternative pathway is critically important for the generation of SLO<sup>18</sup>. However, simultaneous deletion of TNFR1 and RelA precludes formation of all LN and PP in double knockout mice due to a stromal cell defect, even in the presence of LTi cells

expressing normal levels of Lymphotoxin<sup>22</sup>, which suggests that the canonical NF- $\kappa$ B pathway is physiologically important for normal development. It is clear however, that LT $\beta$ R signaling has a profound effect in the generation of most LN and intestinal TLO, a role that cannot be bypassed even in the presence of increased levels of TNF $\alpha$ . While increased TNF $\alpha$  cannot compensate for the lack of LT $\beta$ R in terms of TLO and SLO development, it can partially compensate for lack of LT $\beta$ R-signaling in the development of mLN. Finally, as shown here, increased TNF $\alpha$  expression can correct the splenic defects associated with lack of LT $\beta$ R. These results are in agreement with studies that show that TNF $\alpha$  overexpression can correct splenic defects associated with LT $\alpha$ -deficiency<sup>49, 50</sup>. Taken together, the studies highlight a significant cross-talk between these receptor systems for the development and function of lymphoid structures.

In summary: Our results support a model of LN development in TNF/*Rorc*( $\gamma$ )<sup>-/-</sup> mice where increased expression of TNF $\alpha$  by F4/80<sup>+</sup>CD11b<sup>+</sup> cells is sufficient to promote the homeostasis of lymph node stromal cells up to early postnatal life. After birth the recruitment of lymphoid cells and myeloid cells to the anlagen initiates a series of cross-talk interactions with stromal cells through LT $\alpha$ -LT $\beta$ R signaling that induces the expression of chemokines and cell adhesion molecules to organize specific lymphoid areas and attract further cells to form the proper lymph node structure containing HEVs and lymphatic vasculature. Failure of signaling through LT $\beta$ R in early postnatal life results in the collapse of the anlagen of most lymph nodes with the exception of mesenteric LN that present with a disrupted architecture as shown in TNF/LT $\beta$ R<sup>-/-</sup> mice. Thus, our results show that lymphoid tissue organogenesis can occur in the absence of Ror $\gamma$ <sup>+</sup> LTi cells and suggest that interactions between TNF $\alpha$ -expressing myeloid cells and stromal cells have an important role in this process.

## Materials and Methods

### Mice

TNF<sup>ARE/+</sup> and LT $\beta$ R<sup>-/-</sup> mice have been described<sup>12, 23</sup>. Id2<sup>-/-</sup> mice were a generous gift from Dr. Y. Yokota (University of Fukui, Japan)<sup>6</sup>. C57BL6/J and *Rorc*( $\gamma$ )<sup>-/-</sup> mice were obtained from the Jackson Laboratories and bred in our facility. All mice were housed under specific-pathogen-free conditions in individually ventilated cages at the Mount Sinai School of Medicine Animal Facility. All experiments were performed following institutional guidelines. For timed pregnancies, the day of vaginal plug was considered as E0.5.

### Immunostaining

Sections of frozen tissues were subjected to immunofluorescent staining as described<sup>36</sup> (for details see Supplemental Procedures).

### Cell isolation and flow cytometry

The area of the mesentery corresponding to where mLN are found in WT mice was microdissected from *Rorc*( $\gamma$ )<sup>-/-</sup> and TNF/*Rorc*( $\gamma$ )<sup>-/-</sup> animals and analyzed by flow cytometry (for details see Supplemental Procedures).

### In vivo immunization

*TNF/Rorc(γ)<sup>+/+</sup>* and *TNF/Rorc(γ)<sup>-/-</sup>* mice at 6–8 wks were immunized with ovalbumine (OVA, grade V; Sigma-Aldrich) by intragastric gavage of 100μg of OVA + 20μg cholera toxin (List Biological Laboratories) on seven occasions at 7 day intervals. One week after the last immunization, mice were killed and the mLN was collected for cytokine analysis (for details, please see Supplemental Procedures)

### Analysis of mRNA expression

Total RNA was extracted from mesenteric region using the RNeasy mini Kit (Qiagen) as described<sup>36</sup> (for details see Supplemental Procedures).

### Microarray analysis

Microarrays were done with the Illumina TotalPrep™ RNA Amplification Kit (for details please see Supplemental Procedures).

### Statistics

Statistical analyses were performed using GraphPad Prism (GraphPad Software). Differences among means were evaluated by a 2-tailed *t* test. *P* < 0.05 was considered significant. All results shown represent mean ± SEM

### Transplantation of mesenteric lymph node anlagen

*Rorc(γ)<sup>-/-</sup>* mice (6–8 weeks) were anesthetized with ketamine/xylazine solution. A small incision in the skin and peritoneum was made in order to expose the kidney. A slight pressure to both sides of the incision was applied in order to exteriorize the kidney. A small nick in the kidney capsule was created using a 25 gauge needle, and the mesenteric lymph node anlagen was placed into the kidney capsule pocket created in the nick area. The peritoneum and skin were stitched using 5-0 silk sutures w/ a C-6 19mm needle. Formation of lymph nodes under the kidney capsule was assessed by histology 3 weeks after transplantation.

### Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

### Acknowledgments

This work was supported by NIH grant 2 P01 DK072201-06A1 (to SAL). Work in JC laboratory was supported by the EU FP7 integrated project INFLACARE, a BBSRC project grant and by the College of Medical and Dental Sciences-University of Birmingham, UK.

### References


1. Randall TD, Carragher DM, Rangel-Moreno J. Development of secondary lymphoid organs. *Annu Rev Immunol.* 2008; 26:627–650. [PubMed: 18370924]
2. Neyt K, Perros F, GeurtsvanKessel CH, Hammad H, Lambrecht BN. Tertiary lymphoid organs in infection and autoimmunity. *Trends Immunol.* 2012; 33(6):297–305. [PubMed: 22622061]

3. Kim D, Mebius RE, MacMicking JD, Jung S, Cupedo T, Castellanos Y, et al. Regulation of peripheral lymph node genesis by the tumor necrosis factor family member TRANCE. *J Exp Med*. 2000; 192(10):1467–1478. [PubMed: 11085748]
4. Sun Z, Unutmaz D, Zou YR, Sunshine MJ, Pierani A, Brenner-Morton S, et al. Requirement for RORgamma in thymocyte survival and lymphoid organ development. *Science*. 2000; 288(5475): 2369–2373. [PubMed: 10875923]
5. Eberl G, Marmon S, Sunshine MJ, Rennert PD, Choi Y, Littman DR. An essential function for the nuclear receptor RORgamma(t) in the generation of fetal lymphoid tissue inducer cells. *Nat Immunol*. 2004; 5(1):64–73. [PubMed: 14691482]
6. Yokota Y, Mansouri A, Mori S, Sugawara S, Adachi S, Nishikawa S, et al. Development of peripheral lymphoid organs and natural killer cells depends on the helix-loop-helix inhibitor Id2. *Nature*. 1999; 397(6721):702–706. [PubMed: 10067894]
7. Roozendaal R, Mebius RE. Stromal cell-immune cell interactions. *Annu Rev Immunol*. 2011; 29:23–43. [PubMed: 21073333]
8. Kong YY, Yoshida H, Sarosi I, Tan HL, Timms E, Capparelli C, et al. OPGL is a key regulator of osteoclastogenesis, lymphocyte development and lymph-node organogenesis. *Nature*. 1999; 397(6717):315–323. [PubMed: 9950424]
9. Dougall WC, Glaccum M, Charrier K, Rohrbach K, Brasel K, De Smedt T, et al. RANK is essential for osteoclast and lymph node development. *Genes Dev*. 1999; 13(18):2412–2424. [PubMed: 10500098]
10. De Togni P, Goellner J, Ruddle NH, Streeter PR, Fick A, Mariathasan S, et al. Abnormal development of peripheral lymphoid organs in mice deficient in lymphotoxin. *Science*. 1994; 264(5159):703–707. [PubMed: 8171322]
11. Vondenhoff MF, Greuter M, Goverse G, Elewaut D, Dewint P, Ware CF, et al. LTbetaR signaling induces cytokine expression and up-regulates lymphangiogenic factors in lymph node anlagen. *J Immunol*. 2009; 182(9):5439–5445. [PubMed: 19380791]
12. Futterer A, Mink K, Luz A, Kosco-Vilbois MH, Pfeffer K. The lymphotoxin beta receptor controls organogenesis and affinity maturation in peripheral lymphoid tissues. *Immunity*. 1998; 9(1):59–70. [PubMed: 9697836]
13. Crowe PD, VanArsdale TL, Walter BN, Ware CF, Hession C, Ehrenfels B, et al. A lymphotoxin-beta-specific receptor. *Science*. 1994; 264(5159):707–710. [PubMed: 8171323]
14. Cupedo T, Vondenhoff MF, Heeregrave EJ, De Weerd AE, Jansen W, Jackson DG, et al. Presumptive lymph node organizers are differentially represented in developing mesenteric and peripheral nodes. *J Immunol*. 2004; 173(5):2968–2975. [PubMed: 15322155]
15. Okuda M, Togawa A, Wada H, Nishikawa S. Distinct activities of stromal cells involved in the organogenesis of lymph nodes and Peyer's patches. *J Immunol*. 2007; 179(2):804–811. [PubMed: 17617570]
16. Banks TA, Rouse BT, Kerley MK, Blair PJ, Godfrey VL, Kuklin NA, et al. Lymphotoxin-alpha-deficient mice. Effects on secondary lymphoid organ development and humoral immune responsiveness. *J Immunol*. 1995; 155(4):1685–1693. [PubMed: 7636227]
17. Miyawaki S, Nakamura Y, Suzuka H, Koba M, Yasumizu R, Ikehara S, et al. A new mutation, aly, that induces a generalized lack of lymph nodes accompanied by immunodeficiency in mice. *Eur J Immunol*. 1994; 24(2):429–434. [PubMed: 8299692]
18. Weih F, Caamano J. Regulation of secondary lymphoid organ development by the nuclear factor-kappaB signal transduction pathway. *Immunol Rev*. 2003; 195:91–105. [PubMed: 12969313]
19. Rennert PD, Browning JL, Hochman PS. Selective disruption of lymphotoxin ligands reveals a novel set of mucosal lymph nodes and unique effects on lymph node cellular organization. *Int Immunol*. 1997; 9(11):1627–1639. [PubMed: 9418124]
20. Rennert PD, Browning JL, Mebius R, Mackay F, Hochman PS. Surface lymphotoxin alpha/beta complex is required for the development of peripheral lymphoid organs. *J Exp Med*. 1996; 184(5): 1999–2006. [PubMed: 8920886]
21. Rennert PD, James D, Mackay F, Browning JL, Hochman PS. Lymph node genesis is induced by signaling through the lymphotoxin beta receptor. *Immunity*. 1998; 9(1):71–79. [PubMed: 9697837]

22. Alcamo E, Hacohen N, Schulte LC, Rennert PD, Hynes RO, Baltimore D. Requirement for the NF-kappaB family member RelA in the development of secondary lymphoid organs. *J Exp Med*. 2002; 195(2):233–244. [PubMed: 11805150]
23. Kontoyiannis D, Boulougouris G, Manoloukos M, Armaka M, Apostolaki M, Pizarro T, et al. Genetic dissection of the cellular pathways and signaling mechanisms in modeled tumor necrosis factor-induced Crohn's-like inflammatory bowel disease. *J Exp Med*. 2002; 196(12):1563–1574. [PubMed: 12486099]
24. Eberl G, Littman DR. Thymic origin of intestinal alphabeta T cells revealed by fate mapping of RORgammat+ cells. *Science*. 2004; 305(5681):248–251. [PubMed: 15247480]
25. Tsuji M, Suzuki K, Kitamura H, Maruya M, Kinoshita K, Ivanov, et al. Requirement for lymphoid tissue-inducer cells in isolated follicle formation and T cell-independent immunoglobulin A generation in the gut. *Immunity*. 2008; 29(2):261–271. [PubMed: 18656387]
26. Lorenz RG, Chaplin DD, McDonald KG, McDonough JS, Newberry RD. Isolated lymphoid follicle formation is inducible and dependent upon lymphotoxin-sufficient B lymphocytes, lymphotoxin beta receptor, and TNF receptor I function. *J Immunol*. 2003; 170(11):5475–5482. [PubMed: 12759424]
27. Lochner M, Ohnmacht C, Presley L, Bruhns P, Si-Tahar M, Sawa S, et al. Microbiota-induced tertiary lymphoid tissues aggravate inflammatory disease in the absence of RORgamma t and LTi cells. *J Exp Med*. 2011; 208(1):125–134. [PubMed: 21173107]
28. Picarella DE, Kratz A, Li CB, Ruddle NH, Flavell RA. Insulinitis in transgenic mice expressing tumor necrosis factor beta (lymphotoxin) in the pancreas. *Proc Natl Acad Sci U S A*. 1992; 89(21):10036–10040. [PubMed: 1279667]
29. Viejo-Borbolla A, Martin AP, Muniz LR, Shang L, Marchesi F, Thirunarayanan N, et al. Attenuation of TNF-driven murine ileitis by intestinal expression of the viral immunomodulator CrmD. *Mucosal Immunol*. 2010; 3(6):633–644. [PubMed: 20664576]
30. Ivanov, McKenzie BS, Zhou L, Tadokoro CE, Lepelley A, Lafaille JJ, et al. The orphan nuclear receptor RORgammat directs the differentiation program of proinflammatory IL-17+ T helper cells. *Cell*. 2006; 126(6):1121–1133. [PubMed: 16990136]
31. Veiga-Fernandes H, Coles MC, Foster KE, Patel A, Williams A, Natarajan D, et al. Tyrosine kinase receptor RET is a key regulator of Peyer's patch organogenesis. *Nature*. 2007; 446(7135):547–551. [PubMed: 17322904]
32. Honda K, Nakano H, Yoshida H, Nishikawa S, Rennert P, Ikuta K, et al. Molecular basis for hematopoietic/mesenchymal interaction during initiation of Peyer's patch organogenesis. *J Exp Med*. 2001; 193(5):621–630. [PubMed: 11238592]
33. Benezech C, White A, Mader E, Serre K, Parnell S, Pfeffer K, et al. Ontogeny of stromal organizer cells during lymph node development. *J Immunol*. 2010; 184(8):4521–4530. [PubMed: 20237296]
34. White A, Carragher D, Parnell S, Msaki A, Perkins N, Lane P, et al. Lymphotoxin independent and -independent signals regulate stromal organizer cell homeostasis during lymph node organogenesis. *Blood*. 2007; 110(6):1950–1959. [PubMed: 17526859]
35. Walker JA, Barlow JL, McKenzie AN. Innate lymphoid cells--how did we miss them? *Nat Rev Immunol*. 2013; 13(2):75–87. [PubMed: 23292121]
36. Marinkovic T, Garin A, Yokota Y, Fu YX, Ruddle NH, Furtado GC, et al. Interaction of mature CD3+CD4+ T cells with dendritic cells triggers the development of tertiary lymphoid structures in the thyroid. *J Clin Invest*. 2006; 116(10):2622–2632. [PubMed: 16998590]
37. Rangel-Moreno J, Carragher DM, de la Luz Garcia-Hernandez M, Hwang JY, Kusser K, Hartson L, et al. The development of inducible bronchus-associated lymphoid tissue depends on IL-17. *Nat Immunol*. 2011; 12(7):639–646. [PubMed: 21666689]
38. Kojouharoff G, Hans W, Obermeier F, Mannel DN, Andus T, Scholmerich J, et al. Neutralization of tumour necrosis factor (TNF) but not of IL-1 reduces inflammation in chronic dextran sulphate sodium-induced colitis in mice. *Clin Exp Immunol*. 1997; 107(2):353–358. [PubMed: 9030875]
39. van de Pavert SA, Mebius RE. New insights into the development of lymphoid tissues. *Nat Rev Immunol*. 2010; 10(9):664–674. [PubMed: 20706277]
40. Harmsen A, Kusser K, Hartson L, Tighe M, Sunshine MJ, Sedgwick JD, et al. Cutting edge: organogenesis of nasal-associated lymphoid tissue (NALT) occurs independently of lymphotoxin-


- alpha (LT alpha) and retinoic acid receptor-related orphan receptor-gamma, but the organization of NALT is LT alpha dependent. *J Immunol.* 2002; 168(3):986–990. [PubMed: 11801629]
41. Rangel-Moreno J, Moyron-Quiroz JE, Carragher DM, Kusser K, Hartson L, Moquin A, et al. Omental milky spots develop in the absence of lymphoid tissue-inducer cells and support B and T cell responses to peritoneal antigens. *Immunity.* 2009; 30(5):731–743. [PubMed: 19427241]
  42. Lovas A, Weidemann A, Albrecht D, Wiechert L, Weih D, Weih F. p100 Deficiency is insufficient for full activation of the alternative NF-kappaB pathway: TNF cooperates with p52-RelB in target gene transcription. *PLoS One.* 2012; 7(8):e42741. [PubMed: 22880094]
  43. Luther SA, Lopez T, Bai W, Hanahan D, Cyster JG. BLC expression in pancreatic islets causes B cell recruitment and lymphotoxin-dependent lymphoid neogenesis. *Immunity.* 2000; 12(5):471–481. [PubMed: 10843380]
  44. Martin AP, Canasto-Chibuque C, Shang L, Rollins BJ, Lira SA. The chemokine decoy receptor M3 blocks CC chemokine ligand 2 and CXC chemokine ligand 13 function in vivo. *J Immunol.* 2006; 177(10):7296–7302. [PubMed: 17082648]
  45. Marchesi F, Martin AP, Thirunarayanan N, Devany E, Mayer L, Grisotto MG, et al. CXCL13 expression in the gut promotes accumulation of IL-22-producing lymphoid tissue-inducer cells, and formation of isolated lymphoid follicles. *Mucosal Immunol.* 2009; 2(6):486–494. [PubMed: 19741597]
  46. Halle S, Dujardin HC, Bakocevic N, Fleige H, Danzer H, Willenzon S, et al. Induced bronchus-associated lymphoid tissue serves as a general priming site for T cells and is maintained by dendritic cells. *J Exp Med.* 2009; 206(12):2593–2601. [PubMed: 19917776]
  47. GeurtsvanKessel CH, Willart MA, Bergen IM, van Rijt LS, Muskens F, Elewaut D, et al. Dendritic cells are crucial for maintenance of tertiary lymphoid structures in the lung of influenza virus-infected mice. *J Exp Med.* 2009; 206(11):2339–2349. [PubMed: 19808255]
  48. Muniz LR, Pacer ME, Lira SA, Furtado GC. A critical role for dendritic cells in the formation of lymphatic vessels within tertiary lymphoid structures. *J Immunol.* 2011; 187(2):828–834. [PubMed: 21666055]
  49. Alexopoulou L, Pasparakis M, Kollias G. Complementation of lymphotoxin alpha knockout mice with tumor necrosis factor-expressing transgenes rectifies defective splenic structure and function. *J Exp Med.* 1998; 188(4):745–754. [PubMed: 9705956]
  50. Victoratos P, Lagnel J, Tzima S, Alimzhanov MB, Rajewsky K, Pasparakis M, et al. FDC-specific functions of p55TNFR and IKK2 in the development of FDC networks and of antibody responses. *Immunity.* 2006; 24(1):65–77. [PubMed: 16413924]





**Figure 1. TLO are formed in the ileum *TNF/Rorc(γ)*<sup>-/-</sup> mice**


(a) Representative H&E sections of the Ileum of WT, *Rorc(γ)*<sup>-/-</sup>, *TNF/Rorc(γ)*<sup>+/+</sup> and *TNF/Rorc(γ)*<sup>-/-</sup> mice at 16 wks. Notice the presence of inflammatory infiltrates in the ileum of *TNF/Rorc(γ)*<sup>+/+</sup> and *TNF/Rorc(γ)*<sup>-/-</sup> mice. (b) Ileum sections of indicated mice were stained with anti-B220 antibody to visualize B cell aggregates and DAPI for nuclear staining. Small B cell clusters were found in the ileum of WT but were absent in the ileum of *Rorc(γ)*<sup>-/-</sup> mice. (c) Overexpression of TNF induced the formation of large B cell clusters with few T cells in the ileum of *TNF/Rorc(γ)*<sup>+/+</sup> and *TNF/Rorc(γ)*<sup>-/-</sup> mice. Scale bars = 250 μm, n=4/group.



**Figure 2. Increased expression of TNF induces development SLO in the absence of  $ROR\gamma^+$  LTi cells**


(a) Photograph of the mesentery of WT, *Rorc(γt)<sup>-/-</sup>*, *TNF/Rorc(γt)<sup>+/+</sup>*, and *TNF/Rorc(γt)<sup>-/-</sup>* mice. Photograph of axillary (b) and cervical (c) lymph nodes of WT, *TNF/Rorc(γt)<sup>+/+</sup>* and *TNF/Rorc(γt)<sup>-/-</sup>* mice. (d) Incidence of mesenteric (Mes), axillary (Axi), cervical (Cer), brachial (Bra), inguinal (Ing), para aortic (PA), peripancreatic (Pan), popliteal (Pop), mediastinal (Med) lymph nodes, and Peyer's patches (PP) formed in *TNF/Rorc(γt)<sup>-/-</sup>* mice (n = 80). (e) Lymph nodes from *TNF/Rorc(γt)<sup>+/+</sup>* and *TNF/Rorc(γt)<sup>-/-</sup>* mice at 6 wk of age were analyzed by immunostaining. Note segregation of T and B cell areas, presence of PNA<sup>+</sup> HEV and lymphatic vessels, normal distribution of ER-TR7<sup>+</sup> meshwork and CD35<sup>bright</sup> FDC in mesenteric (mLN) and axillary (aLN) lymph nodes of *TNF/Rorc(γt)<sup>-/-</sup>* mice. These features were similar to the ones observed in the mLN of *TNF/Rorc(γt)<sup>+/+</sup>* mice (n = 5 mice/group). Scale bars = 250μm. (f) OVA-specific IgG and IgA measured in the serum of *TNF/Rorc(γt)<sup>+/+</sup>* (n = 5) and *TNF/Rorc(γt)<sup>-/-</sup>* (n = 4) obtained after 5 rounds of immunization. (g) IFN-γ and (h) IL-17 levels in supernatants of cultured MLN cells 7 weeks after OVA immunization.





**Figure 4. F4/80<sup>+</sup>/CD11b<sup>+</sup> cells produce increased levels of TNF in the mLN of *TNF/RORγ<sup>-/-</sup>* mice**

(a) Flow cytometric analysis of TNF production by CD45<sup>-</sup>, CD45<sup>+</sup>F4/80<sup>-</sup> and CD45<sup>+</sup>F4/80<sup>+</sup> cells isolated from the mLN region of *RORγ<sup>-/-</sup>* and *TNF/RORγ<sup>-/-</sup>* mice at P0.5-P1 stage. (b) Analysis of TNF production by CD45<sup>+</sup>F4/80<sup>hi</sup>CD11b<sup>low</sup> and CD45<sup>+</sup>F4/80<sup>low</sup>CD11b<sup>high</sup> cells. Representative plot of two independent experiments, (n=4–5 animals/group).




**Figure 5.** *TNF/Id2<sup>-/-</sup>* mice lack F4/80<sup>+</sup> cells in the mLN anlagen and do not develop SLO mLN region of *TNF/Id2<sup>+/-</sup>* and *TNF/Id2<sup>-/-</sup>* at P0 stained with CD45, F4/80, LYVE-1 antibodies. Notice the absence of F4/80<sup>+</sup> cells in the MLN region of *Id2<sup>-/-</sup>* and *TNF/Id2<sup>-/-</sup>* mice (dashed lines). Representative staining (n = 3/group). Scale bars; 250 $\mu$ m.











**Figure 7. TNF induces stromal cell activation**

(a) FACS analysis of single-cell suspensions from the mLN region of *Rorc(γt)<sup>-/-</sup>* and *TNF/Rorc(γt)<sup>-/-</sup>* mice at E17.5, P0, P1, and P3 showing the increased expression of ICAM-1/VCAM-1 in the CD45 negative stromal cell population. (b) Expression of ICAM-1/VCAM-1 in stromal cell population in the mLN of WT and *TNF<sup>ARE+/+</sup>* mice at P1. (c) Relative number of ICAM-1<sup>hi</sup>VCAM-1<sup>hi</sup> cells in the CD45 negative stromal cells in the mLN region of *Rorc(γt)<sup>-/-</sup>* and *TNF/Rorc(γt)<sup>-/-</sup>* at E17.5, E18.5, and P0 (n = 4 mice/group), P1 (n = 7 mice/group), and P3 (n = 6 mice/group). (d) Expression of CXCL13, CXCL19 and CCL21 in the stromal (CD45 negative) cell population sorted from the mLN region of *Rorc(γt)<sup>-/-</sup>* and *TNF/Rorc(γt)<sup>-/-</sup>* mice at P0.5-P1 (n = 9–10 anlagen/group).




**Figure 8. TNF-driven formation of most SLO is dependent on LTβR signaling**

(a) Mesentery of *LTβR*<sup>-/-</sup> and *TNF/LTβR*<sup>-/-</sup> mice. (b) With the exception of mLN, SLO were absent in *TNF/LTβR*<sup>-/-</sup> mice (n = 16). (c) Abnormal organogenesis of mLN in *TNF/LTβR*<sup>-/-</sup> mice. mLN of *TNF/LTβR*<sup>+/+</sup> and *TNF/LTβR*<sup>-/-</sup> mice were stained with the indicated antibodies. Notice the lack of distinct T and B cell areas, absence of PNAd<sup>+</sup> HEVs and reduced lymphatic vasculature, lack of CD35<sup>+</sup> FDC, and disorganized ER-TR7 stroma in the MLN of *TNF/LTβR*<sup>-/-</sup> mice. Representative staining (n = 5/group). Scale bars = 250μm.



**Figure 9. Hematopoietic cell influx into the mLN anlagen**

Sections of the mLN region of WT, *Rorc(γt)*<sup>-/-</sup> and *TNF/Rorc(γt)*<sup>-/-</sup> (P0 and P5) mice were stained with anti-CD45, -F4/80 and -LYVE-1 antibodies. Representative staining (n = 3/group). Scale bars = 250μm.



**Figure 10. mLN anlagen from WT and  $TNF/Rorc(\gamma)^{-/-}$  mice promote development of LN-like structures when grafted under the kidney capsule**  
 (a) Schematic representation of the transplantation experiment. (b) mLN anlagen isolated from WT (n = 3) and  $TNF/Rorc(\gamma)^{-/-}$  (n = 14) newborn mice were grafted under the kidney capsule of  $Rorc(\gamma)^{-/-}$  mice. Notice normal segregation of T and B cells and development of PNAd<sup>+</sup> HEV from WT and  $TNF/Rorc(\gamma)^{-/-}$  grafts. Scale bars = 250 $\mu$ m.