

STUDY PROTOCOL**Open Access**

Vitamin D supplementation in older people (VDOP): Study protocol for a randomised controlled intervention trial with monthly oral dosing with 12,000 IU, 24,000 IU or 48,000 IU of vitamin D₃

Inez Schoenmakers^{1*}, Roger M Francis², Elaine McColl³, Thomas Chadwick⁴, Gail R Goldberg¹, Christine Harle³, Alison Yarnall^{2,5}, Jennifer Wilkinson³, Jennie Parker³, Ann Prentice¹ and Terence Aspray^{2,5}

Abstract

The randomised, double blind intervention trial 'Optimising Vitamin D Status in Older People' (VDOP) will test the effect of three oral dosages of vitamin D given for one year on bone mineral density (BMD) and biochemical markers of vitamin D metabolism, bone turnover and safety in older people. VDOP is funded by Arthritis Research UK, supported through Newcastle University and MRC Human Nutrition Research and sponsored by the Newcastle upon Tyne Hospitals NHS Foundation Trust.^a

Background: Vitamin D insufficiency is common in older people and may lead to secondary hyperparathyroidism, bone loss, impairment of muscle function and increased risk of falls and fractures. Vitamin D supplementation trials have yielded conflicting results with regard to decreasing rates of bone loss, falls and fractures and the optimal plasma concentration of 25 hydroxy vitamin D (25OHD) for skeletal health remains unclear.

Method/design: Older (≥ 70 years) community dwelling men and women are recruited through General Practices in Northern England and 375 participants are randomised to take 12,000 international units (IU), 24,000 IU or 48,000 IU of vitamin D₃ orally each month for one year starting in the winter or early spring. Hip BMD and anthropometry are measured at baseline and 12 months. Fasting blood samples are collected at baseline and three-month intervals for the measurement of plasma 25OHD, parathyroid hormone (PTH), biochemical markers of bone turnover and biochemistry to assess the dose-response and safety of supplementation. Questionnaire data include falls, fractures, quality of life, adverse events and outcomes, compliance, dietary calcium intake and sunshine exposure.

Discussion: This is the first integrated vitamin D supplementation trial in older men and women using a range of doses given at monthly intervals to assess BMD, plasma 25OHD, PTH and biochemical markers of bone turnover and safety, quality of life and physical performance. We aim to investigate the vitamin D supplementation and plasma 25OHD concentration required to maintain bone health and to develop a set of biochemical markers that reflects the effect of vitamin D on bone. This will aid future studies investigating the effect of vitamin D supplementation on fracture risk. #ISRCTN 35648481 (assigned 16 August 2012), EudraCT 2011-004890-10.

Keywords: Vitamin D supplementation trial, Bone mineral density, 25 hydroxy vitamin D, Parathyroid hormone, Bone markers, Older people

* Correspondence: Inez.Schoenmakers@mrc-hnr.cam.ac.uk

¹MRC Human Nutrition Research, Elsie Widdowson Laboratory, Fulbourn Road, Cambridge CB1 9NL, UK

Full list of author information is available at the end of the article

Background

Vitamin D status and musculoskeletal health

Vitamin D insufficiency is common in older people and may lead to increased bone resorption, bone loss, impairment of muscle function and an increased risk of falls and fractures. The results of clinical trials assessing the effect of vitamin D supplementation on bone loss, falls and fractures have yielded conflicting results and the blood concentration of 25 hydroxy vitamin D (25OHD) required to maintain optimal skeletal health remains unclear. Unfortunately, there is no universal consensus on the optimal way of assessing vitamin D status or on what constitutes vitamin D repletion at different ages throughout life [1].

Vitamin D status is currently assessed by measurement of plasma 25OHD concentration. In the UK, vitamin D deficiency has been defined as a plasma 25OHD less than 25 nmol/L, the upper end of the range at which vitamin D deficiency osteomalacia and rickets have been observed [2,3]. Recent epidemiological and cross-sectional studies have shown that 25OHD concentrations above 25 nmol/L may also be associated with secondary hyperparathyroidism, higher bone resorption, bone loss, impaired muscle function and a greater risk of falls and fragility fractures [4-8] when compared to concentrations of 40 to 75 nmol/L and above.

Vitamin D status and requirements

A plasma 25OHD concentration of 40 to 50 nmol/L has been suggested as a cut-off for vitamin D insufficiency [2,9-11]. However, some experts have advocated higher concentrations of 25OHD (75 to 150 nmol/L) for the maintenance of bone health and other non-skeletal benefits [12-17]. Few individuals in the UK achieve these concentrations of 25OHD in winter and spring, when more than 80% of 45-year-old men and women have a plasma 25OHD concentration less than 75 nmol/L [18-20]. Vitamin D status is poorer in older people, where approximately 12% of people living in private households and 30% of care home residents have a plasma 25OHD concentration below 25 nmol/L throughout the year, reflecting their reduced exposure to sunlight, low dietary intake of vitamin D and potentially their higher expenditure of vitamin D [14,18,21].

The dietary Reference Nutrient Intake (RNI) for vitamin D in the UK for people over 65 years is currently 400 International Units (IU) (= 10 µg) per day. In the US, the Estimated Average Requirement (EAR) and Recommended Dietary Allowance (RDA) for those over 70 years of age is 600 (15 µg) and 800 IU (20 µg) per day, respectively [11,22]. It is recognised that supplementation is generally necessary to achieve these intakes [23]. Despite this advice, vitamin D supplementation is

uncommon in older people, even in care home residents [18,24-26].

Vitamin D supplementation and fracture risk

Several placebo-controlled intervention trials have examined the effect of vitamin D supplementation on fracture risk. Some meta-analyses suggest that vitamin D supplementation doses of 800 to 1000 IU daily decrease the incidence of falls and fractures in older women, whereas lower doses are ineffective [27-29]. In contrast, other meta-analyses, including the latest Cochrane Review [30], indicate that vitamin D supplementation does not decrease fracture risk, unless combined with calcium, when it reduces the risk of hip and other non-vertebral fractures in care home residents [31,32]. One randomised controlled trial of a single annual oral dose of 500,000 IU vitamin D showed an increase in falls and fractures, particularly in the first three months after administration [33]. Other studies using 300,000 or 600,000 IU per annum have shown mixed results: 100,000 IU administered at four-month intervals was associated with a decrease in fracture risk, whereas 300,000 IU of vitamin D₂ by i.m. injection or 150,000 IU administered orally every three months was associated with no benefit [34-36]. These results suggest that not only the dose but also the frequency of administration may determine the risk and benefit for falls and fractures. The mechanisms of this association remain unclear and are likely to be multifactorial.

The evidence for the effects of vitamin D supplementation on BMD or the prevention of bone loss in older people is conflicting [37,38]. The differences in the reported efficacy of vitamin D may be because the dose of vitamin D in several trials may have been too low or compliance and persistence with supplementation may have been poor, resulting in a failure to achieve a high enough plasma 25OHD to reduce bone loss and obtain an anti-fracture benefit [13,39-41]. Alternatively, the participants in some trials may have been vitamin D replete [41], even if their baseline plasma 25OHD concentrations were lower than those advocated by North American experts [12,15,16]. Moreover, differences in reported efficacy may have been due to differences in the study population, the dose of vitamin D, the route and frequency of administration and/or the use of different forms of vitamin D (vitamin D₂ and vitamin D₃).

Several studies have indicated that the achieved plasma 25OHD concentration, rather than the vitamin D dose given, predicts the effect on bone health [42]. However, measurement of plasma 25OHD, parathyroid hormone (PTH) and other potential markers of vitamin D status have only been performed in a small minority of participants in these intervention studies [41], limiting the ability to explore the relationship between plasma 25OHD

concentrations achieved and bone loss and fracture prevention.

Vitamin D status markers for bone health

Before performing further studies of the effect of vitamin D supplementation on the risk of major outcomes such as fragility fractures or cardiovascular disease [12,13,43,44], there is a need to understand better the relationship between vitamin D intake, plasma 25OHD and bone health in older people and to develop better functional markers of vitamin D status, reflecting vitamin D function at the level of the target organ [3,45]. These functional markers will be instrumental in establishing clearer criteria for vitamin D sufficiency and insufficiency and help to clarify the optimal dose and frequency of vitamin D supplementation. Potential markers of vitamin D status have been reviewed [3]. Plasma PTH has been proposed as a functional marker, because an elevated concentration is a recognised risk factor for bone loss and fractures in older people [6-9,13,46-48], which can potentially be modified by vitamin D supplementation [49,50]. The circulating concentration of 25OHD below which PTH increases outside the normal range may be used to establish a threshold value for vitamin D insufficiency [1]. However, PTH varies within and between people at any given concentration of 25OHD [18,51], as PTH depends on many factors other than vitamin D status [52]. These include stage of life, ethnic background, dietary calcium and phosphate intake, time of day, renal function, serum magnesium, physical inactivity and medication use [3,14,53-55].

Aim

The Optimising Vitamin D Status in Older People (VDOP) trial aims to examine the relationship between vitamin D supplementation at a range of doses and the change in bone mineral density (BMD) in older people living in private households in the North East of England. Vitamin D₃ supplementation is given orally each month, for a year (12,000 IU/month, 24,000 IU/month or 48,000 IU/month, equivalent to 400 IU/day, 800 IU/day and 1,600 IU/day, respectively). We will explore the relationship between the change in BMD and the achieved plasma 25OHD and PTH over 12 months. As BMD is the result of the balance between bone formation and resorption, the effect of the supplement on the change in markers of bone turnover will be explored. Plasma C-terminal telopeptide (CTX), bone specific alkaline phosphatase (bone ALP) and N-terminal propeptide of procollagen type I (P1NP) will be measured, as it has been suggested that these may be the best surrogate bone turnover markers for the effect of pharmacological intervention on fracture incidence [56]. We will also investigate the effect of duration of treatment and season on the dose-response.

Rationale for dosage and dosing regimen

Two of the dosages applied in this trial correspond to the current Scientific Advisory Committee on Nutrition (SACN) RNI (400 IU/day) and the Institute of Medicine (IOM) RDA (800 IU/day) for this age group [2,11]. The highest dose is twice the IOM RDA and well below the IOM Tolerable Upper Intake Level (TUIL) defined by the IOM (4,000 IU/day) [11].

This trial uses a monthly dosage scheme as this is expected to result in improved compliance compared to daily or weekly dosages [57]. Compliance will be encouraged through administration of the dose under direct supervision on four occasions and taking the other doses will be prompted by telephone calls by staff from the research centre.

Methods/design

Study design

The VDOP trial is a single centre parallel group, randomised, double blind interventional trial testing the effects of three dosages of oral vitamin D₃ given each month to men and women 70 years old and older for a year. The primary outcome is the change in BMD at the hip (total hip BMD) from baseline to 12 months post intervention.

Recruitment

Community dwelling men and women 70-years-old and older, are recruited through General Practices (GP) in Tyneside, Wearside and Northumberland, participating in the Northern and Yorkshire Primary Care Research Network (PCRN-NY). Potentially eligible participants are identified from GP practice registers, after initial screening against inclusion and exclusion criteria on electronic medical records. An invitation letter is sent to potential participants, including details of the study and a pre-paid response slip. Those expressing an interest are contacted by the research team, further screening performed to ensure that the participant is eligible and an invitation to participate is then offered, after any discussions the potential participant might wish. If still interested, an appointment is made for the first study visit, when written informed consent is obtained, and a blood sample is collected to perform the final assessment of eligibility (screening for biochemical exclusion criteria: see below). If eligible, baseline visit measurements and questionnaires are completed.

Inclusion criteria

Inclusion criteria are as follows:

1. Ambulatory, community dwelling men and women 70 years old and older
2. Individuals capable of giving informed consent on their own behalf

3. Individuals willing to attend the Study Centre (The Clinical Ageing Research Unit (CARU)) on six occasions and to be contacted by telephone at monthly intervals between study visits over twelve months

Exclusion criteria

Exclusion criteria are as follows:

1. Current antiresorptive or anabolic treatment for osteoporosis
2. Treatment with bisphosphonates for osteoporosis in the past two years
3. Current supplement use of vitamin D (>400 IU/day) or calcium (>500 mg/day) (including use of over-the-counter preparations)
4. Fragility fracture in the previous six months
5. Known primary hyperparathyroidism
6. History of renal stones
7. Hip replacement

Biochemical:

1. Hypercalcaemia (albumin-adjusted serum calcium >2.60 mmol/L)
2. Renal impairment (Stage 4 or 5 Chronic Kidney Disease: estimated glomerular filtration rate (eGFR) below 30 ml/min/1.73m² calculated using the modification of diet in renal disease (MDRD) formula.

There will be no limits to concomitant care or medication use other than defined in the inclusion and exclusion criteria. Participation in the VDOP trial is discontinued when during the course of the trial conditions develop where vitamin D supplementation is contra-indicated.

Randomisation, Investigational Medicinal Product (IMP) and IMP administration

Participants are randomised using a computerised system in a 1:1:1 ratio to three treatment arms (12,000 IU, 24,000 IU or 48,000 IU vitamin D₃ once monthly for 12 months) by a statistician within the Newcastle Clinical Trials Unit (CTU) not otherwise engaged in the study. Permuted blocks of variable length are used to reduce the risk of breach of concealment of allocation. Allocation lists are provided to Apotek Produktion & Laboratorier AB (APL; Stockholm, Sweden) which produces numbered packs of the Investigational Medicinal Product (IMP) per participant for shipping to the Newcastle upon Tyne Hospitals NHS Foundation Trust pharmacy for dispensing. The allocation is not stratified and allocation status is kept concealed from participant

and research staff unless code break is required for safety reasons.

The IMP is composed of vitamin D₃ dissolved in Miglyol as Vigantol® (Merck Sereno GmbH, Darmstadt Germany). One ml of Vigantol® solution contains 0.5 mg cholecalciferol, equivalent to 20,000 IU vitamin D₃. Blinding of Vigantol® oil across the three doses (12,000 IU, 24,000 IU and 48,000 IU) is achieved by further dilution, using Miglyol® 812 oil (Sasol; The Warner Graham Company, Cockeysville, MD USA) and the stabilising agent butylhydroxianisole, both free of vitamin D₃. For the highest dose, only the stabilising agent is added. All three doses of vitamin D are dispensed in 10 ml amber glass bottles (target fill volume of 3 ml including 0.6 ml overage).

MODEPHARMA (London, UK) is responsible for arranging the manufacture of the IMP and project management and assistance relating to the IMP. The formulation, manufacture and release of the IMP is undertaken by APL, a Swedish clinical supplies company licensed for the manufacture and release of IMPs for clinical trial use in the EU.

The IMP is administered orally under direct supervision of a research nurse at baseline and at the third, sixth and ninth month study visits. At the end of each study visit, participants are given two bottles of the study IMP for the two subsequent months before their next scheduled visit to CARU, together with written instructions on how to take the IMP. On months when the participant is not attending the clinic, they are contacted by telephone, to remind them to take the IMP. Participants are directed to return all IMP bottles in their original packaging to the research team. Researchers return these to the trial pharmacist and all returned or unused study IMPs are documented in the Pharmacy File and then destroyed according to Pharmacy Standard Operating Procedures.

Measurements

An overview of the timing of measurements, sample collections and data collection is given in Table 1. Total hip BMD is measured using Dual-Energy X-ray Absorptiometry (DXA) (i-Lunar DXa, GE Healthcare, Madison Wisconsin USA) at baseline and 12 months. The densitometer is calibrated daily prior to participant scanning. The total radiation dose (0.04 mSv) is equivalent to having a chest X-ray. Height and weight, demispan, waist-hip ratio and the Timed Up and Go (TU&G) test are determined [58] and the participant's medical history (including medication and supplement use) reviewed. A standardised physical exam including the assessment of vital signs, blood pressure, and questionnaires assessing bone pain, fracture risk and falls history are performed (adapted from standard clinical care questions and the

Table 1 Schedule of study procedures for VDOP

	Phone call 0	Visit 0	Visit 1	Phone call 1	Phone call 2	Visit 2	Phone call 3	Phone call 4	Visit 3	Phone call 5	Phone call 6	Visit 4	Phone call 7	Phone call 8	Visit 5
Venue	Pre- Screening	Pre- Screening	Screening	Baseline visit	FUP ^a 1	FUP ^a 2	FUP ^a 3	FUP ^a 4	FUP ^a 5	FUP ^a 6	FUP ^a 7	FUP ^a 8	FUP ^a 9	FUP ^a 10	FUP ^a 11
Month	PIC ^a	Home	CARU ^a	CARU ^a	Home	Home	CARU ^a	Home	Home	CARU ^a	Home	Home	CARU ^a	Home	CARU ^a
Identification	X														
Pre-Screening		X													
Informed consent			X												
Venipuncture (screening)				X											
Randomisation					X										
Medical history, Demispan, falls history and fracture risk						X									
Demographics				X											
TU&G ^a test grip strength					X										X
DXA (total hip) weight height				X											X
Venipuncture (safety and biochemistry)			X			X			X			X			X
Urine sample				X			X			X			X		X
Physical exam and vital signs				X			X			X			X		X
Quality of Life			X				X			X			X		X
CalQuest			X							X					X
Sunshine exposure			X				X			X			X		X
Falls diaries		-----	-----	→	X	-----	→	X	-----	→	X	-----	→	X	-----
Administer treatment		X	X	X	X	X	X	X	X	X	X	X	X	X	
Check drug adherence			X	X	X	X	X	X	X	X	X	X	X	X	

^aCARU Clinical Ageing Research Unit, FUP follow-up visit, PIC Patient Identification Centre, TU&G Time up and Go.

Fracture Risk Assessment (FRAX) tool [59]). Blood and timed urine samples are collected after an overnight fast between 8:30 and 11:30 AM. After venipuncture, participants are offered breakfast. Questionnaires are used to collect data on demographics, quality of life (WHOQol-BREF UK version and UK WHOQOL-old; [60,61]), dietary calcium and vitamin D intake (adapted from Calquest; [62]) and sunshine exposure in the UK and during overseas travel (adapted from [63]). Participants are also asked to keep a three-month prospective falls diary. At each visit, information is collected on participant-reported adverse events (AEs), serious adverse events (SAEs) and suspected unexpected serious adverse reactions (SUSARs). In addition, at every visit, safety is assessed specifically by performing full blood count, serum albumin adjusted calcium, creatinine and liver function. Information on clinical fractures during the study is recorded as a safety measure.

Biochemical analyses

Blood samples are collected into tubes containing ethylenediaminetetraacetic acid (EDTA), lithium heparin (LH) or serum-separation gel (BD Diagnostics, Oxford, UK) and one EDTA and one LH tube is immediately placed on ice. Samples for clinical biochemistry and safety measurements are sent to Newcastle upon Tyne hospitals NHS Foundation Trust (NUTH) laboratories and analysed immediately. These include eGFR, full blood count (FBC), urea and electrolytes (U&E) and liver function tests (LFT) which include: serum sodium, potassium, urea, total protein, total bilirubin, alanine transaminase, calcium, phosphate, magnesium, albumin, alkaline phosphatase (ALP) and creatinine. The remaining blood samples are separated within 30 minutes of collection in a refrigerated centrifuge at 1,800 g for 20 minutes and stored at -80°C prior to analysis at MRC Human Nutrition Research, Cambridge, UK. Blood pellets are stored at -20°C for potential genetic analyses. Collected urine is mixed thoroughly and aliquots are analysed for calcium, phosphate and creatinine (Cr) immediately at NUTH laboratories.

All assays are performed in duplicate with the exception of PTH. EDTA plasma is used for analysis of PTH by immunoassay (Immulite, Siemens Healthcare Diagnostics Ltd, Camberley, UK). LH plasma is used for the measurement of 25OHD, BAP (both DiaSorin-Liaison, Stillwater, MN, USA) and CTX (IDS Ltd, Tyne and Wear, UK) according to standardised protocols [64]). Assay performance is monitored using kit and in-house controls and under strict standardisation according to ISO 9001:2000. Quality assurance of 25OHD and PTH assays are performed as part of the Vitamin D External Quality Assessment Scheme (www.deqas.org), and the National External Quality Assessment Scheme (www.ukneqas.org.uk).

Data entry and analyses of questionnaires

An electronic case report form (eCRF) is created for each participant using a standardised computerised system (InferMed's MACRO) at CTU. Biochemistry data are generated electronically and collated in a standardised format. Dietary contents of calcium and vitamin D are derived from food tables. Quality checks of data entry and merging are performed.

Sample size and power calculation

The sample size calculation was based on an unpaired two-group comparison of the absolute change in BMD at the hip within individual participants over the 12 months of the study. This illustrates the power for a primary comparison of interest between any two chosen arms in the three-arm design of this study; given the choices detailed below, the calculations would be identical for any selected pair of arms. Absolute change is used, rather than percentage change, as it is statistically more efficient [65].

Data from a population based study [50] showed that standard deviation (SD) in absolute change of BMD at the hip over 12 months in women 60 to 70 years old, living in Aberdeen, is between 0.012 and 0.015 after supplementation with a placebo or two vitamin D dosages (0, 400 IU and 1,000 IU). As this SD is relatively stable in value but does not include values for two of our three dosage levels, the conservative choice was made to use the largest SD estimate and to apply the two-sample t-test, a two-sided significance level of 0.05 and a power of 80% in the calculations. Aiming to be able to detect a difference of 0.006 g/cm² (equivalent to a moderate standardised effect size of 0.4) between two arms, 100 participants per arm would be needed. Making an allowance for 20% attrition at the 12 month follow-up results in 125 participants per arm (375 in total) being required. This sample size would give 90% power to detect a difference of 0.007 g/cm² between a pair of arms. We believe that a difference in BMD of the order of 0.006 g/cm² against the background of the anticipated yearly decrease in BMD in an untreated population would be clinically significant [50].

Statistical analyses

The primary outcome of this study is the 12 month change in BMD at the hip (total hip BMD).

The secondary outcome measures are the changes in plasma 25OHD concentration and PTH, so as to explore their use as functional markers in relation to change in BMD and biochemical markers of bone turnover. Data are also being obtained on the proportion of participants in each treatment group who fall, the number of falls, quality of life and safety.

Analyses are performed according to a formal pre-planned documented analysis plan. Analysis will be carried

out primarily on an intention-to-treat basis although other exploratory analyses may also be considered. There are no planned interim analyses.

Data with missing observations due to loss to follow-up will be examined to determine both the extent of missingness and whether it is missing at random or is selective. If data are missing to a sufficient extent, the use of appropriate multiple imputation techniques will be considered to allow for this in the analysis.

Analysis of the primary outcome measure (12 month change in BMD at the hip) will be performed using Analysis of Covariance (ANCOVA) to compare the response in the three treatment groups while allowing for the effects of covariates. Baseline BMD at the hip, PTH and 25OHD measures will be included in addition to measures of kidney function, BMI, gender and age. Interactions will also be explored. The inclusion of baseline values as covariates will also enable the examination of possible interactions between effects observed and these values.

A number of secondary analyses will also be undertaken. Alternative clinical indicators (such as change in plasma 25OHD, PTH, falls, physical performance, quality of life measures and other biochemical markers) will be compared between the three treatment groups, using similar techniques to those described above.

The relationship of PTH as a dependent variable with 25OHD and kidney function as independent variables will be examined using regression methods. Further exploratory regression models will explore the relationships of 25OHD and PTH with change in BMD and biochemical markers of bone turnover. The relationship between 25OHD and BMI will also be explored using similar regression methods.

The relationship of 25OHD at different time points as a dependent variable, with season and duration of supplementation as independent variables, will be examined using regression or repeated measures methods.

Ethical considerations

This study has received a favourable opinion from the Sunderland Research Ethics Committee (REC, 12/NE/0050) and R and D approval from the Newcastle upon Tyne Hospitals NHS Foundation Trust (sponsor). The conduct of this study is in accordance with the recommendations for physicians involved in research on human subjects adopted by the 18th World Medical Assembly, Helsinki 1964 and later revisions.

The study involves procedures that may be unfamiliar to participants and are not part of their standard medical care. These procedures are explained in information sheets provided to all potential participants prior to their screening visit and written informed consent is obtained prior to any study procedures. The consent form

specifies the separate study components and includes an opt in/opt out section for genetic analyses and permission to be approached for potential affiliated studies. Participants who lack capacity to consent for themselves are excluded from this study. In the unlikely event that a participant loses capacity after they have provided consent to take part, their case will be reviewed by the Chief Investigator and the legal representative of the participant to decide whether to continue in the study.

The hip BMD DXA measurement involves a low dose of radiation (see measurements).

The latest Cochrane Review suggests that vitamin D supplementation may be associated with a small increased risk of gastrointestinal side effects, hypercalcaemia, renal stones and renal impairment, particularly when given with additional calcium or when active metabolites of vitamin D are used [32]. However, the doses to be used in this study are not generally associated with adverse events [11,22]. As we are using vitamin D alone, without additional calcium supplementation, we do not anticipate that this will pose a significant risk to participants. The highest dose of vitamin D₃ used in this study is less than half the TUIL recently advised by the IOM [11,22].

Reporting and evaluation of adverse events, serious adverse events and suspected unexpected serious adverse reactions

The development or progression of age-related medical conditions is anticipated to occur spontaneously during the study. On the basis of the latest Cochrane Review a small increased risk of side effects may be expected (see above). Any AEs, SAEs or SUSARs are verified against treatment notes/medical records and causality assessed using standard criteria according to Good Clinical Practice (GCP) guidelines [66]. All adverse events judged by the Trial Steering Committee (TSG) as having a reasonable suspected causal relationship to the IMP are considered to be adverse reactions and are reported to the sponsor, the Medicines and Healthcare products Regulatory Agency (MHRA) and REC in accordance with current legislation.

Data protection

Data are handled, computerised and stored in accordance with the Data Protection Act 1998, guidelines for GCP [66,67] and local policy. Data collected on paper are entered on a secure validated clinical data management system (InferMed's MACRO). No participant identifiable data will leave the study site. To preserve anonymity, any data or samples leaving the site will identify participants by a unique study identification code (linked in anonymised form). All identifiable study records and Investigator Site Files are kept at the site in a locked filing cabinet with restricted access and on encrypted and/or pass-word protected computers.

Monitoring

The daily management lies with the Trial Management Group (TMG) which comprises Dr Terry Aspray (Chief Investigator (Chair)), Professor Roger Francis (Co-Investigator), Dr Tom Chadwick (Statistician), Dr Jennifer Wilkinson (Senior Trial Manager), Christine Harle (Trial Manager), Jennie Parker (Assistant Trial Manager), Ian Campbell (Assistant Director of Pharmacy) and Katie Argo (Clinical Trials Pharmacist). The UK CRC registered Newcastle Clinical Trials Unit (NCTU) has delegated authority from the sponsor for obtaining ethical and regulatory approvals, trial management and ensuring that the study is conducted in accordance with the principles of GCP and pharmacovigilance.

The TSG consists of the TMG and Professor David Reid (Independent Chair), Professor Elaine McColl (Co-investigator), Dr Gail Goldberg (Co-investigator), Dr Ann Prentice (Co-investigator), Dr Inez Schoenmakers (Co-investigator), Dr Elina Hypponen (Independent member), Professor Sue Lanham-New (Independent member) and two Lay members (Patient support group representatives).

The Data Monitoring and Ethics Committee (DMEC) comprises Professor Arduino Mangoni (Chair), Dr Nicola Peel and Dr Barbara Gregson (trialist and statistician).

Exit strategy

Participants in this study will discontinue the study supplementation after 12 months of supplementation. They are then free to use over the counter supplements of vitamin D.

All clinically relevant findings will be reported to the participants' GPs by the CI (Dr Terry Aspray) who will also offer advice on clinical management of bone-related health issues through the Freeman Hospital Musculoskeletal Unit.

Discussion and potential impact

This study will provide clarification of the dose of vitamin D required to prevent bone loss in older people in the UK and will influence the formulation of dietary recommendations of vitamin D at the population level. This study is novel and clinically relevant, as it will compare the effect of three different doses of vitamin D on BMD and plasma 25OHD, PTH and biochemical markers of bone turnover in both men and women. This study evaluates the effects of vitamin D as mono-therapy rather than combined with calcium. The data complement current evidence on the balance of risks in using vitamin D with or without calcium supplementation [68]. The study also offers the opportunity to evaluate the impact of different dosages of vitamin D supplement on structural (DXA Scanning) and biochemical markers which allow a better definition of Vitamin D sufficiency for bone health and to determine an adequate vitamin D

intake to maintain musculoskeletal function. We aim to develop a set of biochemical markers that reflect the effect of vitamin D intervention on bone. This will aid future larger studies investigating the effect of vitamin D supplementation on fracture risk.

Although the study is primarily powered to detect an effect of vitamin D supplementation on BMD, it will also provide data on the effect of vitamin D supplementation on other functional outcomes, such as quality of life, physical performance and falls. Further, safety data are collected and analysed for the effects of the different dosages, temporal trends and the cumulative dosage. In addition, this study will provide data on the effect of season on vitamin D status and other markers and improve knowledge of the role of monthly (rather than daily) dosing with vitamin D.

Trial status

The VDOP study commenced recruitment in November 2012; recruitment is scheduled to be completed at end of June 2013, with follow-up for 12 months thereafter.

Endnotes

^aFreeman Hospital, Freeman Road, High Heaton, Newcastle-upon-Tyne, Tyne and Wear, NE7 7DN, United Kingdom.

Abbreviations

25OHD: 25 hydroxy vitamin D; AE: Adverse event; ALP: Alkaline phosphatase; APL: Apotek produktions & laboratorier; AR: Adverse reaction; ARUK: Arthritis research UK; BMD: Bone mineral density; bALP: Bone alkaline phosphatase; CARU: Clinical ageing research unit; CTU: Clinical trials unit; CTX: C-terminal telopeptide Type 1; DMEC: Data monitoring & ethics committee; DXA: Dual energy X-ray absorptiometry; EAR: Estimated average requirement; e-CRF: Electronic case report form; EDTA: Ethylenediaminetetraacetic acid; eGFR: Estimated glomerular filtration rate; FBC: Full blood count; GCP: Good clinical practice; GP: General practitioner; IMP: Investigational medicinal product; IOM: Institute of medicine; IU: International unit; LFT: Liver function test; LH: Lithium heparin; MDRD: Modification of diet in renal disease; MHRA: Medicines and healthcare products regulatory agency; PCRN-NY: Primary care research network – Northern and Yorkshire; PIC: Patient identification centre; P1NP: Terminal propeptide of procollagen type I; PTH: Parathyroid hormone; QC: Quality control; QP: Qualified person; RDA: Recommended daily allowance; REC: Research ethics committee; RN: Reference nutrient intake; SACN: Scientific advisory committee on nutrition; SAE: Serious adverse event; SAR: Serious adverse reaction; SUSAR: Suspected unexpected serious adverse reaction; TMG: Trial management group; TSG: Trial steering group; TU&G: Time up and go; TUIL: Tolerable upper intake level; U&E: Urea and electrolytes; Vitamin D₂: Ergocalciferol; Vitamin D₃: Cholecalciferol; VDOP: Optimising vitamin D status in older people; WHO QOL-OLD: WHO Quality of life measure for older people.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

IS, RMF, EMCC, TC, GG, AP, TA designed the trial; IS, RMF, EMCC, TC, CH, AY, JW, JP and TA are responsible for the conduct of the trial; IS and TA drafted the paper; All authors read and approved the final manuscript.

Funding

This trial is supported by a Clinical Studies Grant (MP/ID19544) from Arthritis Research UK. This work is partly supported by Newcastle University and funding from the core programme of the MRC Nutrition and Bone Health

Group at MRC Human Nutrition Research funded by the UK MRC, grant code U10596037.

Role of the funder

The study design has been internationally peer-reviewed on behalf of the Arthritis Research UK as part of the funding decision process. Reviewer's recommendations were taken into consideration in the design of the trial. Arthritis Research UK has no involvement in the analyses or publication of the results of the trial.

Role of the sponsor

The sponsor has responsibility for the conduct of the trial according to guidelines laid down in GCP, GRP, the Data Protection Act and the Declaration of Helsinki. The sponsor is also responsible for Pharmacovigilance. These responsibilities are delegated to the Principle Investigator Dr T. Aspray and Newcastle CTU.

Author details

¹MRC Human Nutrition Research, Elsie Widdowson Laboratory, Fulbourn Road, Cambridge CB1 9NL, UK. ²Institute for Ageing and Health, Newcastle University, Campus for Ageing and Vitality Nuns Moor, Newcastle upon Tyne NE4 5PL, UK. ³Newcastle Clinical Trials Unit, The Medical School, 4th Floor, William Leech Building Framlington Place, Newcastle upon Tyne NE2 4HH, UK. ⁴Institute of Health and Society, Newcastle University, Baddiley-Clark Building Richardson Road, Newcastle upon Tyne NE2 4 AX, UK. ⁵Bone Clinic Musculoskeletal Unit, Freeman Hospital, Newcastle upon Tyne NE7 7DN, UK.

Received: 7 June 2013 Accepted: 6 September 2013

Published: 17 September 2013

References

- Francis RM: What do we currently know about nutrition and bone health in relation to United Kingdom public health policy with particular reference to calcium and vitamin D? *Br J Nutr* 2008, **99**:155–159.
- Nutrition TSACo: Update on Vitamin D. London: The Stationery Office; 2007.
- Prentice A: Vitamin D deficiency: a global perspective. *Nutr Rev* 2008, **66**:S153–S164.
- Bischoff-Ferrari HA, Conzelmann M, Stahelin HB, Dick W, Carpenter MG, Adkin AL, Theiler R, Pfeifer M, Allum JH: Is fall prevention by vitamin D mediated by a change in postural or dynamic balance? *Osteoporos Int* 2006, **17**:656–663.
- Bischoff-Ferrari HA, Orav EJ, Dawson-Hughes B: Additive benefit of higher testosterone levels and vitamin D plus calcium supplementation in regard to fall risk reduction among older men and women. *Osteoporos Int* 2008, **19**:1307–1314.
- Sahota O, Gaynor K, Harwood RH, Hosking DJ: Hypovitaminosis D and 'functional hypoparathyroidism'—the NoNoF (Nottingham Neck of Femur) study. *Age Ageing* 2001, **30**:467–472.
- Sahota O, Mundey MK, San P, Godber IM, Lawson N, Hosking DJ: The relationship between vitamin D and parathyroid hormone: calcium homeostasis, bone turnover, and bone mineral density in postmenopausal women with established osteoporosis. *Bone* 2004, **35**:312–319.
- Rejnmark L, Vestergaard P, Brot C, Mosekilde L: Increased fracture risk in normocalcemic postmenopausal women with high parathyroid hormone levels: a 16-year follow-up study. *Calcif Tissue Int* 2011, **88**:238–245.
- Lips P: Vitamin D deficiency and secondary hyperparathyroidism in the elderly: consequences for bone loss and fractures and therapeutic implications. *Endocr Rev* 2001, **22**:477–501.
- Boonen S, Rizzoli R, Meunier PJ, Stone M, Nuki G, Syversen U, Lehtonen-Veromaa M, Lips P, Johnell O, Reginster JY: The need for clinical guidance in the use of calcium and vitamin D in the management of osteoporosis: a consensus report. *Osteoporos Int* 2004, **15**:511–519.
- IoMotN A: Dietary Reference Intakes for Calcium and Vitamin D. Washington DC: The National Academies Press; 2011.
- Holick MF: Optimal vitamin D status for the prevention and treatment of osteoporosis. *Drugs Aging* 2007, **24**:1017–1029.
- Bischoff-Ferrari HA, Giovannucci E, Willett WC, Dietrich T, Dawson-Hughes B: Estimation of optimal serum concentrations of 25-hydroxyvitamin D for multiple health outcomes. *Am J Clin Nutr* 2006, **84**:18–28.
- Vieth R, Ladak Y, Walfish PG: Age-related changes in the 25-hydroxyvitamin D versus parathyroid hormone relationship suggest a different reason why older adults require more vitamin D. *J Clin Endocrinol Metab* 2003, **88**:185–191.
- Dawson-Hughes B, Heaney RP, Holick MF, Lips P, Meunier PJ, Vieth R: Estimates of optimal vitamin D status. *Osteoporos Int* 2005, **16**:713–716.
- Vieth R, Bischoff-Ferrari H, Boucher BJ, Dawson-Hughes B, Garland CF, Heaney RP, Holick MF, Hollis BW, Lamberg-Allardt C, McGrath JJ, Norman AW, Scragg R, Whiting SJ, Willett WC, Zittermann A: The urgent need to recommend an intake of vitamin D that is effective. *Am J Clin Nutr* 2007, **85**:649–650.
- Grant WB, Cross HS, Garland CF, Gorham ED, Moan J, Peterlik M, Porojnicu AC, Reichrath J, Zittermann A: Estimated benefit of increased vitamin D status in reducing the economic burden of disease in western Europe. *Prog Biophys Mol Biol* 2009, **99**:104–113.
- Bates CJ, Carter GD, Mishra GD, O'Shea D, Jones J, Prentice A: In a population study, can parathyroid hormone aid the definition of adequate vitamin D status? A study of people aged 65 years and over from the British National Diet and Nutrition Survey. *Osteoporos Int* 2003, **14**:152–159.
- Ruston D, Hoare L, Henderson L, Gregory J, Bates CJ, Prentice A, Birch M, Swan G, Farron M: *The National Diet and Nutrition Survey: Adults Aged 19 to 64 Years. Nutritional Status (Anthropometry and Blood Analytes), Blood Pressure and Physical Activity*. London: The Stationery Office; 2003.
- Hypponen E, Power C: Hypovitaminosis D in British adults at age 45 y: nationwide cohort study of dietary and lifestyle predictors. *Am J Clin Nutr* 2007, **85**:860–868.
- Hirani V, Primatesta P: Vitamin D concentrations among people aged 65 years and over living in private households and institutions in England: population survey. *Age Ageing* 2005, **34**:485–491.
- Ross AC, Manson JE, Abrams SA, Aloia JF, Brannon PM, Clinton SK, Durazo-Arvizu RA, Gallagher JC, Gallo RL, Jones G, Kovacs CS, Mayne ST, Rosen CJ, Shapses SA: The 2011 report on dietary reference intakes for calcium and vitamin D from the Institute of Medicine: what clinicians need to know. *J Clin Endocrinol Metab* 2011, **96**:53–58.
- Department of Health: *Nutrition and Bone Health: with particular reference to calcium and vitamin D*. London: The Stationery Office; 1998.
- Aspray TJ, Stevenson P, Abdy SE, Rawlings DJ, Holland T, Francis RM: Low bone mineral density measurements in care home residents—a treatable cause of fractures. *Age Ageing* 2006, **35**:37–41.
- Meyer HE, Smedshaug GB, Kvaavik E, Falch JA, Tverdal A, Pedersen JL: Can vitamin D supplementation reduce the risk of fracture in the elderly? A randomized controlled trial. *J Bone Miner Res* 2002, **17**:709–715.
- Meyer HE, Falch JA, Sogaard AJ, Haug E: Vitamin D deficiency and secondary hyperparathyroidism and the association with bone mineral density in persons with Pakistani and Norwegian background living in Oslo, Norway, The Oslo health study. *Bone* 2004, **35**:412–417.
- Bischoff-Ferrari HA, Dawson-Hughes B, Staehelin HB, Orav JE, Stuck AE, Theiler R, Wong JB, Egli A, Kiel DP, Henschkowski J: Fall prevention with supplemental and active forms of vitamin D: a meta-analysis of randomised controlled trials. *BMJ* 2009, **339**:b3692.
- Bischoff-Ferrari HA, Willett WC, Wong JB, Giovannucci E, Dietrich T, Dawson-Hughes B: Fracture prevention with vitamin D supplementation: a meta-analysis of randomized controlled trials. *JAMA* 2005, **293**:2257–2264.
- Bischoff-Ferrari HA, Willett WC, Wong JB, Stuck AE, Staehelin HB, Orav JE, Thoma A, Kiel DP, Henschkowski J: Prevention of nonvertebral fractures with oral vitamin D and dose dependency: a meta-analysis of randomized controlled trials. *Arch Intern Med* 2009, **169**:551–561.
- Boonen S, Bischoff-Ferrari HA, Cooper C, Lips P, Ljunggren O, Meunier PJ, Reginster JY: Addressing the musculoskeletal components of fracture risk with calcium and vitamin D: a review of the evidence. *Calcif Tissue Int* 2006, **78**:257–270.
- DIPART (Vitamin D Individual Patient Analysis of Randomized Trials) Group: Patient level pooled analysis of 68 500 patients from seven major vitamin D fracture trials in US and Europe. *BMJ* 2010, **340**:b5463.
- Avenell A, Gillespie WJ, Gillespie LD, O'Connell D: Vitamin D and vitamin D analogues for preventing fractures associated with involutional and post-menopausal osteoporosis. *Cochrane Database Syst Rev* 2009, **2**:CD000227.
- Sanders KM, Stuart AL, Williamson EJ, Simpson JA, Kotowicz MA, Young D, Nicholson GC: Annual high-dose oral vitamin D and falls and fractures in older women: a randomized controlled trial. *JAMA* 2010, **303**:1815–1822.

34. Trivedi DP, Doll R, Khaw KT: Effect of four monthly oral vitamin D3 (cholecalciferol) supplementation on fractures and mortality in men and women living in the community: randomised double blind controlled trial. *BMJ* 2003, **326**:469.
35. Smith H, Anderson F, Raphael H, Maslin P, Crozier S, Cooper C: Effect of annual intramuscular vitamin D on fracture risk in elderly men and women—a population-based, randomized, double-blind, placebo-controlled trial. *Rheumatology (Oxford)* 2007, **46**:1852–1857.
36. Glendenning P, Zhu K, Inderjeeth C, Howat P, Lewis JR, Prince RL: Effects of three-monthly oral 150,000 IU cholecalciferol supplementation on falls, mobility, and muscle strength in older postmenopausal women: a randomized controlled trial. *J Bone Miner Res* 2012, **27**:170–176.
37. Chung M, Balk EM, Brendel M, Ip S, Lau J, Lee J, Lichtenstein A, Patel K, Raman G, Tatsioni A, Terasawa T, Trikalinos TA: Vitamin D and calcium: a systematic review of health outcomes. *Evid Rep Technol Assess (Full Rep)* 2009(183):1–420.
38. Cranney A, Horsley T, O'Donnell S, Weiler H, Puil L, Ooi D, Atkinson S, Ward L, Moher D, Hanley D, Fang M, Yazdi F, Garrity C, Sampson M, Barrowman N, Tsitsvadze A, Mamaladze V: Effectiveness and safety of vitamin D in relation to bone health. *Evid Rep Technol Assess (Full Rep)* 2007(158):1–235.
39. Boonen S, Lips P, Bouillon R, Bischoff-Ferrari HA, Vanderschueren D, Haentjens P: Need for additional calcium to reduce the risk of hip fracture with vitamin D supplementation: evidence from a comparative metaanalysis of randomized controlled trials. *J Clin Endocrinol Metab* 2007, **92**:1415–1423.
40. Bischoff-Ferrari HA: The 25-hydroxyvitamin D threshold for better health. *J Steroid Biochem Mol Biol* 2007, **103**:614–619.
41. Francis RM: The vitamin D paradox. *Rheumatology (Oxford)* 2007, **46**:1749–1750.
42. Bischoff-Ferrari H: Vitamin D: what is an adequate vitamin D level and how much supplementation is necessary? *Best Pract Res Clin Rheumatol* 2009, **23**:789–795.
43. Heaney RP: The case for improving vitamin D status. *J Steroid Biochem Mol Biol* 2007, **103**:635–641.
44. Pearce SH, Cheetham TD: Diagnosis and management of vitamin D deficiency. *BMJ* 2010, **340**:b5664.
45. Brannon PM, Yetley EA, Bailey RL, Picciano MF: Summary of roundtable discussion on vitamin D research needs. *Am J Clin Nutr* 2008, **88**:587S–592S.
46. Arabi A, Baddoura R, El-Rassi R, El-Hajj Fuleihan G: PTH level but not 25 (OH) vitamin D level predicts bone loss rates in the elderly. *Osteoporos Int* 2012, **23**:971–980.
47. Hodzman AB, Hanley DA, Watson PH, Fraher LJ: Parathyroid hormone. In *Principles of Bone Biology*. Edited by Bilezikian JP, Raisz LG, Rodan GA. New York: Academic; 2002:1305–1324.
48. Bischoff-Ferrari HA: Optimal serum 25-hydroxyvitamin D levels for multiple health outcomes. *Adv Exp Med Biol* 2008, **624**:55–71.
49. Malabanan A, Veronikis IE, Holick MF: Redefining vitamin D insufficiency. *Lancet* 1998, **351**:805–806.
50. Macdonald HM, Wood AD, Aucott LS, Black AJ, Fraser WD, Mavroeidi A, Reid DM, Secombe KR, Simpson WG, Thies F: Hip bone loss is attenuated with 1000 IU but not 400 IU daily vitamin D3: a 1 year double-blind RCT in postmenopausal women. *J Bone Miner Res* 2013. doi:10.1002/jbmr.1959.
51. Chapuy MC, Preziosi P, Maamer M, Arnaud S, Galan P, Hercberg S, Meunier PJ: Prevalence of vitamin D insufficiency in an adult normal population. *Osteoporos Int* 1997, **7**:439–443.
52. Durazo-Arvizu RA, Dawson-Hughes B, Sempos CT, Yetley EA, Looker AC, Cao G, Harris SS, Burt VL, Carrquiry AL, Picciano MF: Three-phase model harmonizes estimates of the maximal suppression of parathyroid hormone by 25-hydroxyvitamin D in persons 65 years of age and older. *J Nutr* 2010, **140**:595–599.
53. Patel S, Hyer S, Barron J: Glomerular filtration rate is a major determinant of the relationship between 25-hydroxyvitamin D and parathyroid hormone. *Calcif Tissue Int* 2007, **80**:221–226.
54. Al-ozaizi ZH, Tuck SP, Raj N, Harrop JS, Summers GD, Cook DB, Francis RM, Datta HK: Assessment of vitamin D status in male osteoporosis. *Clin Chem* 2006, **52**:248–254.
55. Sahota O, Mundey MK, San P, Godber IM, Hosking DJ: Vitamin D insufficiency and the blunted PTH response in established osteoporosis: the role of magnesium deficiency. *Osteoporos Int* 2006, **17**:1013–1021.
56. Bouxsein ML, Delmas PD: Considerations for development of surrogate endpoints for antifracture efficacy of new treatments in osteoporosis: a perspective. *J Bone Miner Res* 2008, **23**:1155–1167.
57. Ish-Shalom S, Segal E, Salganik T, Raz B, Bromberg IL, Vieth R: Comparison of daily, weekly, and monthly vitamin D3 in ethanol dosing protocols for two months in elderly hip fracture patients. *J Clin Endocrinol Metab* 2008, **93**:3430–3435.
58. Podsiadlo D, Richardson S: The timed “Up & Go”: a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc* 1991, **39**:142–148.
59. McCloskey EV, Johansson H, Oden A, Kanis JA: From relative risk to absolute fracture risk calculation: the FRAX algorithm. *Curr Osteoporos Rep* 2009, **7**:77–83.
60. Power M, Quinn K, Schmidt S: Development of the WHOQOL-old module. *Qual Life Res* 2005, **14**:2197–2214.
61. Skevington SM, Lotfy M, O'Connell KA: The World Health Organization's WHOQOL-BREF quality of life assessment: psychometric properties and results of the international field trial. A report from the WHOQOL group. *Qual Life Res* 2004, **13**:299–310.
62. Nelson M, Hague GF, Cooper C, Bunker VW: Calcium intake in the elderly: validation of a dietary questionnaire. *J Hum Nutr Diet* 1988, **1**:115–127.
63. Macdonald HM: Contributions of sunlight and diet to vitamin D status. *Calcif Tissue Int* 2013, **92**:163–176.
64. Yan L, Schoenmakers I, Zhou B, Jarjour LM, Smith E, Nigdikar S, Goldberg GR, Prentice A: Ethnic differences in parathyroid hormone secretion and mineral metabolism in response to oral phosphate administration. *Bone* 2009, **45**:238–245.
65. Vickers AJ: The use of percentage change from baseline as an outcome in a controlled trial is statistically inefficient: a simulation study. *BMC Med Res Methodol* 2001, **1**:6.
66. European Medicines Agency: *ICH Topic E 6 (R1): Guideline for Good Clinical Practice*. London: EMEA; 2002.
67. European Medicines Agency: *COMMISSION DIRECTIVE 2005/28/EC: laying down principles and detailed guidelines for good clinical practice as regards investigational medicinal products for human use, as well as the requirements for authorisation of the manufacturing or importation of such products*. Available at: http://ec.europa.eu/health/files/eudralex/vol-1/dir_2005_28/dir_2005_28_en.pdf.
68. Bolland MJ, Grey A, Avenell A, Gamble GD, Reid IR: Calcium supplements with or without vitamin D and risk of cardiovascular events: reanalysis of the Women's Health Initiative limited access dataset and meta-analysis. *BMJ* 2011, **342**:d2040.

doi:10.1186/1745-6215-14-299

Cite this article as: Schoenmakers et al.: Vitamin D supplementation in older people (VDOP): Study protocol for a randomised controlled intervention trial with monthly oral dosing with 12,000 IU, 24,000 IU or 48,000 IU of vitamin D₃. *Trials* 2013 **14**:299.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

