

Meeting abstract

Open Access

Bioactivation of nitroglycerin by the East Asian variant of aldehyde dehydrogenase-2

Matteo Beretta*¹, Antonius CF Gorren¹, Verena M Wenzl¹, Robert Weis², Michael Russwurm³, Doris Koesling³, Kurt Schmidt¹ and Bernd Mayer¹

Address: ¹Department of Pharmacology and Toxicology, Karl Franzens University Graz, 8010 Graz, Austria, ²Department of Pharmaceutical Chemistry, Karl Franzens University Graz, 8010 Graz, Austria and ³Department of Pharmacology and Toxicology, Ruhr University Bochum, 44780 Bochum, Germany

Email: Matteo Beretta* - matteo.beretta@uni-graz.at

* Corresponding author

from 15th Scientific Symposium of the Austrian Pharmacological Society (APHAR) Joint meeting with the Hungarian Society of Experimental and Clinical Pharmacology (MFT) and the Slovenian Pharmacological Society (SDF) Graz, Austria. 19-21 November 2009

Published: 12 November 2009

BMC Pharmacology 2009, 9(Suppl 2):A40 doi:10.1186/1471-2210-9-S2-A40

This abstract is available from: <http://www.biomedcentral.com/1471-2210/9/S2/A40>

© 2009 Beretta et al; licensee BioMed Central Ltd.

Background

The East Asian variant of mitochondrial aldehyde dehydrogenase (ALDH2) exhibits significantly reduced dehydrogenase, esterase and nitroglycerin (GTN) reductase activities [1]. The small molecule Alda-1 was reported to partly restore low acetaldehyde dehydrogenase activity of this variant [2]. In the present study we compared the wild-type enzyme (ALDH2*1) with the East Asian variant (ALDH2*2) regarding GTN bioactivation and effects of Alda-1.

Results

Alda-1 increased acetaldehyde oxidation by ALDH2*1 and ALDH2*2 about 2- and 15-fold, respectively. The effects of this compound on the esterase activities of both enzymes were identical to that of NAD (6- and 10-fold stimulation with ALDH2*1 and ALDH2*2, respectively), but in the presence of the nicotinamide Alda-1 stimulated the esterase activity of ALDH2*1 only 1.7-fold, whereas the ALDH2*2-catalyzed reaction was increased 73-fold. ALDH2*1 exhibited a greater affinity for GTN than ALDH2*2 as well as a 7-fold higher maximal GTN denitration activity. However, bioactivation of the nitrate, measured as soluble guanylate cyclase (sGC) activation, was much more pronounced (8.8 ± 0.2 and 18.2 ± 0.9 $\mu\text{mol cGMP min}^{-1} \text{mg}^{-1}$ in the presence of 100 $\mu\text{g}/100 \mu\text{L}$

ALDH2 and 100 μM GTN). Alda-1 caused 30-70% inhibition of GTN denitration by ALDH2*1 but had no effect on the ALDH2*2-catalyzed reaction and did not affect GTN-induced sGC activation in the presence of either variant.

Conclusion

The present results indicate that Alda-1 stimulates established ALDH2 activities by improving NAD binding but does not restore impaired GTN bioactivation by the East Asian variant. In addition, our data revealed an unexpected discrepancy between GTN reductase activity and sGC activation, suggesting that GTN denitration and bioactivation reflect independent reactions catalyzed by ALDH2.

Acknowledgements

Supported by Fonds zur Förderung der Wissenschaftlichen Forschung in Austria (W901 DK Molecular Enzymology and P20669) and Deutsche Forschungsgemeinschaft (KO1157/4-1).

References

1. Larson HN, Zhou J, Chen Z, Stamler JS, Weiner H, Hurley TD: **Structural and functional consequences of coenzyme binding to the inactive Asian variant of mitochondrial aldehyde dehydrogenase.** *J Biol Chem* 2007, **282**:12940-12950.
2. Chen CH, Budas GR, Churchill EN, Disatnik MH, Hurley TD, Mochly-Rosen D: **Activation of aldehyde dehydrogenase-2 reduces ischemic damage to the heart.** *Science* 2008, **321**:1493-1495.