

## CASE REPORT

# Moyamoya syndrome in a child with HbE $\beta$ -thalassemia

Akmal Zahra<sup>1</sup> | Hanan Al-Abboh<sup>1</sup> | Yousif Habeeb<sup>2</sup> | Adekunle Adekile<sup>1,3</sup> 

<sup>1</sup>Hematology Unit, Department of Pediatrics, Mubarak Al-Kabeer Hospital, Jabriya, Kuwait

<sup>2</sup>Neurology Unit, Department of Pediatrics, Mubarak Al-Kabeer Hospital, Jabriya, Kuwait

<sup>3</sup>Department of Pediatrics, Faculty of Medicine, Kuwait University, Kuwait City, Kuwait

**Correspondence**

Adekunle Adekile, Department of Pediatrics, Faculty of Medicine, Kuwait University, 24923, Safat 13110, Kuwait City, Kuwait.  
Email: adekunle.adekile@ku.edu.kw

**Funding information**

None.

**Abstract**

Moyamoya is a progressive cerebrovascular disease associated with stenosis or occlusion of the arteries of the Circle of Willis. It is uncommon in thalassemia. We present a 9-year-old girl with HbE $\beta$ -thalassemia who presented with headache, vomiting, and episodes of transient hemiparesis with complete occlusion internal carotid arteries.

**KEYWORDS**

cerebral infarcts, Moyamoya, seizures, thalassemia

## 1 | INTRODUCTION

Patients with transfusion- or non-transfusion-dependent thalassemia (TDT or NTDT), suffer from a hypercoagulable state and often present with thromboembolism.<sup>1-3</sup> NTDT is more than 4 times likely to be associated with thromboembolic events than TDT. In the former, the events are mostly venous and relatively rare in the pediatric age group. On the contrary, arterial strokes are more common in TDT than in NTDT (28% vs. 9%, respectively). The risk factors in TDT are the higher rate of iron overload-mediated morbidities like diabetes mellitus, cardiac dysfunction, and arrhythmias.<sup>4-6</sup> Silent cerebral infarcts (SCI) are common in both groups.<sup>3-7</sup>

Moyamoya is an idiopathic, progressive cerebrovascular disease, characterized by bilateral stenosis or occlusion of the arteries of the circle of Willis, typically the supraclinoid internal carotid arteries, followed by extensive collateralization. The patients are prone to thrombosis,

aneurysm, and hemorrhage. It presents an angiographic appearance of tangled, tiny, collateral vasculature that has been compared with a puff of smoke.<sup>8,9</sup>

Moyamoya syndrome (MMS), occurs in a wide range of clinical conditions including hemoglobinopathies, especially sickle cell disease (SCD).<sup>10,11</sup> Few reports of MMS have been reported in TDT,<sup>12,13</sup> but more commonly in NTDT,<sup>14-16</sup> and rarely in E $\beta$ -thalassemia.<sup>17,18</sup>

## 2 | CASE REPORT

A.O., an Indian girl, was first presented at the age of 15 months with pallor, jaundice, and dark urine. She was otherwise well and her development was satisfactory. Apart from jaundice and pallor, the only other positive findings were enlarged liver and spleen (2 cm below the costal margin). There were no skeletal changes. Her initial CBC showed WBC  $5 \times 10^9/L$ , Hb 7 gm/dl, MCV 59 fl,

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Clinical Case Reports* published by John Wiley & Sons Ltd.

platelets  $414 \times 10^9/L$  and reticulocytes 5%. G6PD was normal; HPLC showed HbA 36.3%, Hb A<sub>2</sub> 5.8%, HbF 27.5%, HbE 30.3%.  $\beta$ -globin gene study revealed compound heterozygosity for Codon 41–42 del CTTT and E 26 K (c.79 G > A), thus confirming the diagnosis of E $\beta$ -thalassemia. She was managed as NTDT and maintained on folic acid, with transfusion only for severe anemia. Over the next 1 year, her Hb was  $\sim 7.5$  g/dl and she required transfusion only once. At age 3 years, she was commenced on 20 mg/kg of hydroxyurea (HU) daily, to which she showed a modest response. However, after about one year on HU, she started to develop skeletal changes. She was, therefore, commenced on transfusion every 2 months.

In May 2016, at the age of 5 years, she had haploidentical bone marrow transplantation, with the mother as the donor. However, there was no engraftment and her Hb rapidly dropped to the baseline value. Since September 2016, she has been on regular transfusion every 5–6 weeks and 30 mg/kg/day deferasirox.

She was admitted to the hospital in December 2019 with acute liver injury, elevated transaminases, and direct bilirubin, along with abdominal pain and vomiting. Blood-borne virology and hepatitis screen were negative. Deferasirox was discontinued; she gradually improved with supportive therapy and was discharged after 1 week.

At the outpatient department (OPD) visit in March 2020, she complained of repeated attacks of early morning headache and non-projectile vomiting of recent onset. She reported two episodes of transient right-sided weakness and numbness that lasted for a few minutes. There was no neurologic deficit, the sensorium was unaffected and cranial nerves were normal. The weakness was not progressive. There was no prior history of focal neurological deficit, seizures, cognitive or psychiatric manifestations, and no pertinent family history.

Magnetic resonance imaging (MRI) showed multiple old lacunar infarcts in both cerebral hemispheres, while magnetic resonance angiography (MRA) revealed normal intracranial arteries. The distal segments of both internal carotid arteries (ICAs) were small in caliber. MRA of the neck showed small calibers of both ICAs, with compensatory dilatation of both vertebral arteries.

Electroencephalography showed no specific abnormalities, and echocardiography was normal. Serum ferritin was not elevated, and there was no evidence of iron accumulation in the liver or heart. Thrombophilia screening, inflammatory markers, and antibody studies were all normal.

She was on regular transfusion, low-dose deferasirox (10 mg/kg/day), and prophylactic aspirin. Despite this, 4 months later, she complained of more frequent and more prolonged episodes of weakness in both hands and feet, with associated dysarthria, but no loss of consciousness



**FIGURE 1** Magnetic resonance angiography showing occlusion of both internal carotid arteries (arrow) at the cavernous segment


or spatial perception. These were suspected to be partial seizures, and she was treated with carbamazepine, after which the episodes ceased. She was neurologically normal in between attacks.

Repeat MRI showed no evidence of new ischemic or inflammatory changes; the previously seen cerebral infarcts were evident. However, MRA now showed complete occlusion of both ICAs as they enter the intracranial cavity (Figure 1). The anterior and middle cerebral arteries showed good flow, being supplied from the posterior cerebral arteries through communicating arteries bilaterally. The calibers of the vertebral and basilar arteries and communicating arteries were more than double to compensate for the ICA occlusion (Figure 2).

Three weeks after the imaging studies, the patient had no more morning headache or vomiting, there was reduced frequency of transient ischemic attacks (TIAs) and no weakness. This indicated relative stabilization and/or improvement of her condition.

### 3 | DISCUSSION

E $\beta$ -thalassemia spans the spectrum of NTDT to TDT, and our patient was initially managed as NTDT, but eventually became transfusion dependent. She could, therefore, present with complications associated with the 2 phenotypes. Indeed, she presented with TIA and was eventually found to have several old lacunar infarcts. Therefore, an initial impression of SCI was made but, because of the extension of her symptoms, repeat imaging confirmed complete occlusion of both ICAs, with increased collaterals,



**FIGURE 2** Magnetic resonance angiography showing occlusion (open arrow heads) of both internal carotid arteries. The internal carotid (round arrow heads) caliber is reduced, whereas that of the vertebral arteries (double headed arrow) and basilar arteries (arrow head) are increased

consistent with MMS.<sup>8</sup> She had also shown evidence of deferasirox hepatotoxicity.

Moyamoya syndrome may be asymptomatic but could present with TIA, ischemic or hemorrhagic stroke, and epilepsy. Rarely, patients may develop dystonia, chorea, or dyskinesia.<sup>19,20</sup> Our patient presented with morning headache and transient brief unilateral weakness, which became more frequent and lasted for longer periods of time. She later developed focal seizures, accompanied by dysarthria. Other common causes of abnormal movements such as drug side effects, Sydenham's chorea, and neurodegenerative conditions, vasculitis, autoimmune conditions, infections, thrombophilias, and connective tissue disorders were all ruled out.

She had been on regular transfusion for at least 5 years before her neurological symptoms. This supports the premise that, while transfusion dependency reduces the effect of chronic tissue hypoxia, patients are still at risk of arterial thrombosis.<sup>4,13,21</sup> Moreover, MMS is not uncommon among patients with HbE- $\beta$ -thalassemia. In one study, three of four beta-thalassemia patients with MMS were NTDT and one of them was HbE- $\beta$ -thalassemia.<sup>22</sup> In another recent study among 13 patients of MMS with thalassemia, 6 were TDT and 3 had HbE- $\beta$ -thalassemia.<sup>13</sup> Our patient is, however, much younger than those in the literature.

The complete occlusion of both ICAs in our patient probably prevented the appearance of new vessel formation and the appearance of the classic puff of smoke, especially given the good compensation by communicating vertebral and basilar arteries. This is consistent with stage

II of the Suzuki MMS staging system.<sup>8</sup> The treatment options for ICA occlusion include by-pass surgery,<sup>23</sup> endarterectomy, and stenting.<sup>24</sup>

## 4 | CONCLUSION

There is no curative treatment for MMS; it is, therefore, imperative to search for and treat any underlying condition. The vascular occlusion tends to continue despite any known medical management,<sup>13</sup> as in our patient, who was on aspirin therapy and regular transfusions. She is currently being considered for surgical intervention.

## ACKNOWLEDGMENT

We thank the patient and her family for allowing us to report this case.

## CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

## AUTHOR CONTRIBUTIONS

All authors participated in the care of the patient. AZ wrote the first draft of the manuscript. YH reviewed the MR images and interpreted the findings. All authors critically reviewed and contributed to the manuscript.

## CONSENT

Written informed consent was obtained from the patient's father to publish this report in accordance with the journal's patient consent policy.

## DATA AVAILABILITY STATEMENT

Data sharing not applicable – no new data generated.

## ORCID

Adekunle Adekile  <https://orcid.org/0000-0002-3388-6659>

## REFERENCES

1. Sirachainan N. Thalassemia and the hypercoagulable state. *Thromb Res.* 2013;132(6):637-641.
2. Cappellini MD, Motta I, Musallam KM, Taher AT. Redefining thalassemia as a hypercoagulable state. *Ann N Y Acad Sci.* 2010;1202:231-236.
3. Musallam KM, Beydoun A, Hourani R, et al. Brain magnetic resonance angiography in splenectomized adults with  $\beta$ -thalassemia intermedia. *Eur J Haematol.* 2011;87(6):539-546.
4. Taher A, Isma'eel H, Mehio G, et al. Prevalence of thromboembolic events among 8,860 patients with thalassaemia major and intermedia in the Mediterranean area and Iran. *Thromb Haemost.* 2006;96(4):488-491.
5. Musallam KM, Cappellini MD, Wood JC, et al. Elevated liver iron concentration is a marker of increased morbidity

- in patients with  $\beta$  thalassemia intermedia. *Haematologica*. 2011;96(11):1605-1612.
6. Karimi M, Khanlari M, Rachmilewitz EA. Cerebrovascular accident in beta-thalassemia major (beta-TM) and beta-thalassemia intermedia (beta-TI). *Am J Hematol*. 2008;83(1):77-79.
  7. Pazgal I, Inbar E, Cohen M, Shpilberg O, Stark P. High incidence of silent cerebral infarcts in adult patients with beta thalassemia major. *Thromb Res*. 2016;144:119-122.
  8. Suzuki J, Kodama N. Moyamoya disease—a review. *Stroke*. 1969;14(1):104-109.
  9. Suzuki J. Cerebrovascular "Moyamoya" disease. Disease showing abnormal net-like vessels in base of brain. *Arch Neurol*. 1969;20(3):288-299.
  10. Fujimura M, Bang OY, Kim JS. Moyamoya disease. *Front Neurol Neurosci*. 2016;40:204-220.
  11. Dobson SR, Holden KR, Nietert PJ, et al. Moyamoya syndrome in childhood sickle cell disease: a predictive factor for recurrent cerebrovascular events. *Blood*. 2002;99(9):3144-3150.
  12. Inati A, Tourjuman O, Bizri D, et al. A rare case of Moyamoya syndrome in a  $\beta$ -thalassemia major patient. *Blood Cells Mol Dis*. 2013;51(3):149-150.
  13. Das S, Dubey S, Acharya M, et al. Thalassemia and Moyamoya syndrome: unfurling an intriguing association. *J Neurol*. 2019;266(11):2838-2847.
  14. Sanefuji M, Ohga S, Kira R, Yoshiura T, Torisu H, Hara T. Moyamoya syndrome in a splenectomized patient with beta-thalassemia intermedia. *J Child Neurol*. 2006;21(1):75-77.
  15. Göksel BK, Ozdogu H, Yildirim T, Oğuzkurt L, Asma S. Beta-thalassemia intermedia associated with Moyamoya syndrome. *J Clin Neurosci*. 2010;17(7):919-920.
  16. Oberoi S, Bansal D, Singh P, Marwaha RK. Stroke in a young boy with  $\beta$ -thalassemia intermedia secondary to moyamoya syndrome. *J Pediatr Hematol Oncol*. 2010;32(7):568-570.
  17. Parker TM, Ward LM, Johnston DL, Ventureya E, Klaassen RJ. A case of Moyamoya syndrome and hemoglobin E/beta-thalassemia. *Pediatr Blood Cancer*. 2009;52(3):422-424.
  18. Doctor PN, Choudhari A, Verma M, Merchant RH. Moyamoya syndrome in hemoglobin E-beta thalassemia: a rare presentation and association. *J Postgrad Med*. 2018;64(4):240-242.
  19. Ahn ES, Scott RM, Robertson RL Jr, Smith ER. Chorea in the clinical presentation of moyamoya disease: results of surgical revascularization and a proposed clinicopathological correlation. *J Neurosurg Pediatr*. 2013;11(3):313-319.
  20. Baik JS, Lee MS. Movement disorders associated with moyamoya disease: a report of 4 new cases and a review of literatures. *Mov Disord*. 2010;25(10):1482-1486.
  21. Borgna Pignatti C, Carnelli V, Caruso V, et al. Thromboembolic events in beta thalassemia major: an Italian multicenter study. *Acta Haematol*. 1998;99(2):76-79.
  22. Oberoi S, Bansal D, Singh P, Marwaha RK. Stroke in a young boy with beta-thalassemia intermedia secondary to moyamoya syndrome. *J Pediatr Hematol Oncol*. 2010;32(7):568-570.
  23. Baaj AA, Agazzi S, Sayed ZA, Toledo M, Spetzler RF, van Loveren H. Surgical management of moyamoya disease: a review. *Neurosurg Focus*. 2009;26(4):E7.
  24. Hokari M, Isobe M, Asano T, et al. Treatment strategy for bilateral carotid stenosis: 2 cases of carotid endarterectomy for the symptomatic side followed by carotid stenting. *J Stroke Cerebrovasc Dis*. 2014;23(10):2851-2856.

**How to cite this article:** Zahra A, Al-Abboh H, Habeeb Y, Adekile A. Moyamoya syndrome in a child with HbE $\beta$ -thalassemia. *Clin Case Rep*. 2022;10:e05536. doi:[10.1002/ccr3.5536](https://doi.org/10.1002/ccr3.5536)