

Prevalence of Extended-Spectrum and Metallo β -Lactamase Production in AmpC β -Lactamase Producing *Pseudomonas aeruginosa* Isolates From Burns

Roya Rafiee¹; Fereshteh Eftekhari^{1*}; Seyyed Ahmad Tabatabaei²; Dariush Minaee Tehrani¹

¹Department of Microbiology, Faculty of Biological Sciences, Shahid Beheshti University, Tehran, IR Iran

²Department of Pediatrics, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

*Corresponding author: Fereshteh Eftekhari, Department of Microbiology, Faculty of Biological Sciences, Shahid Beheshti University, Tehran, IR Iran. Tel: +98-2129903208, Fax: +98-2122431664, E-mail: Feftekhari@sbu.ac.ir

Received: December 15, 2013; Revised: June 09, 2014; Accepted: June 21, 2014

Background: *Pseudomonas aeruginosa* is one of the most common causes of nosocomial infections. Resistance of *P. aeruginosa* to β -lactam antibiotics may be the result of acquired resistance through mutation and over production of various antibiotic inactivating enzymes. This research aimed to determine the prevalence of extended-spectrum β -lactamases (ESBL) and metallo β -lactamase (MBL) production as well as the presence of their related genes among AmpC β -lactamase producing *P. aeruginosa* isolated from burns.

Objectives: The current study aimed to determine the prevalence of class A ESBL and MBL production in relation to the presence of their related genes among AmpC β -lactamase producing *P. aeruginosa* isolated from burns.

Materials and Methods: The antimicrobial susceptibility of 51 *P. aeruginosa* isolates from patients with burns was examined against 13 antibiotics by the disc diffusion method. Minimum inhibitory concentrations (MIC) for imipenem and ceftazidime were measured by the microdilution method. AmpC production was detected by AmpC disc and the modified three-dimensional extract tests. ESBL phenotype was confirmed by the double disc synergy test (DDST). Presence of β -lactamase genes was detected by specific primers and polymerase chain reaction (PCR).

Results: All isolates were multidrug resistant. AmpC, ESBL and MBL production were observed in 35 (68.6%), 20 (39.2%) and 19 (37.3%) isolates, respectively. Overall, 43 isolates (84.3%) carried β -lactamase genes, out of which 31 (60.8%) harbored bla_{AmpC} , 20 (39.2%) had bla_{TEM} and 11 (21.6%) carried bla_{PER-1} genes. Among the AmpC producers, two isolates (6.5%) carried $bla_{AmpC} + bla_{ESBL}$, 13 (41.9%) had $bla_{AmpC} + bla_{MBL}$ and six (19.4%) produced the three enzymes.

Conclusions: A high prevalence of multiple β -lactamase production was observed among the AmpC producers (60%), of which the majority co-produced AmpC and MBL. The current study results showed correlation between β -lactamase production and the presence of antibiotic resistance genes in the isolates.

Keywords: *Pseudomonas aeruginosa*; Beta-Lactamase; bla_{AmpC} ; bla_{ESBL} ; bla_{MBL}

1. Background

Pseudomonas aeruginosa is an important opportunistic pathogen with the ability to propagate on medical devices, hospital environments and even in disinfectants (1). Due to the intrinsic resistance to many antimicrobial agents, *P. aeruginosa* infections present serious therapeutic challenges both in the community and health centers. In addition, the organism can acquire resistance elements to multiple classes of antibacterial agents, even during the course of treatment (2, 3). *P. aeruginosa* isolates from burn wounds are often responsible for systemic sepsis, graft loss, prolonged hospital stay and mortality (4).

Several mechanisms are responsible for resistance to β -lactam antibiotics in *P. aeruginosa* including: genetic mutations that lead to stable over expression of chromosome-mediated AmpC cephalosporinases, acquisition of transferable β -lactamase genes, overproduction of efflux systems, and reduced permeability (5). Among the numerous β -lactamases, Ambler class A extended-spectrum

β -lactamases (ESBLs) and class B metallo β -lactamases (MBLs) are reported as rapidly growing enzymes in clinical isolates of *P. aeruginosa*. ESBLs are capable of hydrolyzing penicillins, cephalosporins, and aztreonam (except for cephamycins or carbapenems). These enzymes are inhibited by β -lactamase inhibitors such as clavulanic acid (6, 7). Among more than 200 different known ESBLs in Gram-negative bacteria, 32 are detected in *P. aeruginosa* belonging to class A (TEM, SHV, CTX-M, PER, VEB, GES, BEL), and class D (OXA type) β -lactamases (8).

AmpC β -lactamases hydrolyze cephamycins (e.g. ceftoxitin and cefotetan), oxyimino-cephalosporins (e.g. ceftazidime, cefotaxime, and ceftriaxone), and monobactams (e.g. aztreonam) (9). In *P. aeruginosa*, decreased susceptibility to the extended-spectrum cephalosporins such as ceftazidime mostly results from over expression of the naturally occurring AmpC β -lactamases (10). Among the carbapenems, intermediate susceptibility to imipenem is widely associated with the modification of the outer

membrane protein OprD, production of MBLs and over expression of the cephalosporinases associated with decreased outer membrane permeability. Recently, a novel resistance mechanism was shown in *P. aeruginosa*, where peculiar AmpC β -lactamases with expanded-spectrum activity toward imipenem may also contribute to decreased imipenem susceptibility (11).

2. Objectives

The current study aimed to determine the prevalence of ESBL and MBL production among AmpC β -lactamase producing *P. aeruginosa* isolated from burns and evaluate the presence of their related antibiotic resistance genes.

3. Materials and Methods

3.1. Bacterial Isolates

Fifty one non-duplicate *P. aeruginosa* isolates were randomly selected from a collection of burn isolates provided by Shahid Motahari Hospital in 2011. The antibiotic susceptibility patterns and MBL production were previously determined using disc diffusion and the double disc synergy tests, respectively (12). The isolates were maintained in brain heart infusion broth (Oxoid, UK) containing 10% dimethyl sulfoxide and stored at -20°C until use. Controls for polymerase chain reaction (PCR) tests were: *P. aeruginosa* PAO1 containing bla_{AmpC} (provided by Dr. Abdi, Al-Zahra University, Iran), DNA from *Klebsiella pneumoniae* 7881 harboring bla_{TEM}, bla_{SHV} and bla_{CTX-M}, *P. aeruginosa* KOAS carrying bla_{PER-1} and *P. aeruginosa* 10.2 harboring bla_{VEB-1} were obtained from Pasteur Institute, Tehran, Iran. *P. aeruginosa* ATCC 27853 was also used as the susceptible control strain.

3.2. Antimicrobial Susceptibility Testing

Antibiotic susceptibility testing of the isolates was carried out by the disc diffusion method, according to the Clinical and Laboratory Standards Institute guidelines (CLSI) (5). The used antibiotic discs (MAST Group LTD, Merseyside, UK) were: ceftazidime (30 μ g), aztreonam (30 μ g), carbenicillin (100 μ g), piperacillin (100 μ g), ticarcillin (75 μ g), cotrimoxazole (25 μ g), amikacin (30 μ g), cefepime (30 μ g), ciprofloxacin (5 μ g), tobramycin (10 μ g), meropenem (10 μ g), imipenem (10 μ g), ceftazidime (30 μ g), and piperacillin/tazobactam (100/10 μ g).

3.3. Determination of Minimum Inhibitory Concentrations

Minimum inhibitory concentrations (MICs) for imipenem and ceftazidime were determined by the microdilution assay as recommended by the CLSI (13).

3.4. Phenotypic Detection of AmpC Production

Initial screening for AmpC β -lactamase production was

performed by the AmpC disc test. The results showed a zone diameter of < 18 mm for ceftazidime (14). Briefly, a blank disc moistened with sterile saline, was inoculated with a few colonies of the test strain. The disc was then placed next to a ceftazidime disc (30 μ g) on the surface of a Muller Hinton agar plate (Leofilchem, Italy) previously inoculated with a lawn of *Escherichia coli* ATCC 25922. The plate was incubated overnight at 37°C. An indentation of the ceftazidime inhibition zone adjacent to the disc containing the test strain was an indication of AmpC β -lactamase production.

AmpC production was also determined using a modified three-dimensional extract test (TDET) (15). Briefly, bacterial colonies from overnight cultures on Mueller Hinton agar were transferred to a pre-weighed sterile micro centrifuge tube. The tube was weighed again to obtain 10 to 15 mg of bacterial wet weight per sample. The bacteria were then suspended in peptone water (Merck, Germany) and centrifuged at 3000 rpm (Sigma 1-13 micro-centrifuge, Germany) for 15 minutes. Crude cell extracts were prepared by 10 cycles of alternate freeze-thawing at -78°C (dry ice/ethanol bath) and 37°C. The extracts were then used to screen for AmpC using the following method. Linear slits (3 cm) were cut into plates inoculated with each test organism using a sterile scalpel blade, 3 mm away from a ceftazidime disc in an outward radial direction. Approximately 40 to 50 μ L of the prepared extract was loaded into each slit and the plates were kept upright for 5 to 10 minutes followed by incubation at 37°C overnight. AmpC production was recorded when clear distortion of ceftazidime inhibition zones were observed (Figure 1).

3.5. Phenotypic Detection of ESBL

ESBL production was detected by the conventional double disc synergy test (DDST) using ceftazidime (30 μ g) and ceftazidime discs (30 μ g) with or without clavulanic acid (10 μ g) as recommended by the CLSI. An increase of \geq 5 mm in the inhibition zones of either cephalosporin in combination with clavulanic acid compared to the cephalosporin alone was interpreted as ESBL production (13).

3.6. PCR Amplification of AmpC and ESBL Genes

DNA extraction was carried out by boiling. Briefly, a loopful of an overnight grown culture of each test isolate as well as the control strains were suspended in 500 μ L of sterile double distilled water, boiled at 100°C for 10 minutes and were centrifuged at 12'000 \times g for 10 minutes. The supernatants were then used as DNA template for PCR amplification tests.

PCR amplifications were performed using specific primers for Ambler class A (bla_{TEM}, bla_{SHV}, bla_{CTX-M}, bla_{VEB-1}, bla_{PER-1}) and class C (bla_{AmpC}) β -lactamase genes (Table 1). AmpC amplifications were initially performed using a pair of external primers (PreAmpC-PA1 and PostAmpC-PA2) yielding a 1243-base pair product encompassing the entire AmpC gene of *P. aeruginosa*, but excluding its

promoter sequences. Another pair of primers (AmpC-PA-A and AmpC-PA-B) was also used as internal primers for AmpC gene yielding a 1063 bp product (Table 1).

PCR experiments were carried out in 25 μ L volume reaction mixtures containing 10 pM of each primer, 200 μ M dNTP, 1.4 mM MgCl₂, 1 μ L of crude DNA template and 1U of Taq DNA polymerase in the reaction buffer provided by the manufacturer (CinnaGen, Tehran, Iran). Amplifications were carried out in thermocycler (Techne, UK) using the following program: five minutes denaturation at 95°C followed by 30 cycles of 95°C for one minute, one minute at the annealing temperature (43 to 63°C for ES-

BLs, 62°C and/or 65°C for AmpC), and one minute at 72°C with a final extension period of 10 minutes at 72°C.


3.7. Statistical Analyses

Comparisons of β -lactamase phenotypes and presence of the related genes were performed using the chi-square test. Correlation between antibiotic resistance and β -lactamase gene carriage was analyzed by the nonparametric Mann-Whitney U test. All statistical analyses were carried out using SPSS software version 19. A P Value of < 0.05 was considered statistically significant.

Table 1. Primers Used for PCR Detection of ESBLs and AmpC Genes

β -lactamase (Primer)	Sequence (5' to 3')	Product Size (bp)	Reference
Class A			
TEM-F	GAGTATTCAACATTTCCTGTC	851	(10)
TEM-R	TAATCAGTGAGGCACCTATCTC		
SHV-F	AAGATCCACTATCGCCAGCAG	231	(10)
SHV-R	ATTCAGTTCGGTTCCAGCGG		
CTX-M-F	CGCTTGGCATGTGCAG	550	(16)
CTX-M-R	ACCGGATATCGTTGGT		
PER-1-F	ATGAATGTCATTATAAAAGC	920	(10)
PER-1-R	AATTTGGGCTTAGGGCAGAA		
VEB-1-F	CGACTTCCATTCCCGATGC	643	(10)
VEB-1-R	GGACTCTGCAACAATACGC		
Class C			
PreAmpC-PA1	ATGCAGCCAACGACAAAGG	1243	(11)
PostAmpC-PA2	CGCCCTCGCGAGCGCGCTTC		
ampC-PA-A	CTCCACACTGCTGTTCCGCC	1063	(11)
ampC-PA-B	TTGGCCAGGATCACCAGTCC		

Figure 1. Modified Three-Dimensional Test for Phenotypic Detection of AmpC β -Lactamase



A; clear distortion of inhibition zone indicates AmpC β -lactamase production, B, C and D; absence of distortion indicates negative test. E; negative control

4. Results

Disc diffusion results showed that all test isolates were resistant to three or more antibiotic classes; therefore, they were identified as multidrug resistant (MDR). Antibiotic resistance rates shown in Figure 2 were: piperacilin, carbenicillin, tobramycin and ceftazidime (100%), imipenem, ticarcillin, aztreonam and cotrimoxazole (98%), amikacin, cefepime and ciprofloxacin (96%), meropenem (94%), ceftazidime (92%) and piperacillin/tazobactam (88%). The MIC results showed 88.2% resistance to ceftazidime and 98% to imipenem, confirming the results obtained by the disc diffusion method. There was no significant association between antibiotic resistance patterns and β -lactamase production. AmpC production was observed in 35 isolates (68.6%) using the AmpC disc test and only 18 (51.4%) by the TDET test suggesting the superiority of the AmpC disc test to TDET. ESBL production was observed in 20 isolates (39.2%) and 19 isolates (37.3%) were MBL producers. Of the 35 AmpC producers, two (5.7%) had the ESBL phenotype, 13 (37.1%) produced MBL and six (17.1%) were positive for all three enzymes.

PCR amplification results showed that 43 isolates (84.3%) carried β -lactamase genes. Bla_{AmpC} was present in 31 isolates (60.8%), two of which were susceptible to ceftazidime (Table 2). There was a significant correlation between the AmpC disc test phenotype and AmpC gene carriage ($P < 0.05$), while no relationship was observed between TDET and presence of the AmpC gene. Among the class A β -lactamases, 20 isolates (39.2%) harbored bla_{TEM} and 11 (21.6%) carried bla_{PER-1} genes. SHV, CTX-M and VEB-1 genes were not detected in any of the isolates. All MBL producing isolates harbored their related genes (12). Overall, there was correlation between β -lactamase productions with the presence of their related genes in the test isolates ($P < 0.05$).

Table 2. Distribution of β -Lactamase Genes Among *Pseudomonas aeruginosa* Burn Isolates


β -Lactamases	NO. (%) Isolates	Ceftazidime ^a	Imipenem ^a
AmpC alone	10 (19.6)	≥ 256 (1/10)	16-64 (0/10)
AmpC + ESBL ^b	2 (3.9)	> 256 (0/2)	64 (0/2)
AmpC + MBL ^c	13 (25.5)	> 256 (1/13)	16-64 (0/13)
AmpC + ESBL + MBL	6 (11.8)	32 to ≥ 256 (0/6)	16-64 (0/6)

^a MIC (μ g/mL) (NO. Susceptible) (MIC, Minimum Inhibitory Concentration)

^b Extended-Spectrum β -Lactamase

^c Metallo- β -Lactamase

Figure 2. Antibiotic Resistance Patterns of *Pseudomonas aeruginosa* Isolates from Burns



CAZ, ceftazidime; IPM, imipenem; MEM, meropenem; PRL, piperacilin; PTZ, piperacillin-tazobactam; TC, ticarcillin; PY, carbenicillin; AK, amikacin; TN, tobramycin; ATM, azteonam; CPM, cefepime; CIP, ciprofloxacin; TS, cotrimoxazole; FOX, ceftioxin.

5. Discussion

Multiple β -lactamase producing *P. aeruginosa* cause major therapeutic failure and pose significant clinical challenges if remain undetected (17). Among more than 800 β -lactamases identified in Gram-negative bacteria, at least 120 were detected in *P. aeruginosa*, among which, AmpC, ESBLs, and MBLs are clinically significant (8). The Current

CLSI guidelines do not describe a method for detection of AmpC β -lactamases. AmpC disc test was originally introduced to detect plasmid-mediated AmpC β -lactamases (18). However, Black et al. reported the detection of chromosomally mediated inducible AmpC β -lactamases in a number of bacteria including *P. aeruginosa*, by the AmpC disc test (19). In the present study, using the AmpC disc test, AmpC production (68.6%) was higher among the *P. aeruginosa* isolates from burns compared to those of other reports (17.3-59.4%) (17, 20-23). On the other hand, only half of the isolates which were positive in AmpC disc test showed AmpC production by TDET. Basak et al. also showed that the AmpC disc test was superior to TDET in screening for AmpC production (20).

The current study results showed a significant association between the AmpC phenotype determined by the AmpC disc test and AmpC gene carriage ($P < 0.05$). The rate of AmpC gene carriage (60.8%) in the present study was similar to a report from Taiwan, where presentation of AmpC gene in *P. aeruginosa* was 59.6% (16). However, in a study conducted in Iran, Fazeli et al. reported 100% AmpC gene carriage among 72 pediatric *P. aeruginosa* isolates from various clinical specimens, but did not show the AmpC phenotype (24). Gene presence does not necessarily mean its expression and depends on environmental conditions. Therefore, the current study believes that it is important to associate the enzyme phenotype with the presence of resistance genes.

Detection of ESBLs in AmpC producing Gram-negative bacteria is often a problem. High level expression of AmpC can prevent recognition of ESBLs leading to false negative results (25, 26). ESBL production in our isolates (39.2%) was higher compared to the other reports from Iran (2.2-23.3%) and other countries (1.81-17%) (16, 27-31). Coproduction of AmpC and ESBL (3.9%) in the isolates of this study was similar to the findings of Upadhyay et al. (3.3%), but much lower than several other reports (24.5-26%) (17, 22, 23). Finally, similar to a report from Brazil, the isolates of the present study did not harbor SHV and CTX-M genes (32).

In the past 30 years, extensive use of carbapenems to treat AmpC and ESBL producing *P. aeruginosa*, has led to increased levels of bacterial resistance mostly due to production of MBLs (9, 33, 34). Coproduction of AmpC and MBL was detected in 25.5% of the isolates which was low compared to the other studies worldwide (45.5-46.6%) (22, 23). Alarmingly, 11.8% of the isolates of the current study produced AmpC along with ESBL and MBL. Detection of multiple β -lactamases in highly resistant bacteria could be useful for the selection of suitable antibiotic therapy and avoiding treatment failure as well as reducing mortality rates in hospitalized patients.

In conclusion, the present study showed a high prevalence of AmpC production among *P. aeruginosa* isolates from burn wounds (68.6%). More importantly, multiple β -lactamase production was observed in 60% of the AmpC producers, the majority of which also produced

MBL (25.5%). Co-production of AmpC with ESBL occurred in 3.9% of the isolates and 11.9% produced the three β -lactamases.

Acknowledgements

The authors wish to thank Shahid Beheshti University Research Council for providing a special grant to support this research, and also Shahid Motahari Hospital microbiology laboratory for providing the bacterial isolates.

Authors' contributions

The funder organization had no role in the design and conducting the study, collection, management and analysis of the data, preparation, review, and approval of the manuscript. Roya Rafiee (Ph.D student) carried out the research, collected the data and wrote a draft of the manuscript, Fereshteh Eftekhari (Ph.D supervisor) was responsible for designing the research project, interpretation of the results and final preparation of the manuscript, Seyyed Ahmad Tabatabaei (Ph.D adviser) provided some bacterial isolates and advised the choice of antibiotics, Dariush Minaee-Tehrani (Ph.D adviser) provided technical advice and support.

Funding/Support

This research was financially supported by Shahid Beheshti University Research Council.

References

- Fluit AC, Verhoef J, Schmitz FJ. Antimicrobial resistance in European isolates of *Pseudomonas aeruginosa*. European SENTRY Participants. *Eur J Clin Microbiol Infect Dis*. 2000;**19**(5):370-4.
- Lister PD, Wolter DJ, Hanson ND. Antibacterial-resistant *Pseudomonas aeruginosa*: clinical impact and complex regulation of chromosomally encoded resistance mechanisms. *Clin Microbiol Rev*. 2009;**22**(4):582-610.
- Selim S, El Alfy S, Al Ruwaili M, Abdo A, Al Jaouni S. Susceptibility of imipenem-resistant *Pseudomonas aeruginosa* to flavonoid glycosides of date palm (*Phoenix dactylifera* L.) tamar growing in Al Madinah, Saudi Arabia. *Afr J Biotechnol*. 2012;**11**(2):416-22.
- Coetzee E, Rode H, Kahn D. *Pseudomonas aeruginosa* burn wound infection in a dedicated paediatric burns unit. *S Afr J Surg*. 2013;**51**(2):50-3.
- Livermore DM. Multiple mechanisms of antimicrobial resistance in *Pseudomonas aeruginosa*: our worst nightmare? *Clin Infect Dis*. 2002;**34**(5):634-40.
- Bush K, Jacoby GA, Medeiros AA. A functional classification scheme for beta-lactamases and its correlation with molecular structure. *Antimicrob Agents Chemother*. 1995;**39**(6):1211-33.
- Paterson DL, Bonomo RA. Extended-spectrum beta-lactamases: a clinical update. *Clin Microbiol Rev*. 2005;**18**(4):657-86.
- Zhao WH, Hu ZQ. Beta-lactamases identified in clinical isolates of *Pseudomonas aeruginosa*. *Crit Rev Microbiol*. 2010;**36**(3):245-58.
- Jacoby GA. AmpC beta-lactamases. *Clin Microbiol Rev*. 2009;**22**(1):161-82.
- Weldhagen GF, Poirel L, Nordmann P. Ambler class A extended-spectrum beta-lactamases in *Pseudomonas aeruginosa*: novel developments and clinical impact. *Antimicrob Agents Chemother*. 2003;**47**(8):2385-92.
- Rodriguez-Martinez JM, Poirel L, Nordmann P. Extended-spectrum cephalosporinases in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother*. 2009;**53**(5):1766-71.
- Salimi F, Eftekhari F. Coexistence of AmpC and extended spectrum β -lactamases in metallo- β -lactamase producing *Pseudomonas aeruginosa* burn isolates in Tehran. *Jundishapur J Microbiol*. 2013;**6**(8).
- Cockerill FR, Clinical. *Performance standards for antimicrobial susceptibility testing: twenty-second informational supplement; provides updated tables for... M02-A11 and M07-A9*. National Committee for Clinical Laboratory Standards; 2012.
- Singhal S, Mathur T, Khan S, Upadhyay DJ, Chugh S, Gaiand R, et al. Evaluation of methods for AmpC beta-lactamase in Gram negative clinical isolates from tertiary care hospitals. *Indian J Med Microbiol*. 2005;**23**(2):120-4.
- Manchanda V, Singh NP. Occurrence and detection of AmpC beta-lactamases among Gram-negative clinical isolates using a modified three-dimensional test at Guru Tegh Bahadur Hospital, Delhi, India. *J Antimicrob Chemother*. 2003;**51**(2):415-8.
- Lin SP, Liu MF, Lin CF, Shi ZY. Phenotypic detection and polymerase chain reaction screening of extended-spectrum beta-lactamases produced by *Pseudomonas aeruginosa* isolates. *J Microbiol Immunol Infect*. 2012;**45**(3):200-7.
- Upadhyay S, Sen MR, Bhattacharjee A. Presence of different beta-lactamase classes among clinical isolates of *Pseudomonas aeruginosa* expressing AmpC beta-lactamase enzyme. *J Infect Dev Ctries*. 2010;**4**(4):239-42.
- Black JA, Moland ES, Hossain A, Lockhart TJ, Olson LB, Thamson K S, editors. Prevalence of plasmid-mediated AmpC β -lactamases in *Klebsiella pneumoniae* (KP), *Klebsiella oxytoca* (KO), *Proteus mirabilis* (PM), and *Salmonella* (S) isolates from 42 ICU and 21 non-ICU sites in the United States.; Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAC)..2003;
- Black JA, Moland ES, Thomson KS. AmpC disk test for detection of plasmid-mediated AmpC beta-lactamases in Enterobacteriaceae lacking chromosomal AmpC beta-lactamases. *J Clin Microbiol*. 2005;**43**(7):3110-3.
- Basak S, Khodke M, Bose S, Mallick SK. Inducible Amp C beta-lactamase producing *Pseudomonas aeruginosa* isolated in a rural hospital of central India. *J Clin Diagn Res*. 2009;**3**:19-7.
- Shahid M, Malik A, Agrawal M, Singhal S. Phenotypic detection of extended-spectrum and AmpC beta-lactamases by a new spot-inoculation method and modified three-dimensional extract test: comparison with the conventional three-dimensional extract test. *J Antimicrob Chemother*. 2004;**54**(3):684-7.
- Kumar V, Sen MR, Nigam C, Gahlot R, Kumari S. Burden of different beta-lactamase classes among clinical isolates of AmpC-producing *Pseudomonas aeruginosa* in burn patients: A prospective study. *Indian J Crit Care Med*. 2012;**16**(3):136-40.
- Sreeshma P, Champa H, Sunil RP, Subbannayya K. Detection of extended spectrum β -lactamase, AmpC β -lactamase and metallo β -lactamase in clinical isolates of *Pseudomonas aeruginosa*. *J Pharm Biomed Sci*. 2013;**33**(33):1506-15.
- Fazeli H, Sadighian H, Esfahani BN, Pourmand MR. Identification of class-1 integron and various β -lactamase classes among clinical isolates of *Pseudomonas aeruginosa* at children's medical center hospital. *J Med Bacteriol*. 2012;**1**(2):25-36.
- Laghawe A, Jaitly N, Thombare V. The Simultaneous Detection of the ESBL and the AmpC β -Lactamases in Gram Negative Bacilli. *J Clin Diag Res*. 2012;**6**(4suppl2):660-3.
- Thomson KS. Controversies about extended-spectrum and AmpC beta-lactamases. *Emerg Infect Dis*. 2001;**7**(2):333-6.
- Glupczynski Y, Bogaerts P, Deplano A, Berhin C, Huang TD, Van Eldere J, et al. Detection and characterization of class A extended-spectrum-beta-lactamase-producing *Pseudomonas aeruginosa* isolates in Belgian hospitals. *J Antimicrob Chemother*. 2010;**65**(5):866-71.
- Imani Folooodi AA, Rostami Z, Shapouri R. Antimicrobial resistance and ESBL prevalence in *Pseudomonas aeruginosa* strains isolated from clinical specimen by phenotypic and genotypic methods. *J Ardabil Univ Med Sci*. 2010;**10**(37):189-98.
- Jabalamel F, Mirsalehian A, Sotoudeh N, Jabalamel L, Aligholi M, Khoramian B, et al. Multiple-locus variable number of tandem repeats (VNTR) fingerprinting (MLVF) and antibacterial resistance profiles of extended spectrum beta lactamase (ESBL) producing *Pseudomonas aeruginosa* among burnt patients in Tehran. *Burns*. 2011;**37**(7):1202-7.

30. Shacheraghi F, Shakibaie MR, Noveiri H. Molecular Identification of ESBL genes blaGES-1, blaVEB-1, blaCTX-M blaOXA-1, blaOXA-4, blaOXA-10 and blaPER-1 in *Pseudomonas aeruginosa* strains isolated from burn patients by PCR, RFLP and sequencing techniques. *Int J Biol life Sci.* 2010;**3**(6):138-42.
31. Shahcheraghi F, Nikbin VS, Feizabadi MM. Prevalence of ESBLs genes among multidrug-resistant isolates of *Pseudomonas aeruginosa* isolated from patients in Tehran. *Microb Drug Resist.* 2009;**15**(1):37-9.
32. Zaranza AV, Morais FC, do Carmo MS, de Mendonça Marques A, Andrade-Monteiro C, Ferro TF, et al. Antimicrobial Susceptibility, Biofilm Production and Adhesion to HEp-2 Cells of *Pseudomonas aeruginosa* Strains Isolated from Clinical Samples. *J Biomat Nanobiotechnol.* 2013;**4**(1):98-106.
33. Bush K. Carbapenemases: Partners in crime. *J Glob Antimicrob Res.* 2013;**1**(1):7-16.
34. Livermore DM, Woodford N. Carbapenemases: a problem in waiting? *Curr Opin Microbiol.* 2000;**3**(5):489-95.