

# The Enigmatic Canal-Associated Neurons Regulate *Caenorhabditis elegans* Larval Development Through a cAMP Signaling Pathway

Jason Chien,\* Fred W. Wolf,<sup>†</sup> Sarah Grosche,\*<sup>1</sup> Nebeyu Yosef,\*<sup>2</sup> Gian Garriga,<sup>\*,3</sup> and Catarina Mörck\*

\*Department of Chemistry and Molecular Biology, University of Gothenburg, Sweden 405 30, <sup>†</sup>Department of Molecular and Cell Biology, University of California, Merced, California 95343, and <sup>3</sup>Department of Molecular and Cell Biology, University of California, Berkeley, California 94720-3204

ORCID ID: 0000-0003-3641-9439 (G.G.)

**ABSTRACT** *Caenorhabditis elegans* larval development requires the function of the two Canal-Associated Neurons (CANs): killing the CANs by laser microsurgery or disrupting their development by mutating the gene *ceh-10* results in early larval arrest. How these cells promote larval development, however, remains a mystery. In screens for mutations that bypass CAN function, we identified the gene *kin-29*, which encodes a member of the Salt-Inducible Kinase (SIK) family and a component of a conserved pathway that regulates various *C. elegans* phenotypes. Like *kin-29* loss, gain-of-function mutations in genes that may act upstream of *kin-29* or growth in cyclic-AMP analogs bypassed *ceh-10* larval arrest, suggesting that a conserved adenylyl cyclase/PKA pathway inhibits KIN-29 to promote larval development, and that loss of CAN function results in dysregulation of KIN-29 and larval arrest. The adenylyl cyclase *ACY-2* mediates CAN-dependent larval development: *acy-2* mutant larvae arrested development with a similar phenotype to *ceh-10* mutants, and the arrest phenotype was suppressed by mutations in *kin-29*. *ACY-2* is expressed predominantly in the CANs, and we provide evidence that the *acy-2* functions in the CANs to promote larval development. By contrast, cell-specific expression experiments suggest that *kin-29* acts in both the hypodermis and neurons, but not in the CANs. Based on our findings, we propose two models for how *ACY-2* activity in the CANs regulates KIN-29 in target cells.

**KEYWORDS** *C. elegans*; KIN-29; salt-inducible kinase (SIK); cAMP; canal-associated neuron

**T**HE nematode *Caenorhabditis elegans* requires only three neurons for survival: the M4 motor neuron and the two Canal-Associated Neurons (CANs). The M4 neuron is located in the pharynx—the *C. elegans* feeding organ—and is required for peristaltic movements that move food along the pharynx (Avery and Horvitz 1987, 1989). The CANs are two bilaterally symmetric neurons that are born in the head and migrate posteriorly to the middle of the worm during embryogenesis. After the CANs have completed their migration, each neuron extends two axons: one axon grows anteriorly

toward the head, and the other grows posteriorly toward the tail (White *et al.* 1986; Wu *et al.* 2011). If the CANs are killed by laser microsurgery or if the neurons fail to differentiate, the worms arrest their development early in larval development (Forrester and Garriga 1997; Forrester *et al.* 1998). How the CANs regulate larval development is unknown.

Phenotypic analysis of mutants with CAN defects also reveals their role in larval development. The CANs express two differentiation markers, the homeodomain transcription factors CEH-10 and CEH-23 (Wang *et al.* 1993; Svendsen and McGhee 1995). Loss of *ceh-10* also results in larval arrest, which is thought to result from the failure of the CANs to differentiate (Forrester and Garriga 1997; Forrester *et al.* 1998). The posteriorly directed migrations of many cells and growth cones require the gene *vab-8* (variable abnormal) (Wightman *et al.* 1996; Wolf *et al.* 1998). In *vab-8* null mutants, the CANs fail to migrate posteriorly, and their posterior axons fail to extend or extend a short distance. The posterior body of older *vab-8* mutant larvae and adults becomes thin

Copyright © 2019 by the Genetics Society of America

doi: <https://doi.org/10.1534/genetics.119.302628>

Manuscript received October 11, 2019; accepted for publication October 13, 2019; published Early Online October 16, 2019.

Available freely online through the author-supported open access option.

<sup>1</sup>Present address: CeMM Research Center for Molecular Medicine of the Austrian Academy of Sciences, 1090 Vienna, Austria.

<sup>2</sup>Present address: Faculty of Medicine Research Program Unit Stem Cells and Metabolism, University of Helsinki, 00014 Finland.

<sup>3</sup>Corresponding author: Department of Molecular and Cell Biology, 16 Barker Hall, University of California, Berkeley, CA 94720-3204. E-mail: [garriga@berkeley.edu](mailto:garriga@berkeley.edu)

and develops abnormally. This withered tail (Wit) phenotype is thought to result from the lack of CAN function in the posterior of the mutant animals (Wightman *et al.* 1996)—a hypothesis that is supported by a correlation in different mutants between the severity of the defect in the extension of the CAN posterior axon and the penetrance of the Wit phenotype (Forrester and Garriga 1997).

In an attempt to reveal the function of the CANs, we mutagenized *ceh-10* or *vab-8* mutants and screened for mutations that can suppress the mutant larval arrest or Wit phenotypes without suppressing their CAN neuron defects. In our screens, we identified three alleles of *kin-29*, which encodes a serine/threonine kinase that is a member of the Salt-Inducible Kinase (SIK) family involved in the regulation of feeding and fasting states (Koo *et al.* 2005; Dentin *et al.* 2007; Wang *et al.* 2008; Choi *et al.* 2011).

The three mammalian SIKs are inhibited by a conserved G-protein Coupled Receptor (GPCR) pathway that activates adenylyl cyclase (ACY) and Protein Kinase A (PKA) (Wang *et al.* 1999; Takemori *et al.* 2002; Okamoto *et al.* 2004; van der Linden *et al.* 2008). Here, we report that mutations that cause an increase of cAMP levels or the activation of PKA rescue the *ceh-10* larval arrest phenotype. We also provide evidence that ACY-2 is the adenylyl cyclase that generates the cAMP necessary for CAN-dependent larval development. ACY-2 is expressed in the CANs and in a few other neurons (Korswagen *et al.* 1998). We found that, when expressed in the CANs, *acy-2* partially rescued the mutant *acy-2* larval arrest phenotype. Furthermore, CAN-specific RNAi of *acy-2* induced larval arrest. Together, these findings suggest that ACY-2 produces cAMP in the CANs. To address where KIN-29 acts when CAN function is defective, we performed cell-specific expression experiments that suggest that KIN-29 functions in the hypodermis and neurons, but not in the CANs. Our observations are consistent with a model where cAMP produced by ACY-2 in the CANs negatively regulates KIN-29 in neurons and hypodermal cells to promote larval development. We propose that cAMP diffuses from the CANs through gap junctions to inhibit KIN-29 through PKA to promote proper larval development.

## Materials and Methods

### C. elegans genetics

Worms were cultured as previously described (Brenner 1974). All strains were maintained at 20°, unless otherwise noted. The following mutant alleles were used: LGI: *gsa-1(ce94)*, *lin-35(n745)*, *mef-2(gv1)* LGII: *pde-4(ce268)* LGIII: *acy-1(pk1279)*, *ceh-10(gm58)*, *rpf-3(pk1426)* LGIV: *eri-1(mg366)* LGV: *acy-2(pk465)*, *ergo-1(gg100)*, *nre-1(hd20)*, *rde-1(ne219)*, *vab-8(e1017)* LGX: *hda-4(oy59)*, *kin-2(ce179)*, *kin-29(gm112)*, *kin-29(jehm1)*, *kin-29(jehm2)*, *kin-29(gk288)*, *lin-15B(n744)*, *lin-15B(hd126)*.

Transgenes: *gms18[Pceh-23::GFP]*; pRF4 [*rol-6(su1006)*] (Zinovyeva and Forrester 2005).

The double and triple mutants created in the genetic interaction studies were sequenced to confirm that all mutations were present.

### *ceh-10* and *vab-8* suppressor screens

*ceh-10(gm58)/ht2*; *gms18* or *vab-8(e1017)* *gms18* worms were mutagenized for 4 hr by incubation in 0.05 M ethyl methane sulfonate (EMS). Worms were washed in M9 buffer (22 mM KH<sub>2</sub>PO<sub>4</sub>, 42 mM Na<sub>2</sub>HPO<sub>4</sub>, 85.5 mM NaCl, and 1 mM MgSO<sub>4</sub>) and placed on a large culture dish; 2 h later, L4 hermaphrodites were transferred to new plates in groups of 10 worms for the *vab-8* suppressor screen or 5 worms for the *ceh-10* suppressor screen. F1 progeny were picked individually to new plates 5–6 days later, and, on days 9–13, the F2 progeny were screened for ability to rescue *ceh-10* larval arrest or *vab-8* withered tail (Wit) phenotype. More than 40,000 mutagenized genomes were screened in the *vab-8* suppressor screen, and ~20,000 genomes were screened in the *ceh-10* suppressor screen. Of the five mutations that suppressed the *vab-8* Wit phenotype, only *gm112* did not suppress the CAN migration defect. Of the four mutations that suppressed *ceh-10* larval lethality, only *jehm1* and *jehm2* were *kin-29* alleles. Neither of the other two suppressor mutations were *mef-2* alleles. All suppressors were outcrossed at least three times to the wild-type N2 strain.

### Mutant identification

For identification of *kin-29(gm112)*, we used a combination of SNP mapping, RNAi interference, and sequencing. The Hawaiian isolate CB4856 was used for SNP mapping (Wicks *et al.* 2001), which placed *gm112* between SNPs in the *H01M10.1* and *pccb-1* genes. Genes located between *H01M10.1* and *pccb-1* were tested for suppression of *vab-8* Wit and *ceh-10* larval arrest by feeding worms bacteria expressing double-stranded RNA specific to a single gene. RNAi clones were obtained from the Ahringer RNAi library (Kamath *et al.* 2003) or the *C. elegans* ORFeome library (Rual *et al.* 2004) and were verified by sequencing. The experiments were performed as previously described (Timmons *et al.* 2001).

RNAi against *kin-29* rescued both *vab-8* Wit and *ceh-10* larval arrest phenotypes. The mutant *kin-29* genes were sequenced by amplifying fragments covering the entire *kin-29* gene by PCR.

### DNA plasmid constructs and transgenic lines

**Pges-1: *kin-29cDNA*** was generated by PCR amplifying 3323 bp of the *ges-1* promoter using wild type genomic DNA as template with the following primers: 5'-ctcgagctaagc ttaatgaagtattttc -3' (*XhoI* site underlined) and 5'-ggatccct gaattcaaagataagatatgt-3' (*BamHI* site underlined). The PCR product was cloned into pCR2.1-TOPO (Invitrogen), cut out with *XhoI* and *BamHI*, and ligated into pBluescriptKS-. The 2468-bp *kin-29cDNA* was amplified from *Pkin-29::kin-29cDNA::GFP* (a kind gift from Piali Sengupta) with the following primers:

5'-ggatccatggctgcgccacggcgcc-3' (*Bam*HI site underlined) and 5'-gcggccgctcactccgagctccagcttg-3' (*Not*I site underlined). The PCR product was cloned into pCR2.1-TOPO (Invitrogen), cut out with *Bam*HI and *Not*I and ligated into the pBluescriptKS- vector containing the promoter of *ges-1* (*Pges-1*). A 744-bp fragment of the *unc-54* 3'UTR was generated by PCR amplification using wild type genomic DNA as template with the following primers: 5'-gcggccgcatctcgcgccctgcctc-3' (*Not*I site underlined) and 5'-gcggccgcaaacagttatgtttggtat-3' (*Not*I site underlined). The PCR product was cloned into pCR2.1-TOPO (Invitrogen), cut out with *Not*I and ligated into *Pges-1::kin-29cDNA*;pBluescriptKS- creating *Pges-1::kin-29cDNA::unc-54 3'UTR*. The plasmid was injected into *ceh-10;kin-29* at 25 ng/μl together with 2 ng/μl *Pmyo-2::mCherry*.

***Pges-1***: *GFP* was generated by PCR amplifying 3323 bp of the *ges-1* promoter using wild type genomic DNA as template with the following primers: 5'-ggatccctaagcttaaatgaagttatttc-3' (*Bam*HI site underlined) and 5'-ccatggctgaattcaaaagataagatgt-3' (*Nco*I site underlined). The PCR product was cloned into pCR2.1-TOPO (Invitrogen), cut out with *Bam*HI and *Nco*I and ligated into pPD95.77. The plasmid was injected into wild-type worms at 25 ng/μl together with 40 ng/μl pRF4 [*rol-6(su1006)*].

***Phlh-1***: *kin-29cDNA* was generated by PCR amplifying 3052 bp of the *hlh-1* promoter using wild-type genomic DNA as template with the following primers: 5'-ctgcagcagaattctgtgaaataagc-3' (*Pst*I site underlined) and 5'-ggatccttctgaaaaattattgaaaat-3' (*Bam*HI site underlined). The PCR product was cloned into pCR2.1-TOPO (Invitrogen), cut out with *Pst*I and *Bam*HI and ligated into pBluescriptKS-. The *kin-29cDNA* and *unc-54* 3'UTR was amplified, cloned, cut and ligated as described for the *Pges-1::kin-29cDNA* construct (see above). The plasmid was injected into *ceh-10; kin-29* at 25 ng/μl together with 2 ng/μl *Pmyo-2::mCherry*.

***Phlh-1***: *GFP* was generated by cutting out *Phlh-1* from pCR2.1-TOPO (Invitrogen) (see above) with *Pst*I and *Bam*HI and ligate the fragment into pPD95.77. The plasmid was injected into wild-type worms at 25 ng/μl together with 40 ng/μl pRF4 [*rol-6(su1006)*].

***Pelt-3***: *kin-29cDNA* was generated by PCR amplifying 1964 bp of the *elt-3* promoter using wild-type genomic DNA as template with the following primers: 5'-ctgcagtgtgacacgtgttttccaggtc-3' (*Pst*I site underlined) and 5'-ggatccgaagttgaaataccaggtagc-3' (*Bam*HI site underlined). The PCR product was cloned into pCR2.1-TOPO (Invitrogen), cut out with *Pst*I and *Bam*HI and ligated into pBluescriptKS-. The *kin-29cDNA* and *unc-54* 3'UTR were amplified, cloned, cut and ligated as described for the *Pges-1::kin-29cDNA* construct (see above). The plasmid was injected into *ceh-10; kin-29* at 25 ng/μl together with 2 ng/μl *Pmyo-2::mCherry*.

***Pelt-3***: *GFP* was generated by cutting out *Pelt-3* from pCR2.1-TOPO (Invitrogen) (see above) with *Pst*I and *Bam*HI, and the fragment ligated into pPD95.77. The plasmid was injected into wild-type worms at 25 ng/μl together with 40 ng/μl pRF4 [*rol-6(su1006)*].

***Prab-3***: *kin-29cDNA* was generated by PCR amplifying 1329 bp of the *rab-3* promoter using wild-type genomic DNA as template with the following primers: 5'-ctgcagcgaagctataatagtttttc-3' (*Pst*I site underlined) and 5'-ggatccggtcttctctgtttccgcc-3' (*Bam*HI site underlined). The PCR product was cloned into pCR2.1-TOPO (Invitrogen), cut out with *Pst*I and *Bam*HI and ligated into pBluescriptKS-. The *kin-29cDNA* and *unc-54* 3'UTR was amplified, cloned, cut and ligated as described for the *Pges-1::kin-29cDNA* construct (see above). The plasmid was injected into *ceh-10; kin-29* at 10 ng/μl together with 2 ng/μl *Pmyo-2::mCherry*.

***Prab-3***: *GFP* was generated by cutting out *Prab-3* from pCR2.1-TOPO (Invitrogen) (see above) with *Pst*I and *Bam*HI and ligating the fragment into pPD95.77. The plasmid was injected into wild-type animals at 10 ng/μl together with 40 ng/μl pRF4 [*rol-6(su1006)*].

***Pkin-29***: *kin-29cDNA* was generated by PCR amplifying 1400 bp of the *kin-29* promoter using wild-type genomic DNA as template with the following primers: 5'-ctgcagctattactgtaacacctttac-3' (*Pst*I site underlined) and 5'-ggatccctgcagtgtgtgtggcgccg-3' (*Bam*HI site underlined). The PCR product was cloned into pCR2.1-TOPO (Invitrogen), cut out with *Pst*I and *Bam*HI and ligated into pBluescriptKS-. The *kin-29cDNA* and *unc-54* 3'UTR were amplified, cloned, cut, and ligated as described for the *Pges-1::kin-29cDNA* construct (see above). The plasmid was injected into wild type worms at 25 ng/μl together with 2 ng/μl *Pmyo-2::mCherry*.

***Pkin-29***: *kin-29::GFP* was generated by cutting out *Pkin-29* from pCR2.1-TOPO (Invitrogen) (see above) with *Pst*I and *Bam*HI and ligating the fragment into pPD95.77. The plasmid was injected into wild-type animals at 25 ng/μl together with 40 ng/μl pRF4 [*rol-6(su1006)*].

***Pkin-29***: *kin-29SER517ALA* was generated by modifying *Pkin-29::kin-29cDNA* using PCR-based mutagenesis (Quickchange II XL site-directed mutagenesis kit, Stratagene). The following primers were used: 5'-caaagagtgaacccgagctgcccggtgaaactcttctgcc-3' and its reverse complement. The plasmid was injected into wild-type worms at 25 ng/μl together with 2 ng/μl *Pmyo-2::mCherry*.

***Pceh-23\_L***: *acy-2* fragment (*sense/anti-sense*) *Pceh-23\_L::acy-2* fragment (*sense*) was generated with a Gibson assembly cloning kit (NEB) by assembly of the following two DNA fragments: (1) *Pceh-23\_L*, which was amplified from *Pceh-23\_L::unc-53cDNA* with the primers:

5'- ggtactccagccgactccatgatgattcgcgccgcatTTTcaattttaata-3' and 5'- ctactctccctgtttccagcttatggctgcagttttctaccggtaccctca-3'; and (2) 1252 bp *acy-2* genomic fragment, which was amplified from N2 genomic DNA with the primers: 5'- tgagggtaccgta gaaaaactgcagccataagctggaacagggagagtag-3' and 5'- tatt taaatttgaaaatcgcgccgcaatcatatggagtcggctggagtag-3'.

**Pacy-2:** *acy-2* genomic was generated with a Gibson assembly cloning kit (NEB) by assembly of the following two DNA fragments: (1) 1200 bp *Pacy-2*, which was amplified from wild-type genomic DNA template with the primers: 5'- gctgtctactgccaatacgtc-3' and 5'- tgcgcgctggaattcagg-3'; and (2) *acy-2* genomic backbone, which was amplified from *Pceh-23\_L::acy-2(genomic)* with the primers: 5'- cctgaattc caggcgcgcaatgtcgacagtgtggaatgtcgacg-3' and 5'- gacg tatttggcagtagacagccccagctttgttcccttagtg-3'. *Pacy-2::acy-2 genomic* was injected into wild-type worms at 20 ng/ $\mu$ l with 3 ng/ $\mu$ l *Pmyo-2::GFP*.

*Pceh-23\_L: acy-2 fragment(anti-sense)* was generated with a Gibson assembly cloning kit (NEB) by assembly of the following two DNA fragments: (1) *Pceh-23\_L*, which was amplified from *Pceh-23\_L::unc-53cDNA* with the primers: 5'- ctactctccctgtttccagcttatggcgccgcatTTTcaattttaata-3'; and 5'- ggtactccagccgactccatgatgattctgcagttttctaccggtaccctca-3' and (2) 1252 bp *acy-2* genomic fragment, which was amplified from N2 genomic DNA with the primers: 5'- tgagggtaccg gtagaaaaactgcagaatcatatggagtcggctggagtag-3' and 5'- tatt taaatttgaaaatcgcgccgcaatcatatggagtcggctggagtag-3'.

*Pceh-23: acy-2 fragment(sense)* and *Pceh-23::acy-2 fragment(anti-sense)* were injected together into wild-type worms at 20 ng/ $\mu$ l each with 3 ng/ $\mu$ l *Pmyo-2::GFP*.

**Pceh-23\_L: acy-2(genomic)** was generated with a Gibson assembly cloning kit (NEB) by assembly of the following two DNA fragments: (1) *Pceh-23\_L*, which was amplified from *Pceh-23\_L::GFP* with the primers: 5'- gactccaaaattttc caaacttaactataaatcaaaagaatagaccgaga-3'; and 5'- cgtcga catttccatcactgtcgacatctgcagttttctaccggtaccctca-3' and (2) *acy-2* genomic DNA, which was amplified from N2 genomic DNA with the primers: 5'- tgagggtaccgtagaaaaactgcagatgtc gacagtgtggaatgtcgacg-3' and 5'- tctcggctctattctttgattta taagtaagtttgaaaattttggagtag-3'. *Pceh-23\_L::acy-2(genomic)* was injected into wild-type worms at 20 ng/ $\mu$ l together with 3 ng/ $\mu$ l *Pmyo-2::GFP*.

Germline transformation was performed by direct injection of various plasmid DNAs into the gonads of adult wild-type animals as described (Mello *et al.* 1991).

### Survival assay

Eggs were transferred to fresh NGM plates and allowed to hatch. The newly hatched L1 larvae were transferred to new plates and the stage of the worms were studied after 24, 48, and 72 hr. At least three biological replicates were performed for each strain. Error bars show the 95% confidence interval determined by Z-tests. The *P*-values were calculated using Fisher's exact test.

### cAMP feeding

8-Br-cAMP (Tocris) was mixed with fresh growing *Escherichia coli* (OP50) bacteria (grown overnight in Luria Broth medium with shaking at 37°); 75  $\mu$ l of bacteria-cAMP mix was seeded onto small NGM plates and the plates allowed to dry for 1 hr. Worms were transferred to the plates, and immediately another batch of 75  $\mu$ l bacteria-cAMP mix was added on top of the worms. For survival studies, eggs from mothers previously grown on cAMP plates were transferred to new cAMP plates and allowed to hatch and develop for 72–96 hr; the stage of the worms was then determined.

### Fluorescence microscopy

Worms were anesthetized in 10 mM levamisole. A Zeiss Axioskop two microscope was used to examine worms. Images were collected using an ORCA-ER CCD camera (Hamamatsu) and Openlab imaging software (Improvision).

### Data availability statement


The authors affirm that all data necessary for confirming the conclusions of this article are represented fully within the article and its tables and figures.

## Results


### Mutations in *kin-29* rescued phenotypes caused by defective CANs

The two bilaterally symmetrical *C. elegans* CANs are generated in the head, and migrate toward the tail to occupy positions near the center of the embryo (Sulston 1983). Each CAN extends an anterior axon to the head and a posterior axon to the tail (White *et al.* 1986; Wu *et al.* 2011) (Figure 1A). Normal morphogenesis and larval development require CAN neuron function (Forrester and Garriga 1997; Forrester *et al.* 1998). In mutants lacking *vab-8* function, for example, the CAN cell bodies usually fail to migrate, and either lack, or have short, posterior axons (Hedgecock *et al.* 1987; Manser and Wood 1990; Wightman *et al.* 1996) (Figure 1C). The lack of CAN function in the posterior of *vab-8* mutants is thought to result in thinning of the posterior body leading to the Wit phenotype (Figure 1D). In *ceh-10(gm58)* mutants, the CANs cannot be detected using Nomarski optics or a CAN differentiation marker, and the worms arrest as early larvae (Figure 1, G and H) (Forrester and Garriga 1997; Forrester *et al.* 1998). Because laser killing of the CANs also results in larval arrest (Forrester and Garriga 1997), the developmental arrest phenotype of *ceh-10* mutants is thought to result from loss of CAN function. It is unclear, however, whether the CANs are absent in *ceh-10(gm58)* mutants or whether they are present but fail to differentiate.

To investigate how the CANs regulate larval development, we carried out two suppressor screens. In the first screen, we mutagenized *vab-8(e1017)* mutants and screened for suppressor mutations that rescued the Wit phenotype without rescuing the CAN migration or axon extension defects, and



**Figure 1** Mutations in *kin-29* rescue phenotypes caused by defective CANs. (A), (C), (E), and (G) Fluorescence photomicrographs and (B), (D), (F), and (I) Nomarski microscopy of worms containing the *Pceh-23::GFP* transgene, which is expressed in the CANs and in tail and head neurons. (A–J) Anterior is to the left and dorsal is up, the Bar, represents 100  $\mu$ m. (A) In wild-type worms, the CAN cell bodies are located in the middle of the worm and each neuron extends axons both anteriorly and posteriorly. (B) The body morphology of a wild-type worm. (C) In *vab-8* mutants, the CANs fail to migrate and are located in the head among the other neurons that express GFP. The CAN axons fail to extend to the tail (the arrowhead indicates where one of the axon's projection stops). (D) The posterior body, as indicated by the arrows, is much thinner in *vab-8* mutants (the Withered Tail or Wit phenotype). (E) In *vab-8; kin-29* double mutants, the CAN migration and extension defects are not rescued, (F) but the Wit phenotype is rescued. (G) In *ceh-10* mutants, the CANs fail to express *Pceh-23::GFP*. (H) *ceh-10* mutants arrest their development during the L1-L2 larval stage. (I) In *ceh-10; kin-29* double mutants, the CANs still are undetectable (J), but the larval arrest is rescued. (K) The structure of *kin-29* cDNA and the different mutant alleles used in this study. (L) and (M) Quantification of survival past the L3 larval stage. The number of animals scored for each genotype is indicated above each bar. Error bars show



**Figure 2** Mutations that upregulate cAMP levels and PKA activity and reduce the function transcription factor MEF-2 rescue the *ceh-10* mutant larval arrest phenotype. Quantification of survival past the L3 larval stage of wild type, single, double, and triple mutant strains containing the *ceh-10* mutation. The number of animals scored for each genotype is indicated above each bar. Error bars show the 95% confidence interval determined by Z-tests. \*  $P < 0.00001$  (Fisher's exact test).

identified *kin-29(gm112)*. Because this mutation also suppressed the larval arrest phenotype of *ceh-10* mutants (see below), we mutagenized *ceh-10(gm58)* mutants and screened for mutations that suppressed the larval arrest phenotype but did not restore the CANs based on our inability to detect the cells using Nomarski optics or a *Pceh-23::gfp* reporter transgene. In this screen, we isolated four suppressed strains. Two of these strains contained the *kin-29(jehm1)* or *kin-29(jehm2)* mutations (Figure 1, E, F, I, and J).

*kin-29* encodes a serine/threonine kinase that is homologous to the SIKs that are related to the AMPK/SNF1 family of kinases (Lanjuin and Sengupta 2002). Sequencing of the *kin-29* gene from the different mutants revealed that the *jehm1* allele is a missense mutation that changes a conserved glutamate in the kinase domain to lysine (E112K), that the *jehm2* allele is a nonsense mutation that changes a conserved tryptophan in the kinase domain to an amber stop codon (W172STOP), and that the *gm112* allele is a nonsense mutation that changes a glutamine to an amber stop codon (Q447STOP) (Figure 1K). We also analyzed the *kin-29(gk288)* allele isolated by the International *C. elegans* Gene Knockout Consortium. The 575-bp deletion removes most of the kinase domain and results in a downstream frameshift. All of the *kin-29* mutant alleles rescued the *vab-8* Wit phenotype (data not shown), and, by scoring the ability of animals to develop past the third larval (L3) stage, all of the *kin-29* mutant alleles also suppressed the *ceh-10* mutant larval arrest phenotype (Figure 1L). These findings indicate that the

morphological and larval arrest phenotypes caused by CAN dysfunction or loss require *kin-29* function. In all of the studies described below, we used the *kin-29(gm112)* allele.

#### ***kin-29* functions in neurons and hypodermal cells to mediate CAN function**

To determine where the *kin-29* mutations act to suppress the *ceh-10* larval arrest phenotype, we expressed a *kin-29* cDNA from cell-specific promoters in *ceh-10; kin-29* double mutants, and asked whether *kin-29* expression in specific cell types produced the larval arrest phenotype of the *ceh-10* single mutant. We tested expression in intestine, body-wall muscle, hypodermal cells, and neurons—cell types known to express *kin-29* (Maduzia *et al.* 2005). For intestinal expression we used the *ges-1* promoter (Aamodt *et al.* 1991), for body-wall muscle expression we used the *hlh-1* promoter (Qadota *et al.* 2007), and for hypodermal expression we used the *elt-3* promoter (Gilleard *et al.* 1999). For neuronal expression we used the *rab-3* promoter, which is expressed in all neurons except the CANs (Stefanakis *et al.* 2015). To confirm that the promoters used to drive *kin-29* in these cells were indeed specific, we also fused the promoters to the GFP gene, studied the expression of the transgenic animals at different developmental stages, and found that the promoters drove expression in the predicted cells. Only when neurons or hypodermal cells expressed the *kin-29* cDNA was the *ceh-10* larval arrest phenotype restored, suggesting that deregulated *kin-29* activity in either neurons (other than the CANs) or

the 95% confidence interval determined by Z-tests. (L) *kin-29* mutant alleles rescue the larval arrest phenotype of *ceh-10* mutants. (M) Tissue-specific expression of *kin-29* cDNA in *ceh-10; kin-29* mutants. *kin-29* cDNA was expressed from an intestinal specific (*Pges-1*), a body wall muscle specific (*Phlh-1*), a hypodermal specific (*Pelt-3*) and a pan-neuronal promoter (*Prab-3*). The number of animals for *ceh-10; kin-29; Pelt-3::kin-29* and *ceh-10; kin-29; Prab-3::kin-29* were small because these animals arrested development and could not be propagated. The arrested transgenic animals were identified based on the presence of the cotransforming marker.

hypodermal cells is sufficient to arrest larval development (Figure 1M).

Because all known CAN promoters require *ceh-10* function, we were unable to confirm that *kin-29* does not act in the CANs to suppress *ceh-10* lethality. However, we were able to express a *kin-29* cDNA in the CANs of *vab-8*; *kin-29* double mutants since the CANs are present in these animals. To ensure specific expression in the CANs, we used a part of the *ceh-23* promoter that drives expression only in the CANs (*Pceh-23\_L*) (Wenick and Hobert 2004). CAN-specific expression of *kin-29* did not restore the Wit phenotype ( $N = 50$ ), consistent with the hypothesis that KIN-29 can act in neurons and the hypodermis to mediate the effects of CAN function.


#### Loss of the MEF-2 MADS domain transcription factor also rescues *ceh-10* mutant larval arrest

In *C. elegans* and cultured cells, SIKs phosphorylate and inhibit class II histone deacetylases, which act either upstream of, or in a complex with, the transcription factor MEF2 to regulate gene transcription (Miska *et al.* 1999; Lanjuin and Sengupta 2002; Chan *et al.* 2003; van der Linden *et al.* 2008; Cohen *et al.* 2009). Loss of the *C. elegans* homologs of MEF2 (*mef-2*) and the class IIa HDACs (*hda-4*) suppress several *kin-29* mutant phenotypes: small body size, long lifespan, slow growth, hyper-foraging, and chemoreceptor gene regulation (van der Linden *et al.* 2007). If the sole activity of KIN-29 in suppressing the larval arrest phenotype of *ceh-10* mutants is to inhibit the function of a HDA-4/MEF-2 repressive complex, then *hda-4* and *mef-2* mutants should exhibit a larval arrest phenotype similar to *ceh-10* mutants, but both *hda-4* and *mef-2* mutants are viable and fertile.

To determine whether HDA-4 and MEF-2 function differently in the regulation of morphogenesis and larval development, we asked whether mutations in these genes interacted with a *ceh-10* mutation. Although an *hda-4* mutation had no effect on the *ceh-10* larval arrest phenotype, both the *mef-2(gv1)* mutation and *mef-2(RNAi)* suppressed larval arrest (Figure 1L and Figure 2). The *ceh-10*; *mef-2*; *kin-29* triple mutant has survival rates similar to the *ceh-10*; *kin-29* and *ceh-10*; *mef-2* double mutants. These findings indicate that the functional relationship between *kin-29* and *mef-2* differs in chemoreceptor regulation and CAN-dependent larval development.

#### Mutations that upregulate the cAMP-dependent PKA pathway suppressed *ceh-10* mutant larval arrest

cAMP-dependent protein kinase A (PKA) inhibits KIN-29 and its SIK homologs at both the transcriptional and post-translational levels (Takemori *et al.* 2002; Okamoto *et al.* 2004; Berdeaux *et al.* 2007; van der Linden *et al.* 2008; Wang *et al.* 2008). In *C. elegans*, PKA consists of two subunits, the catalytic subunit KIN-1 and the regulatory subunit KIN-2. PKA is activated by cAMP that is produced by adenylyl cyclases. One of these, ACY-1, can be activated by the heterotrimeric G protein GSA-1 (Berger *et al.* 1998). Loss of *acy-1*, *gsa-1*, *kin-1*, and *kin-2* result in embryonic or larval lethality.



**Figure 3** Feeding *ceh-10* mutants synthetic cAMP rescues the larval arrest phenotype. Quantification of survival past the L3 larval stage of *ceh-10* mutants grown on normal plates seeded with bacteria mixed with different concentrations of 8-Br-cAMP. The number of animals scored for each genotype is indicated above each bar. Error bars show the 95% confidence interval determined by Z-tests. \*  $P < 0.00001$  (Fisher's exact test).

If the proteins encoded by these genes inhibit KIN-29 function, then, like loss-of-function mutations in *kin-29*, gain-of-function mutations in these genes might also suppress the larval arrest phenotype of *ceh-10* mutants. *kin-2(ce179)* mutants express a PKA holoenzyme that is hypersensitive to low levels of cAMP, and *gsa-1(ce94)* mutants express a  $G\alpha$  protein that constitutively activates PKA (Korswagen *et al.* 1997; Schade *et al.* 2005; Charlie *et al.* 2006). Similar to the *kin-2(ce179)* and *gsa-1(ce94)* mutations suppressed the *ceh-10* larval arrest phenotype without suppressing the CAN defects of *ceh-10* mutants (Figure 2 and data not shown).

We also asked whether elevating the levels of cAMP could suppress the *ceh-10* larval arrest phenotype. The gene *pde-4* encodes a cAMP phosphodiesterase that is homologous to human cAMP phosphodiesterase 4D (Charlie *et al.* 2006). cAMP phosphodiesterases convert cAMP to 5'-AMP and thus lower cAMP levels (Sunahara *et al.* 1996). The *pde-4(ce268)* mutation disrupts the PDE-4 catalytic domain and reduces PDE-4 function, which is predicted to increase cAMP levels (Charlie *et al.* 2006). This mutation suppressed *ceh-10* mutant lethality (Figure 2).

#### Hyperactive KIN-29 results in larval arrest

We asked if we could phenocopy the larval arrest phenotype of *ceh-10* mutants by introducing a hyperactive version of KIN-29 into wild-type hermaphrodites. We created a construct with a mutation in the conserved PKA phosphorylation

**Table 1** *C. elegans* adenylyl cyclases

Gene	Homolog	Expression	Phenotype
<i>acy-1</i>	AC9	Neurons in head and tail ganglia, ventral nerve cord, weak in CANs, vulva and body wall muscles	Larval arrest, paralysis Moorman and Plasterk (2002)
<i>acy-2</i>	AC2	Few neurons in head ganglia and ventral nerve cord, strong expression in CANs	Larval arrest, clear Korswagen <i>et al.</i> (1998)
<i>acy-3</i>	AC5	Support cells of ciliated neurons, head and tail ganglia, two pairs of neurons in the retrovesicular ganglia, spermatheca	Reduced fat content Ashrafi <i>et al.</i> (2003), improved mitochondrial function, antioxidant defense and lifespan Vatner <i>et al.</i> (2015)
<i>acy-4</i>	AC5,6	Gonad sheath cells, spermatheca	Sterile adults Govindan <i>et al.</i> (2009)

site (Ser 517-Ala) (Takemori *et al.* 2002; van der Linden *et al.* 2008) and expressed it using the *kin-29* promoter, *Pkin-29::kin-29cDNA(Ser517-Ala)*. As controls, we made two constructs lacking the mutation, *Pkin-29::kin-29cDNA* and *Pkin-29::GFP*. The control constructs were injected into wild-type worms, creating several stable transgenic lines lacking obvious phenotypes. The construct with the *kin-29* promoter driving GFP showed the same expression pattern as described previously by Lanjuin and Sengupta, (2002): it is expressed broadly in the nervous system, including the CANs, in body wall muscle cells, and in the hypodermis, persisting through development. We injected the mutated construct into wild-type hermaphrodites, and 72% ( $N = 58$ ) of the transgenic worms arrested as early larvae. The surviving transgenic worms failed to produce lines. This finding is consistent with the hypothesis that *KIN-29* is a PKA target.

#### **ACY-2 functions in the CANs to produce essential levels of cAMP**

Our results suggest that the CANs signal to the hypodermis and other neurons, activating PKA in these tissues. PKA represses *KIN-29*, which allows larval development to proceed. Because the *pde-4* mutation rescued the *ceh-10* larval arrest phenotype to a similar degree as the *kin-29* mutations, we asked if exogenous cAMP could also rescue *ceh-10* loss. To explore this possibility, we fed *ceh-10* mutants a synthetic version of cAMP, 8-Br-cAMP, which is a cell-permeable cAMP analog that is resistant to hydrolysis by phosphodiesterases (Sandberg *et al.* 1991). We tested different concentrations and found that 5 mM 8-Br-cAMP gave the best rescue, with 75% of the *ceh-10* mutants developing past the L3 stage (Figure 3). The *ceh-10* mutants could be maintained for generations on 8-Br-cAMP, but, within hours of removing the worms from 8-Br-cAMP, the animals became sick and arrested development.


cAMP is synthesized from ATP by adenylyl cyclases (ACs). *C. elegans* has four ACs: *ACY-1*, *ACY-2*, *ACY-3*, and *ACY-4*. With the exception of *ACY-1*, which is broadly expressed in neurons and body wall muscles (Moorman and Plasterk 2002), the other three ACs have relatively restricted expression patterns. For example, *ACY-2* is only expressed in a few neurons in head ganglia and in the CANs (Korswagen *et al.* 1998) (Table 1). Loss-of-function mutations in *acy-1* or *acy-2* result in larval arrest (Korswagen *et al.* 1998; Moorman and Plasterk 2002), which prompted us to examine if *acy-1* and

*acy-2* mutants arrest development in a similar way to *ceh-10* mutants. The *acy-1(pk1279)*, *acy-2(pk465)* and *ceh-10(gm58)* mutants were maintained as balanced strains. To score the arrested larvae, we picked newly hatched worms that lacked the balancer chromosome and scored their phenotypes after 72 hr. We noted that the arrested larvae displayed three different phenotypes: normal (Figure 4A), morphological defective (Figure 4B), or clear (Figure 4C). Most of the *acy-1* arrested worms appeared normal, while *acy-2* and *ceh-10* mutants displayed the morphological defective and clear phenotypes at similar frequencies (Figure 4D). We then asked if *acy-1* and *acy-2* mutants could be rescued by the same mutations and treatments as *ceh-10*. The *pde-4* mutation was previously shown to partly suppress the *acy-1* mutant (Charlie *et al.* 2006), and we found that it also rescued the *acy-2* mutant (Figure 4E). Feeding *acy-2* mutants with 8-Br-cAMP also rescued larval arrest (Figure 4E). These findings are not unexpected for an adenylyl cyclase mutant. The *kin-29* or *mef-2* mutations, however, suppressed the *acy-2* but not the *acy-1* mutant phenotypes, consistent with the hypothesis that cAMP produced by *ACY-2* negatively regulates *KIN-29* to promote larval development (Figure 4E, and data not shown).

Mutations in *kin-29* and *mef-2* rescued the *acy-2* and *ceh-10* mutant defects slightly better than *kin-29* and *mef-2* RNAi (Figure 2 and Figure 5A and data not shown). It is noteworthy that many of the RNAi-treated *acy-2* and *ceh-10* mutant worms became visibly sick hours after being transferred to plates with bacteria that did not express *kin-29* or *mef-2* dsRNA (data not shown), suggesting that the activities of *KIN-29* and *MEF-2* need to be continuously provided for *acy-2* and *ceh-10* worms to survive. These observations imply that CANs need to constantly signal, presumably by producing cAMP that acts in the nervous system and hypodermis.

The expression pattern of *ACY-2* suggests that it could act in the CANs to promote larval development. To test this hypothesis, we expressed *ACY-2* specifically in the CANs of *acy-2* mutants. Our attempt to isolate an *acy-2* cDNA was unsuccessful, possibly because *acy-2* is expressed in only a few cells (Korswagen *et al.* 1998). Instead, we expressed the wild-type *acy-2* genomic DNA fused to GFP under the control of the CAN-specific promoter *Pceh-23\_L* (Wenick and Hobert 2004). We generated two independent, nonintegrated transgenic lines (*Pceh-23\_L::acy-2(genomic)::GFP*), and only the CANs and a single tail neuron expressed these transgenes






**Figure 4** The *kin-29* and *acy-2* mutants have similar phenotypes. (A) An arrested *ceh-10* larva with normal body morphology. The arrowheads mark the width of the intestine. (B) A *ceh-10* mutant with a morphological defective body in which the internal cells appear abnormal. (C) An arrested *ceh-10* mutant with a Clr phenotype. Note that the intestine is much thinner compared to the intestine in the worm in (A) (see arrowheads). (D) Quantification of *acy-1*, *acy-2*, and *ceh-10* mutants that arrest either with a normal body, a morphological defective body or with a Clr phenotype. (E) Quantification of survival past the L3 larval stage of *acy-2* single mutants, *acy-2*; *kin-29*, *mef-2*; *acy-2* and *pde-4*; *acy-2* double mutants and *acy-2* mutants fed with 5 mM 8-Br-cAMP. The number of animals scored for each genotype in (E) is indicated above each bar. Error bars show the 95% confidence interval determined by Z-tests. \*  $P < 0.00001$ , NS, Not Significant (Fisher's exact test). For (D) significance was determined for each phenotypic class.

(not shown). Because the transgenes were extrachromosomal and can be lost meiotically, we compared transgenic and nontransgenic worms originating from the same transgenic mother. Both transgenic lines rescued the larval arrest phenotype, with >40% of the transgenic worms developing past L3 stage (Figure 5B). Those progeny that had lost the transgenes arrested development. We also generated transgenic lines that express *acy-2* from its endogenous promoter (*Pacy-2::acy-2(genomic)*). We observed a more robust rescue compared to CAN-specific expression of *ACY-2* with >90% of the transgenic worms developing past the L3 stage. This finding suggests either that the endogenous promoter drives higher levels of *acy-2* in the CANs or that neurons other than the CANs are also important for *acy-2* mutant larvae to develop.

As an alternative test of this hypothesis, we performed CAN-specific *acy-2* RNAi to ask if we could phenocopy the *acy-2* mutant phenotype. We generated transgenes that expressed both *acy-2* sense and *acy-2* antisense RNA driven from the CAN promoter to generate *acy-2* dsRNA in the CANs. These transgenes were expressed in an *ergo-1* mutant to sensitize the background to RNAi effects (Pavelec *et al.* 2009). To obtain viable transgenic lines, we grew the worms on plates with bacteria that produced dsRNA that targeted *kin-29*, transferred transgenic worms to plates with control bacteria that did not express *kin-29* dsRNA and scored survival in the next generation. As a control, we subjected *acy-2(pk465)* mutants to the same protocol. When the *Pceh-23\_L::acy-2(RNAi)* transgenic worms and the *acy-2* mutants were transferred to plates with control bacteria, both strains arrested development



**Figure 5** Larval development requires *acy-2* function in the CANs. (A) Quantification of survival past the L3 larval stage of *acy-2* mutant worms or wild-type worms carrying transgenes that express *acy-2* dsRNA specifically in the CANs [*acy-2*(CAN RNAi)]. The worms were initially grown on bacteria that produce *kin-29* dsRNA (*kin-29* RNAi) and then transferred to plates with bacteria that do not express *kin-29* dsRNA. (B) Quantification of survival past the L3 larval stage of *acy-2* mutant worms either lacking or carrying the extrachromosomal *PCAN::acy-2* transgene that expresses *acy-2* specifically in the CANs (two lines). As control *ACY-2* was expressed from its endogenous promoter. The number of animals scored for each genotype in (E–G) is indicated above each bar. Error bars show the 95% confidence interval determined by Z-tests. \*  $P < 0.0001$  (Fisher’s exact test).

at similar frequencies (Figure 5A). These findings further support the hypothesis that *ACY-2* can act in the CANs to promote larval development.

## Discussion

### A conserved pathway mediates the essential function of the CANs

The function of the CANs is mysterious. It has been proposed that the CANs regulate the function of the excretory canal cell, which is involved in osmoregulation (Hedgecock *et al.* 1987; Forrester and Garriga 1997). This hypothesis is based on the accumulation of fluid in the pseudocoelom, the Clear (Clr) phenotype, in animals missing their CANs. In screens for mutations that bypass the requirement for the CANs in larval development, we identified the gene *kin-29*, which encodes a SIK homolog, and showed that the CANs regulate a conserved cAMP pathway that inhibits *kin-29*.

The *KIN-29*/SIK pathway mediates diverse functions that range from transcriptional regulation of *C. elegans* chemoreceptors to lipid metabolism in adipocytes (van der Linden *et al.* 2007; Henriksson *et al.* 2012). In *C. elegans*, *kin-29* functions in sensory neurons to regulate body size, entry into the dauer stage, and foraging behavior (Lanjuin and Sengupta 2002; Maduzia *et al.* 2005). Although the CANs have no obvious role in any of these processes, the essential nature of the cell makes testing its role in other processes difficult. In this context, it is noteworthy that the analogous cell in the nematode *Pristionchus pacificus* expresses the gene *dauerless*, which antagonizes dauer development. Although killing the CAN-like cell in this species does not cause larval arrest, it makes the animals more sensitive to dauer pheromone (Mayer *et al.* 2015).


Suppression of the *ceh-10* larval arrest phenotype by mutations in genes that act in the *kin-29* pathway suggests that

the adenylyl cyclase *ACY-2* acting through PKA inhibits *KIN-29* activity, allowing larval development to proceed. It is unclear whether the CANs and this pathway regulate larval development directly or a physiological state that allows development to proceed.

SIKs inhibit the function of class IIa histone deacetylases, which can interact with the MEF2 transcription factor to repress the transcription of target genes (Di Giorgio and Brancolini 2016). The sole *C. elegans* member of the class IIa HDAC family is *HDA-4*. Mutations in either *hda-4* or *mef-2* suppress the effects of *kin-29* mutations on chemoreceptor gene transcription, consistent with the inhibition of *HDA-4*/MEF2 repressor activity by *KIN-29* (van der Linden *et al.* 2007). If *KIN-29* inhibits *HDA-4*/MEF2 repressor functions in larval development as it does in chemoreceptor regulation, *hda-4* and *mef-2* mutations should cause larval arrest, which they do not. The *mef-2* mutation, however, suppressed the larval arrest phenotype of the *ceh-10* mutant. One model to explain these observations is that *KIN-29* activates MEF2, either directly or indirectly. Stimulation of cortical neurons by BDNF results in the activation of MEF2 transcriptional targets. In these cells, SIK1 phosphorylates the class IIa histone deacetylase HDAC5, resulting in HDAC5 export from the nucleus and allowing MEF2 to function as a transcriptionally activator (Finsterwald *et al.* 2013).

### Models

cAMP has traditionally been described as an intracellular “secondary messenger” that is released in response to signals from “first messengers.” How then can cAMP produced in the CANs regulate *KIN-29* in other cell types? We can think of two explanations. One possibility is that there is a cAMP pathway that functions in the CANs and results in the CANs providing a signal to neurons and hypodermal cells that regulates the *KIN-29* pathway (Figure 6A). *KIN-29* could be



**Figure 6** Models for CAN function in larval development. (A) In this model, cAMP promotes the release of a CAN signal (black squares) that acts through a GPCR/Adenylyl Cyclase (AC)/PKA/KIN-29 pathway in the target cells. An uncharacterized pathway (thick black arrows) mediates the release of the signal by cAMP. (B) In this model, cAMP synthesized in the CANs enters the target cells through gap junctions to regulate PKA and KIN-29.

regulated in neurons and hypodermal cells by cAMP. Suppression of the *ceh-10* developmental arrest by the *gsa-1* and *kin-2* gain-of-function mutations and the *pde-4* reduction-of-function mutation could result from decreased activity of *kin-29* in the nervous system and hypodermis. The ability of CAN-specific expression of *acy-2* to rescue the *acy-2* mutant phenotype suggests that the sole adenylyl cyclase functioning in this pathway is not *ACY-2*. It is possible that *ACY-1* provides this cyclase function in the nervous system and hypodermal cells, but our finding that *ceh-10* and *acy-1* mutants arrest development with distinct phenotypes does not support this idea (Figure 4D). Although the more complex phenotypes of the *ceh-10* mutants could reflect the function of the gene in other cells, laser killing of the CANs can result in a clear phenotype (Forrester and Garriga 1997)—a phenotype displayed by *ceh-10* and *acy-2* mutants, but not by *acy-1* mutants. Mutations in, or RNAi of, the remaining two adenylyl cyclase genes, *acy-3* and *acy-4*, do not produce a larval arrest phenotype (Table 1) (Ashrafi *et al.* 2003; Govindan *et al.* 2009; Vatner *et al.* 2015).

A specific version of this model proposes that cAMP is the signal from the CANs. This model is similar to the role played by cAMP in the slime mold *Dictyostelium discoideum*. Under

nutrient-limiting conditions, *Dictyostelium* cells release cAMP through the ABC transporter AbcB3 (Miranda *et al.* 2015). The released cAMP binds to a GPCR in the secreting and surrounding cells, leading to activation of a signal transduction pathway that promotes aggregation and subsequent differentiation (Loomis 2014). In mammals, a few ABC transporters are known to export cAMP (Chen *et al.* 2001; van Aubel *et al.* 2002; Wielinga *et al.* 2003). *C. elegans* has 14 predicted AbcB3 homologs, but none have been shown to be expressed in the CANs. In addition, none of the *C. elegans* GPCRs are obvious homologs of the *D. discoideum* cAMP receptor.

A second possibility is cAMP could diffuse between the CAN and other cells via gap junctions, intercellular channels that allow passive transport of ions and small molecules (Figure 6B). Vertebrate gap junctions are hemichannels that consist of connexin (Cx) proteins (Elfgang *et al.* 1995). Several studies have shown that cAMP diffuses between cells via connexins; for example, it is well established that cAMP passes through the Cx26, Cx32, Cx36, Cx43, Cx45, and Cx47 gap junction channels in HeLa cells (Bedner *et al.* 2003, 2006; Hernandez *et al.* 2007; Chandrasekhar *et al.* 2013). It is noteworthy that White *et al.* (1986) described the CAN as other

than having “a few unconvincing gap junctions to the excretory cell, no other synapses can be assigned to CAN”. Yet, the CANs express the innexin (INX) genes *inx-7* and *inx-13* (Bhattacharya *et al.* 2019), and an *inx-13* mutation results in larval arrest (Johnsen *et al.* 2000). *C. elegans* gap junctions are assembled from INX proteins (Phelan *et al.* 1998). The CAN-specific RNAi of *inx-13* also produced larval arrest (not shown), which suggests that the gap junctions between the CANs and other cells promote larval development. However, loss of neither *kin-29* nor *mef-2* suppressed the larval arrest phenotype of an *inx-13* mutant (not shown). The lack of suppression may reflect the expression of *inx-13* in other cell types, where it could function to promote larval development. The ALA, CAN, and PVD neuronal processes run together in the lateral fascicule, and the site of *KIN-29* neuronal function could be the ALA and PVD neurons.

A major concern with this model is the suppression of *ceh-10* mutants by mutations predicted to increase endogenous cAMP levels when the CAN is missing. Reduction of *pde-4* function, for example, significantly reduces larval arrest by *ceh-10* loss. Low levels of cAMP produced by cyclases in the target cells of this mutant could rise to levels sufficient to inhibit *KIN-29* in *pde-4* mutants.

Our findings provide a framework that can be used in future experiments to address the role of the CANs in larval development. In particular, the observation that loss of *mef-2* can bypass the need for the CANs suggests that the transcriptional targets of *MEF-2* could provide important insights into how the CANs regulate larval development.

## Acknowledgments

The authors thank Dawna Sweeney and Ranjan Devkota for assistance in the laboratory and valuable discussions. We thank Piali Sengupta for the plasmid containing *Pkin-29::kin-29cDNA::GFP*. Some of the nematode strains used in this study were provided by the *Caenorhabditis* Genetics Center, which is funded by the National Institutes of Health (NIH) National Center for Research Resources (NCRR). The *kin-29(gk288)* strain was provided by the *C. elegans* Gene Knockout Facility at the Oklahoma Medical Research Foundation (funded by the NIH) and the *C. elegans* Reverse Genetics Core Facility at the University of British Columbia (funded by the Canadian Institute for Health Research, Genome Canada, Genome BC, the Michael Smith Foundation, and the NIH). This work was supported by the Swedish Research Council grant 621-2012-3075 and the Olle Engkvist grant 2014/568 to C.M., and NIH grant NS32057 to G.G.

## Literature Cited

Aamodt, E. J., M. A. Chung, and J. D. McGhee, 1991 Spatial control of gut-specific gene expression during *Caenorhabditis elegans* development. *Science* 252: 579–582. <https://doi.org/10.1126/science.2020855>

Ashrafi, K., F. Y. Chang, J. L. Watts, A. G. Fraser, R. S. Kamath *et al.*, 2003 Genome-wide RNAi analysis of *Caenorhabditis elegans* fat regulatory genes. *Nature* 421: 268–272. <https://doi.org/10.1038/nature01279>

Avery, L., and H. R. Horvitz, 1987 A cell that dies during wild-type *C. elegans* development can function as a neuron in a *ced-3* mutant. *Cell* 51: 1071–1078. [https://doi.org/10.1016/0092-8674\(87\)90593-9](https://doi.org/10.1016/0092-8674(87)90593-9)

Avery, L., and H. R. Horvitz, 1989 Pharyngeal pumping continues after laser killing of the pharyngeal nervous system of *C. elegans*. *Neuron* 3: 473–485. [https://doi.org/10.1016/0896-6273\(89\)90206-7](https://doi.org/10.1016/0896-6273(89)90206-7)

Bedner, P., H. Niessen, B. Odermatt, K. Willecke, and H. Harz, 2003 A method to determine the relative cAMP permeability of connexin channels. *Exp. Cell Res.* 291: 25–35. [https://doi.org/10.1016/S0014-4827\(03\)00323-9](https://doi.org/10.1016/S0014-4827(03)00323-9)

Bedner, P., H. Niessen, B. Odermatt, M. Kretz, K. Willecke *et al.*, 2006 Selective permeability of different connexin channels to the second messenger cyclic AMP. *J. Biol. Chem.* 281: 6673–6681. <https://doi.org/10.1074/jbc.M511235200>

Berdeaux, R., N. Goebel, L. Banaszynski, H. Takemori, T. Wandless *et al.*, 2007 SIK1 is a class II HDAC kinase that promotes survival of skeletal myocytes. *Nat. Med.* 13: 597–603. <https://doi.org/10.1038/nm1573>

Berger, A. J., A. C. Hart, and J. M. Kaplan, 1998 G alphas-induced neurodegeneration in *Caenorhabditis elegans*. *J. Neurosci.* 18: 2871–2880. <https://doi.org/10.1523/JNEUROSCI.18-08-02871.1998>

Bhattacharya, A., U. Aghayeva, E. G. Berghoff, and O. Hobert, 2019 Plasticity of the electrical connectome of *C. elegans*. *Cell* 176: 1174–1189.e16. <https://doi.org/10.1016/j.cell.2018.12.024>

Brenner, S., 1974 The genetics of *Caenorhabditis elegans*. *Genetics* 77: 71–94.

Chan, J. K., L. Sun, X. J. Yang, G. Zhu, and Z. Wu, 2003 Functional characterization of an amino-terminal region of HDAC4 that possesses MEF2 binding and transcriptional repressive activity. *J. Biol. Chem.* 278: 23515–23521. <https://doi.org/10.1074/jbc.M301922200>

Chandrasekhar, A., E. A. Kalmykov, S. R. Polusani, S. A. Mathis, S. N. Zucker *et al.*, 2013 Intercellular redistribution of cAMP underlies selective suppression of cancer cell growth by connexin26. *PLoS One* 8: e82335. <https://doi.org/10.1371/journal.pone.0082335>

Charlie, N. K., A. M. Thomure, M. A. Schade, and K. G. Miller, 2006 The Dunce cAMP phosphodiesterase PDE-4 negatively regulates G alpha(s)-dependent and G alpha(s)-independent cAMP pools in the *Caenorhabditis elegans* synaptic signaling network. *Genetics* 173: 111–130. <https://doi.org/10.1534/genetics.105.054007>

Chen, Z. S., K. Lee, and G. D. Kruh, 2001 Transport of cyclic nucleotides and estradiol 17-beta-D-glucuronide by multidrug resistance protein 4. Resistance to 6-mercaptopurine and 6-thioguanine. *J. Biol. Chem.* 276: 33747–33754. <https://doi.org/10.1074/jbc.M104833200>

Choi, S., W. Kim, and J. Chung, 2011 *Drosophila* salt-inducible kinase (SIK) regulates starvation resistance through cAMP-response element-binding protein (CREB)-regulated transcription coactivator (CRTC). *J. Biol. Chem.* 286: 2658–2664. <https://doi.org/10.1074/jbc.C110.119222>

Cohen, T. J., T. Barrientos, Z. C. Hartman, S. M. Garvey, G. A. Cox *et al.*, 2009 The deacetylase HDAC4 controls myocyte enhancing factor-2-dependent structural gene expression in response to neural activity. *FASEB J.* 23: 99–106. <https://doi.org/10.1096/fj.08-115931>

Dentin, R., Y. Liu, S. H. Koo, S. Hedrick, T. Vargas *et al.*, 2007 Insulin modulates gluconeogenesis by inhibition of the coactivator TORC2. *Nature* 449: 366–369. <https://doi.org/10.1038/nature06128>

- Di Giorgio, E., and C. Brancolini, 2016 Regulation of class IIa HDAC activities: it is not only matter of subcellular localization. *Epigenomics* 8: 251–269. <https://doi.org/10.2217/epi.15.106>
- Elfngang, C., R. Eckert, H. Lichtenberg-Frate, A. Butterweck, O. Traub *et al.*, 1995 Specific permeability and selective formation of gap junction channels in connexin-transfected HeLa cells. *J. Cell Biol.* 129: 805–817. <https://doi.org/10.1083/jcb.129.3.805>
- Finsterwald, C., A. Carrard, and J. L. Martin, 2013 Role of salt-inducible kinase 1 in the activation of MEF2-dependent transcription by BDNF. *PLoS One* 8: e54545. <https://doi.org/10.1371/journal.pone.0054545>
- Forrester, W. C., and G. Garriga, 1997 Genes necessary for *C. elegans* cell and growth cone migrations. *Development* 124: 1831–1843.
- Forrester, W. C., E. Perens, J. A. Zallen, and G. Garriga, 1998 Identification of *Caenorhabditis elegans* genes required for neuronal differentiation and migration. *Genetics* 148: 151–165.
- Gilleard, J. S., Y. Shafi, J. D. Barry, and J. D. McGhee, 1999 ELT-3: a *Caenorhabditis elegans* GATA factor expressed in the embryonic epidermis during morphogenesis. *Dev. Biol.* 208: 265–280. <https://doi.org/10.1006/dbio.1999.9202>
- Govindan, J. A., S. Nadarajan, S. Kim, T. A. Starich, and D. Greenstein, 2009 Somatic cAMP signaling regulates MSP-dependent oocyte growth and meiotic maturation in *C. elegans*. *Development* 136: 2211–2221. <https://doi.org/10.1242/dev.034595>
- Hedgecock, E. M., J. G. Culotti, D. H. Hall, and B. D. Stern, 1987 Genetics of cell and axon migrations in *Caenorhabditis elegans*. *Development* 100: 365–382.
- Henriksson, E., H. A. Jones, K. Patel, M. Peggie, N. Morrice *et al.*, 2012 The AMPK-related kinase SIK2 is regulated by cAMP via phosphorylation at Ser358 in adipocytes. *Biochem. J.* 444: 503–514. <https://doi.org/10.1042/BJ20111932>
- Hernandez, V. H., M. Bortolozzi, V. Pertegato, M. Beltramello, M. Giarin *et al.*, 2007 Unitary permeability of gap junction channels to second messengers measured by FRET microscopy. *Nat. Methods* 4: 353–358. <https://doi.org/10.1038/nmeth1031>
- Johnsen, R. C., S. J. Jones, and A. M. Rose, 2000 Mutational accessibility of essential genes on chromosome I(left) in *Caenorhabditis elegans*. *Mol. Gen. Genet.* 263: 239–252. <https://doi.org/10.1007/s004380051165>
- Kamath, R. S., A. G. Fraser, Y. Dong, G. Poulin, R. Durbin *et al.*, 2003 Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature* 421: 231–237. <https://doi.org/10.1038/nature01278>
- Koo, S. H., L. Flechner, L. Qi, X. Zhang, R. A. Screaton *et al.*, 2005 The CREB coactivator TORC2 is a key regulator of fasting glucose metabolism. *Nature* 437: 1109–1111. <https://doi.org/10.1038/nature03967>
- Korswagen, H. C., J. H. Park, Y. Ohshima, and R. H. Plasterk, 1997 An activating mutation in a *Caenorhabditis elegans* Gs protein induces neural degeneration. *Genes Dev.* 11: 1493–1503. <https://doi.org/10.1101/gad.11.12.1493>
- Korswagen, H. C., A. M. van der Linden, and R. H. Plasterk, 1998 G protein hyperactivation of the *Caenorhabditis elegans* adenylyl cyclase SGS-1 induces neuronal degeneration. *EMBO J.* 17: 5059–5065. <https://doi.org/10.1093/emboj/17.17.5059>
- Lanjuin, A., and P. Sengupta, 2002 Regulation of chemosensory receptor expression and sensory signaling by the KIN-29 Ser/Thr kinase. *Neuron* 33: 369–381. [https://doi.org/10.1016/S0896-6273\(02\)00572-X](https://doi.org/10.1016/S0896-6273(02)00572-X)
- Loomis, W. F., 2014 Cell signaling during development of Dictyostelium. *Dev. Biol.* 391: 1–16. <https://doi.org/10.1016/j.ydbio.2014.04.001>
- Maduzia, L. L., A. F. Roberts, H. Wang, X. Lin, L. J. Chin *et al.*, 2005 *C. elegans* serine-threonine kinase KIN-29 modulates TGFbeta signaling and regulates body size formation. *BMC Dev. Biol.* 5: 8. <https://doi.org/10.1186/1471-213X-5-8>
- Manser, J., and W. B. Wood, 1990 Mutations affecting embryonic cell migrations in *Caenorhabditis elegans*. *Dev. Genet.* 11: 49–64. <https://doi.org/10.1002/dvg.1020110107>
- Mayer, M. G., C. Rodelsperger, H. Witte, M. Riebesell, and R. J. Sommer, 2015 The orphan gene dauerless regulates dauer development and intraspecific competition in nematodes by copy number variation. *PLoS Genet.* 11: e1005146. <https://doi.org/10.1371/journal.pgen.1005146>
- Mello, C. C., J. M. Kramer, D. Stinchcomb, and V. Ambros, 1991 Efficient gene transfer in *C. elegans*: extrachromosomal maintenance and integration of transforming sequences. *EMBO J.* 10: 3959–3970. <https://doi.org/10.1002/j.1460-2075.1991.tb04966.x>
- Miranda, E. R., E. A. Nam, A. Kuspa, and G. Shaulsky, 2015 The ABC transporter, AbcB3, mediates cAMP export in *D. discoideum* development. *Dev. Biol.* 397: 203–211. <https://doi.org/10.1016/j.ydbio.2014.11.006>
- Miska, E. A., C. Karlsson, E. Langley, S. J. Nielsen, J. Pines *et al.*, 1999 HDAC4 deacetylase associates with and represses the MEF2 transcription factor. *EMBO J.* 18: 5099–5107. <https://doi.org/10.1093/emboj/18.18.5099>
- Moorman, C., and R. H. Plasterk, 2002 Functional characterization of the adenylyl cyclase gene *sgs-1* by analysis of a mutational spectrum in *Caenorhabditis elegans*. *Genetics* 161: 133–142.
- Okamoto, M., H. Takemori, and Y. Katoh, 2004 Salt-inducible kinase in steroidogenesis and adipogenesis. *Trends Endocrinol. Metab.* 15: 21–26. <https://doi.org/10.1016/j.tem.2003.11.002>
- Pavelec, D. M., J. Lachowicz, T. F. Duchaine, H. E. Smith, and S. Kennedy, 2009 Requirement for the ERI/DICER complex in endogenous RNA interference and sperm development in *Caenorhabditis elegans*. *Genetics* 183: 1283–1295. <https://doi.org/10.1534/genetics.109.108134>
- Phelan, P., J. P. Bacon, J. A. Davies, L. A. Stebbings, M. G. Todman *et al.*, 1998 Innexins: a family of invertebrate gap-junction proteins. *Trends Genet.* 14: 348–349. [https://doi.org/10.1016/S0168-9525\(98\)01547-9](https://doi.org/10.1016/S0168-9525(98)01547-9)
- Qadota, H., M. Inoue, T. Hikita, M. Koppen, J. D. Hardin *et al.*, 2007 Establishment of a tissue-specific RNAi system in *C. elegans*. *Gene* 400: 166–173. <https://doi.org/10.1016/j.gene.2007.06.020>
- Rual, J. F., J. Ceron, J. Koreth, T. Hao, A. S. Nicot *et al.*, 2004 Toward improving *Caenorhabditis elegans* phenome mapping with an ORFeome-based RNAi library. *Genome Res.* 14: 2162–2168. <https://doi.org/10.1101/gr.2505604>
- Sandberg, M., E. Butt, C. Nolte, L. Fischer, M. Halbrugge *et al.*, 1991 Characterization of Sp-5,6-dichloro-1-beta-D-ribofuranosylbenzimidazole-3',5'-monophosphorothioate (Sp-5,6-DCl-cBiMPS) as a potent and specific activator of cyclic-AMP-dependent protein kinase in cell extracts and intact cells. *Biochem. J.* 279: 521–527. <https://doi.org/10.1042/bj2790521>
- Schade, M. A., N. K. Reynolds, C. M. Dollins, and K. G. Miller, 2005 Mutations that rescue the paralysis of *Caenorhabditis elegans ric-8* (synembryon) mutants activate the G alpha(s) pathway and define a third major branch of the synaptic signaling network. *Genetics* 169: 631–649. <https://doi.org/10.1534/genetics.104.032334>
- Stefanakis, N., I. Carrera, and O. Hobert, 2015 Regulatory logic of pan-neuronal gene expression in *C. elegans*. *Neuron* 87: 733–750. <https://doi.org/10.1016/j.neuron.2015.07.031>
- Sulston, J. E., 1983 Neuronal cell lineages in the nematode *Caenorhabditis elegans*. *Cold Spring Harb. Symp. Quant. Biol.* 48: 443–452. <https://doi.org/10.1101/SQB.1983.048.01.049>
- Sunahara, R. K., C. W. Dessauer, and A. G. Gilman, 1996 Complexity and diversity of mammalian adenylyl cyclases. *Annu. Rev. Pharmacol. Toxicol.* 36: 461–480. <https://doi.org/10.1146/annurev.pa.36.040196.002333>
- Svendsen, P. C., and J. D. McGhee, 1995 The *C. elegans* neuro-nally expressed homeobox gene *celh-10* is closely related to

- genes expressed in the vertebrate eye. *Development* 121: 1253–1262.
- Takemori, H., Y. Katoh, N. Horike, J. Doi, and M. Okamoto, 2002 ACTH-induced nucleocytoplasmic translocation of salt-inducible kinase. Implication in the protein kinase A-activated gene transcription in mouse adrenocortical tumor cells. *J. Biol. Chem.* 277: 42334–42343. <https://doi.org/10.1074/jbc.M204602200>
- Timmons, L., D. L. Court, and A. Fire, 2001 Ingestion of bacterially expressed dsRNAs can produce specific and potent genetic interference in *Caenorhabditis elegans*. *Gene* 263: 103–112. [https://doi.org/10.1016/S0378-1119\(00\)00579-5](https://doi.org/10.1016/S0378-1119(00)00579-5)
- van Aubel, R. A., P. H. Smeets, J. G. Peters, R. J. Bindels, and F. G. Russel, 2002 The MRP4/ABCC4 gene encodes a novel apical organic anion transporter in human kidney proximal tubules: putative efflux pump for urinary cAMP and cGMP. *J. Am. Soc. Nephrol.* 13: 595–603.
- van der Linden, A. M., K. M. Nolan, and P. Sengupta, 2007 KIN-29 SIK regulates chemoreceptor gene expression via an MEF2 transcription factor and a class II HDAC. *EMBO J.* 26: 358–370. <https://doi.org/10.1038/sj.emboj.7601479>
- van der Linden, A. M., S. Wiener, Y. J. You, K. Kim, L. Avery *et al.*, 2008 The EGL-4 PKG acts with KIN-29 salt-inducible kinase and protein kinase A to regulate chemoreceptor gene expression and sensory behaviors in *Caenorhabditis elegans*. *Genetics* 180: 1475–1491. <https://doi.org/10.1534/genetics.108.094771>
- Vatner, D. E., L. Yan, L. Lai, C. Yuan, L. Mouchiroud *et al.*, 2015 Type 5 adenylyl cyclase disruption leads to enhanced exercise performance. *Aging Cell* 14: 1075–1084. <https://doi.org/10.1111/acel.12401>
- Wang, B., J. Goode, J. Best, J. Meltzer, P. E. Schilman *et al.*, 2008 The insulin-regulated CREB coactivator TORC promotes stress resistance in *Drosophila*. *Cell Metab.* 7: 434–444. <https://doi.org/10.1016/j.cmet.2008.02.010>
- Wang, B. B., M. M. Muller-Immergluck, J. Austin, N. T. Robinson, A. Chisholm *et al.*, 1993 A homeotic gene cluster patterns the anteroposterior body axis of *C. elegans*. *Cell* 74: 29–42. [https://doi.org/10.1016/0092-8674\(93\)90292-X](https://doi.org/10.1016/0092-8674(93)90292-X)
- Wang, Z., H. Takemori, S. K. Halder, Y. Nonaka, and M. Okamoto, 1999 Cloning of a novel kinase (SIK) of the SNF1/AMPK family from high salt diet-treated rat adrenal. *FEBS Lett.* 453: 135–139. [https://doi.org/10.1016/S0014-5793\(99\)00708-5](https://doi.org/10.1016/S0014-5793(99)00708-5)
- Wenick, A. S., and O. Hobert, 2004 Genomic cis-regulatory architecture and trans-acting regulators of a single interneuron-specific gene battery in *C. elegans*. *Dev. Cell* 6: 757–770. <https://doi.org/10.1016/j.devcel.2004.05.004>
- White, J. G., E. Southgate, J. N. Thomson, and S. Brenner, 1986 The structure of the nervous system of the nematode *Caenorhabditis elegans*. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 314: 1–340. <https://doi.org/10.1098/rstb.1986.0056>
- Wicks, S. R., R. T. Yeh, W. R. Gish, R. H. Waterston, and R. H. Plasterk, 2001 Rapid gene mapping in *Caenorhabditis elegans* using a high density polymorphism map. *Nat. Genet.* 28: 160–164. <https://doi.org/10.1038/88878>
- Wielinga, P. R., I. van der Heijden, G. Reid, J. H. Beijnen, J. Wijnholds *et al.*, 2003 Characterization of the MRP4- and MRP5-mediated transport of cyclic nucleotides from intact cells. *J. Biol. Chem.* 278: 17664–17671. <https://doi.org/10.1074/jbc.M212723200>
- Wightman, B., S. G. Clark, A. M. Taskar, W. C. Forrester, A. V. Maricq *et al.*, 1996 The *C. elegans* gene *vab-8* guides posteriorly directed axon outgrowth and cell migration. *Development* 122: 671–682.
- Wolf, F. W., M. S. Hung, B. Wightman, J. Way, and G. Garriga, 1998 *vab-8* is a key regulator of posteriorly directed migrations in *C. elegans* and encodes a novel protein with kinesin motor similarity. *Neuron* 20: 655–666. [https://doi.org/10.1016/S0896-6273\(00\)81006-5](https://doi.org/10.1016/S0896-6273(00)81006-5)
- Wu, Y., A. Ghitani, R. Christensen, A. Santella, Z. Du *et al.*, 2011 Inverted selective plane illumination microscopy (iSPIM) enables coupled cell identity lineaging and neurodevelopmental imaging in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. USA* 108: 17708–17713. <https://doi.org/10.1073/pnas.1108494108>
- Zinovyeva, A. Y., and W. C. Forrester, 2005 The *C. elegans* Frizzled CFZ-2 is required for cell migration and interacts with multiple Wnt signaling pathways. *Dev. Biol.* 285: 447–461. <https://doi.org/10.1016/j.ydbio.2005.07.014>

Communicating editor: O. Hobert