

Once-daily, oral levofloxacin monotherapy for low-risk neutropenic fever in cancer patients: a pilot study in China

Lixian He^a, Caicun Zhou^b, Su Zhao^c, Heng Weng^d and Guowang Yang^e

This pilot study assesses the safety and efficacy of once-daily, oral levofloxacin monotherapy in Chinese patients with low-risk febrile neutropenia. In this prospective, single-arm, open-label, multicenter clinical trial, 46 adult Chinese patients with solid tumors and low-risk febrile neutropenia were included. Patients received oral levofloxacin monotherapy (500 mg orally/day) until day 12, followed by 7 days of follow-up (day 19). Body temperature was measured three times per day. On days 2, 3, 5–7, 9, 12, and 19, disease symptoms and vital signs were recorded, adverse drug reactions were assessed, and blood samples were collected to determine the whole-blood cell count and the absolute neutrophil count. Blood cultures and chest radiographs were performed simultaneously until negative results were found. Oral levofloxacin was effective and well tolerated in 97.6% of patients irrespective of the cancer type and cause of fever. Body temperature began to decline in 24.4, 68.3, and 90.2% of patients, respectively, at 12, 24, and 48 h after initiating levofloxacin therapy. On days 5 and 7, 95.1 and 97.6% of the patients had complete defervescence, respectively. The median time for absolute neutrophil count recovery to at least 1500/mm³ after initiation of treatment

Introduction

Patients with chemotherapy-induced neutropenia often develop fever of unknown origin and some may be at risk for serious infections [1–3]. A subset of patients with febrile neutropenia can be identified as being at a low risk for the development of serious clinical complications and subsequent hospitalization [4,5]. Outpatient management for adult cancer patients with low-risk febrile neutropenia includes the administration of oral antimicrobial agents, which is shown to be safe and effective in reducing fever and to compare favorably with hospital treatment [4,6,7]. Fluoroquinolone antibacterial agents are important clinical tools used for the prevention and management of infections in cancer patients with neutropenia [2,3,8,9]. Levofloxacin is a broad-spectrum antimicrobial agent of the fluoroquinolone drug class that has shown excellent tissue penetration [10]. Compared with other fluoroquinolones (e.g. ciprofloxacin), levofloxacin has shown greater activity against Gram-positive bacteria and less activity against Gram-negative bacteria [10]. Levofloxacin reduces the incidence of fever and different types of infections in cancer

was 3 days. Only one patient reported mild diarrhea. This pilot study showed that oral levofloxacin quickly and effectively reduced fever, initiated neutrophil recovery, and was well tolerated in Chinese low-risk febrile neutropenic patients with solid tumors. Further study is needed to compare patient data of levofloxacin with the standard amoxicillin/ciprofloxacin protocol in this population for both safety and efficacy. *Anti-Cancer Drugs* 26:359–362 Copyright © 2015 Wolters Kluwer Health, Inc. All rights reserved.

Anti-Cancer Drugs 2015, 26:359–362

Keywords: cancer, febrile neutropenia, fluoroquinolone, levofloxacin, low risk, solid tumor

^aZhongshan Hospital, Fudan University, ^bShanghai Pulmonary Hospital, Tongji University, Shanghai, ^cTongji Medical College Huazhong University of Science & Technology, The Central Hospital of Wuhan, Wuhan, ^dFuzhou Pulmonary Hospital of Fujian, Fuzhou and ^eBeijing Hospital of T.C.M Affiliated to Capital University of Medicine Sciences, Beijing, China

Correspondence to Caicun Zhou, Shanghai Pulmonary Hospital, Tongji University, 507 Zhengmin Road, Yangpu District, Shanghai 200433, China
Tel: +86 133 018 25532; fax: +86 21 556 60346;
e-mail: caicunzhou@yahoo.com.cn

Received 1 July 2014 Revised form accepted 1 November 2014

patients with neutropenia when compared with placebo [11–14]. However, differences have been shown between the microbiological spectrum in febrile neutropenic patients in Asia compared with that in western countries [15]. In addition, quinolone-resistant bacteria are prevalent in China [16–18]. Therefore, it is clinically important to understand the safety and efficacy of fluoroquinolone antibiotics such as levofloxacin within a Chinese population of cancer patients at low risk for infection. Accordingly, our goal was to carry out an initial prospective study to observe the ability of levofloxacin to treat infection and reduce fever in Chinese cancer patients with febrile neutropenia. This pilot study aimed to evaluate the safety and efficacy of once-daily, oral levofloxacin in the treatment of low-risk neutropenic fever in Chinese cancer patients.

Patients and methods

This pilot study was designed as a prospective, single-arm, open-label, multicenter clinical trial. The study was approved by the Ethics Committees of Zhongshan Hospital of Fudan University, Fudan, China, and the Respiratory Hospital of Tongji University (Shanghai, China). All enrolled patients provided signed informed consent. Eligible patients were more than 18 years of age. All patients had pathologically diagnosed cancer with solid tumors, were receiving chemotherapy, and had a

This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 3.0 License, where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially.

body temperature of more than 38°C but less than 40°C when taken orally. Low risk was defined as patients having an oral temperature of at least 38.5°C twice within 24 h and less than or equal to 7 days of fever, as described previously [19]. The low-risk patients had an absolute neutrophil count (ANC) less than 1500/mm³, serum creatinine levels less than 2 mg/dl, serum aminotransferase levels four times the upper limit of normal, and a Multinational Association for Supportive Care in Cancer risk index score of at least 21 [8]. Patients with evidence of active infection, including mycobacteria, fungi, viruses, protozoa, or HIV, or those who had been treated with antibiotics less than 4 days before the start of the study, had used study drugs within 30 days before enrollment, were allergic to quinolones, or had cancer-related or unrelated fever before receiving chemotherapy were excluded. A total of 46 patients were enrolled, but the efficacy population was based on 41 patients per protocol. Patients were treated with oral levofloxacin (Daiichi Sankyo Co., Ltd, Japan) 500 mg once daily (750 mg levofloxacin is not commercially available in China) at the same time every day (immediately after breakfast) for 12 days. No other antibiotics were administered during the study period. Use of antifungal and antiviral drugs was also prohibited. Granulocyte colony-stimulating factor, blood transfusions, and intravenous infusions were administered as needed. Levofloxacin treatment was discontinued after 12 days (day 12). Patients were then followed for an additional 7-day period (day 19) without treatment. Body temperature was measured three times per day. On days 1–3, 5, 7, 9, 12, and 19, disease symptoms and vital signs were evaluated and recorded. Adverse drug reactions were assessed simultaneously. Blood samples were collected on the same day for whole-blood cell count, and ANC was counted simultaneously. Blood cultures and chest radiographs were performed at the same time until negative results were found. Urine and sputum cultures were obtained when signs of infection were clinically indicated to identify the site of infection. Some patients who responded well and quickly to levofloxacin treatment were permitted to discontinue treatment earlier than day 12, but these patients were still followed for an additional 7 days without treatment. Patients were withdrawn if their fever persisted or worsened, if ANC remained high, or if a levofloxacin-associated adverse reaction occurred. The primary endpoints were the effectiveness of levofloxacin as evaluated by the proportion of patients with fever defervescence on days 5 and 7, the median time to starting defervescence, and the median time to complete defervescence. Defervescence was defined as a body temperature of less than or equal to 37°C or returning to the baseline value. Other primary endpoints included the proportion of patients who did not respond and were treated in the intensive care unit, received other antibiotics, and deaths. The secondary endpoints included neutrophil recovery indicated by time for ANC returning to at least 1500/mm³

and for white blood cell (WBC) counts to return to at least 3000/mm³ or at least 5000/mm³. No resistance data were collected. Safety endpoints included the incidence of adverse drug reactions and the proportion of patients who were withdrawn from the study because of adverse drug reactions. Continuous data were summarized by medians and ranges, and categorical data were expressed by frequencies and percentages. The response rates were described as the proportion of patients who responded and reported with a 95% confidence interval. All statistical analyses were carried out using SAS software version 9.2 (SAS Institute Inc., Cary, North Carolina, USA).

Results

Most patients (30/41; 73.2%) had been diagnosed with primary lung cancer. Other cancer types were mammary cancer ($n=5$), gastric cancer ($n=3$), gallbladder cancer ($n=1$), thyroid carcinoma ($n=1$), and renal carcinoma ($n=1$). The median Multinational Association for Supportive Care in Cancer risk index score was 23 (range: 21–26). Before levofloxacin therapy, the median ANC was 590/mm³ and the median WBC count was 1410/mm³. The median body temperature was 38.7°C and the median duration of levofloxacin therapy was 10 days. Among 41 patients in the effectiveness population, the median time of starting defervescence was 24 h and the median time to complete fever reduction was 3 days (Table 1). The proportions of patients who showed a reduction in fever within 12, 24, and 48 h were 24.4, 68.3, and 90.2%, respectively, and the percentages with complete defervescence within 5 and 7 days were 95.1 and 97.6%, respectively (Table 1). Forty patients in the efficacy population responded to levofloxacin therapy, indicating a response rate of 97.6% (95% confidence interval: 87.4–99.6%, Table 2). The median time for improvement in ANC to at least 1500/mm³ was 3 days, and that for recovery of WBC counts to at least 3000/mm³ and at least 5000/mm³ were 3 and 4 days, respectively, after initiation of treatment. Only one patient required administration of granulocyte colony-stimulating factor. Overall, 37 patients were diagnosed to have fever of

Table 1 Time to defervescence after receiving oral levofloxacin therapy

Time to defervescence	<i>N</i> = 41
Starting defervescence [<i>n</i> (%)] (h)	
≤ 12	10 (24.4)
≤ 24	28 (68.3)
≤ 48	37 (90.2)
Time to starting defervescence (h)	
Median (range)	24 (0–72)
Complete defervescence [<i>n</i> (%)]	
≤ 12 h	2 (4.9)
≤ 1 day	10 (24.4)
≤ 2 days	17 (41.5)
≤ 5 days	39 (95.1)
≤ 7 days	40 (97.6)
Time to complete defervescence (days)	
Median (range)	3 (0–14)

Table 2 Response to therapy on the basis of the nature of febrile episode and type of infection

Type of infection	Number of patients	Number of response (%)
Unexplained fever	37	37
Clinically diagnosed infection by chest radiograph		
Pneumonia	2	1
Confirmation of microbial infection		
<i>Escherichia coli</i> (urine culture)	1	1
<i>Klebsiella pneumoniae</i> (sputum culture)	1	1
Total	41	40 (97.6%)

unknown origin; two patients had confirmed infection, including one with *Klebsiella pneumoniae* found by sputum culture and one with *Escherichia coli* found by urine culture; two patients were clinically diagnosed to have infection on the basis of chest radiograph (Table 2). None of the 40 patients who responded to treatment experienced a new infection at day 7 of the follow-up period. Of the 46 patients in the safety population, only one patient reported an adverse event, which was drug-related diarrhea, but the symptoms were mild and the patient remained in the study.

Discussion and conclusion

The present study showed that once-daily, oral levofloxacin effectively treated infection and quickly reduced fever among Chinese patients with low-risk febrile neutropenia. Defervescence started at a median of 24 h, with complete fever reduction in median 3 days. Among 41 patients in the effectiveness group, 95.1 and 97.6% had complete defervescence within 5 and 7 days, respectively. Neutrophil recovery was indicated by a 3-day median time for improvement in ANC to at least $1500/\text{mm}^3$ and WBC counts returning to at least $3000/\text{mm}^3$ and at least $5000/\text{mm}^3$ in 3 and 4 days, respectively. These results in a Chinese population are especially important because of the prevalence of quinolone-resistant bacteria in China [16–18].

Recent studies measured antimicrobial resistance of Gram-positive isolates from 18 hospitals to commonly used antimicrobial agents according to the Clinical Laboratory Standards Institute guidelines, showing that nationwide resistance rates in China ranged from 45 to 81% for MRSA and MRSE, respectively, and were lower for vancomycin and linezolid resistance [16]. Macrolide resistance was more serious than that reported for Western countries [16]. In another Chinese study, a high proportion of extended-spectrum β -lactamase and/or plasmid-mediated *AmpC*-producing Enterobacteriaceae such as *Klebsiella* spp. were shown to carry the quinolone-resistance gene (*qnr*) [18].

Our results are also important when considering monotherapy versus combined therapy for addressing neutropenic fever. Both the IDSA and the NCCN recommend applying antibiotic combination therapy for the treatment

of low-risk febrile neutropenia, namely, ciprofloxacin in combination with amoxicillin/clavulanate, which shows anti-Gram-positive activity [19,20]. However, in our experience, combined therapies potentially increase side effects, resulting in lower treatment compliance and increased medical costs. Levofloxacin has a broad antibacterial spectrum, potent anti-Gram-positive activity, a long half-life, and is usually well tolerated [21]. The bioavailability of levofloxacin is as high as 99%, and once-daily oral levofloxacin is shown to be effective in treating febrile neutropenia [14,22]. Oral levofloxacin administered as monotherapy has also been used to successfully treat other diseases such as community-acquired pneumonia, chronic bronchitis, and enteric fever [21,23]. Good results have also been achieved with oral administration of the standard amoxicillin/ciprofloxacin combination. Results of first-day step-down administration of oral amoxicillin/ciprofloxacin on an outpatient basis for children with low-risk chemotherapy-induced neutropenia showed that outcomes were noninferior to the usual emergent hospitalization and intravenous antimicrobial therapy [24]. A comparison between monotherapy with once-daily oral moxifloxacin and twice-daily ciprofloxacin plus amoxicillin/clavulanic acid showed that oral moxifloxacin was safe and efficacious in low-risk febrile neutropenic patients who were discharged early and treated as outpatients; effectiveness and survival were similar in the oral monotherapy group and the combination therapy group [25]. However, the limited number of studies that compare levofloxacin monotherapy with standard intravenous therapies indicates the need for additional studies to evaluate the efficacy and safety of levofloxacin monotherapy compared with other treatments. In addition, the differences between the microbiological spectrum in febrile neutropenic patients in Asia and those in Western countries [15] as well as the prevalence of quinolone-resistant bacteria in China [18] make it important to examine the ability of levofloxacin monotherapy to treat infection and reduce fever in Chinese cancer patients with low-risk febrile neutropenia. The results of the present study have shown that levofloxacin is safe and effective in treating Chinese low-risk febrile neutropenic patients. Theoretically, the presence of infection is related to the neutropenia. For example, an ANC of less than $1000/\text{mm}^3$ predicts risk for infection; an ANC of less than $500/\text{mm}^3$ predicts increased susceptibility to infection; and an ANC of $100/\text{mm}^3$ indicates severe infection or bacteremia, but the infection may not be related to the types and sites of cancer [26]. Although our inclusion criteria used ANC less than $1500/\text{mm}^3$, other studies had different inclusion criteria, including ANC less than $500/\text{mm}^3$ or less than $1000/\text{mm}^3$ with a predicted decrease to $500/\text{mm}^3$ within the subsequent 48 h [8,26]. These differences may help to explain the differences between our results and those of other studies. As low-risk neutropenic patients with ANC less than $1500/\text{mm}^3$ were selected for the present pilot study, this may, in fact, contribute toward the rapid antibiotic response and better recovery of ANC.

The results of this pilot study are limited because we did not include a comparator treatment arm without determining whether neutrophil recovery or defervescence with levofloxacin was better than that achieved with the standard amoxicillin/ciprofloxacin regimen. We also did not consider the effects of different chemotherapy regimens and may not have had a sufficient sample size to fully support our conclusion.

In conclusion, oral levofloxacin quickly and effectively reduced fever and was well tolerated in Chinese low-risk febrile neutropenic patients with solid tumors. Although this pilot study showed the safety and efficacy of oral levofloxacin in Chinese cancer patients with febrile neutropenia, additional prospective, multicenter, large-cohort studies are needed to compare the patient data of levofloxacin treatment with those of amoxicillin/ciprofloxacin in this population, investigating both neutrophil recovery and defervescence as indicators of effective treatment for low-risk febrile neutropenia.

Acknowledgements

The authors would like to acknowledge Prof. Wen Fuqiang from West China Hospital, Sichuan University, Department of Respiratory Disease; Prof. Lin Zhuli from The First Affiliated Hospital of Guangzhou University of Chinese Medicine, Department of Oncology; Prof. Shen Huahao from The Second Affiliated Hospital, Zhengjiang University, Department of Respiratory Disease; Prof. Zhou Jianying from The First Affiliated Hospital, Zhengjiang University, Department of Respiratory Disease; and Prof. Zhou Jianying and Prof. Zhang Shucai from Beijing Chest Hospital, Capital Medical University, Department of Respiratory Disease for their contributions to this study.

This study was financially supported by Daiichi Sankyo Co. Ltd, Tokyo, Japan.

Conflicts of interest

There are no conflicts of interest.

References

- Picazo JJ. Association for Health Research and Development (ACINDES). Management of the febrile neutropenic patient. *Int J Antimicrob Agents* 2005; **Suppl 2**:S120–S122, discussion S133–S140.
- Gafter-Gvili A, Fraser A, Paul M, Leibovici L. Meta-analysis: antibiotic prophylaxis reduces mortality in neutropenic patients. *Ann Intern Med* 2005; **142** (Pt 1):979–995.
- Leibovici L, Paul M, Cullen M, Bucaneve G, Gafter-Gvili A, Fraser A, Kern WV. Antibiotic prophylaxis in neutropenic patients: new evidence, practical decisions. *Cancer* 2006; **107**:1743–1751.
- Kern WV. Risk assessment and treatment of low-risk patients with febrile neutropenia. *Clin Infect Dis* 2006; **42**:533–540.
- Klastersky J, Paesmans M, Rubenstein EB, Boyer M, Elting L, Feld R, et al. The Multinational Association for Supportive Care in Cancer risk index: a multinational scoring system for identifying low-risk febrile neutropenic cancer patients. *J Clin Oncol* 2000; **18**:3038–3051.
- Innes H, Marshall E. Outpatient therapy for febrile neutropenia. *Curr Opin Oncol* 2007; **19**:294–298.
- Carstensen M, Sørensen JB. Outpatient management of febrile neutropenia: time to revise the present treatment strategy. *J Support Oncol* 2008; **6**:199–208.
- Rolston KV, Frisbee-Hume SE, Patel S, Manzullo EF, Benjamin RS. Oral moxifloxacin for outpatient treatment of low-risk, febrile neutropenic patients. *Support Care Cancer* 2010; **18**:89–94.
- Bliotiotis IA, Michalopoulos A, Kasiakou SK, Samonis G, Christodoulou C, Chrysanthopoulou S, Falagas ME. Ciprofloxacin vs an aminoglycoside in combination with a beta-lactam for the treatment of febrile neutropenia: a meta-analysis of randomized controlled trials. *Mayo Clin Proc* 2005; **80**:1146–1156.
- Langtry HD, Lamb HM. Levofloxacin. Its use in infections of the respiratory tract, skin, soft tissues and urinary tract. *Drugs* 1998; **56**:487–515.
- Bucaneve G, Micozzi A, Menichetti F, Martino P, Dionisi MS, Martinelli G, et al. Gruppo Italiano Malattie Ematologiche dell'Adulto (GIMEMA) Infection Program. Levofloxacin to prevent bacterial infection in patients with cancer and neutropenia. *N Engl J Med* 2005; **353**:977–987.
- Reuter S, Kern WV, Sigge A, Döhner H, Marre R, Kern P, von Baum H. Impact of fluoroquinolone prophylaxis on reduced infection-related mortality among patients with neutropenia and hematologic malignancies. *Clin Infect Dis* 2005; **40**:1087–1093.
- Cullen M, Steven N, Billingham L, Gaunt C, Hastings M, Simmonds P, et al. Simple Investigation in Neutropenic Individuals of the Frequency of Infection after Chemotherapy +/- Antibiotic in a Number of Tumours (SIGNIFICANT) Trial Group. Antibacterial prophylaxis after chemotherapy for solid tumors and lymphomas. *N Engl J Med* 2005; **353**:988–998.
- Cornely OA, Wicke T, Seifert H, Bethe U, Schwonzen M, Reichert D, et al. Once-daily oral levofloxacin monotherapy versus piperacillin/tazobactam three times a day: a randomized controlled multicenter trial in patients with febrile neutropenia. *Int J Hematol* 2004; **79**:74–78.
- Chen CY, Tang JL, Hsueh PR, Yao M, Huang SY, Chen YC, et al. Trends and antimicrobial resistance of pathogens causing bloodstream infections among febrile neutropenic adults with hematological malignancy. *J Formos Med Assoc* 2004; **103**:526–532.
- Li Y, Lv Y, Xue F, Zhang XZ, Hu XJ, et al. Antimicrobial susceptibility surveillance of gram-positive bacteria from Ministry of Health National Antimicrobial Resistance Investigation Net (Mohnarin) 2011-2012. *Chin J Clin Pharmacol* 2013; **30**:251–259.
- Li Y, Lv Y, Xue F, Zhang XZ, Hu XJ, et al. Antimicrobial susceptibility surveillance of gram-negative bacteria from Mohnarin 2011-2012. *Chin J Clin Pharmacol* 2013; **30**:260–277.
- Lee CC, Lui G, Ip M, Ling TK, Lee N. Frequent detection of plasmid-mediated quinolone resistance (qnr) genes in multidrug-resistant Enterobacteriaceae blood isolates, Hong Kong. *Eur J Clin Microbiol Infect Dis* 2012; **31**:3183–3189.
- National Comprehensive Cancer Network. *NCCN Clinical Practice Guidelines in Oncology: prevention and treatment of cancer related infections. Version 1.2012*. Fort Washington, Pennsylvania: National Comprehensive Cancer Network; Available at: http://www.nccn.org/professionals/physician_gls/pdf/infections.pdf. [Accessed: 20 February 2012].
- Freifeld AG, Bow EJ, Sepkowitz KA, Boeckh MJ, Ito JI, Mullen CA, et al. Infectious Diseases Society of America. Clinical practice guideline for the use of antimicrobial agents in neutropenic patients with cancer: 2010 update by the Infectious Diseases Society of America. *Clin Infect Dis* 2011; **52**: e56–e93.
- Noreddin AM, Elkhatib WF, Cunnion KM, Zhanel GG. Cumulative clinical experience from over a decade of use of levofloxacin in community-acquired pneumonia: critical appraisal and role in therapy. *Drug Healthc Patient Saf* 2011; **3**:59–68.
- Galan-Herrera JF, Poo JL, Rosales-Sanchez O, Fuentes-Fuentes E, Cariño L, Burke-Fraga V, et al. Bioavailability of two oral formulations of a single dose of levofloxacin 500 mg: an open-label, randomized, two-period crossover comparison in healthy Mexican volunteers. *Clin Ther* 2009; **31**:1796–1803.
- Urueta-Robledo J, Ariza H, Jardim JR, Caballero A, García-Calderón A, Amábilis-Cuevas CF, et al. MOX-CB Study Group. Moxifloxacin versus levofloxacin against acute exacerbations of chronic bronchitis: the Latin American Cohort. *Respir Med* 2006; **100**:1504–1511.
- Brack E, Bodmer N, Simon A, Liebundgut K, Kühne T, Niggli FK, Ammann RA. First-day step-down to oral outpatient treatment versus continued standard treatment in children with cancer and low-risk fever in neutropenia. A randomized controlled trial within the multicenter SPOG 2003 FN study. *Pediatr Blood Cancer* 2012; **59**:423–430.
- Kern WV, Marchetti O, Drgona L, Akan H, Aoun M, Akova M, et al. Oral antibiotics for fever in low-risk neutropenic patients with cancer: a double-blind, randomized, multicenter trial comparing single daily moxifloxacin with twice daily ciprofloxacin plus amoxicillin/clavulanic acid combination therapy – EORTC infectious diseases group trial XV. *J Clin Oncol* 2013; **31**:1149–1156.
- Chamilos G, Bamias A, Efstathiou E, Zorzou PM, Kastiris E, Kostis E, et al. Outpatient treatment of low-risk neutropenic fever in cancer patients using oral moxifloxacin. *Cancer* 2005; **103**:2629–2635.