

A validated high-performance thin layer chromatography method for estimation of lornoxicam and paracetamol in their combined tablet dosage form

Abstract

Introduction: Lornoxicam (LORN) a nonsteroidal antiinflammatory drug of oxamic class is marketed in combination with Paracetamol, a common analgesic for acute inflammatory disease of joints. **Materials and Methods:** LORN and Paracetamol (PCM) were estimated at 280 nm by densitometry using silica gel 60 F₂₅₄ as stationary phase and a premix of toluene: chloroform: methanol: formic acid (3:5:1.5:0.2 v/v/v/v) as mobile phase. The method was found linear in a range of 160–560 nanograms/spot for LORN and 10 000–35 000 nanograms/spot for PCM with a correlation coefficient >0.99 for both. **Result:** PCM and LORN were well resolved with R_f 0.57 ± 0.02 and 0.75 ± 0.02, respectively. **Conclusions:** The developed high-performance thin layer chromatography method was found to be simple, specific, precise, and reproducible and can be used for the routine estimation of LORN and PCM in the combined tablet dosage form, available in market.

Key words: High-performance thin layer chromatography; lornoxicam and paracetamol; simultaneous validation

INTRODUCTION

Lornoxicam (LORN) is a nonsteroidal anti-inflammatory drug of the oxamic class with the analgesic, anti-inflammatory and antipyretic properties having chemical name (3E)-6-chloro-3-[hydroxy(pyridin-2-ylamino)methylene]-2-methyl-2,3-dihydro-4H-thieno[2,3-e][1,2]thiazin-4-one 1,1-dioxide. It has been found to be effective in Inflammatory diseases of the joints, osteoarthritis, pain following surgery, and sciatica.^[1] Unlike other oxamic, it has shorter elimination half-life of 3–5 h.^[2] Paracetamol (PCM) common analgesic have chemical name *N*-(4-hydroxyphenyl)acetamide. PCM/acetaminophen is used for relief in fevers, aches, and pains associated with many parts of the body. It has weak antiinflammatory properties. It is combined with LORN in the tablet dosage form.^[3-6]

Analytical techniques such as spectrophotometry,^[7,8] high-performance liquid chromatography (HPLC),^[9-11] high-performance thin layer chromatography (HPTLC),^[12] LC/MS/MS,^[13] etc. are reported for the detection of LORN and PCM individually in plasma and bulk pharmaceutical formulation alone, and also some spectrophotometry^[14,15] and HPTLC^[16] methods for the estimation of LORN and PCM in their combination have been reported. Hence, our aim is to develop a new accurate simple and rapid HPTLC method for the estimation of LORN and PCM in the combined tablet dosage form.

MATERIALS AND METHODS

Instrumentation

Chromatographic separation of drugs were performed on Merck TLC plates

Sanjay L. Borisagar,
H. U. Patel¹, C. N. Patel¹,
Ujjawal P. Jayswal

Department of Quality Assurance
and ¹Pharmaceutical Chemistry,
Sarvajani Pharmacy College,
Mehsana, Gujarat, India

Address for correspondence:

Mr. Sanjay L. Borisagar,
Department of Quality Assurance,
Shree Sarvajani Pharmacy College,
Mehsana, Gujarat - 384 001, India.
E-mail: sanjay86_pharma@yahoo.
com, borisagarsanjay@gmail.com

Access this article online

Website: www.phmethods.org

DOI: 10.4103/2229-4708.84440

Quick response code



precoated with silica gel 60 F₂₅₄ (20×10 cm with 250 mm layer thickness, E. Merck, Germany). The samples were applied onto the plates as a band with the width of 4 mm using Camag 100 µl sample syringe (Hamilton, Switzerland) with an applicator (AS-30, Desaga, Switzerland). Linear ascending development was carried out in a twin trough glass chamber (20×10 cm). Densitometric scanning was performed using the TLC scanner (CD 60, Desaga, Switzerland) and operated by software (proquant). Electronic balance (ACCULAB Model ALC-210.4 Huntington valley, PA), Sonicator (EN 30 US, Entertech Fastclean, Mumbai, India).

Materials

LORN and PCM working standards were obtained as a gift samples from Cirex pharmaceutical Ltd (Gundla, Mandal district, A.P., India) and Cadila Pharmaceutical Ltd (Dholka, Ahmedabad, Gujarat, India), respectively. LORN and PCM combined tablet (Claiming 8 mg of LORN and 500 mg of PCM per tablet) of two different brands Lornasafe-plus (B.No. LFS045, Mankind Pharma Ltd., New Delhi) and Lorsaid-P (B.No. LOS10001, Piramal Healthcare Ltd., Mumbai) were collected from market and analyzed for the LORN and PCM content by the proposed method. All the other chemicals and reagents were of analytical grade.

Selection of detection wavelength

The wavelength was selected at 280 nm, at which, LORN shows high absorbance. This could be used to compensate for relatively low concentration of LORN compared to PCM in the marketed formulation. In the tablet dosage form PCM and LORN were found in the ratio of 500:8 mg/tab. Hence, the selected wavelength was convenient to obtain good response peaks for both the drugs.

Preparation of solution

Standard stock solution of both drugs were prepared by dissolving 1000 mg of PCM and 16 mg of LORN in Acetonitrile to obtain 100 ml stock solution (10 000:160 µg/ml) and further diluted to get final linear concentrations.

Chromatographic condition

A premix of toluene: chloroform: methanol: formic acid (3:5:1.5:0.2 v/v/v/v), respectively was optimized for thin layer chromatography plate development. The chamber was saturated with the mobile phase at room temperature for 30 min. A run distance was kept

about 67 mm and 10 ml of the mobile phase was used for single development. The dosing speed of nitrogen applicator was kept 150 nl/sec with a predosage volume of 5 µl. Samples were applied as bands of 4 mm width with the gaps of 10 mm in between. Developed plates were dried at room temperature for 5 min. Detection was done at 280 nm using the deuterium lamp in the absorption reemission mode. The slit dimension of detection was kept to be 0.4 mm × 0.02 mm.

Method validation

This optimized HPTLC method was then validated for the parameters listed below as per International Conference on Harmonisation (ICH) guidelines.^[17]

Linearity

Different concentrations of LORN (160 to 560 ng/band) and PCM (10 000 to 35 000 ng/band) were applied on the TLC plate and peak area were measured in densitometer. The calibration curves were constructed by plotting the peak areas *vs* concentrations for both drugs and the regression equation was calculated. Each response was an average of three determinations.

Precision

Precision of the method was determined in the terms of intraday and interday variation (%RSD). Intraday precision (%RSD) was assessed by analyzing standard drug solutions (240:15 000 ng/spot, 320:20 000 ng/spot and 400:25 000 ng/spot of LORN:PCM) within the calibration range, three times on the same day. Interday precision (%RSD) was assessed by analyzing drug solutions within the calibration range on three different days over a period of a week.

Accuracy

To check accuracy of the method, recovery studies were carried out by addition of standard drug solution to preanalyzed sample solution at three different levels 50%, 100%, and 150 %. Mean percentage recovery was determined.

Specificity

The absence of any secondary spot having spectra different from LORN and PCM in the typical constituted placebo chromatogram of the tablet preparation, which may interfere with LORN and PCM peak, indicates the specificity of the analytical method.

Limit of detection and limit of quantitation

The limit of detection (LOD) and limit of quantitation (LOQ) were obtained by calculating using the standard formula as per the ICH guidelines,

$$\text{LOD}=3.3\times(\sigma/S), \text{LOQ}=10\times(\sigma/S).$$

Where σ is Standard deviation of the response and S is slope of the calibration curve.

Formulation analysis

Ten combined tablets of two different LORN and PCM tablet samples Lornasafe-plus and Lorsaid-P were finely powdered and powder equivalent to 1000 mg of PCM and 16 mg of LORN was dissolved in acetonitrile to obtain 100 ml stock solution (10 000:160 $\mu\text{g/ml}$). It was sonicated and filtered through the milipore filter and further diluted with acetonitrile to get final concentration.

RESULTS AND DISCUSSION

Development of the optimum mobile phase

Different mobile phases were tried to resolve LORN and PCM. The optimum results were obtained with mobile phase consisting of toluene: chloroform: methanol: formic acid (3:5:1.5:0.2 $v/v/v/v$). The R_f values of LORN and PCM peak were observed about 0.75 ± 0.02 and 0.57 ± 0.02 , respectively. The representative densitogram is given in [Figure 1].

Validation of the developed stability-indicating method

Linearity

The response for the drugs was found to be linear in the concentration range 160–560 ng/band for LORN and 10 000–35 000 ng/band for PCM with correlation coefficient of 0.995 and 0.997, respectively. The linear regression equation obtained are $y = 0.901(x$

+ 335.4 and $y = 0.062(x) + 1023$ for LORN and PCM, respectively [Table 1].

Precision

The % RSD values for intraday precision study were found to be not more than 1.95% and 1.44% for LORN and PCM, respectively and for interday precision were found to be not more than 1.98% and 1.95% for LORN and PCM, respectively, thus confirming precision of the method [Table 2].

Accuracy

Excellent recoveries were obtained at each level of added concentration. The results obtained ($n = 3$ for each 80%, 100%, 150% level) indicated the mean recovery for LORN 98.67–99.06% and for PCM 99.14–101.77% [Table 3].

Limit of detection

The LOD as calculated by standard formula as given in ICH guidelines was found to be 63 ng/band and 1478 ng/band for LORN and PCM, respectively.

Limit of quantitation

The LOQ as calculated by standard formula as given in ICH guidelines was found to be 192 ng/band and 4480 ng/band for LORN and PCM, respectively.

Table 1: Calibration data for linearity

Amount in nanogram/spot		AUC ^a ± %RSD ^b at 280 nm (n=3)	
LORN	PCM	LORN	PCM
160	10 000	490 ± 2.00	1622 ± 1.74
240	15 000	544 ± 1.95	1951 ± 1.44
320	20 000	628 ± 1.69	2347 ± 1.43
400	25 000	704 ± 1.94	2600 ± 1.18
480	30 000	795 ± 1.97	2923 ± 1.07
560	35 000	844 ± 1.72	3165 ± 1.15

^a AUC: Area under the curve. ^b %RSD- % relative standard deviation.
LORN: Lornoxicam, PCM: Paracetamol




Figure 1: HPTLC densitogram of Paracetamol (PCM) and Lornoxicam (LORN).

Table 2: System precision of the analytical method


Amount in nanogram/spot		AUC ± %RSD (n=3) intraday		AUC ± %RSD (n=3) interday	
LORN	PCM	LORN	PCM	LORN	PCM
240	15 000	557 ± 1.95	1986 ± 1.44	558 ± 1.98	1963 ± 1.81
320	20 000	641 ± 1.69	2391 ± 1.43	614 ± 1.82	2366 ± 1.95
400	25 000	722 ± 1.94	2562 ± 1.18	689 ± 1.97	2605 ± 1.90

LORN: Lornoxicam, PCM: Paracetamol

Table 3: Recovery study of the analytical method

Test concentration		Amount spiked		%Recovery ± %RSD	
LORN	PCM	LORN	PCM	LORN	PCM
160	10 000	80	15 000	98.78 ± 1.57	99.14 ± 1.22
160	10 000	160	20 000	99.06 ± 1.27	101.77 ± 1.43
160	10 000	240	25 000	98.67 ± 1.82	99.61 ± 1.36

LORN: Lornoxicam, PCM: Paracetamol

**Figure 2:** Chromatogram of constituted tablet placebo**Table 4: Summary of validation**

Validation parameter	LORN	PCM
Regression equation	$y=0.901(x) + 335.4$	$y=0.062(x) + 1023$
Linearity	0.995	0.997
Precision	1.69–1.95	1.18–1.44
Recovery	98.67–99.06 %	99.14–101.77 %
LOD	63 ng/band	1478 ng/band
LOQ	192 ng/band	4480 ng/band
Specificity	Specific	Specific
%Assay (lorsaid-P)	98.83%	100.18%
(lornasafe-plus)	98.23%	99.45%

LORN: Lornoxicam, PCM: Paracetamol, LOD: Limit of detection, LOQ: Limit of quantitation

Formulation analysis

The percentage Assay of LORN and PCM in two different tablet samples Lornasafe-plus and Lorsaid-P were calculated and recorded in Table 4.

Specificity

The specificity of the method was ascertained by the

absence of any other peak of placebo [Figure 2]. The validation summary is given in Table 4.

CONCLUSION

The developed HPTLC method was found to be simple, specific, precise, accurate, and reproducible and can be used for the routine estimation of LORN and PCM in the combined tablets dosage form available in market. The developed method is found to be less sensitive but more accurate and specific than the previously published method.

ACKNOWLEDGMENT

Authors are thankful to Cirex pharmaceutical Ltd (Gundla, Mandal district, A.P., India) and Cadila Pharmaceutical Ltd (Dholka, Ahmedabad, Gujarat, India) for providing the gift sample of LORN and PCM standard.

REFERENCES

- Balfour JA, Fitton A, Barradell LB. A review of its pharmacology and therapeutic potential in the management of painful and inflammatory conditions. *Drugs* 1996;51:639-57.
- Skjodt NM, Davies NM. Clinical pharmacokinetics of lornoxicam. A short half-life oxiam. *Clin Pharmacokinet* 1998;34:421-8.
- Indian Pharmacopoeia. Vol 3. New Delhi: The Controller of Publications; 2007. p. 1514.
- British Pharmacopoeia. London, UK: General Medicine Council; 1998. p. 743.
- United State Pharmacopoeia-28, National formulary-23. Asian edition. US Pharmacopoeia Convention; 2005. p. 16-23.
- Hinz, B, Cheremina, O, Brune K. Acetaminophen (Paracetamol) is a selective cyclooxygenase-2 inhibitor in man. *FASEB J* 2008;22:383-90.
- Nemutlu E, Demircan S, Kir S. Determination of Lornoxicam in pharmaceutical preparations by zero and first order derivative UV Spectrophotometric methods. *Pharmazie* 2005;60:421-5.
- Schmid J, Fedorcak A, Koss FW. A sensitive, specific colorimetric method for determining Paracetamol in human plasma. *Arzneimittelforschung* 1980;30:996-8.
- Mattimarad. Rapid RP HPLC Method for Quantitative Determination of Lornoxicam in tablets. *J Clin Pharm* 2010;1:118-24.
- Lau SN, Critchley JA. The estimation of Paracetamol and its major metabolites in both plasma and urine by a single high-performance liquid chromatography assay. *J Pharm Biomed Anal* 1994;12:1563-72.
- Lotfi M, Frida D. Simultaneous LC determination of Paracetamol and related compounds in pharmaceutical formulations using a carbon-based column. *J Pharm Biomed Anal* 2002;27:851-60.
- Nadia M, Mostaf. Stability indicating method for the determination of Paracetamol in its pharmaceutical preparations by TLC densitometric method. *J Saudi Chem Soc* 2010;14:341-4.
- Zeng YL, Chen XY, Zhang YF, Zhong DF. Determination of Lornoxicam in human plasma by LC/MS/MS. *Yao Xue Xue Bao* 2004;39:132-5.
- Lakshmi S, Lakshmi KS, Tintu T. Simultaneous spectrophotometric estimation of Paracetamol and Lornoxicam in Tablet Dosage Form. *Int J Pharm Pharma Sci* 2010;2:112-5.
- Bhavsar KC, Gaikwad PD, Bankar VH, Pawar SP. Development and validation of UV spectrophotometric method for simultaneous estimation of Paracetamol and Lornoxicam in Tablet Dosage Form. *Int J Pharm Tech* 2010;2:429-39.
- Dhara JP, Vivek PP. Simultaneous Determination of Paracetamol and Lornoxicam in Tablets by Thin Layer Chromatography Combined with Densitometry. *Int J ChemTech Res* 2010;2:1929-32.
- Q2 (R1) - Validation of analytical methods, Text and methodology, International conference on harmonization of technical requirements for registration of pharmaceuticals for human use; 2005.

How to cite this article: Borisagar SL, Patel HU, Patel CN, Jayswal UP. A validated high-performance thin layer chromatography method for estimation of lornoxicam and paracetamol in their combined tablet dosage form. *Pharm Methods* 2011;2:83-7.
Source of Support: Nil, **Conflict of Interest:** None declared.

Author Help: Online submission of the manuscripts

Articles can be submitted online from <http://www.journalonweb.com>. For online submission, the articles should be prepared in two files (first page file and article file). Images should be submitted separately.

1) **First Page File:**

Prepare the title page, covering letter, acknowledgement etc. using a word processor program. All information related to your identity should be included here. Use text/rtf/doc/pdf files. Do not zip the files.

2) **Article File:**

The main text of the article, beginning with the Abstract to References (including tables) should be in this file. Do not include any information (such as acknowledgement, your names in page headers etc.) in this file. Use text/rtf/doc/pdf files. Do not zip the files. Limit the file size to 1024 kb. Do not incorporate images in the file. If file size is large, graphs can be submitted separately as images, without their being incorporated in the article file. This will reduce the size of the file.

3) **Images:**

Submit good quality color images. Each image should be less than **4096 kb (4 MB)** in size. The size of the image can be reduced by decreasing the actual height and width of the images (keep up to about 6 inches and up to about 1800 x 1200 pixels). JPEG is the most suitable file format. The image quality should be good enough to judge the scientific value of the image. For the purpose of printing, always retain a good quality, high resolution image. This high resolution image should be sent to the editorial office at the time of sending a revised article.

4) **Legends:**

Legends for the figures/images should be included at the end of the article file.