



Contents lists available at ScienceDirect

Journal of Traditional and Complementary Medicine

journal homepage: <http://www.elsevier.com/locate/jtcm>

Original article

A randomized ethnomedicinal survey of snakebite treatment in southwestern parts of Bangladesh

Md. Nazmul Hasan^{a, b}, Nur Kabidul Azam^c, Md. Nasir Ahmed^c, Akinori Hirashima^{a, *}^a Laboratory of Pesticide Chemistry, Division of Molecular Biosciences, Department of Bioscience and Biotechnology, Faculty of Agriculture, Kyushu University, Fukuoka 812-8581, Japan^b Department of Genetic Engineering and Biotechnology, Faculty of Biological Science and Technology, Jessore University of Science & Technology, Jessore 7408, Bangladesh^c Tech B Herbal Solution, Bheramara Kushtia, Bangladesh

ARTICLE INFO

Article history:

Received 9 January 2015

Received in revised form

16 March 2015

Accepted 25 March 2015

Available online 21 April 2015

Keywords:

Medicinal plants

Ethnomedicine

Snakebite

Antivenom

Bangladesh

ABSTRACT

Snakebite is the single most important toxin-related injury, causing substantial mortality in many parts of the Africa, Asia and the Americas. Incidence of snakebite is usually recorded in young people engaged in active physical work in rural areas. The various plant parts used to treat snakebite included whole plant, leaves, barks, roots and seeds. Most bites in Bangladesh are recorded between May and October with highest number in June. Lower and upper limbs are most common sites of snakebite, but it may happen in other sites as well. Snake venom (蛇毒 *shé dú*) has been the cause of innumerable deaths worldwide. However, antiserum does not provide enough protection against venom induced hemorrhage, necrosis, nephrotoxicity and hypersensitivity reactions. Informed consent was obtained from the practitioners prior to interviews. After the survey, it is concluded that the medicinal plants used by tribal medicinal practitioners in Bangladesh for treatment against snakebite are *Acyranthes aspera* L. (土牛膝 *tǔ niú xī*), *Amaranthus Viridis* L. (野苋菜 *yě xiàng cài*), *Asparagus racemosus* Willd (總序天冬 *zǒng xù tiān dōng*) and *Emblca officinalis* Gaertn (油柑 *yóu gān*), while the non-tribal communities used 35 plant species among them, most of the plants reported as new species used against snakebite in the belonging family. The plants present a considerable potential for discovery of novel compounds with fewer side effects for treatment of antsnake venom and can, at least in Bangladesh, become a source of affordable and more easily available drugs.

Copyright © 2015, Center for Food and Biomolecules, National Taiwan University. Production and hosting by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Snakes and snakebite are mythical attars in the mind of people of the world. Based on the geographical position and climatic conditions, Bangladesh is a disaster prone country. Due to this reason, snakebite is a serious but neglected public health issue in Bangladesh. A great diversity of locales for snakes, as well as others wild species have developed in Bangladesh in the junction of Indo-Malayan, Indo-Chinese and Indo-Himalayan regions which provides excellent opportunities to make welcoming habitat for those dangerous beasts. Biting occurs generally when individuals are at

work, engaging in activities such as cultivation, fishing, plantation, wood collection, tending crops and gardening. On the other hand, when the victims are walking on rural footpaths or while sleeping in the floor, bites are fairly common. For the period of monsoon, due to the rainfall snakebite occurrences are being increased because snakes leave their shelter. As most of the houses in rural area of Bangladesh are made of mud not bricks, the snakes sometimes live in the holes of the muddy floors and suitable places in the home where people stored grains and keep flock.

In Bangladesh, particularly in rural areas, snakebite is a major cause of mortality and morbidity and it has a noteworthy effect on human health and economy through treatment related expenses and loss of productivity. Most of the cases, the victim of snakebite is a poor, young and active individual. The snakebite occurrence by division in Bangladesh presented in Table 1. The number of fatalities attributed to snakebites varies greatly by geographical area. South Asia, Southeast Asia, and sub-Saharan Africa report the highest

* Corresponding author.

E-mail address: ahirasim@agr.kyushu-u.ac.jp (A. Hirashima).

Peer review under responsibility of The Center for Food and Biomolecules, National Taiwan University.

Table 1
Snakebite distribution by division in Bangladesh.

Division	Number of snakebites	Annual incidence per 1000,000 person-years (95% CI)
Barisal	22	2667.70 (1787.20–3829.50)
Chittagong	9	397.80 (211.80–680.30)
Dhaka	22	440.00 (285.00–649.90)
Khulna	20	936.20 (104.40–750.20)
Rajshahi	20	472.70 (288.00–680.30)
Sylhet	5	321.60 (104.40–750.20)
Over all	98	623.40 (513.40–789.20)

number of bites and the high incidence is in the Entropic and other equatorial and subtropical regions.^{1–3} Each year tens of thousands of people die because of snakebites. The global incidence of snakebites, envenomation and mortality per year have been shown in Table 2.^{1–6}

Treatment of snakebite in Bangladesh was mostly subjected by old-fashioned snake charmers (Ozha). Folks used to be content with their traditional approaches of tight tourniquet, multiple incisions at bite site, application of herbal products and different rituals. The outcome was determined by chance. Even the medical experts were not well aware of the scientific approaches of management of snakebite.⁷ The treatment for snakebite is as variable as the bite itself. The only available treatment in modern science is the usage of antivenom against snakebite, which was developed by Albert Calmette, a French scientist of the Pasteur Institute in 1895, against the Indian Cobra (*Naja naja*).

The traditional therapists are the first line defense against of this illness in Bangladesh. The success of these therapists is ambiguously understood, sometimes partially by their unknown *materia medica* and occult mystical nature of their practice, but direct testimony from victims confirms success of their treatments. Although their practice was ignored by biomedicine, they serve more snakebite accident fatalities than modern practitioners. The healers, especially the elderly and spirit inspired, are reputed to have an inherent expertise to handle these cases. They are accepted as healthcare providers with a thoughtful socio-cultural understanding of their communities in both ethnic groups.⁸ Other alternative treatment involves the usage of folk traditional medicines (民間傳統藥 *mín jiān chuán tōng yào*) in snakebites. Various plants have been used against snakebite in folk traditional medicine. Traditional medicinal knowledge has been a means towards the discovery of many modern medicines.⁹ Bangladesh has a rich history of several traditional medicinal systems, among which the most notable ones are the Ayurvedic, Unani and the folk medicinal systems. Folk medicine is practiced by the traditional medical practitioners (generally known as Kavirajes by the mainstream community) who utilize simple formulations of medicinal plants in

Table 2
Global incidence of snakebite occurrences, envenomation and mortality per year.

Location	Total reported cases	Envenomation	Death/year	Reference
Europe	25,000	800	30	1–6
Middle East	20,000	15,000	100	
North USA and Canada	45,000	6,500	15	
Central and South USA	3000,000	150,000	5,000	
Asia	4 million	2 million	100,000	
Africa	1 million	500,000	20,000	
Oceania	10,000	3000	200	

most of their preparations. The southern part of Bangladesh posse's hills, mountains and costal area of Bay of Bengal, whereas the western part posses forests and hills as well. These regions of Bangladesh still remain undeveloped in different sectors including education, treatment, and transportation. Snakebite incidents and mortality rate is higher in this part of Bangladesh. Due to illiteracy and lacking of modern treatment, general public are used to depend on traditional or folk medicines. For proper documentation and finding effective folk medicine, these areas were randomly selected for this study. It was the objective of the present study to conduct a completely randomized survey of Kavirajes of south-western Bangladesh to learn more about medicinal plants used for treatment of snakebite. The expectation was that the medicinal plants used by the Kavirajes could prove to be a useful source for further scientific studies leading to discovery of more efficacious drugs.

2. Materials and methods

The present survey was carried out between September 2011 to July 2012, among Kavirajes of southwestern parts in Bangladesh which includes twelve districts (Fig. 1), namely Bagerhat, Barisal, Dinajpur, Jessore, Jenidaha, Khulna, Kurigram, Kushtia, Pabna, Pirojpur, Rajbari, Rajshahi and three tribal communities of Bangladesh namely Garo (Netrokona), Rakahine (Chittagong hill tracks region) and Santal (Rajshahi). Informed consent was obtained from the Kavirajes prior to the survey.

For this specific ethnomedicinal survey the number of visits were made to the Kavirajes to gain their confidence. Actual surveys were conducted with the help of a semi-structured open ended questionnaires and the guided field-walk method as described by Martin and Maundu.^{10,11} A total of 21 kavirajes (35–65 years) were interviewed during the survey including medicine men (*Ojha*). Kavirajes were asked specifically as to whether they know about anti-snake venom and whether they treat the snakebite on a regular basis. Kavirajes were selected based on their affirmative answer to both questions. The Kavirajes mentioned the plants with which they treated snakebite to the interviewers and took the interviewers to spots from where they collected the plants. All interviews were conducted in the Bangla language, which was spoken by both Kavirajes and the interviewers. After that, it was converted into English by the author ownself. The plants were shown along with provision of local names and the parts used. Plant specimens were collected and dried in the field and later brought back to Dhaka for complete identification at the Bangladesh National Herbarium. The collected plants were cross-checked by neighboring herbalists and traditional medicinal healers. For each species the proportion of informants who independently reported its use in snakebite was assessed.¹² Each medicinal practice was cross checked with at least 2 to 3 informants.

3. Results

Bangladesh is a country of rich of flora and fauna, where 722 numbers of medicinal plants have already been discovered and they have much variety diversity based on locality in size, shape, flowering etc. The names of a total 38 of plant species were obtained from 21 Kavirajes of the twelve districts surveyed, most of the cases used plants belong to the same area or nearby area of the healer which were naturally distributed. Among them 4 plant species were obtained from the 3 tribal community areas (Garo, Rakhain and Santal). The plant species belonged to 25 families. The Fabaceae family contributed four plants; Amaranthaceae and Solanaceae contributed three plants followed by the Acanthaceae, Apocynaceae, Asteraeae, Euphorbiaceae, Lamiaceae and Rubiaceae




Fig. 1. Map of Bangladesh highlighting ethnomedical surveyed areas.

families with two plants each. The results are summarized in Tables 3 and 4.

Among 38 plant species, two of them (*Rauwolfia serpentina* L. Benth ex-Kurz (印度蛇木 yìn dù shé mù) and *Aristolochia indica* L. (馬兜鈴 mǎ dòu líng)) were used very commonly in five districts, which are Bagerhat, Jessore, Ishwardi, Pabna and Rajshahi. Whole plants as well as plant parts like leaves, barks, roots and seeds were used for treatment. It was observed that in several instances, a single plant part (like seeds of *Cajanus cajan* L. Millsp) was used. Different plants species and the different communities use parts of the plants based on the individual Kabirajes because of their earned knowledge from their ancestor only. Overall, the maximum number of plants (21 out of 38) for treatment of snake bite was obtained from the Kavirajes of Bagerhat, Khulna and Rajbari and these are *Aegle marmelos* (L.) Corr. (木橘 mù jú), *Aerva sanguinolenta* (絹毛莧 juān máo xiàn), *Agaricus albolutescens* Zeller, *Anisomeles malabarica* (L.) R.Br. ex Sims, *Aristolochia indica* L. (馬兜鈴 mǎ dòu líng), *Cecropia peltata* L. (號角樹 hào jiǎo shù), *Clitoria ternatea* L. (蝶豆 dié dòu), *Cycas revoluta* Thunb. (蘇鐵 sū tiě), *Datura metel* L. (南洋金花 nán yáng jīn huā), *Emilia sonchifolia* (L.) DC. ex Wight (紫背草 zǐ biè cǎo), *Euphorbia milii* "Lutea" Hort (鐵海棠 tiě hǎi táng), *Ipomoea*

aquatic Forsk. (蕹菜 wèng cài), *Justicia adhatoda* L. (鴨嘴花 yā zuǐ huā), *Justicia gendarussa* L. (車桑子 chē sāng zǐ), *Leucas aspera* (Wild.) Link (蜂窩草 fēng wō cǎo), *Piper longum* L. (萹荳 biān dòu), *Rauwolfia serpentina* (L.) Benth ex Kurz (印度蛇木 yìn dù shé mù), *Sansevieria trifasciata* Prain (虎尾蘭 hǔ wěi lán), *Tylophora indica* (Burm.f) Merr., *Uraria picta* (Jacq.) DC. (兔尾草 tù wěi cǎo), *Wissadula periplocifolia* (L.) C. Presl ex Thwaites.

4. Discussion

There is a wide varieties of venome among diverse snake species but in general, snake venom (蛇毒 shé dú) contains enzymes that digest proteins which cause animals to go into shock and cause damage to body tissues and internal organs that ultimately affect the nerve functions which lead to paralysis including stopping breathing as well as heart beat which is the eventual cause of death. Snakebites in rural areas are commonly treated with plant extracts.^{13–17} Traditional healers have reputation of treating difficult snakebite cases and are trusted by their patients. In both study areas, cases of deaths in victims attended by traditional healers were very rare, (less than 3%). In a Colombian study healers

Table 3
Medicinal plants used for the treatment of snakebite by folk medicinal practitioners in the twelve districts, Bangladesh.

SL. no.	Scientific name	Family	Local name	Part(S) used	Area of information collection
1	<i>Aegle marmelos</i> (L.) Corr. (木橘 mù jú)	Rutaceae	Bel	root	Khulna
2	<i>Aerva sanguinolenta</i> (縞毛茛 jiǎn máo xiàn)	Amaranthaceae	Bish korali	leaf	Rajbari
3	<i>Agaricus albolutescens</i> Zeller	Agaricaceae	Bang chata	whole plant	Bagerhat
4	<i>Amaranthus Viridis</i> L. (野苋菜 yě xiàng cài)	Amaranthaceae	Gandhori ara	whole plant	Jessore
5	<i>Anisomeles malabarica</i> (L.) R.Br. ex Sim	Lamiaceae	Raaz-moni	whole plant	Chalna, Khulna
6	<i>Anogeissus latifolia</i>	Combretaceae	Doya		Jhenidaha
7	<i>Anthocephalus chinensis</i> (Lam.) A. Rich. Ex Walp.	Rubiaceae	Kodom	leaf, bark	Pirojpur
8	<i>Aristolochia indica</i> L. (馬兜鈴 mǎ dòu líng)	Aristolochiaceae	Iche; Ishwarmul	leaf; root	Jessore; Ishwardi upazilla, Pabna; Rajshahi; Jhenidaha
9	<i>Cajanus cajan</i> (L.) Millsp.	Fabaceae	Arhor kalai	seeds	Kurigram
10	<i>Cassia sophera</i> L.	Fabaceae	Thon-thoni	leaf	Dinajpur
11	<i>Cecropia peltata</i> L. (號角樹 hào jiǎo shù)	Moraceae	Jongli udal	whole plant	Bagerhat
12	<i>Cissus pedata</i> Lam.	Vitaceae	Goala	leaf	Jessore
13	<i>Clitoria ternatea</i> L. (蝶豆 dié dòu)	Fabaceae	Aparajita	flower, seed	Chalna, Khulna
14	<i>Couroupita guianensis</i> Aubl.	Lecythidaceae	Naglingam	leaf, bark	Gaurnadi upazilla, Barisal
15	<i>Curcuma longa</i> L. (薑黃 jiāng huáng)	Zingiberaceae	Holud	rhizome	Bheramara, Kushtia
16	<i>Cycas revoluta</i> Thunb. (蘇鐵 sū tiē)	Cycadaceae	Cycas	whole plant	Bagerhat
17	<i>Datura metel</i> L. (南洋金花 nán yáng jīn huā)	Solanaceae	Dhutura	leaf, root, fruit, seed	Bagerhat
18	<i>Eclipta prostrata</i> L. (鱧腸草 lǐ cháng cǎo)	Asteraceae	Kalo keshari shak	leaves	Kurigram
19	<i>Emilia sonchifolia</i> (L.) Wight (紫背草 zǐ biè cǎo)	Asteraceae	Chadho	whole plant	Bagerhat
20	<i>Euphorbia milii</i> "Lutea" Hort (鐵海棠 tiě hǎi táng)	Euphorbiaceae	Dudh Kata	whole plant	Bagerhat
21	<i>Ipomoea aquatic</i> Forsk. (蘿菜 luó cài)	Convolvulaceae	Kolmi Shak	whole plant; leaves	Bagerhat; Kurigram
22	<i>Justicia adhatoda</i> L. (鴨嘴花 yā zuǐ huā)	Acanthaceae	Bashok	leaf, root, flower	Khulna
23	<i>Justicia gendarussa</i> L. (車桑子 chē sāng zǐ)	Acanthaceae	Rikfol	leaf	Rajbari
24	<i>Leucas aspera</i> (Wild.) Link (蜂窩草 fēng wō cǎo)	Lamiaceae	Dondo kolosh	leaf stalk, leaf; young stems	Rajbari; Kurigram
25	<i>Morinda citrifolia</i> L. (諾麗果 nuò lì guǒ)	Rubiaceae	Boro Chad	root	Bheramara, Kushtia
26	<i>Piper longum</i> L. (胡椒 hú jiāo)	Piperaceae	Pipul	root, flower, fruit, sap	Khulna
27	<i>Polyalthia longifolia</i> (Sonn.) Thwaites (PL)	Annonaceae	Devdaru	whole plant	Bheramara, Kushtia
28	<i>Rauwolfia canescens</i> L.	Apocynaceae	Boro chanda	root	Pabna
29	<i>Rauwolfia serpentina</i> (L.) Benth ex Kurz (印度蛇木 yìn dù shé mù)	Apocynaceae	Shorpogondha; coto chada; choto chondro	leaf, root, flower, seed	Bagerhat; Ishwardi upazilla, Pabna; Rajshahi; Jhenidaha
30	<i>Sansevieria trifasciata</i> Prain (虎尾蘭 hǔ wěi lán)	Agavaceae	Bagha-chokro	whole plant	Chalna, Khulna
31	<i>Solanum torvum</i> Swartz (水茄 shuǐ qié)	Solanaceae	Tit begun	root, fruit	Pirojpur
32	<i>Solanum capsicoides</i> (刺茄 cì qié)	Solanaceae	Tit begun	seed	Barisal
33	<i>Tylophora indica</i> (Burm.f) Merr.	Asclepiadaceae	Ishil mul	leaf	Rajbari
34	<i>Uraria picta</i> (Jacq.) DC. (兔尾草 tù wěi cǎo)	Fabaceae	Rahu Chondal	whole plant	Bagerhat
35	<i>Wissadula periplocifolia</i> (L.) C. Presl ex Thwaites	Malvaceae	Naag-moni	leaf, root	Chalna, khulna

interviewed reported only 4.4% death in cases they handled.¹⁸ Medicinal plants have been used for many years to treat a great variety of diseases including envenomations by animal bites.^{19–23} These plants play a key role in world health, as they are a rich source of many natural inhibitors and pharmacologically active compounds. Many of these substances structurally resemble to biological compounds, and this similarity is the basis of their physiological action.²⁴ Its is already proved that the methanolic extract and fractions of *Serjania erecta* rich in flavonoids and tannins exhibited as powerful inhibitors of the hemorrhagic and clotting activity, possibly due to interaction with metalloproteases and thrombin-like enzymes.²⁵ Studies of several plants (*Heliconia curtispatha*, *Pleopeltis percussa*, *Brownea rosademonte*, *Bixa orellana* (紅木 hóng mù), *Trichomanes elegans*, *Struthanthus orbiculareis* and *Casearia sylvestris* (林生腳骨脆 lín shēng jiǎo gǔ cuì) describe the

inhibitory effect of all or part of the coagulant activity of snake venoms from *B. asper*, *B. jararacussu*, *B. pirajai*, *B. neuwiedi*, *B. moojeni* and *C. d. terrificus*.^{26–28} The present study documents 38 species of ethnomedicinal plants from 25 families used by the tribal healers and medical practitioners in non-tribal communities. Among them, most of the plants are reported as new species in the belonging family (about use, bioactive compounds and mode of action). They mainly use leaf, root, bark, rhizome, stem, fruit, flower, leaf stalk and whole plant as antidote against snakebite. The reported plants are administered as decoction, extracts, paste and juice. Some of these plants like *Emblia officinalis* (油柑 yóu gān) and *Rauwolfia serpentina* L. Benth ex Kurz (印度蛇木 yìn dù shé mù) have been reported to have antisnake-venom activity in various ethnomedicinal studies.^{29,30} The reported plants may contain flavonoids, tannins or some other new lead compounds which

Table 4
Medicinal plants used for the treatment of snakebite by the three major tribal medicinal practitioners in Bangladesh.

SL. no.	Scientific name	Family	Local name	Part (S) used	Tribe name and district where the information was collected
1	<i>Achyranthes aspera</i> L. (土牛膝 tǔ niú xī)	Amaranthaceae	Chaim-per-on; Mimang-khache	root, leaf, stem; seed	Rakhain tribe, Chittagong Hill Tracks; Garo Tribe, Netrokona
2	<i>Amaranthus Viridis</i> L. (野苋菜 yě xiàng cài)	Amaranthaceae	Gandhori ara	whole plant	Santal tribe, Rajshahi
3	<i>Asparagus racemosus</i> Wild. (總序天冬 zǒng xù tiān dōng)	Liliaceae	Mimang-thamache	root	Garo tribe, Netrokona
4	<i>Emblia officinalis</i> Gaertn (油柑 yóu gān)	Euphorbiaceae	Chele-chibong	fruit, bark, root	Rakhain tribe, Chittagong Hill Tracks

possess the strong inhibitory activity against the hemorrhagic and clotting activity and possibly it acts via the same mode of action described above by Farnandes et al. (2011) as anti-hemorrhage and/or anti-coagulant. Therefore, this survey created a prodigious scope for pharmacologist to find out the active compounds, mode of action and the appropriate uses of those plants.

Pharmacological studies have revealed that the extracts and fractions from some of plants used in traditional medicine are able to antagonize the activity of various crude venoms and purified toxins.^{17,27,28,31–38} Medicinal plant extracts have been shown to antagonize the activity of some venoms and toxins. Several plant species are popularly known as antsnake venom and have been scientifically investigated, such as *Eclipta sp.* (鱧腸草 *lǐ cháng cǎo*), *Curcuma longa* (薑黃 *jiāng huáng*), *Hibiscus esculentus* (秋葵 *qiū kuí*), *Casearia sp.*, *Musa paradisiaca* (香蕉 *xiāng jiāo*), *Mucuna pruriens* (黎豆 *lí dòu*), *Bauhinia forficata* (羊蹄甲 *yáng tí jiǎ*), *Annoma senegalensis*, *Mikania glomerata*, *Piper sp.* (胡椒 *hú jiāo*), *Cordia verbenacea* (破布子 *pò bù zǐ*), *Pentaclethra macroloba* and others.^{19,22,38,39} Many active ingredients from snake venoms have been purified. Their molecular structures have been identified and characterized in the most advanced laboratories.^{40–45} Many other plants such as *Aristolochia albidia* (廣防己 *guǎng fáng jǐ*); *Vitex negundo* (黃荊 *huáng jīng*), *Emblica officinalis* (油柑 *yóu gān*), *Strychnos nux vomica* (馬錢子 *mǎ qián zǐ*), *Hemidesmus indicus* (菝葜 *bá qiā*) and *Mimosa pudica* (含羞草 *hán xiū cǎo*) were reported for the antsnake-venom activity.^{46–50} Pure substances from plants shown to protect mice from ophitoxaemia are generally nitrogen-free, low-molecular weight compounds: phenolics, phytosterols (β -amyirin and sitosterol) and triterpenoids but exceptions are found in 12-methoxy-4-methylvoachalotine, an alkaloid. Proposed views advanced in indicate that these micromolecules interact with macromolecular targets; receptors and enzymes; resulting in venom-inactivation, analgesic and anti-inflammatory action.^{19,21,36,38,40,41,51,52}

5. Conclusion

Recently the World Health Organization (WHO) estimated that 80% people worldwide rely on herbal medicines for some aspect. WHO has shown great interest in documenting the use of medicinal plants used by tribes from different parts of the world. Many developing countries have intensified their efforts in documenting the ethnomedical data and scientific research on medicinal plants.

From just a brief survey of the literature, it appears that the plants used by the Kavirajes in twelve districts and three tribal medical practitioners of Bangladesh present considerable potential in the treatment of snakebite. The consensus among users indicates that plants have protective activity when administered for snakebite. Particularly plants like *Rauwolfia serpentina* L. (印度蛇木 *yīn dù shé mù*), *Emblica officinalis* Linn (油柑 *yóu gān*), *Aristolochia indica* L. (馬兜鈴 *mǎ dòu líng*) and *Morinda citrifolia* L. (諾麗果 *nuò lì guǒ*) have a high consensus agreement regarding their use in snakebite. Thus the present investigation can provide leads for specific venom inhibitory compounds from the reported thirty-eight medicinal plants that could be used in combined therapy with antiserum in the near future.

Conflicts of interest

All contributing authors declare no conflicts of interest.

Ethics committee approval

The Institutional Ethics Committee of Jessore University of Science and Technology, Bangladesh approved the present study.

Acknowledgment

The authors thankful to Md. Mizanur Rahman, Beauty Akter Rupa and Md. Zobaer Ahmed Bhuiyan, Department of Biotechnology and Genetic Engineering, University of Development Alternative, Dhaka, Bangladesh for their contribution in this work. The authors also thankful to Mr. Mafuzur Rahman, Doctoral student, Environmental System Engineering Department, Nagaoka University of Technology, Nagaoka, Niigata, Japan for his contribution in preparing the map.

References

- Kasturiratne A, Wickramasinghe AR, De Silva N, Gunawardena NK, Pathmeswaran A. The global burden of snakebite, a literature analysis and modeling based on regional estimates of envenoming and deaths. *PLoS Med.* 2008;5:218.
- Chippaux JP. Snakebites: appraisal of the global situation. *Bulletin-WHO.* 1998;76:515–524.
- Gutierrez JM, Bruno L, Guillermo L, Alexandra R, Fernando C, Yamileth A. Trends in snake bite envenomation therapy: scientific, technological and public health considerations. *Curr Pharm Des.* 2007;3:2935–2950.
- Bawaskar HS. Snake venoms and antivenoms: critical supply issues. *J Assoc Physicians India.* 2004;52:11–13.
- Nhachi C, Kasilo O. Snake poisoning in rural Zimbabwe – a prospective study. *J Appl Toxicol.* 1994;14:191–193.
- Sutherland SK. Antivenom use in Australia. Premedication, adverse reactions and the use of venom detection kits. *Med J Aust.* 1992;157:734–739.
- Faiz MA. Snake bite in Bangladesh. *The Orion.* 2006;23:322.
- Bethwell O, Daniel PK. Kenyan medicinal plants used as antivenin: a comparison of plant usage. *J Ethnobiol Ethnomed.* 2006;2:7. <http://dx.doi.org/10.1186/1746-4269-2-7>.
- Cotton CM. *Ethnobotany: Principle and Application.* New York: John Wiley and Sons; 1996:399.
- Martin GJ. *Ethnobotany: A 'People and Plants' Conservation Manual.* London: Chapman and Hall; 1995:268.
- Alexiades MN. *Collecting Ethnobotanical Data. Selecting Guidelines for Ethnobotanical Research.* The New York Botanical Garden Press; 1996:53–94.
- Trotter R, Logan M. Informant consensus: a new approach for identifying potentially effective medicinal plants. In: Etkin NI, ed. *Plants in Indigenous Medicine and Diet: Biobehavioural Approaches.* Abingdon: Routledge; 1986: p.91–112.
- Mebs D. Notes on the traditional use of plants to treat snake bite in northern Papua New Guinea. *Toxicon.* 2000;38:299–302.
- Houghton PJ, Osibogun IM. Flowering plants used against snakebite. *J Ethnopharmacol.* 1993;39:1–29.
- Asuzu IU, Harvey AL. The antsnake venom activities of *Parkia biglobosa* (Mimosaceae) stem bark extract. *Toxicon.* 2003;42:763–768.
- Yang LC, Wang F, Liu MA. Study of an endothelin antagonist from a Chinese anti-snake venom medicinal herb. *J Cardiovasc Pharmacol.* 1998;31:249–250.
- Mors WB. Plants against snakebites. *Mem Ins Oswaldo Cruz.* 1991;86(suppl 2): 193.
- Otero R, Fonnegra R, Jimenez SL, et al. Snakebites and ethnobotany in the northwest region of Colombia: part I: traditional use of plants. *J Ethnopharmacol.* 2000;71:493–504.
- Martz W. Plant with a reputation against snakebite. *Toxicon.* 1992;30: 1131–1142.
- Mors WB, Nascimento MC, Pereira BM, Pereira NA. Plant natural products active against snakebite – the molecular approach. *Phytochemistry.* 2000;55: 627–642.
- Soares AM, Januario AH, Lourenco MV, Pereira AM, Pereira PS. Neutralizing effects of Brazilian plants against snake venoms. *Drugs Future.* 2004;29: 1105–1117.
- Soares AM, Tielci FK, Marcussi S, et al. Medicinal plants with inhibitory properties against snake venoms. *Curr Med Chem.* 2005;12:2625–2641.
- Soares AM, Marcussi S, Fernandes RS, et al. Medicinal plant extracts and molecules as the source of new antsnake venom drugs. In: Rahman A, Reitz AB, Choudhary MI, eds. **Karachi-Pakistan: Bentham Science Publishers; 2009:309–346. Front Med Chem.**
- Havsteen B. Flavonoids, a class of natural products of high pharmacological potency. *Biochem Pharmacol.* 1983;32:1141–1148.
- Fernandes RS, Costa TR, Marcussi S, et al. Neutralization of pharmacological and toxic activities of *Bothrops jararacussu* snake venom and isolated myotoxins by *Serjania erecta* methanolic extract and its fraction. *J Venom Anim Toxins Incl Trop Dis.* 2011;17.
- Núñez V, Otero R, Barona J, et al. Neutralization of the edema-forming, defibrinating and coagulant effects of *Bothrops asper* venom by extracts of plants used by healers in Colombia. *Braz J Med Biol Res.* 2004;37:969–977.
- Borges MH, Soares AM, Rodrigues VM, et al. Effects of aqueous extract of *Casearia sylvestris* (Flacourtiaceae) on actions of snake and bee venoms and on

- activity of phospholipases A2. *Comp Biochem Physiol B Biochem Mol Biol*. 2000;127:21–30.
28. Borges MH, Soares AM, Roddrigues VM, et al. Neutralization of proteases from *Bothrops* snake venoms by the aqueous extract from *Casearia sylvestris* (Flacourtiaceae). *Toxicon*. 2001;39:1863–1869.
 29. Sarkhel S, Chakravarty AK, Das R, Gomes A, Gomes A. A snake venom-neutralizing factor from the root extract of *Emblia officinalis* Linn. *Orient Pharm Exp Med*. 2011;11:25–33.
 30. Dey A, De JN. Ethnomedicinal aspects of *Rauwolfia serpentina* (L.) benth ex kurz. in India, Nepal and Bangladesh. *J Med Plants Res*. 2011;5:144–150.
 31. Biondo R, Pereira AM, Marcussi S, Pereira PS, Franca SC, Soares AM. Inhibition of enzymatic and pharmacological activities of some snake venoms and toxins by *Mandevilla velutina* (Apocynaceae) aqueous extract. *Biochimie*. 2003;85:1017–1025.
 32. Biondo R, Soares AM, Bertoni BW, Franca SC, Pereira AMS. Direct organogenesis of *Mandevilla illustris* (Vell) Woodson and effects of its aqueous extract on the enzymatic and toxic activities of *Crotalus durissus terrificus* snake venom. *Plant Cell Reports*. 2004;22:549–552.
 33. Januario AH, Santos SL, Marcussi S, et al. Neo-clerodane diterpenoid, a new metalloprotease snake venom inhibitor from *Baccharis trimera* (Asteraceae): anti-proteolytic and anti-hemorrhagic properties. *Chem-Biol Interact*. 2004;150:243–251.
 34. www.kent.ac.uk/sac/staff-profiles/profiles/staff_pdfs/alexiaades_miguel/thesis.pdf.
 35. Da-Silva JO, Coppede JS, Fernandes VC, et al. Antihemorrhagic, antinucleolytic and other antiophidian properties of the aqueous extract from *Pentaclethra macroloba*. *J Ethnopharmacol*. 2005;100:145–152.
 36. Mariorano VA, Marcussi S, Daher MA, et al. Antiophidian properties of the aqueous extract of *Mikania glomerata*. *J Ethnopharmacol*. 2005;102:364–370.
 37. Oliveira CZ, Maiorano VA, Marcussi S, et al. Anticoagulant and anti-fibrinolytic properties of the aqueous extract from *Bauhinia forficata* against snake venoms. *J Ethnopharmacol*. 2005;98:213–216.
 38. Ticli FK, Hage LI, Cambraia RS, et al. Rosmarinic acid, a new snake venom phospholipase A2 inhibitor from *Cordia verbenacea* (Boraginaceae): antiserum action potentiation and molecular interaction. *Toxicon*. 2005;46:318–327.
 39. Diogo LC, Fernandes RS, Marcussi S, et al. Inhibition of snake venoms and phospholipases A2 by extracts from native and genetically modified *Eclipta alba*: isolation of active coumestans. *Basic Clin Pharmacol Toxicol*. 2009;104:293–299.
 40. Ferreira LA, Henriques OB, Andreoni AA, et al. Antivenom and biological effects of *ar*-turmerone isolated from *Curcuma longa* (Zingiberaceae). *Toxicon*. 1992;30:1211–1218.
 41. Selvanayagam ZE, Gnanavendhan SG, Balakrishna K, et al. Ehretianone, a novel quinonoid xanthene from *Ehretia buxifolia* with antisnake venom activity. *J Nat Prod*. 1996;59:664–667.
 42. Abubakar MS, Sulf MI, Pateh UU, Abdurahman EM, Haruna AK, Jahun BM. *In vitro* snake venom detoxifying action of the leaf extract of *Guiera senegalensis*. *J Ethnopharmacol*. 2000;69:253–257.
 43. Batina MF, Cintra AC, Veronese EL, et al. Inhibition of the lethal and myotoxic activities of *Crotalus durissus terrificus* venom by *Tabernaemontana catharinensis*: identification of one of the active components. *Planta Medica*. 2000;66:424–428.
 44. Aguiy J, Guerranti R, Pagani R, Marinello E. Blood chemistry of rats pretreated with *Mucuna pruriens* seed aqueous extract MP101UJ after *Echis carinatus* venom challenge. *Phytother Res*. 2001;15:712–714.
 45. Mahanta M, Mukherjee AK. Neutralization of lethality, myotoxicity, and toxic enzymes of *Naja kaouthia* venom by *Mimosa pudica* root extracts. *J Ethnopharmacol*. 2001;75:55–60.
 46. Haruna AK, Choudhury MK. *In vivo* antisnake venom activity of a furanoid diterpene from *Aristolochia alba* Duch (aristolochiaceae). *Indian J Pharm Sci*. 1995;57:222–224.
 47. Alam MI, Gomes A. Snake venom neutralization by Indian medicinal plants (*Vitex negundo* and *Emblia officinalis*) root extracts. *J Ethnopharmacol*. 2003;86:75–80.
 48. Chatterjee I, Chakravarty AK, Gomes A. Antisnake's venom activity of ethanolic seed extract of *Strychnos nux vomica* linn. *Indian J Exp Biol*. 2004;42:468–475.
 49. Chatterjee I, Chakravarty AK, Gomes A. *Daboia russellii* and *Naja kaouthia* venom neutralization by lupeol acetate isolated from the root extract of Indian *Sarsaparilla hemidesmus indicus* R.Br. *J Ethnopharmacol*. 2006;106:38–43.
 50. Girish KS, Mohanakumari HP, Nagaraju S, Vishwanath BS, Kemparaju K. Hyaluronidase and protease activities from Indian snake venoms: neutralization by *Mimosa pudica* root extract. *Fitoterapia*. 2004;75:378–380.
 51. Alam MI, Gomes A. Adjuvant effect and antiserum action potentiation by a (herbal) compound 2-hydroxy-4-methoxy benzoic acid isolated from the root extract of the Indian medicinal plant 'Sarsaparilla' (*Hemidesmus indicus* R.Br.). *Toxicon*. 1998;36:1423–1431.
 52. Vilegas JHY, Lançasa FM, Vilegas W, Pozettib GL. Steroids and Furocoumarins from Brazilian medicinal plants of *Dorstenia* genus (Moraceae). *J Braz Chem Soc*. 1997;8:529–535.