

RESEARCH

Open Access

# Increased prevalence of the *pfdhfr*/*phdhps* quintuple mutant and rapid emergence of *pfdhps* resistance mutations at codons 581 and 613 in Kisumu, Kenya

Maroya D Spalding<sup>1,4</sup>, Fredrick L Eyase<sup>2</sup>, Hoseah M Akala<sup>2</sup>, Sheryl A Bedno<sup>3</sup>, Sean T Prigge<sup>4</sup>, Rodney L Coldren<sup>3</sup>, William J Moss<sup>1,4</sup>, Norman C Waters<sup>5\*</sup>

## Abstract

**Background:** Anti-malarial drug resistance in Kenya prompted two drug policy changes within a decade: sulphadoxine-pyrimethamine (SP) replaced chloroquine (CQ) as the first-line anti-malarial in 1998 and artemether-lumefantrine (AL) replaced SP in 2004. Two cross-sectional studies were conducted to monitor changes in the prevalence of molecular markers of drug resistance over the period in which SP was used as the first-line anti-malarial. The baseline study was carried out from 1999-2000, shortly after implementation of SP, and the follow-up study occurred from 2003-2005, during the transition to AL.

**Materials and methods:** Blood was collected from malaria smear-positive, symptomatic patients presenting to outpatient centers in Kisumu, Kenya, during the baseline and follow-up studies. Isolates were genotyped at codons associated with SP and CQ resistance. *In vitro* IC<sub>50</sub> values for antifolates and quinolones were determined for isolates from the follow-up study.

**Results:** The prevalence of isolates containing the *pfdhfr* N511/C59R/S108N/*phdhps* A437G/K540E quintuple mutant associated with SP-resistance rose from 21% in the baseline study to 53% in the follow-up study ( $p < 0.001$ ). Isolates containing the *pfdhfr* I164L mutation were absent from both studies. The *phdhps* mutations A581G and A613S/T were absent from the baseline study but were present in 85% and 61%, respectively, of isolates from the follow-up study. At follow-up, parasites with mutations at five *phdhps* codons, 436, 437, 540, 581, and 613, accounted for 39% of isolates. The CQ resistance-associated mutations *pfcr* K76T and *pfmdr1* N86Y rose from 82% to 97% ( $p = 0.001$ ) and 44% to 76% ( $p < 0.001$ ), respectively, from baseline to follow-up.

**Conclusions:** During the period in which SP was the first-line anti-malarial in Kenya, highly SP-resistant parasites emerged, including isolates harboring *phdhps* mutations not previously observed there. SP continues to be widely used in Kenya; however, given the highly resistant genotypes observed in this study, its use as a first-line anti-malarial should be discouraged, particularly for populations without acquired immunity to malaria. The increase in the *pfcr* K76T prevalence, despite efforts to reduce CQ use, suggests that either these efforts are not adequate to alleviate CQ pressure in Kisumu, or that drug pressure is derived from another source, such as the second-line anti-malarial amodiaquine.

## Background

In Kenya, a sub-Saharan African nation of 34.7 million people, approximately 70% of the population is at risk for malaria infection, and malaria accounts for approximately 20% of admissions to health facilities and 30-50% of

outpatient clinic visits [1]. Effective treatment of malaria in Kenya has been hampered by high levels of drug resistance, both to chloroquine (CQ), which was the first-line anti-malarial for uncomplicated malaria in Kenya until 1998 [2], and to its replacement, sulphadoxine-pyrimethamine (SP). Widespread treatment failures were observed within five years of implementing SP as the first-line anti-malarial, and in 2004 the recommended treatment for

\* Correspondence: norman.waters@us.army.mil

<sup>5</sup>Walter Reed Army Institute of Research, Silver Spring, Maryland, USA  
Full list of author information is available at the end of the article

uncomplicated malaria was changed to the artemisinin combination therapy (ACT), artemether-lumefantrine (AL) [3]. More than two years elapsed before the training of health care workers and distribution of AL were completed in 2006 [3]. Although no longer the official first-line anti-malarial, SP remains available and is frequently used in the private sector [4]. It also plays an important role in the prevention of malaria in pregnancy when used for intermittent presumptive treatment of malaria in pregnant women (IPTp) [5] and has been included in several trials of intermittent presumptive treatment in infants (IPTi) [6-11].

The efficacy of SP depends on the inhibition of two parasite enzymes necessary for the synthesis and recycling of folate: dihydrofolate reductase (DHFR), the target of pyrimethamine, and dihydropteroate synthase (DHPS), the target of sulphadoxine. The key event in the development of pyrimethamine resistance is a mutation at codon 108 that changes serine to asparagine, resulting in partial pyrimethamine resistance. Further mutation at codons 51 (N51I) and/or 59 (C59R) increases the severity of pyrimethamine resistance [12,13]. An additional mutation, I164L, is associated with high-level SP resistance and has been found in the presence of triple mutant *pfdhfr* in South American and Asian isolates [14,15]. This mutation appears to be rare in Africa [16], but has been detected at low prevalence in isolates from Kisumu and Western Kenya [17-19]. In the background of double or triple mutant *pfdhfr*, mutations that enhance resistance to antifolates arise in the *pfdhps* gene. Five mutations in *pfdhps* (S436A/F, A437G, K540E, A581G, and A613S/T) have been implicated in sulphadoxine resistance [20,21]. *In vivo*, the *pfdhps* double mutant A437G/K540E is associated with clinical SP failure in Africa [13]. Mutations at codons 581 and 613 have begun to emerge in Africa in the background of the A437G mutation [22,23]. These mutations are widespread in South America and South East Asia, and are associated with high-level SP resistance [15,24]. Together, the *pfdhps* double mutant A437G/K540E and the *pfdhfr* triple mutant S108N/N51I/C59R form the *pfdhfr/pfdhps* quintuple mutant, which is associated with SP treatment failure [13].

Chloroquine resistance is associated with mutations in two transporters, the chloroquine resistance transporter (PfCRT) and the transmembrane glycoprotein Pgh1, a homolog of the mammalian multidrug resistance transporter encoded by the gene *pfmdr1*. The *pfcr* mutation K76T is sufficient for CQ resistance [25]. The *pfmdr1* mutation N86Y is linked with CQ resistance and mefloquine sensitivity [26,27]. In contrast, quinine and mefloquine resistance are associated with parasites that are wild-type at codon 86, but have increased *pfmdr1* copy number [26]. Among other common *pfmdr1* polymorphisms, the single

mutation N1042D and the allele combination S1034C/N1042D/D1286Y are associated with increased *in vitro* resistance to quinine and increased susceptibility to mefloquine and artemisinin [28]. *In vivo*, treatment with AL selects for parasites that are wild-type at codons 86 and 1246 and mutant at codon 184 [29,30]. In Africa, where CQ has exerted the most drug pressure on these alleles, the *pfcr* K76T haplotype and single copy *pfmdr1* with the N86Y haplotype predominate. While both markers are associated with CQ resistance, *pfcr* K76T is more strongly associated with *in vivo* CQ resistance than either *pfmdr1* N86Y or both markers together [31]. Research in Malawi has shown that fitness costs imparted by the K76T mutation result in the return of CQ sensitivity in the absence of CQ pressure. In 1993, CQ was replaced as the first-line anti-malarial in Malawi. By 2001, the K76T mutation had completely disappeared, and in 2006, CQ was demonstrated to again be effective *in vivo* [32,33].

In 1999, shortly after SP became the first-line anti-malarial in Kenya, the U.S. Army initiated a study of the prevalence of molecular markers of CQ and SP resistance in Kisumu, Kenya, a malaria endemic area near Lake Victoria. To evaluate changes in molecular markers of SP and CQ resistance in the presence of continued SP selection pressure, and presumably decreased CQ selection pressure, a follow-up study was conducted from 2003-2005. During this time period, the national first-line anti-malarial policy changed from SP to AL due to widespread SP treatment failures, although AL was not fully implemented until after completion of the follow-up study [3]. Here, changes in the prevalence of drug resistance-associated *pfdhfr*, *pfdhps*, *pfcr*, and *pfmdr1* haplotypes in the malaria-endemic area of Kisumu, Kenya are compared between the baseline and follow-up studies.

## Methods

### Study site and *Plasmodium falciparum* patient isolates

The goal of this study was to compare the prevalence of molecular markers of drug resistance between the period directly following adoption of SP as the first-line anti-malarial in 1998 and approximately five years later. A baseline study of resistance markers was conducted from 1999-2000 and was reported previously [34]. The findings of a second study conducted from August 2003 through July 2005 are described here and compared to the baseline study results. During both study periods, *P. falciparum* isolates were collected from persons in Kisumu, Kenya, a city located on the shore of Lake Victoria in a region where malaria is holoendemic. Patients of all ages who presented at the outpatient clinics of New Nyanza Provincial Hospital, Kisumu, Kenya, tested positive for malaria by thick smear, and who had not taken anti-malarial drugs within 24 hours were eligible.

Informed consent was obtained from all patients, or their parents or guardians. Venous blood for parasite genotyping and *in vitro* drug testing was collected from study subjects prior to anti-malarial treatment. The studies and protocols were approved by the Institutional Review Boards at the Walter Reed Army Institute of Research (WRAIR) and the Kenya Medical Research Institute (KEMRI).

### Genotyping

Blood spots from venous blood collected prior to treatment were dried on filter paper and DNA was isolated by Chelex extraction. To control for variation between the studies, genotype analyses for the baseline and follow-up studies were conducted under identical laboratory conditions by the same personnel, and laboratory strains were included as controls. Parasite genotypes at *pfdhfr* codons 51, 59, 108 (detection of S108N only), and 164, *pfdhps* codons 436, 437, 540, 581, and 613, and *pfcr* codon 76 were determined by mutation-specific nested PCR. The full-length *pfdhfr* and *pfdhps* genes were amplified in a primary PCR reaction according to the methods of Duraisingh [35] and Wang [36], respectively. Allele specific PCR analyses were performed according to published methods [13]. The genotype of *pfcr* codon 76 was determined by nested PCR according to the methods described by Djimde and colleagues [31]. The genotypes of *pfmdr1* codons 86, 184, 1034, 1042, and 1246 were analysed by PCR-RFLP [31]. Genotypes were initially classified as wild-type, mutant, or mixed wild-type and mutant. Isolates were classified as mixed genotype when strong bands of similar intensity were detected for both the wild-type and mutant genotypes. To determine changes in the mutation prevalence at each locus, isolates were grouped as wild-type (pure wild-type only) or mutant (pure and mixed mutant). For each codon, isolates missing data for that codon were excluded from the mutation prevalence analysis.

### *pfdhfr/pfdhps* haplotype analysis

The prevalence of the *pfdhfr* triple mutant allele N511/C59R/S108N, the *pfdhps* double mutant allele A437G/K540E, and the *pfdhfr/pfdhps* quintuple mutant were determined using a modification of the method described by Kublin and colleagues [13]. This method is particularly useful for mixed genotype isolates, such as those observed here. In these isolates, it is difficult to ascertain haplotype because the observed haplotype is a combination of the haplotypes of all the parasite lines in the infection; the method of Kublin and co-workers accounts for this when classifying haplotypes [13].

*Pfdhfr* and *pfdhps* haplotypes were first classified as wild-type, single mutant, double mutant (mixed or pure), or, in the case of *pfdhfr*, triple mutant (mixed or

pure) (Additional file 1). These haplotypes were then used to determine the *pfdhfr/pfdhps* haplotype (Additional file 2). Although in the published method mutations at codons 51 or 59 were always accompanied by mutant codon 108, several of the Kisumu isolates had mutations at codons 51 and/or 59 but not 108. Thus, in these samples, mutation at codon 108 could not be considered a prerequisite for further *pfdhfr* mutations, and the method was modified to classify isolates with mixed mutations at two *pfdhfr* loci as single mutant (Additional file 1). The published method was further modified by creating a double mutant mixed *pfdhps* category to account for the high number of mixed haplotypes in the samples (Additional file 1).

Due to the high prevalence of *pfdhps* mutations at codons 436, 581, and 613 in the follow-up study, expanded *pfdhps* haplotypes that included these codons were constructed. Using the *pfdhps* A437G/K540E haplotypes (Additional file 1), it was determined whether isolates that were wild-type, single mutant, or double mutant at codons 437 and 540 also carried mutations (mixed and pure mutant combined) at codons 436, 581, and 613. Isolates for which data were missing for any of *pfdhfr* codons 51, 59, or 108 or *pfdhps* codons 436, 437, 540, 581, and 613 were excluded from all haplotype analyses.

### *pfcr* K76T allele frequency estimation

The frequency of the K76T allele in the baseline and follow-up studies was estimated using the formulas described by Hill and Babiker, assuming a conditional Poisson distribution for the number of clones [37].

### *In vitro* drug susceptibility assays

Red blood cells from venous blood samples taken from smear positive patients were resuspended in folate-free RPMI (Gibco) supplemented with 10% human serum and buffered with HEPES and NaHCO<sub>3</sub>. The culture suspension was added to 96-well plates containing serial dilutions of the anti-malarial drugs sulphadoxine, dapsone, pyrimethamine, chlorcycloguanil, proguanil, chloroquine, mefloquine, quinine, and amodiaquine. The reference strains D2 and W6 were included as controls. Inhibition of parasite growth was assessed by <sup>3</sup>H-hypoxanthine uptake as previously described [38]. The *in vitro* IC<sub>50</sub> cutoff values used to classify parasites as drug sensitive or resistant were based on published values determined by the correlation of *in vitro* IC<sub>50</sub> values with *in vivo* treatment failure (for CQ) and by the statistical definition of resistance as an IC<sub>50</sub> value greater than two standard deviations above the mean IC<sub>50</sub> of sensitive parasites (all other drugs). IC<sub>50</sub> values considered discriminative for resistance were 100 nM (32 ng/ml) for CQ [39]; 800 nM (259.5 ng/ml) for quinine, with 500 nM

(162.2 ng/ml) considered a marker of intermediate susceptibility [40]; 30 nM (11.4 ng/ml) for mefloquine; 80 nM (28.5 ng/ml) for amodiaquine [41]; 2 μM (497.4 ng/ml) for pyrimethamine with intermediate levels of susceptibility denoted by IC<sub>50</sub> values between 500 nM (124.5 ng/ml) and 2 μM [42]; 32 μM (10 μg/ml) for sulphadoxine [43]; and 12 μM for dapson (3 μg/ml) [44].


### Statistical analysis

Changes in prevalence by study period were analysed by Fischer's exact test if one or more cells contained five or fewer samples, and by the  $\chi^2$  test for all other analyses. For all analyses, statistical significance was determined at the  $p < 0.05$  level. Statistical analyses were performed with Stata Statistical Software: Release 9.0 (StataCorp, College Station, TX).

### Results

Mutations at four codons in *pfdhfr* and at five codons in *pfdhps* were successfully analysed for 38 to 40 out of 40 total samples from the baseline study (1999-2000) and 220 to 243 out of 250 total samples from the follow-up study (2003-2005). The prevalence of mutant haplotypes (mixed and pure mutant) at each locus was compared between the study periods. In the baseline study [34], the prevalence of patient isolates containing the *pfdhfr* mutations N51I and C59R was high, totaling 97.5% and 100%, respectively, while the prevalence of the mutation S108N was slightly lower at 87.5%. In the follow-up study, the mutation prevalence at codons 51 and 59 was 99.6% and 97.0%, respectively, and did not change significantly from the baseline study ( $p = 0.14$  and  $p = 0.28$ , respectively) (Figure 1A). The prevalence of the S108N mutation increased significantly to 99.5% ( $p < 0.001$ ). Thus, after approximately five years of SP use in Kisumu, over 95% of symptomatic patients had parasites with mutations at *pfdhfr* codons 51, 59, and 108 in either mixed or pure form. The highly drug resistant mutation I164L was not detected in either study period.


The mutations A437G and K540E are the *pfdhps* mutations most strongly associated with SP treatment failure in Africa [13]. The *pfdhps* mutations S436A, A581G, and A613S/T combined with A437G increase the IC<sub>50</sub> of sulfa drugs in *in vitro* culture relative to A437G alone [20,21], although their role in *in vivo* SP failure is not well understood. The prevalence of A437G and K540E increased from 37.5% to 98.8% ( $p < 0.001$ ) and from 57.5% to 96.3% ( $p < 0.001$ ), respectively, between the baseline and follow-up study periods (Figure 1B). The prevalence of S436A/F increased from 23.7% to 68.6% between the study periods ( $p < 0.001$ ), following the trend of increasing mutation prevalence observed for codons 437 and 540 (Figure 1B). The mutations A581G and A613S/T were not detected in



**Figure 1** Prevalence of patient isolates with mutations in (A) *pfdhfr* and (B) *pfdhps* from the baseline study (study period 1) and follow-up study (study period 2).

the initial study period; however, these mutations were present in 85.1% ( $p < 0.001$ ) and 60.7% ( $p < 0.001$ ) of isolates from the second study period, respectively.


Changes in prevalence were also analysed for the three *pfdhfr* and *pfdhps* haplotypes that have been associated with clinical failure of SP: triple mutant *pfdhfr* N51I/C59R/S108N, double mutant *pfdhps* A437G/K540E, and the *pfdhfr/pfhps* quintuple mutant formed from these combined haplotypes. Haplotype analyses were performed on 38 of the 40 isolates from the baseline study and 200 of the 250 isolates from the follow-up study that had complete genotype data for *pfdhfr* codons 51, 59, and 108 and *pfdhps* codons 436, 437, 540, 581, and 613. The prevalence of the *pfdhfr* triple mutant (mixed and pure mutant combined) decreased from 81.5% in the baseline study to 68.0% in the follow-up study, with a coincident rise in the prevalence of *pfdhfr* double mutants from 18.4% to 32.0% ( $p < 0.001$ ) (Figure 2A). The percentage of isolates containing the *pfdhps* double mutant (mixed and pure combined) increased from 23.7% in the baseline study period to 76.5% in the follow-up period ( $p < 0.001$ ) (Figure 2B). The prevalence of parasites wild-type at *pfdhps* codons 437 and 540 decreased between the two study periods from 31.6% to 0.5%. The *pfdhfr/pfhps* quintuple mutant, in either



mixed or pure form, is the most clinically relevant molecular marker for SP resistance. The prevalence of mixed and pure quintuple mutant parasites increased from 21.1% to 53.5% in the three to five year interval between studies ( $p < 0.001$ ) (Figure 2C).

Based on the high prevalence of *pfdhps* mutations at codons other than 437 and 540 in the follow-up study, *pfdhps* genotypes were constructed based on whether

isolates that were wild-type, single, or double mutant at codons 437 and 540 had additional mutations at codons 436, 581, and 613. In the baseline study, isolates that were wild-type at codons 436, 581, and 613 and single or double mutant at codons 437 and 540 accounted for 34.2% and 15.8% of isolates, respectively. Only 7.9% of isolates that were A437G/K540E double mutant also had an additional mutation (Figure 3A). In the follow-up study, isolates that were A437G/K540E double mutant with additional mutations in *pfdhps* predominated, such that 14.5% of isolates had one additional mutation, 17.5% had two additional mutations, and 38.5% had three additional mutations (Figure 3A). Only 5.5% contained the A437G/K540E double mutant without additional mutations in *pfdhps*. An additional 17.5% of isolates were single mutant with respect to codons 437 and 540 and carried mutations at codons 436, 581, and 613. In the baseline study, A437G/K540E double mutant isolates were found with one additional mutation, S436A/F; however, in the follow-up study, the majority of isolates with one additional mutation carried A581G (Figure 3B). Among double mutant isolates with



**Table 1 *In vitro* geometric mean IC<sub>50</sub> data for antifolates in the follow-up study**


Drug	N	Mean IC <sub>50</sub> (ng/ml)	Range (ng/ml)	Percent resistant
pyrimethamine	43	171	21-2028	16.3 (53.5) <sup>a</sup>
chlorcycloguanil	41	2	0.5-11	
proguanil	41	929	129-3376	
sulphadoxine	41	9310	890-48,153	51.2
dapsone	27	648	16-129	14.8

<sup>a</sup> percent resistant using the intermediate level of susceptibility of 500 nM (124.5 ng/ml) as the cutoff

two additional mutations, S436A/F and A581G most commonly occurred together (Figure 3C).

In addition to evaluating changes in parasite genotypes associated with drug resistance, *in vitro* IC<sub>50</sub> values for the common antifolates pyrimethamine, proguanil, chlorcycloguanil, and dapsone were determined for between 27 and 43 isolates from the follow-up study (Table 1). The geometric mean IC<sub>50</sub> of pyrimethamine was 171.0 ng/ml and the geometric mean IC<sub>50</sub> of sulphadoxine was 9.3 µg/ml.

To evaluate whether changes in quinoline resistance occurred after SP replaced chloroquine as the first-line anti-malarial, the prevalence of mutations in the genes *pfprt* and *pfmdr1* were compared between the baseline and follow-up studies. Mutations at *pfprt* codon 76 and at *pfmdr1* codons 86, 184, 1034, 1042, and 1246 were successfully analysed in 36 to 40 out of 40 samples from the 1999-2000 study period and 243 to 247 out of 250 samples from the follow-up period (with the exception of *pfmdr1* codon 1246, for which 90 samples were genotyped). The prevalence of the *pfprt* mutation K76T in patient isolates increased between the study groups from 82.1% to 97.1% (p = 0.001) (Figure 4). In the *pfmdr1* gene, significant changes in mutation prevalence were observed at codons 86, 1034, and 1246. The



**Table 2 *In vitro* geometric mean IC<sub>50</sub> data for quinolines in the follow-up study**

Drug	N	Mean IC <sub>50</sub> (ng/ml)	Range (ng/ml)	Percent resistant
chloroquine	60	15.3	3-110	15
mefloquine	62	12.0	1-48	51.6
quinine	61	47.7	14-313	1.6 (4.9) <sup>a</sup>
amodiaquine	54	7.4	3-24	0

<sup>a</sup> percent resistant using the intermediate level of susceptibility of 500 nM (162.2 ng/ml) as the cutoff

prevalence of the N86Y mutation increased from 43.6% to 75.7% (p < 0.001), while at codons 1034 and 1246 mutation prevalence decreased from 100% to 6.1% (p < 0.001) and 31.5% to 4.4% (p < 0.001), respectively. The mutation prevalence at codon 184 remained at 100% for both the baseline and follow-up studies, and the prevalence of mutant codon 1042 was similar between the baseline and follow-up studies (90.0% and 84.2%, respectively; p = 0.475) (Figure 4).

The prevalence of wild-type *pfprt* 76 has been increasing in coastal Kenya [45] and in Tanzania [46]; therefore, it was important to evaluate whether Kisumu isolates deviated from this trend due to an increase in the frequency of the K76T allele or as a consequence of the higher prevalence of polyclonal infections in the follow-up study compared to the baseline study. The frequency of the K76T allele in the baseline and follow-up studies was estimated using the method of Hill and Babiker [37]. The frequency of K76T was estimated to be 77.8% (63.7%-88.5%) in the baseline study and 85.0% (81.8%-87.9%) in the follow-up study.

IC<sub>50</sub> values for 54 to 62 samples from the follow-up study were obtained for CQ, mefloquine, quinine, and amodiaquine (AQ) (Table 2). Geometric mean IC<sub>50</sub> values for CQ, mefloquine, quinine, and AQ were 15.3, 12.0, 47.7, and 7.4 ng/ml, respectively. The prevalence of *in vitro* resistance to CQ, mefloquine, and quinine was also determined. The CQ resistance level is of particular interest because the cutoff of 100 nM for resistant parasites was associated with *in vivo* treatment failure in Cameroon [39], in contrast to other resistance cut-offs that were determined statistically. *In vitro* resistance to CQ was observed in 15% of the isolates; quinine *in vitro* resistance was 1.6%, with 4.9% showing intermediate levels of susceptibility; and 51.6% of isolates were resistant to mefloquine.

## Discussion

The first-line anti-malarial treatment regimen for uncomplicated malaria in Kenya changed from CQ to SP in 1998 and from SP to AL in 2004. In both instances, the change was necessitated by widespread treatment failure. The prevalence of SP and chloroquine

resistance-associated genotypes and *in vitro* resistance was compared between the period just after SP implementation (1999-2000) and a period five to seven years later (2003-2005) in patients with febrile illness presenting to outpatient clinics in Kisumu, Kenya. Malaria in Kisumu and in other areas surrounding Lake Victoria is endemic. For the 12 months ending in June 2004, the entomological inoculation rate in Kisumu district was 31.1 infective bites per person per year [47], and in the time period 2003-2006, the median *P. falciparum* parasite rate for children 2 to 10 years of age (PfPR<sub>2-10</sub>) was 71% in Kisumu [48]. Despite changes in government anti-malarial policy, as late as 2010 SP was used to treat malaria in 37% of households surveyed in Kisumu, compared to 32% that used ACT [4].

The prevalences of *pfdhfr* codons N51I, C59R, and S108N were near saturation in the baseline study [34] and remained relatively stable between the baseline and follow-up studies. In contrast, the prevalence of mutations at all five *pfdhps* codons analysed increased dramatically over the same period. It thus appears that a significant proportion of isolates carried *pfdhfr* mutations prior to 1998, and use of SP as the first-line anti-malarial in Kenya resulted in the development of highly mutant *pfdhps*. Pre-existing mutations in *pfdhfr* may have derived from a combination of two sources: the use of SP as a second-line anti-malarial in Kenya prior to 1998 and the ongoing treatment of persons with HIV/AIDS with co-trimoxazole, a bacterial DHFR/DHPS inhibitor used to treat respiratory tract infections and prevent opportunistic infections. Co-trimoxazole shows cross-resistance with pyrimethamine and sulphadoxine in *in vitro* *P. falciparum* culture [49,50] and may play a role in the development of mutations in *pfdhfr* and *pfdhps*.

One goal of this study was to determine whether the prevalence of the clinically relevant *pfdhfr/pfdhps* quintuple mutant, which is associated with SP treatment failure in Africa [13,51], changed during the course of SP use as the first-line anti-malarial in Kenya. Between the baseline and follow-up studies, the prevalence of the quintuple mutant doubled to 53.5% and may be even higher. Due to absence of MOI data in these studies, *pfdhfr/pfdhps* haplotypes were deduced according to the method described by Kublin and colleagues [13]. Because this method conservatively classifies isolates that are pure mutant at one *pfdhfr* allele and mixed at two other *pfdhfr* alleles as double mutant, the prevalence of the *pfdhfr* triple mutant may be systematically underestimated in populations where mixed infections are common. In the follow-up study, many more isolates had mixed *pfdhfr* genotypes than at baseline, and the prevalence of the *pfdhfr* triple mutant decreased between the baseline and follow-up studies. This

observation contrasts with other findings showing that the population of SP-resistant parasites expanded over this period, and may be explained by more severe systematic underestimation in the follow-up study compared to the baseline. Thus, the prevalence of the *pfdhfr* triple mutant and the *pfdhfr/pfdhps* quintuple mutant in the follow-up study may actually be higher than estimated here.

Although the prevalence of the *pfdhfr* triple mutant appeared to decrease over time, increases in the prevalence of the *pfdhps* A437G/K540E double mutant led to a rise in the *pfdhfr/pfdhps* quintuple mutant prevalence. The very high prevalence of mutant codons 437 and 540 is consistent with other reports that these mutations are common in western Kenya [19,52]. Notably, the data presented here show that SP use in Kenya was not just associated with the expansion of A437G and K540E, but also the progressive accumulation of mutations in *pfdhps* at codons 436, 581, and 613. This process appears to be rapid, as demonstrated by the emergence and spread of the A581G and A613S/T mutations in the short time between the baseline and follow-up studies.

Mutations at *pfdhps* codons 436, 581, and 613 are associated with increased levels of *in vitro* resistance when they occur with the A437G mutation [20,43]. Among isolates in the follow-up study, over half had the mutations S436A/E, A581G, and A613S/T in addition to the A437G/K540E single or double mutant; A437G was present in either mixed or pure form in all single mutant isolates. Thus, at follow-up the majority of isolates had parasite genotypes associated with extremely high *in vitro* sulphadoxine resistance. Although the effect of these mutations *in vivo* is not well-studied, under SP selection, parasites containing the quintuple mutant and A581G have been found to have a selective advantage *in vivo* over quintuple mutant parasites [53]. This selective advantage is reflected in the swift development and expansion of *pfdhps* mutations between the baseline and follow-up studies.

This is the first study to report the presence of the A581G mutation in Kenya. Although relatively rare in sub-Saharan Africa, this mutation has been documented in the neighboring countries of Uganda and Tanzania [19]. In the background of mutant *pfdhfr* in Tanzania, a similar increase in the prevalence of A437G and K540E accompanied by the emergence and rapid spread of A581G was observed over the six year period following SP implementation [46]. Notably, in this and other East African studies that reported the presence of A581G, all or most samples were wild-type at codon 613 [46,53,54]. This contrasts sharply with the findings reported here, where mutant codon 613 was present in over half of patient isolates in the follow-up study. It thus appears that the mutation A581G is becoming increasingly



common in East Africa, while Kisumu may be a regional hotspot for the mutation A613S/T.

Although progressive mutations in *pfdhps* were observed, the *pfdhfr* mutation I164L, which is associated with high levels of chlorocycloguanil and pyrimethamine resistance in South America and South East Asia, was absent from isolates in both study periods. This mutation was detected in isolates from Kisumu in 2002 [17], prior to the start of the follow-up study, and in western Kenya in 2004 [18]. The absence of the I164L mutation in isolates from the follow-up study, conducted from 2003-2005, indicates that parasites harboring this mutation in 2002 did not become widespread in Kisumu. Taken together, these findings are consistent with evidence that the I164L mutation is rare in sub-Saharan Africa, outside local hotspots in south-west Uganda and Rwanda [16,19,55].

The change from CQ to SP as the first-line anti-malarial in 1998 was expected to decrease CQ use in Kenya, possibly leading to restoration of CQ sensitive parasites. In Malawi, discontinuing the use of CQ led to the reversion of codon 76 to wild-type over a seven year period, and CQ sensitivity *in vivo* was restored after 12 years [32,33]. Trends of increasing prevalence of wild-type codon 76 have recently been reported in other areas of East Africa. In Tanzania, the prevalence of wild-type *pfert* codon 76 increased over a six year period after anti-malarial policy replaced CQ with SP [46]. Surveillance in coastal Kenya from 1993-2006 showed that the wild-type codon 76 prevalence also increased, although at a slower rate than in Malawi [45]. Therefore, it was surprising to observe a statistically significant increase in the prevalence of isolates containing *pfert* K76T from 82% to 97%, as well as a significant increase in the prevalence of *pfmdr1* N86Y, also associated with chloroquine resistance, from 44% to 76%. To analyse whether the apparent increase in the prevalence of mutant codon 76 arose from the higher prevalence of polyclonal infections observed in the follow-up study, the method of Hill and Babiker was used to estimate the frequency of the K76T allele. Even accounting for polyclonal infections, there was an apparent although not statistically significant increase in the frequency of K76T. Thus, the K76T mutation does not appear to be abating in Kisumu, as it is in coastal Kenya and in other areas in East Africa.

The persistence of the *pfert* K76T mutation in isolates from a high transmission setting such as Kisumu indicates that selection for the mutant codon is ongoing. The *pfert* K76T mutation is associated with amodiaquine as well as chloroquine treatment failure [56]. Although AQ was officially the second-line therapy used in Kenya during the study period, it was available at 95% of drug retail outlets, whereas SP was available at only 29% of outlets [57]. By comparison, CQ was stocked by 15% of drug retail outlets [57]. It thus appears that widespread use of

AQ in the private sector may be high enough to exert selective pressure on the parasite population.

Specific *pfmdr1* genotypes have been linked to altered susceptibility to artemisinin, a component of the current first-line anti-malarial therapy AL. *In vitro*, the allele combination S1034C/N1042D/D1246Y is associated with increased artemisinin susceptibility [28], and *in vivo*, treatment with AL selects for the N86, Y184F, and D1246 alleles [29,30]. In this study, the prevalence of alleles associated with artemisinin sensitivity (S1034C and D1246Y) decreased significantly; however, this decline is not likely due to selective pressure induced by AL as completion of the follow-up study predated widespread use of AL. Whether the prevalence of *pfmdr1* mutations continues to change under increasing AL pressure will be of interest.

## Conclusions

During the period in which SP was used as the first-line anti-malarial in Kenya, the prevalence of clinically relevant molecular markers of SP and CQ resistance increased in Kisumu, a city located on the shore of Lake Victoria in a malaria endemic region. Mutations at *pfdhps* codons 581 and 613, which have not previously been observed in Kenya, became highly prevalent during this period. In light of these highly SP-resistant genotypes and reports that SP remains widely used in Kenya [4], it is important to discourage use of SP to treat symptomatic malaria, particularly for populations that are not semi-immune. Furthermore, whether the *pfdhfr/pfdhps* quintuple mutant combined with the *pfdhps* mutations S436A/E, A581G, and A613S/T adversely impacts the efficacy of IPTp and IPTi should be evaluated. The continued rise in the prevalence of mutations associated with CQ resistance indicates that continued drug pressure, either from CQ or amodiaquine use, is preventing restoration of CQ-susceptible parasites in Kisumu. That these findings contrast with a study covering the same period in coastal Kenya [45] highlights the importance of surveillance and the heterogeneity in drug resistance that may occur within a country.

## Additional material

### Additional file 1: Classification of *pfdhfr* and *pfdhps* genotypes.

White boxes indicate wild-type genotype, gray boxes indicate mixed mutant genotype, and black boxes indicate pure mutant genotype. Adapted from [13].

### Additional file 2: Classification of *pfdhfr/pfdhps* haplotypes using the genotype classifications in Additional file 1. Table showing *pfdhfr/pfdhps* haplotype classifications.

## Acknowledgements

This work was funded by the U.S. Army, Military Infectious Disease Research Program, and the Department of Defense-Global Emerging Infections



System. MDS was supported by a Johns Hopkins Malaria Research Institute Pre-doctoral Fellowship. The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or reflecting the views of the Department of the Army or the Department of Defense.

#### Author details

<sup>1</sup>Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA. <sup>2</sup>Global Emerging Infections System (GEIS), United States Army Research Unit-Kenya Walter Reed/KEMRI project, Kisumu, Kenya. <sup>3</sup>Department of Preventive Medicine and Biometrics, Uniformed Services University of the Health Sciences, Bethesda, Maryland, USA. <sup>4</sup>Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA. <sup>5</sup>Walter Reed Army Institute of Research, Silver Spring, Maryland, USA.

#### Authors' contributions

MDS participated in the design of the study, analysed the data, and wrote the manuscript. FE and HA performed the molecular analyses. RLC, SAB, and NCW conceived of and coordinated the studies. RLC and NCW wrote the protocol. STP assisted in writing the manuscript. WJM helped design the study, analyse the data, and write the manuscript. All authors read and approved the final manuscript.

#### Competing interests

The authors declare that they have no competing interests.

Received: 10 August 2010 Accepted: 24 November 2010

Published: 24 November 2010

#### References

- Kenya Medical Research Institute Kenya Malaria Fact Sheet. [http://www.kemri.org/index.php/help-desk/search/diseases-a-conditions/29-malaria/113-kenya-malaria-fact-sheet].
- Shretta R, Omumbo J, Rapuoda B, Snow RW: **Using evidence to change antimalarial drug policy in Kenya.** *Trop Med Int Health* 2000, **5**:755-764.
- Amin A, Zurovac D, Kangwana B, Greenfield J, Otieno D, Akhwale W, Snow R: **The challenges of changing national malaria drug policy to artemisinin-based combinations in Kenya.** *Malar J* 2007, **6**:72.
- Watsierah CA, Jura WG, Oyugi H, Abong'o B, Ouma C: **Factors determining anti-malarial drug use in a peri-urban population from malaria holoendemic region of western Kenya.** *Malar J* 2010, **9**:295.
- ter Kuile FO, van Eijk AM, Filler SJ: **Effect of sulfadoxine-pyrimethamine resistance on the efficacy of intermittent preventive therapy for malaria control during pregnancy: a systematic review.** *JAMA* 2007, **297**:2603-2616.
- Kobbe R, Kreuzberg C, Adjei S, Thompson B, Langefeld I, Thompson PA, Abruquah HH, Kreuels B, Ayim M, Busch W, Marks F, Amoah K, Opoku E, Meyer CG, Adjei O, May J: **A randomized controlled trial of extended intermittent preventive antimalarial treatment in infants.** *Clin Infect Dis* 2007, **45**:16-25.
- Gosling RD, Carneiro I, Chandramohan D: **Intermittent preventive treatment of malaria in infants: how does it work and where will it work?** *Trop Med Int Health* 2009, **14**:1003-1010.
- Grobusch MP, Lell B, Schwarz NG, Gabor J, Dornemann J, Potschke M, Oyakhromire S, Kiessling GC, Necek M, Langin MU, Klein Klouwenberg P, Klopfer A, Naumann B, Altun H, Agnandji ST, Goesch J, Decker M, Salazar CL, Supan C, Kombila DU, Borchert L, Koster KB, Pongratz P, Adegnikna AA, Glasenapp I, Issifou S, Kreamsner PG: **Intermittent preventive treatment against malaria in Gabon—a randomized, double-blind, placebo-controlled trial.** *J Infect Dis* 2007, **196**:1595-1602.
- Macete E, Aide P, Aponte JJ, Sanz S, Mandomando I, Espasa M, Sigauque B, Dobano C, Mabunda S, DgeDge M, Alonso P, Menendez C: **Intermittent preventive treatment for malaria control administered at the time of routine vaccinations in Mozambican infants: a randomized, placebo-controlled trial.** *J Infect Dis* 2006, **194**:276-285.
- Mockenhaupt FP, Reither K, Zanger P, Roepcke F, Danquah I, Saad E, Ziniel P, Dzisi SY, Frempong M, Agana-Nsiire P, Amoo-Sakyi F, Otchwemah R, Cramer JP, Anemana SD, Dietz E, Bienze U: **Intermittent preventive treatment in infants as a mean of malaria control: a randomized, double-blind, and placebo-controlled trial in northern Ghana.** *Antimicrob Agents Chemother* 2007, AAC.00513-00507.
- Schellenberg D, Menendez C, Kahigwa E, Aponte J, Vidal J, Tanner M, Mshinda H, Alonso P: **Intermittent treatment for malaria and anaemia control at time of routine vaccinations in Tanzanian infants: a randomised, placebo-controlled trial.** *Lancet* 2001, **357**:1471-1477.
- Nzila-Mounda A, Mberu EK, Sibley CH, Plowe CV, Winstanley PA, Watkins WM: **Kenyan *Plasmodium falciparum* field isolates: correlation between pyrimethamine and chlorocycloguanil activity *in vitro* and point mutations in the dihydrofolate reductase domain.** *Antimicrob Agents Chemother* 1998, **42**:164-169.
- Kublin JG, Dzinjalama FK, Kamwendo DD, Malkin EM, Cortese JF, Martino LM, Mukadam RA, Rogerson SJ, Lescano AG, Molyneux ME, Winstanley PA, Chimpeni P, Taylor TE, Plowe CV: **Molecular markers for failure of sulfadoxine-pyrimethamine and chlorproguanil-dapsone treatment of *Plasmodium falciparum* malaria.** *J Infect Dis* 2002, **185**:380-388.
- Foot SJ, Galatis D, Cowman AF: **Amino acids in the dihydrofolate reductase-thymidylate synthase gene of *Plasmodium falciparum* involved in cycloguanil resistance differ from those involved in pyrimethamine resistance.** *Proc Natl Acad Sci USA* 1990, **87**:3014-3017.
- Plowe CV, Cortese JF, Djimde A, Nwanyanwu OC, Watkins WM, Winstanley PA, Estrada-Franco JG, Mollinedo RE, Avila JC, Cespedes JL, Carter D, Doumbo OK: **Mutations in *Plasmodium falciparum* dihydrofolate reductase and dihydropteroate synthase and epidemiologic patterns of pyrimethamine-sulfadoxine use and resistance.** *J Infect Dis* 1997, **176**:1590-1596.
- Lynch C, Pearce R, Pota H, Cox J, Abeku TarekgnÅ A, Rwakimari J, Naidoo I, Tibenderana J, Roper C: **Emergence of a *dhfr* mutation conferring high level drug resistance in *Plasmodium falciparum* populations from southwest Uganda.** *J Infect Dis* 2008, **197**:1598-1604.
- McCollum AM, Poe AC, Hamel M, Huber C, Zhou Z, Shi YP, Ouma P, Vulule J, Bloland P, Slutsker L, Barnwell JW, Udhayakumar V, Escalante AA: **Antifolate resistance in *Plasmodium falciparum*: multiple origins and identification of novel *dhfr* alleles.** *J Infect Dis* 2006, **194**:189-197.
- Oesterholt MJ, Alifrangis M, Sutherland CJ, Omar SA, Sawa P, Howitt C, Gouagna LC, Sauerwein RW, Bousema T: **Submicroscopic gametocytes and the transmission of antifolate-resistant *Plasmodium falciparum* in Western Kenya.** *PLoS ONE* 2009, **4**:e4364.
- Drug Resistance Maps. [http://www.drugresistancemaps.org/].
- Triglia T, Menting JGT, Wilson C, Cowman AF: **Mutations in dihydropteroate synthase are responsible for sulfone and sulfonamide resistance in *Plasmodium falciparum*.** *Proc Natl Acad Sci USA* 1997, **94**:13944-13949.
- Triglia T, Wang P, Sims PF, Hyde JE, Cowman AF: **Allelic exchange at the endogenous genomic locus in *Plasmodium falciparum* proves the role of dihydropteroate synthase in sulfadoxine-resistant malaria.** *Embo J* 1998, **17**:3807-3815.
- Alker A, Mwapasa V, Purfield A, Rogerson S, Molyneux M, Kamwendo D, Tadesse E, Chaluluka E, Meshnick S: **Mutations associated with sulphadoxine-pyrimethamine and chlorproguanil resistance in *Plasmodium falciparum* isolates from Blantyre, Malawi.** *Antimicrob Agents Chemother* 2005, **49**:3919-3921.
- Geburu-Woldearegai T, Hailu A, Grobusch MP, Kun JFJ: **Molecular surveillance of mutations in dihydrofolate reductase and dihydropteroate synthase genes of *Plasmodium falciparum* in Ethiopia.** *Am J Trop Med Hyg* 2005, **73**:1131-1134.
- Biswas S, Escalante A, Chaiyaroj S, Angkasekwinai P, Lal AA: **Prevalence of point mutations in the dihydrofolate reductase and dihydropteroate synthetase genes of *Plasmodium falciparum* isolates from India and Thailand: a molecular epidemiologic study.** *Trop Med Int Health* 2000, **5**:737-743.
- Fidock DA, Nomura T, Talley AK, Cooper RA, Dzekunov SM, Ferdig MT, Ursos LM, Sidhu AB, Naude B, Deitsch KW, Su XZ, Wootton JC, Roepe PD, Wellem TE: **Mutations in the *P. falciparum* digestive vacuole transmembrane protein PfCRT and evidence for their role in chloroquine resistance.** *Mol Cell* 2000, **6**:861-871.
- Price RN, Uhlemann AC, Brockman A, McGready R, Ashley E, Phaipun L, Patel R, Laing K, Looareesuwan S, White NJ, Nosten F, Krishna S: **Mefloquine resistance in *Plasmodium falciparum* and increased *pfmdr1* gene copy number.** *Lancet* 2004, **364**:438-447.

27. Mockenhaupt FP, Ehrhardt S, Eggelte TA, Agana-Nsiire P, Stollberg K, Mathieu A, Markert M, Otchwemah RN, Bienzle U: **Chloroquine-treatment failure in northern Ghana: roles of *pfprt* T76 and *pfmdr1* Y86.** *Ann Trop Med Parasitol* 2005, **99**:723-732.
28. Sidhu ABS, Valderramos SG, Fidock DA: ***pfmdr1* mutations contribute to quinine resistance and enhance mefloquine and artemisinin sensitivity in *Plasmodium falciparum*.** *Molecular Microbiology* 2005, **57**:913-926.
29. Dokomajilar C, Nsohya SL, Greenhouse B, Rosenthal PJ, Dorsey G: **Selection of *Plasmodium falciparum* *pfmdr1* alleles following therapy with artemether-lumefantrine in an area of Uganda where malaria is highly endemic.** *Antimicrob Agents Chemother* 2006, **50**:1893-1895.
30. Humphreys GS, Merinopoulos I, Ahmed J, Whitty CJ, Mutabingwa TK, Sutherland CJ, Hallett RL: **Amodiaquine and artemether-lumefantrine select distinct alleles of the *Plasmodium falciparum* *mdr1* gene in Tanzanian children treated for uncomplicated malaria.** *Antimicrob Agents Chemother* 2007, **51**:991-997.
31. Djimdé A, Doumbo OK, Cortese JF, Kayentao K, Doumbo S, Diourte Y, Dicko A, Su X-z, Nomura T, Fidock DA, Wellemes TE, Plowe CV, Coulibaly D: **A molecular marker for chloroquine-resistant *falciparum* malaria.** *N Engl J Med* 2001, **344**:257-263.
32. Kublin JG, Cortese JF, Njunju EM, Mukadam GRA, Wirima JJ, Kazembe PN, Djimde AA, Kouriba B, Taylor TE, Plowe CV: **Reemergence of chloroquine-sensitive *Plasmodium falciparum* malaria after cessation of chloroquine use in Malawi.** *J Infect Dis* 2003, **187**:1870-1875.
33. Laufer MK, Thesing PC, Eddington ND, Masonga R, Dzinjalama FK, Takala SL, Taylor TE, Plowe CV: **Return of chloroquine antimalarial efficacy in Malawi.** *N Engl J Med* 2006, **355**:1959-1966.
34. Mbaisi A, Liyala P, Eyase F, Achilla R, Akala H, Wangui J, Mwangi J, Osuna F, Alam U, Smoak BL, Davis JM, Kyle DE, Coldren RL, Mason C, Waters NC: **Drug susceptibility and genetic evaluation of *Plasmodium falciparum* isolates obtained in four distinct geographical regions of Kenya.** *Antimicrob Agents Chemother* 2004, **48**:3598-3601.
35. Duraisingh M, Curtis J, Warhurst D: ***Plasmodium falciparum*: Detection of polymorphisms in the dihydrofolate reductase and dihydropteroate synthase genes by PCR and restriction digestion.** *Exp Parasitol* 1998, **89**:1-8.
36. Wang P, Brooks DR, Sims PFG, Hyde JE: **A mutation-specific PCR system to detect sequence variation in the dihydropteroate synthetase gene of *Plasmodium falciparum*.** *Molecular and Biochemical Parasitology* 1995, **71**:115-125.
37. Hill WG, Babiker HA: **Estimation of numbers of malaria clones in blood samples.** *Proc Biol Sci* 1995, **262**:249-257.
38. Webster HK, Boudreau EF, Pavanand K, Yongvanitchit K, Pang LW: **Antimalarial drug susceptibility testing of *Plasmodium falciparum* in Thailand using a microdilution radioisotope method.** *Am J Trop Med Hyg* 1985, **34**:228-235.
39. Ringwald P, Basco LK: **Comparison of *in vivo* and *in vitro* tests of resistance in patients treated with chloroquine in Yaounde, Cameroon.** *Bull World Health Organ* 1999, **77**:34-43.
40. Basco LK, Le Bras J: ***In vitro* activity of halofantrine and its relationship to other standard antimalarial drugs against African isolates and clones of *Plasmodium falciparum*.** *Am J Trop Med Hyg* 1992, **47**:521-527.
41. Pradines B, Tall A, Fusai T, Spiegel A, Hienne R, Rogier C, Trape JF, Le Bras J, Parzy D: ***In vitro* activities of Benflumetol against 158 Senegalese isolates of *Plasmodium falciparum* in comparison with those of standard antimalarial drugs.** *Antimicrob Agents Chemother* 1999, **43**:418-420.
42. Basco LK, Ramiliarisoa O, Bras JL: ***In vitro* Activity of pyrimethamine, cycloguanil, and other antimalarial drugs against African isolates and clones of *Plasmodium falciparum*.** *Am J Trop Med Hyg* 1994, **50**:193-199.
43. Wang P, Lee C-S, Bayoumi R, Djimde A, Doumbo O, Swedberg G, Dao LD, Mshinda H, Tanner M, Watkins WM, Sims PF, Hyde JE: **Resistance to antifolates in *Plasmodium falciparum* monitored by sequence analysis of dihydropteroate synthetase and dihydrofolate reductase alleles in a large number of field samples of diverse origins.** *Mol Biochem Parasitol* 1997, **89**:161-177.
44. Mberu EK, Nzila AM, Nduati E, Ross A, Monks SM, Kokwaro GO, Watkins WM, Hopkins Sibley C: ***Plasmodium falciparum*: *in vitro* activity of sulfadoxine and dapsone in field isolates from Kenya: point mutations in sulfadiazine synthase may not be the only determinants in sulfa resistance.** *Exp Parasitol* 2002, **101**:90-96.
45. Mwai L, Ochong E, Abdirahman A, Kiara S, Ward S, Kokwaro G, Sasi P, Marsh K, Borrmann S, Mackinnon M, Nzila A: **Chloroquine resistance before and after its withdrawal in Kenya.** *Malar J* 2009, **8**:106.
46. Alifrangis M, Lusingu JP, Mmbando B, Dalgaard MB, Vestergaard LS, Ishengoma D, Khalil IF, Theander TG, Lemnge MM, Bygbjerg IC: **Five-year surveillance of molecular markers of *Plasmodium falciparum* antimalarial drug resistance in Korogwe District, Tanzania: accumulation of the 581G mutation in the *P. falciparum* dihydropteroate synthase gene.** *Am J Trop Med Hyg* 2009, **80**:523-527.
47. Ndenga B, Githeko A, Omukunda E, Munyekenye G, Atieli H, Wamai P, Mbogo C, Minakawa N, Zhou G, Yan G: **Population dynamics of malaria vectors in western Kenya highlands.** *J Med Entomol* 2006, **43**:200-206.
48. Okiro EA, Alegana VA, Noor AM, Snow RW: **Changing malaria intervention coverage, transmission and hospitalization in Kenya.** *Malar J* 2010, **9**:285.
49. Khalil I, Ronn AM, Alifrangis M, Gabar HA, Satti GM, Bygbjerg IC: **Dihydrofolate reductase and dihydropteroate synthase genotypes associated with *in vitro* resistance of *Plasmodium falciparum* to pyrimethamine, trimethoprim, sulfadoxine, and sulfamethoxazole.** *Am J Trop Med Hyg* 2003, **68**:586-589.
50. Iyer JK, Milhous WK, Cortese JF, Kublin JG, Plowe CV: ***Plasmodium falciparum* cross-resistance between trimethoprim and pyrimethamine.** *The Lancet* 2001, **358**:1066-1067.
51. Mockenhaupt F, Bousema T, Eggelte T, Schreiber J, Ehrhardt S, Wassilew N, Otchwemah R, Sauerwein R, Bienzle U: ***Plasmodium falciparum* *dhfr* but not *dhps* mutations associated with sulphadoxine-pyrimethamine treatment failure and gametocyte carriage in northern Ghana.** *Trop Med Int Health* 2005, **10**:901-908.
52. Naidoo I, Roper C: **Following the path of most resistance: *dhps* K540E dispersal in African *Plasmodium falciparum*.** *Trends Parasitol* 2010, **26**:447-456.
53. Harrington WE, Mutabingwa TK, Muehlenbachs A, Sorensen B, Bolla MC, Fried M, Duffy PE: **Competitive facilitation of drug-resistant *Plasmodium falciparum* malaria parasites in pregnant women who receive preventive treatment.** *Proc Natl Acad Sci USA* 2009, **106**:9027-9032.
54. Gesase S, Gosling RD, Hashim R, Ord R, Naidoo I, Madebe R, Mosha JF, Joho A, Mandia V, Mrema H, Mapunda E, Savael Z, Lemnge M, Mosha FW, Greenwood B, Roper C, Chandramohan D: **High resistance of *Plasmodium falciparum* to sulphadoxine/pyrimethamine in northern Tanzania and the emergence of *dhps* resistance mutation at codon 581.** *PLoS ONE* 2009, **4**:e4569.
55. Karema C, Imwong M, Fanello CI, Stepniewska K, Uwimana A, Nakeesathit S, Dondorp A, Day NP, White NJ: **Molecular correlates of high-level antifolate resistance in Rwandan children with *Plasmodium falciparum* malaria.** *Antimicrob Agents Chemother* 2010, **54**:477-483.
56. Ochong E, Van den Broek I, Keus K, Nzila A: **Short report: Association between chloroquine and amodiaquine resistance and allelic variation in the *Plasmodium falciparum* multiple drug resistance 1 gene and the chloroquine resistance transporter gene in isolates from the upper Nile in southern Sudan.** *Am J Trop Med Hyg* 2003, **69**:184-187.
57. Amin A, Snow R: **Brands, costs and registration status of antimalarial drugs in the Kenyan retail sector.** *Malar J* 2005, **4**:36.

doi:10.1186/1475-2875-9-338

**Cite this article as:** Spalding et al.: Increased prevalence of the *pfdhfr/pfdhps* quintuple mutant and rapid emergence of *pfdhps* resistance mutations at codons 581 and 613 in Kisumu, Kenya. *Malaria Journal* 2010 **9**:338.