

Corrigendum

Clin Pediatr Endocrinol 2016; 25(2): 71–76. doi: 10.1297/cpe.25.71 PMID: PMC4860518

Short Communication

Growth standard charts for Japanese children with mean and standard deviation (SD) values based on the year 2000 national survey

Tsuyoshi Isojima^{1,2}, Noriko Kato³, Yoshiya Ito^{1,4}, Susumu Kanzaki^{1,5}, and Mitsunori Murata^{1,6}

¹ The Japanese Society for Pediatric Endocrinology, Kyoto, Japan

² Department of Pediatrics, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan

³ Department of Early Childhood and Elementary Education, Jumonji University, Niiza, Japan

⁴ Faculty of Nursing, Japanese Red Cross Hokkaido College of Nursing, Kitami, Japan

⁵ Division of Pediatrics and Perinatology, Tottori University Faculty of Medicine, Yonago, Japan

⁶ Health Center, Wayo Women's University, Chiba, Japan

Received: December 2, 2015/ Accepted: December 29, 2015

Corresponding author: Noriko Kato, Department of Early Childhood and Elementary Education, Jumonji University, 2-1-28 Sugasawa, Niiza, Saitama 352-8510, Japan

E-mail: kato@niph.go.jp

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License <<http://creativecommons.org/licenses/by-nc-nd/4.0/>>.

The authors and publisher would like to apologize for the mistake in the following equation.

In p. 72, in lines 8–12 of the right column:

The SD score (Z-score) of each measurement (y value) could be calculated from the L, M and S curves, using values appropriate for the age and gender, with the following equation:

$$Z = \frac{[(y/M)^{L-1}] / (L \times S)}{L}, \text{ or } Z = \ln(y/M)/S \text{ if } L = 0.$$

should have been

The SD score (Z-score) of each measurement (y value) could be calculated from the L, M and S curves, using values appropriate for the age and gender, with the following equation:

$$Z = \frac{[(y/M)^{L-1}] / (L \times S)}{L}, \text{ or } Z = \ln(y/M)/S \text{ if } L = 0.$$