



Phytochemical content, especially spermidine derivatives, presenting antioxidant and antilipoxygenase activities in Thai bee pollens

Phanthiwa Khongkarat¹, Preecha Phuwapraisirisan² and Chanpen Chanchao³

¹Program in Biotechnology, Faculty of Science, Chulalongkorn University, Patumwan, Bangkok, Thailand

²Center of Excellence in Natural Products, Department of Chemistry, Faculty of Science, Chulalongkorn University, Patumwan, Bangkok, Thailand

³Department of Biology, Faculty of Science, Chulalongkorn University, Patumwan, Bangkok, Thailand

ABSTRACT

Background. Bee pollen (BP) is full of useful nutrients and phytochemicals. Its chemical components and bioactivities depend mainly on the type of floral pollen.

Methods. Monofloral BP from *Camellia sinensis* L., *Mimosa diplotricha*, *Helianthus annuus* L., *Nelumbo nucifera*, *Xyris complanata*, and *Ageratum conyzoides* were harvested. Crude extraction and partition were performed to yield solvent-partitioned extracts of each BP. Total phenolic content (TPC) was assayed by the Folin-Ciocalteu method, while the flavonoid content (FC) was measured by the aluminium chloride colorimetric method. Antioxidant capacity was measured by the (i) 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity, (ii) 2,2'-azino-bis (3-ethylbenzthiazoline-6-sulphonic acid) (ABTS) scavenging activity and its Trolox equivalent antioxidant capacity (TEAC), and (iii) ferric reducing antioxidant power (FRAP). All samples were tested for lipoxygenase inhibitory (LOXI) activity. The most active sample was enriched by silica gel 60 column chromatography (SiG60-CC) and high performance liquid chromatography (HPLC), observing the chemical pattern of each fraction using thin layer chromatography. Chemical structure of the most active compound was analyzed by proton nuclear magnetic resonance and mass spectrometry.

Results. Dichloromethane (DCM)-partitioned BP extracts of *H. annuus* L. and *M. diplotricha* (DCMMBP) showed a very high TPC, while DCMMBP had the highest FC. In addition, DCMMBP had the strongest DPPH and ABTS radical scavenging activities (as a TEAC value), as well as FRAP value. Also, DCMMBP (60 µg/mL) gave the highest LOXI activity (78.60 ± 2.81%). Hence, DCMMBP was chosen for further enrichment by SiG60-CC and HPLC. Following this, the most active fraction showed higher antioxidant and LOXI activities with an EC₅₀ for DPPH and ABTS of 54.66 ± 3.45 µg/mL and 24.56 ± 2.99 µg/mL (with a TEAC value of 2,529.69 ± 142.16 µmole TE/g), respectively, and a FRAP value of 3,466.17 ± 81.30 µmole Fe²⁺/g and an IC₅₀ for LOXI activity of 12.11 ± 0.36 µg/mL. Triferuloyl spermidines were revealed to be the likely main active components.

Conclusions. TPC, FC, and spermidine derivatives played an important role in the antioxidant and antilipoxygenase activities in *M. diplotricha* bee pollen.

Submitted 31 January 2022

Accepted 6 May 2022

Published 25 May 2022

Corresponding author

Chanpen Chanchao,
chanpen.c@chula.ac.th

Academic editor

Mohd Adnan

Additional Information and
Declarations can be found on
page 17

DOI 10.7717/peerj.13506

© Copyright

2022 Khongkarat et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Agricultural Science, Biotechnology

Keywords Bee pollen, Bioactivity, Flavonoid content, Phytochemical, Total phenolic content

INTRODUCTION

People nowadays have a longer expected lifespan due to advances in medicine and technology. In addition, people pay more attention to their health. Hence, a lot of nutrient supplements have been introduced, which are mostly from natural products. Bee pollen (BP) is one of the natural bee products that is widely used for nutritional and medical applications. It is produced by foraging bees by mixing the floral pollens with some nectar or honey, enzymes, wax, and bee secretion (*Chantarudee et al., 2012*). Bee pollens are a good protein source because they contain a high level of well-balanced proteins (those with all the essential amino acids necessary for the human body). Furthermore, BPs contain other required nutrients, including unsaturated fatty acids, vitamins, minerals, and trace elements. As a result, it is commonly used as a natural dietary supplement (*Sommano et al., 2020*).

Additionally, BP contains a variety of beneficial chemicals, particularly phenolic and flavonoid components, such as quercetin, kaempferol, caffeic acid, and naringenin (*Saric et al., 2009*). As a result, many bioactivities of BP have been discovered, such as antimicrobial and anti-inflammatory activities (*Saral et al., 2022*). The phenolic compounds from rape BP showed *in vitro* antioxidant and tyrosinase inhibition (TYRI) properties and could inhibit melanogenesis. These activities were attributed to rutin, the major flavonoid in the extract (*Sun et al., 2017*). Also, the phenol and flavonoid compounds in BP from Nigeria had an α -amylase inhibitory activity, indicating the potential therapeutic potential of BP in the management of diabetes (*Daudu, 2019*). In addition, a monoamine oxidase (MAO) inhibitory activity was reported for chestnut BP (*Yildiz et al., 2014*), and so its potential use in the treatment of depressive disorders, Parkinson's disease, and Alzheimer's disease. The relationship between MAO inhibition and the total phenolic content (TPC) as well as the antioxidant capacity of BP was subsequently reported (*Yildiz et al., 2014*). Hence, the TPC and flavonoid contents (FC) from BP, which mainly depended on the floral origin, have the potential to be used as a natural antioxidant agent and enzyme inhibitor.

The antioxidant and lipoxygenase inhibition (LOXI) properties were found to be associated with anti-inflammatory, anti-cancer, and anti-aging effects (*Eshwarappa et al., 2016*). The antioxidant properties can protect cells from the oxidative damage of free radical molecules, which are unstable molecules with one or more unpaired electrons (*Lobo et al., 2010*). The antioxidant can react with other molecules, including the intracellular proteins, lipids, and DNA. After such damage, the cells are in an oxidative stress that can induce the development of many chronic diseases, such as cancer, autoimmune disorders, aging, cataracts, rheumatoid arthritis, and cardiovascular and neurodegenerative diseases (*Pham-Huy, He & Pham-Huy, 2010*).

Therefore, to help prevent oxidative damage to cells, antioxidant agents that can donate electrons or hydrogen atoms are important in order to reduce the free radical level (*Shahidi & Zhong, 2015*). The lipoxygenases (LOXs) are an important family of enzymes in inflammatory and immune responses. They catalyze the conversion of polyunsaturated fatty acids and arachidonic acid to inflammatory eicosanoids, including hydroperoxy-eicosatetraenoic acid and leukotrienes (*Mashima & Okuyama, 2015*). The overexpression

of LOX promotes the development of several inflammation-related diseases, such as arthritis, asthma, cancer, and allergic diseases (Wang et al., 2021). This can be prevented via the inhibition of the LOX pathway and the targeting of LOX with LOXIs is a promising therapeutic target for treating a wide spectrum of human diseases (Eshwarappa et al., 2016).

The chemical composition of BP mainly depends on its floral and geological/geographical origins. In Thailand, some dominant floral origins are different from those in other countries (Chamchumroon et al., 2017), while the bioactivities of some BPs have not been reported before. This led to our interest in the bioactivities of different types of Thai monofloral BPs. Six such samples were collected and then sequentially extracted and partitioned with three organic solvents of different polarities. The enrichment of the active compounds, using bioactivity and chemical profile screening of the fractions, was performed. The structure of the isolated active compounds was analyzed by proton nuclear magnetic resonance spectroscopy ($^1\text{H-NMR}$) and the molecular weight (MW) was measured by MALDI-TOF mass spectrometry (MS). The benefit of this work is the discovery of potential new antioxidant and LOXI compounds and the promotion of BP as a nutraceutical food to be developed for use in the pharmaceutical industry. This may encourage the growth of Thai apiaries, resulting in an improved income for bee farmers.

MATERIALS & METHODS

Sample collection

Six monofloral *Apis mellifera* BP samples were collected from five provinces in Thailand (Table 1) and stored at room temperature (25 °C) until used. The BP was identified by palynological analysis and sequence analysis of the ITS-2 region of the rRNA genes, as previously reported (Khongkarat et al., 2022). Both analyses were consistent and confirmed that the six *A. mellifera* BP samples were from *Camellia sinensis* L. (BP1), *Mimosa diplotricha* (BP2) (Chiangmai province), *Helianthus annuus* L. (BP3) (Lopburi province), *Nelumbo nucifera* (BP4) (Nakhon Sawan province), *Xyris complanata* (BP5) (Udon Thani), and *Ageratum conyzoides* (BP6) (Lamphun).

Crude extraction and partition

All collected samples were extracted as previously described (Umthong et al., 2011). In brief, each BP sample (140 g) was extracted with 800 mL of methanol (MeOH) by shaking at 100 rpm, 15 °C, for 18 h. After that, the supernatant was collected and the residue was extracted two more times each with 800 mL of MeOH. The three supernatants were pooled and dried under reduced pressure to obtain the respective MeOH crude extract (MCE). Each MCE was further partitioned by hexane, dichloromethane (DCM), and MeOH as follows. At first, the MCE was dissolved in MeOH (250 mL) and then mixed with an equal volume of hexane in a separating funnel and left to phase separate. The upper hexane phase was collected and the lower MeOH phase was further partitioned two more times in the same manner, with the three hexane extracts being pooled and dried under reduced pressure to yield the respective hexane-partitioned BP extract (HX); designated as HXCBP, HXMBP, HXHBP, HXNBP, HXXBP, and HXABP for BP1–6, respectively. Next, the residual MeOH phase was added to an equal volume of water and subsequently partitioned three times with

Table 1 Detail of sample collection.

Type of BP	Sample code	Collecting time	Collecting site (province)	Geographical location	
				Longitude	Latitude
<i>Camellia sinensis</i> L. BP	BP1	February, 2018	Chiangmai	99°03'45.3"E	18°46'37.3"N
<i>Mimosa diplotricha</i> BP	BP2	February, 2018	Chiangmai	99°03'45.3"E	18°46'37.3"N
<i>Helianthus annuus</i> L. BP	BP3	February, 2018	Lopburi	101°01'07.5"E	14°51'31.7"N
<i>Nelumbo nucifera</i> BP	BP4	February, 2018	Nakhon Sawan	100°15'01.4"E	15°41'02.4"N
<i>Xyris complanata</i> BP	BP5	February, 2018	Udon Thani	102°45'28.8"E	17°21'12.7"N
<i>Ageratum conyzoides</i> BP	BP6	February, 2018	Lamphun	98°53'42.6"E	18°04'49.2"N

DCM in the same manner as above (except the DCM phase was the lower layer), with the pooled DCM extracts being evaporated as above to provide the DCM-partitioned extracts; designated as DCMCBP, DCMMBP, DCMHBP, DCMNBP, DCMXBP, and DCMABP for BP1–6, respectively. Finally, the residual MeOH phases were evaporated as above to obtain the MeOH-partitioned extracts; designated as MTCBP, DCMMBP, MTHBP, MTNBP, MTXBP, and MTABP for BP1–6, respectively. All partitioned extracts were kept at -20°C in the dark until used.

Determination of the TPC

The TPC of each partitioned extract was evaluated as previously reported ([Marghitas et al., 2009](#)) based on the Folin-Ciocalteu method. The method was adapted to be performed in a 96 well plate. Firstly, 125 μL of 0.2 N Folin-Ciocalteu reagent was added to 25 μL of the diluted partitioned extract or gallic acid solution in dimethyl sulfoxide (DMSO) and mixed for 5 min. Then, 100 μL of 7.5% (w/v) sodium carbonate solution was added per well and incubated at room temperature for 2 h. The absorbance at a wavelength of 700 nm was measured using a microplate reader against DMSO as a blank. All reactions were performed in triplicate. The results were expressed as mg of gallic acid equivalents (GAE)/g of partitioned extract using a standard graph for gallic acid in the range of 0.02–0.1 mg/mL.

Determination of the FC

The FC of each partitioned extract was measured as previously described ([Marghitas et al., 2009](#)) based on the aluminium chloride (AlCl_3) colorimetric method but with adaptation for use in a 96 well microplate reader. Briefly, 25 μL of the diluted partitioned extract solution dissolved in DMSO was mixed with 100 μL of distilled water. Then, 7.5 μL of 5% (w/v) sodium nitrite solution was added and incubated at room temperature for 5 min, followed by the addition of 7.5 μL of 10% (w/v) AlCl_3 . After 6 min of incubation at room temperature, 50 μL of 1 M sodium hydroxide was added and the absorbance at wavelength of 510 nm was measured. Each assay was performed in triplicate. The FC was expressed as mg of quercetin equivalents (QE)/g of partitioned extract using a standard graph for quercetin (0.1–1 mg/mL).

Determination of the antioxidant capacity

The DPPH free radical scavenging activity assay

The potential of the partitioned extracts was determined as previously described (Khongkarat et al., 2020). Five different concentrations of each sample were prepared in DMSO. For each concentration, 20 μL of the sample was mixed with 80 μL of 0.15 mM DPPH in MeOH and incubated at room temperature for 30 min. The absorbance was measured at a wavelength of 517 nm (A_{517}) using a microplate reader. Ascorbic acid (vitamin C) was used as the standard reference. Each assay was performed in triplicate. The free radical scavenging activity was calculated from Eq. (1):

$$\% \text{ DPPH radical scavenging activity} = [(A - B)/(A)] \times 100, \quad (1)$$

where A is the A_{517} of the negative control and B is the A_{517} of the treatment.

The % inhibition (Y axis) was plotted against the extract concentrations (X axis) and the effective concentration at 50% (EC_{50}) was obtained from the graph.

Determination of the Trolox equivalent antioxidant capacity (TEAC)

The TEAC assay followed the described method (Suriyatem et al., 2017). The stock $\text{ABTS}^{\bullet+}$ solution was prepared by reacting 7 mM ABTS solution with 2.45 mM potassium persulphate solution in distilled water at a 1:1 (v/v) ratio in the dark at room temperature for 16 h before use. The working $\text{ABTS}^{\bullet+}$ solution was prepared by diluting the stock $\text{ABTS}^{\bullet+}$ solution (one mL) with ethanol (35 mL) to get an absorbance at 734 nm (A_{734}) of 0.700 ± 0.025 . After that, 0.3 mL of each partitioned extract at different concentrations dissolved in DMSO was mixed with 2.7 mL of the prepared $\text{ABTS}^{\bullet+}$ solution, left for 6 min in the dark and then the A_{734} was read. The percentage of inhibition was calculated according to Eq. (2),

$$\% \text{ ABTS radical scavenging activity} = [(A - B)/(A)] \times 100, \quad (2)$$

where A is the A_{734} of the negative control and B is the A_{734} of the treatment.

The % inhibition (Y axis) was plotted against the respective extract concentration (X axis) and the EC_{50} value was obtained from the graph. The results were then compared with the Trolox standard curve (0–0.2 mM) and the results are expressed as μmole Trolox equivalents (TE)/g partitioned extract.

Determination of the ferric reducing antioxidant power (FRAP)

The FRAP assay was performed as previously described (Sun et al., 2017). The FRAP reagent was freshly prepared by mixing 100 mL of 300 mM acetate buffer (pH 3.6), 10 mL of 10 mM 2,4,6-tris(2pyridyl)-s-triazine solution in 40 mM hydrochloric acid, and 10 mL of 20 mM ferric chloride solution. Later, 100 μL of the sample at a desired concentration was added to three mL of the FRAP reagent and incubated at 37 °C in a water bath for 6 min. The absorbance was measured at a wavelength of 593 nm. Aqueous solutions of ferrous sulfate (0–2,000 μM) were used to establish the standard graph and the reducing capacity was expressed as μmole of Fe^{2+} /g extract. Each assay was performed in triplicate.

***In vitro* TYRI activity**

The *in vitro* TYRI activity was determined as described (Khongkarat et al., 2020). The reaction mixture contained 120 μL of 2.5 mM L-DOPA in 80 mM phosphate buffer (pH 6.8), 30 μL of 80 mM phosphate buffer (pH 6.8), and 10 μL of partitioned extract or kojic acid (positive control) at different concentrations in DMSO. After mixing, the reaction was pre-incubated at 25 °C for 10 min. Then, 40 μL of 165 units (U)/mL mushroom TYR in phosphate buffer was added and incubated at 25 °C for 5 min. The absorbance at 475 nm (A_{475}) was measured using a microplate reader. The TYRI activity was calculated as the IC_{50} value. Each assay was performed in triplicate and the percentage TYRI was calculated from Eq. (3);

$$\text{Percentage of tyrosinase inhibition} = \frac{\{(A - B) - (C - D)\}}{(A - B)} \times 100, \quad (3)$$

where: A is the A_{475} after incubation without extract, B is the A_{475} after incubation without an extract and tyrosinase, C is the A_{475} after incubation with an extract and tyrosinase, and D is the A_{475} after incubation with an extract, but without tyrosinase.

The % TYRI (Y axis) was plotted against the respective extract concentration (X axis) and the IC_{50} value was obtained from the graph.

***In vitro* LOXI activity**

The LOXI activity of each partitioned extract was determined as described (Sethiya & Mishra, 2014) with some modifications. Firstly, 220 μL of 0.2 M borate buffer pH 9.0, 30 μL of the extract in DMSO, and 250 μL of 20,000 U/mL LOX from soybean in 0.2 M borate buffer pH 9.0 were mixed and incubated at 25 °C for 5 min. After that, 1,000 μL of 0.6 mM linoleic acid solution was added and the absorbance at 234 nm (A_{234}) was measured. Nordihydroguaiaretic acid (NDGA) was used as the reference standard. Each assay was performed in triplicate. The percentage of LOXI was calculated from Eq. (4);

$$\text{Percentage of LOXI} = \frac{\{(A - B) - (C - D)\}}{(A - B)} \times 100, \quad (4)$$

where A is the A_{234} after incubation without an extract, B is the A_{234} after incubation without an extract and LOX, C is the A_{234} after incubation with an extract and LOX, and D is the A_{234} after incubation with an extract but without LOX.

The % LOXI (Y axis) was plotted against the respective extract concentration (X axis) and the IC_{50} value was obtained from the graph.

Enrichment of active fractions

Among the mentioned bioactivities, the most active extract for antioxidants and *in vitro* LOXI activity was further fractionated following the bioactivity as in previous studies (Teerasripreecha et al., 2012).

Fractionation by silica gel 60 column chromatography (SiG60-CC; 500-mL size)

The most active extract for antioxidant and *in vitro* LOXI activities was selected for fractionation by SiG60-CC. Briefly, the 500-mL column was packed with fine SiG60 (Merck, for CC). The partitioned extract (5.64 g) was dissolved in 20 mL of MeOH and

combined with 20 g of rough SiG60 (Merck, for CC). After drying, it was poured over the surface of the packed SiG60 column and then eluted with 6.5 L of DCM, 8.5 L of 7% (v/v) MeOH in DCM, and 3.5 L of MeOH, respectively. Eluted fractions (250 mL each) were collected, and the solvent was removed by evaporation under reduced pressure at a maximum temperature of 40–45 °C. The pattern of chemical compounds in each fraction was profiled by thin layer chromatography (TLC; see below). Fractions with the same TLC pattern were pooled together and tested for antioxidant and *in vitro* LOXI activities.

Chemical profiling by TLC

A sample was spotted onto the starting line of a 5 × five cm² TLC plate (silica immobile phase) using a capillary tube and then dried at room temperature. It was then resolved in one direction using 7% (v/v) MeOH: DCM as the mobile phase. The resolved compounds on the TLC plate were visualized under UV light at 254 nm or by dipping in 3% (v/v) anisaldehyde in MeOH and heating over a hot plate.

Enrichment by HPLC

To separate (enrich) the compounds of a similar polarity in the mixture (extract) the HPLC method reported by *Lv et al. (2015)* was further developed and modified. The optimal operating condition was found using a SB-PHENYL column (5 μm, 9.4 × 250 mm), loading 10 × 20 μL aliquots of the respective sample (100 mg/mL in MeOH) with a column temperature of 25 °C, and eluting in an isocratic mobile phase (two mL/min) of milli Q H₂O and MeOH ranging from 0:100 to 60:40 (v/v) H₂O: MeOH. The eluted fractions were detected by UV-visible spectroscopy at 254 nm (A₂₅₄). The retention time of the extract was determined.

Chemical structure analysis by ¹H-NMR and MS

Among the selected fractions from the SiG60-CC (500 mL size) and HPLC fractions, the most active fraction for both activities was evaporated and analyzed as reported (*Do et al., 2021*). Briefly, the evaporated sample was dissolved in an appropriate deuterated solvent (methanol-d₄, Merck) at a ratio of 5–20 mg of compound to 600 μL of the solvent. Next, it was transferred to an NMR tube and shaken until totally dissolved. The ¹H-NMR spectrum was recorded using a Jeol JNM-ECZ 500MHz operated at 500 MHz for ¹H-NMR nuclei with tetramethylsilane as the internal standard. The chemical shift in δ (ppm) was assigned with reference to the signal from the residual protons in the deuterated solvents, while the chemical shift and J coupling value were determined using the MestReNova version 12.0.3 software. The MW of the active fractions was analyzed using a microTOF focus II MS with electrospray ionization in the positive mode and α-cyano-4-hydroxycinnamic acid as the matrix.

Data analysis

All experiments were done in triplicate. Numerical data are reported as the mean ± one standard deviation (SD), determined in the Microsoft Excel 2019 software (*Khongkarat et al., 2022*). One-way ANOVA and *T*-test were used to test for significant differences. Tukey's and Dunnett T3 test (*p* < 0.05) was applied for pairwise multiple comparisons

(Durovic *et al.*, 2022). All statistical analyses were performed using the IBM SPSS statistics version 22 for windows software.

The overall procedure of BP screening and enrichment of the antioxidant and enzyme inhibitory activities from the most active extract is summarized schematically in Fig. 1.

RESULTS

The partitioned extracts of BPs

After partitioning the MCEs of the six different types of BPs with organic solvents from the lowest to the highest in polarity (hexane, DCM, and MeOH), a total of 18 partitioned extracts were obtained. These partitioned extracts were then evaporated and weighed, and the character of each extract was observed, with the results summarized in Supplement 1. The MeOH-partitioned extracts had the highest yield (above 40%). Only the DCM-partitioned extracts exhibited a sticky solid form, whereas the MeOH- and hexane-partitioned extracts exhibited an oil form. The TPC and FC of the MeOH-, DCM- and hexane-partitioned extracts of all six types of BP were then determined.

Determination of the TPC and FC

The TPC and FC were determined from the calibration curves of gallic acid ($y = 8.5519x - 0.0133$; $R^2 = 0.9991$) and quercetin ($y = 0.6342x + 0.0083$; $R^2 = 0.9982$), respectively, with the TPC and FC of each extract shown in Table 2. The effect of the partition solvent on the TPC and FC was significantly evident, with the highest TPC and FC being found in the DCM-partitioned extracts of all six BP samples, followed by the MeOH-partitioned extracts, while the hexane-partitioned extracts had the lowest TPC and no FC. The TPC of these BP extracts varied between 7.20 ± 0.25 and 53.26 ± 0.85 mg GAE/g, being highest in DCMHBP (53.26 ± 0.85 mg GAE/g). The FC varied between 3.09 ± 0.59 and 104.13 ± 3.80 mg QE/g, being highest in DCMMBP (104.13 ± 3.80 mg QE/g). The FC of the hexane-partitioned extracts could not be determined due to the absorption being too low (below the range of the quercetin standard curve). Since the DCM-partitioned extract of each type of BP had the significantly highest TPC and FC, they were used in the subsequent screening for antioxidant and enzyme inhibitory activities.

Antioxidant activity of the partitioned extracts

The results for the three different antioxidant assays of the DCM-partitioned extracts from the six types of BP are shown in the Table 3 along with that for ascorbic acid as the standard reference for the DPPH and ABTS assays. The EC_{50} values ranged from 176.85 ± 8.31 to $2,305.98 \pm 59.53$ $\mu\text{g/mL}$. Only the DCMMBP extract showed a significantly strong DPPH radical scavenging ability (EC_{50} of 176.85 ± 8.31 $\mu\text{g/mL}$), although this was still almost 2.6-fold less effective than ascorbic acid (EC_{50} of 68.00 ± 1.13 $\mu\text{g/mL}$), while the DCMNBP extract had the weakest DPPH scavenging capacity.

For the ABTS assay, the EC_{50} values (Table 3) ranged from 195.59 ± 11.44 to 875.04 ± 49.84 $\mu\text{g/mL}$. Again, only DCMMBP showed a significantly strong ABTS radical scavenging ability (EC_{50} of 195.59 ± 11.44 $\mu\text{g/mL}$), although this was still over four-fold less effective than ascorbic acid (EC_{50} of 44.54 ± 0.19 $\mu\text{g/mL}$), while DCMNBP demonstrated the weakest ABTS scavenging capacity.




Figure 1 Summary of the extraction, screening, and enrichment procedures for the selected BP.

Full-size DOI: [10.7717/peerj.13506/fig-1](https://doi.org/10.7717/peerj.13506/fig-1)

Table 2 The TPC and FC of the partitioned BP extracts.

Partitioned extract	CBP	HBP	MBP	NBP	XBP	ABP
<i>TPC (mg GAE/g):</i>						
MT	29.45 ± 0.94 ^b	20.19 ± 0.75 ^b	24.04 ± 0.37 ^b	11.84 ± 0.23 ^a	18.16 ± 0.16 ^b	26.55 ± 0.37 ^b
DCM	42.44 ± 0.25 ^c	53.26 ± 0.85 ^c	47.82 ± 0.39 ^c	16.83 ± 0.04 ^b	47.91 ± 0.36 ^c	39.33 ± 0.66 ^c
HX	10.45 ± 0.24 ^a	9.69 ± 0.47 ^a	7.20 ± 0.25 ^a	12.48 ± 0.41 ^a	10.03 ± 0.57 ^a	11.90 ± 0.18 ^a
<i>FC (mg QE/g):</i>						
MT	8.86 ± 0.44 ^a	3.09 ± 0.59 ^a	6.39 ± 0.27 ^a	–	4.27 ± 0.16 ^a	16.95 ± 0.44 ^a
DCM	31.07 ± 2.94 ^b	56.13 ± 2.52 ^b	104.13 ± 3.80 ^b	6.74 ± 0.45	58.44 ± 1.36 ^b	24.64 ± 0.44 ^b
HX	–	–	–	–	–	–

Notes.

Data are shown as the mean ± 1SD derived from three replicates. Means within a column with a different superscript letter are significantly different [$p < 0.05$; one-way ANOVA and Post Hoc (Tukey) test (TPC) or Independent-Samples *T*-Test (FC)].

Table 3 Antioxidant activity of the partitioned extracts.

Sample	DPPH EC ₅₀ (μg/mL)	ABTS EC ₅₀ (μg/mL)	ABTS (μmole TE/g)	FRAP (μmole Fe ²⁺ /g)
DCMCBP	1,952.99 ± 8.44 ^c	563.85 ± 23.48 ^{cd}	163.92 ± 10.24 ^{bc}	412.11 ± 11.04 ^c
DCMHBP	2,291.18 ± 36.99 ^d	681.1 ± 28.83 ^{cd}	164.80 ± 7.60 ^{bc}	395.17 ± 13.00 ^{bc}
DCMMBP	176.85 ± 8.31 ^b	195.59 ± 11.44 ^b	296.95 ± 16.87 ^d	1,058.92 ± 28.78 ^e
DCMNBP	>4,000 ^e	>2,000 ^e	35.71 ± 2.98 ^a	141.83 ± 5.61 ^a
DCMXBP	1,970 ± 3.34 ^c	494.04 ± 25.41 ^c	176.58 ± 10.91 ^c	469.61 ± 15.86 ^d
DCMABP	2,305.98 ± 59.53 ^d	875.04 ± 49.84 ^d	136.22 ± 13.60 ^b	351.97 ± 15.67 ^b
Ascorbic acid	68.00 ± 1.13 ^a	44.54 ± 0.19 ^a	–	–

Notes.

Data are shown as the mean ± 1 SD. Within a column, EC₅₀ means with a different superscript letter are significantly different ($p < 0.05$; one-way ANOVA and Post Hoc (Dunnnett T3) test). Likewise, for the mean ABTS and FRAP μmole TE g⁻¹ and μmole Fe²⁺ g⁻¹ values, means within a column with a different superscript letter are significantly different [$p < 0.05$; one-way ANOVA and Post Hoc (Tukey) test].

Furthermore, the ABTS assay was also evaluated in terms of the TEAC, determined from the Trolox calibration curve ($y = -3.0279x + 0.5931$; $R^2 = 0.9978$). The TEAC values of the BP extracts ranged from 35.71 ± 2.98 to 296.95 ± 16.87 μmole TE/g (Table 3), where DCMMBP had the significantly highest TEAC value and DCMNBP the lowest.

The reducing power, in terms of the FRAP value, of the DCM-partitioned extracts was calculated from the calibration curve of FeSO₄ ($y = 0.0004x - 0.0292$; $R^2 = 0.9947$). The significantly highest antioxidant activity was found in DCMMBP ($1,058.92 \pm 28.78$ μmole Fe²⁺/g) (Table 3), while the lowest was found in DCMNBP (141.83 ± 5.61 μmole Fe²⁺/g).

Overall, from the three antioxidant assays (DPPH, ABTS, and FRAP), DCMMBP showed the strongest antioxidant activity. However, its activity was lower than ascorbic acid (positive standard). To increase its activity, DCMMBP was further enriched (fractionated) by chromatographic methods.

Enzyme inhibitory activity

In vitro TYRI activity

Due to the large number of fractions to be screened, all the partitioned extracts were initially screened for TYRI activity at a single final extract concentration of 200 μg/mL.

Table 4 The TYRI and LOXI activities (%/IC₅₀) of the partitioned extracts.

Sample	TYRI (% at 200 μg/mL)/IC ₅₀	LOXI (% at 60 μg/mL)/IC ₅₀
DCMCBP	58.88 ± 0.36/101.13 ± 3.18 ^c	25.80 ± 4.73/–
DCMHBP	60.41 ± 0.45/42.63 ± 1.65 ^b	28.58 ± 4.96/–
DCMMBP	10.05 ± 0.86/–	78.60 ± 2.81/32.55 ± 1.31 ^b
DCMNBP	0.00 ± 0.00/–	15.64 ± 3.28/–
DCMXPB	52.95 ± 0.09/179.97 ± 3.77 ^d	28.27 ± 1.82/–
DCMABP	51.76 ± 0.93/193.87 ± 5.06 ^e	24.47 ± 2.95/–
Kojic acid	–/9.61 ± 0.47 ^a	–
NDGA	–	–/22.41 ± 2.35 ^a

Notes.

The IC₅₀ values are shown as the mean ± 1SD. Within a column, means with a different superscript letter are significantly different ($p < 0.05$; one-way ANOVA and Post Hoc Tukey test for TYRI activity and Independent-Samples T-Test for LOXI).

The obtained A₄₇₅ was converted to the TYRI activity (%) and the results are presented as the mean ± SD in Table 4. At this extract concentration, DCMCBP, DCMHBP, DCMXPB, and DCMABP showed a more than 50% TYRI activity, and so they were further evaluated at different concentrations and their derived IC₅₀ values are shown in Table 4. Of these extracts, DCMHBP had the significantly highest TYRI activity (IC₅₀ 42.63 ± 1.65 μg/mL) but this was significantly (some 4.4-fold) less effective than kojic acid (IC₅₀ of 9.61 ± 0.47 μg/mL).

***In vitro* LOXI activity**

The partitioned extracts were also initially screened for LOXI activity at a single final extract concentration of 60 μg/mL, with the LOXI activity (%) presented as the mean ± SD in Table 4. At this extract concentration, only DCMMBP provided a high *in vitro* LOXI activity (78.60 ± 2.81%) and so was further evaluated at different concentrations, revealing an IC₅₀ value of 32.55 ± 1.31 μg/mL (Table 4), some 1.4-fold less significantly effective than NDGA (22.41 ± 2.35 μg/mL).

**Antioxidant and LOXI activities of compounds from DCMMBP
Fractionation of DCMMBP by SiG60-CC**

Since the DCMMBP provided the best LOXI and antioxidant activities, the sample (5.64 g) was further enriched using SiG60-CC. A total of 74 fractions were collected, but after pooling fractions with a similar TLC plate profile, five different fractions (DCMMBP1–5) were obtained (Supplement 2). The TLC profile DCMMBP1 (lane 2, R_f of 0.90) revealed one major band under UV light and a few bands after staining with anisaldehyde, while DCMMBP2 (lane 3, R_f of 0.26, 0.38, and 0.59) showed two bands under UV light and one more major band after staining. For DCMMBP3 (lane 4, R_f of 0.21), it showed only one major band under UV light and so it might be a pure compound, while DCMMBP 4 (lane 5, R_f of 0.00 and 0.21) and DCMMBP 5 (lane 6, R_f of 0.00 and 0.21) each showed a major band at the base line and a minor band under UV light. When the TLC profiles of each fraction were compared to the original fraction (lane 1, R_f of 0.23), the compound in fraction 3 was found to be the main active compound. Their respective weights and characteristics are recorded in Supplement 1.

All five pooled fractions were separately screened for their antioxidant and LOXI activities. An antioxidant activity of more than 50% was found in DCMMBP3 ($81.21 \pm 1.38\%$) and DCMMBP4 ($69.52 \pm 1.52\%$) at a concentration of $500 \mu\text{g/mL}$ based on the DPPH assay. The ABTS assay revealed more 50% activity for the DCMMBP2 ($92.87 \pm 4.00\%$), DCMMBP3 ($99.32 \pm 0.01\%$), DCMMBP4 ($97.97 \pm 0.57\%$), and DCMMBP5 ($72.12 \pm 1.49\%$). However, at a concentration of $20 \mu\text{g/mL}$, a LOXI activity of over 50% was only found in DCMMBP3 ($73.35 \pm 0.85\%$). Therefore, these fractions were further investigated at various concentrations and their respective EC_{50} and IC_{50} values derived.

The data for the DPPH assay are shown in [Table 5](#), where the EC_{50} value of DCMMBP3 and DCMMBP4 was 54.66 ± 3.45 and $184.84 \pm 5.47 \mu\text{g/mL}$, respectively, and DCMMBP3 was significantly more effective than ascorbic acid ($68.00 \pm 1.13 \mu\text{g/mL}$) and the parental DCMMBP extract ($176.85 \pm 8.31 \mu\text{g/mL}$; [Table 3](#)).

For the ABTS assay, the obtained IC_{50} values are shown in [Table 5](#). The EC_{50} value and $\mu\text{mole TE/g}$ for the ABTS assay of DCMMBP3 was $24.56 \pm 2.99 \mu\text{g/mL}$ and $2,529.69 \pm 142.16 \mu\text{mole TE/g}$, respectively, which was significantly stronger than that for ascorbic acid ($44.54 \pm 0.19 \mu\text{g/mL}$) and the parental DCMMBP extract ($195.59 \pm 11.44 \mu\text{g/mL}$ and $296.95 \pm 16.87 \mu\text{mole TE/g}$, respectively).

For the FRAP assay ([Table 5](#)), DCMMBP3 was $3,466.17 \pm 81.30 \mu\text{mole Fe}^{2+}/\text{g}$, which was over three-fold higher than that the parental DCMMBP extract ($1,058.92 \pm 28.78 \mu\text{mole Fe}^{2+}/\text{g}$; [Table 3](#)).

For the LOXI activity, the IC_{50} values are reported in [Table 5](#). The IC_{50} value for the LOXI activity of DCMMBP3 was $12.11 \pm 0.36 \mu\text{g/mL}$, which was significantly (1.85- and 2.69-fold) more effective than that of NDGA (IC_{50} of $22.41 \pm 2.35 \mu\text{g/mL}$) and the parental DCMMBP extract (IC_{50} of $32.55 \pm 1.31 \mu\text{g/mL}$; [Table 4](#)), respectively. Therefore, DCMMBP3 was further enriched by HPLC.

Fractionation of DCMMBP3 by HPLC

To optimize the separation, the HPLC was eluted with an isocratic gradient of 0:100 to 60:40 (v/v) H_2O : MeOH, which separated DCMMBP3 into five peaks, with the two main peaks eluting at a retention time of 15.352 and 20.182 min, respectively, ([Fig. 2](#)). These two fractions (DCMMBP3-1 and DCMMBP3-2) were defined as compounds 1 and 2, respectively. Their weight and characters are summarized in [Supplement 1](#). Compounds 1 and 2 were tested for their antioxidant and LOXI activities, where compounds 1 and 2 showed similar antioxidant and LOXI activities ([Table 5](#)). The structure of compounds 1 and 2 was characterized by $^1\text{H-NMR}$ and MS analyses, and found to share the same structure, consistent with their bioactivities.

Structural identification of compounds 1 and 2

After fractionation of DCMMBP3 by HPLC, compounds 1 and 2 were obtained. The $^1\text{H-NMR}$ spectra of compounds 1 ([Supplement 3](#)) and 2 ([Supplement 4](#)) revealed essentially similar signals to each other. Their MS spectra also showed identical pseudomolecular ions $[\text{M}+\text{Na}]^+$ and $[\text{M}+\text{H}]^+$ as well as mass fragment ion $[\text{M}+\text{Na}-177]^+$ ([Supplement 5–Supplement 7](#)) at 696, 674, and 522, respectively. The above mentioned spectroscopic

Table 5 Antioxidant and LOXI activities of the respective fractions after SiG60 and HPLC chromatography.

Sample	DPPH EC ₅₀ ($\mu\text{g/mL}$)	ABTS EC ₅₀ ($\mu\text{g/mL}$)	ABTS ($\mu\text{mol TE/g}$)	FRAP ($\mu\text{mol Fe}^{2+}/\text{g}$)	LOX IC ₅₀ ($\mu\text{g/mL}$)
<i>After SiG60-CC:</i>					
DCMMBP1	–	–	86.63 \pm 6.84 ^a	304.50 \pm 33.46 ^a	–
DCMMBP2	–	125.81 \pm 12.97 ^d	878.09 \pm 73.48 ^c	1,038.11 \pm 34.60 ^c	–
DCMMBP3	54.66 \pm 3.45 ^a	24.56 \pm 2.99 ^a	2,529.69 \pm 142.16 ^c	3,466.17 \pm 81.30 ^c	12.11 \pm 0.36 ^a
DCMMBP4	184.84 \pm 5.47 ^c	66.1 \pm 3.55 ^c	1,482.24 \pm 63.00 ^d	1,363.11 \pm 47.30 ^d	–
DCMMBP5	–	318.66 \pm 8.66 ^c	456.82 \pm 45.31 ^b	657.56 \pm 23.52 ^b	–
Ascorbic acid	68.00 \pm 1.13 ^b	44.54 \pm 0.19 ^b	–	–	–
NDGA	–	–	–	–	22.41 \pm 2.35 ^b
<i>After HPLC:</i>					
DCMMBP3-1	53.05 \pm 2.60 ^a	25.38 \pm 0.67 ^a	2,527.63 \pm 7.99 ^a	3,477.33 \pm 52.07 ^a	11.31 \pm 0.46 ^a
DCMMBP3-2	57.7 \pm 2.77 ^a	28.29 \pm 1.05 ^b	2,339.16 \pm 34.62 ^b	3,091.78 \pm 44.48 ^b	11.57 \pm 0.93 ^a
Ascorbic acid	68.00 \pm 1.13 ^b	44.54 \pm 0.19 ^c	–	–	–
NDGA	–	–	–	–	22.41 \pm 2.35 ^b

Notes.

Data are shown as the mean \pm 1SD. Means within a column with a different superscript letter are significantly different ($p < 0.05$; one-way ANOVA plus for samples after SiG60-CC: Post Hoc (Tukey) test for DPPH and FRAP data, Dunnett T3 test for ABTS data, independent samples T-Test for IC₅₀ values; for post-HPLC data: Tukey test for DPPH and ABTS EC₅₀ data, independent samples T-Test for ABTS $\mu\text{mol TE/g}$ and FRAP values; and Dunnett test for LOX IC₅₀ values).

data suggested that compounds 1 and 2 were isomeric compounds. Compound 1 exhibited chemical shifts of the methylene hydrocarbon ($-\text{CH}_2-$) at 1.43 and 1.87 ppm and of a methylene hydrocarbon connected to a nitrogen atom at 3.18 ppm, thus indicating the presence of a spermidine moiety. A cluster of singlet methoxy peaks at around 3.75 ppm together with the chemical shifts between 6.73 and 7.48 ppm suggested the presence of a phenolic compound containing methoxy groups. The chemical shifts at 5.95 and 6.40 ppm with small coupling constants of 12.9 Hz indicated *cis*-olefinic protons, while those of 6.40 and 7.40 ppm with large coupling constants of 15.8 Hz suggested *trans*-olefinic protons. These signals in the aromatic region suggested the presence of *cis*- and *trans*-feruloyl moieties (Fig. 3A). To determine the ratio of *cis*- and *trans*-feruloyl moieties in the whole structure, the signal integration of olefinic protons was analyzed. A ratio of 1:2 suggested that compound 1 was comprised one *cis*-feruloyl and two *trans*-feruloyls in the molecule. Therefore, three possible structures of compound 1 were established; N^1, N^5, N^{10} -tri-(*Z, E, E*)-, N^1, N^5, N^{10} -tri-(*E, Z, E*)-, and N^1, N^5, N^{10} -tri-(*E, E, Z*)-feruloyl spermidine, respectively, as shown in Figs. 3C–3E. An attempt to determine exactly which structure matched to the spectroscopic data was unsuccessful due to the severely overlapped olefinic protons. In nature, the *cis*-isomer is less stable and gradually isomerizes into the *trans*-congener. Compound 2 was nearly identical to compound 1, except for the lack of *cis*-olefinic protons (Fig. 3B). This data suggested that compound 2 is acyl spermidine containing all *trans*-feruloyl moieties. Therefore, the structure of 2 was established as N^1, N^5, N^{10} -tri-(*E, E, E*)-feruloyl spermidine, as shown in Fig. 3F.




Figure 2 The HPLC chromatogram of DCMMBP3 showing the elution of DCMMBP3-1 and DCMMBP3-2 at a retention time of 15.352 and 20.182 min, respectively.

Full-size  DOI: [10.7717/peerj.13506/fig-2](https://doi.org/10.7717/peerj.13506/fig-2)

DCMMBP3 fraction (Mixture of compounds 1 and 2)

$^1\text{H-NMR}$ (500 MHz, methanol- d_4) δ 7.54–7.33 (m, 5H), 7.14–6.62 (m, 17H), 6.61–6.43 (m, 1H), 6.43–6.31 (m, 3H), 3.81 (qd, $J = 7.5, 3.5$ Hz, 20H), 3.76–3.70 (m, 3H), 3.60–3.48 (m, 4H), 3.48–3.39 (m, 2H), 3.32 (d, $J = 5.8$ Hz, 12H), 3.26–3.13 (m, 1H), 1.95–1.78 (m, 3H), 1.74–1.51 (m, 7H), 1.50–1.36 (m, 1H), and 1.31–1.23 (m, 1H).

DISCUSSION

Different types of BP can provide different amounts and types of secondary metabolites, depending on their floral origin, which can result in a wide range of bioactivities between different BPs (*Rzepecka-Stojko et al., 2015*). In our work, six monofloral *A. mellifera* BPs were evaluated because many types of monofloral BP are harvested in Thailand due to the large amount of diverse monocrops that are cultivated. Moreover, the floral origin of monofloral BP is easier to identify, which makes it easier to control the quality and pharmaceutical properties of the BP compared to that of multifloral BP (*Campos et al., 2010*). Since bee products are edible, the food safety aspect should be considered, including the risks associated with pesticides, toxic metals, mycotoxins, allergens, and pollen grains of inedible plants (*Vegh et al., 2021*). There is a risk bias in certain plant groups (*Helianthus*, *Rosaceae*, *Raxinus*, and *Sophora*) that can contain high concentrations of toxic metals




Figure 3 $^1\text{H-NMR}$ (500 MHz, MeOD-D_4) in the range of 7.6–5.7 ppm of (A) compound 1 and (B) compound 2. The chemical structures of (C–F) spermidine derivatives. $^1\text{H-NMR}$ (500 MHz, MeOD-D_4) in the range of 7.6–5.7 ppm of (A) compound 1 and (B) compound 2. The chemical structures of (C) N^1,N^5,N^{10} -tri-(*Z,E,E*)-feruloyl spermidine, (D) N^1,N^5,N^{10} -tri-(*E,Z,E*)-feruloyl spermidine, (E) N^1,N^5,N^{10} -tri-(*E,E,Z*)-feruloyl spermidine, and (F) N^1,N^5,N^{10} -tri-(*E,E,E*)-feruloyl spermidine.

Full-size DOI: 10.7717/peerj.13506/fig-3

(Kostic *et al.*, 2015). In addition, BPs from pine, canola, kiwi, willow, corn poppy, rose, lotus, and camellia contain allergens that can induce anaphylaxis in some people (Yang *et al.*, 2019).

The most practical method to determine the principal botanical origin(s) of BP is palynological analysis using microscopic examination (Kieliszek *et al.*, 2018), but molecular analysis can also be used for minor components. Partitioning of BP with different polar organic solvents can separate the compounds based upon their polarity. This process makes further enrichment easier by screening for only the active extracts to undergo further fractionation. Of the different solvent extracts, the DCM-partitioned extracts of the six BPs in this study all showed the highest TPC and FC. Among these six DCM-partitioned BP extracts, DCMMBP and DCMHBP had the highest amount of both TPC and TFC. This result was consistent with Chantarudee *et al.* (2012) who reported that the DCM-partitioned extracts of corn (*Zea mays*) BP had the highest antioxidant activity compared to that of the MeOH- and hexane-partition extracts. Moreover, Bittencourt *et al.* (2015) reported that DCM partitioning of Brazilian BP enriched its antioxidant activity. As in previous reports, the active component for the antioxidant, TYRI, and LOXI activities in natural products were found to be phenolic and flavonoid compounds (Rebiai & Lanez, 2012; Sroka, Sowa

& Drys, 2017; Kim et al., 2015). Therefore, the DCM-partitioned extracts of the BPs that provided the highest TPC and TFC were selected to study for their antioxidant, TYRI, and LOXI activities.

In this work, the *in vitro* LOXI and TYRI activities were assayed using soybean LOX and mushroom TYR since both of these enzymes show structural and functional similarities with human enzymes. Therefore, they are commonly used as a model for human enzyme inhibition (Moita et al., 2014; Chang, 2009). The highest TYRI activity was found in DCMHBP followed by in DCMMBP, while the strongest LOXI activity was found in DCMMBP. Both activities were also correlated with the very high TPC and FC found in DCMHBP and DCMMBP. It was previously reported that phenol or flavonoid compounds can chelate the copper and iron ions in the active sites of TYR and LOX enzymes, respectively, and so it is highly possible that an antioxidant compound, especially a phenolic compound, can also inhibit 5-LOX (Obaid et al., 2021; Ratnasari, Walters & Tsopmo, 2017).

By comparison to the polyamine derivatives separated from *Microdesmis keayana* roots (Zamble et al., 2006), the two main active components in our work are likely to be N^1 , N^5 , N^{10} -triferuloyl spermidine or Keayanidine C. This finding agrees with the report on the phenolic profile in monofloral BP in Brazil, which also found N^1 , N^5 , N^{10} -triferuloyl spermidine in *Mimosa scabrella* BP (De-Melo et al., 2018). However, in our work, the $^1\text{H-NMR}$ of compound 1 showed *cis*-olefinic hydrogen and *trans*-olefinic hydrogen at a 1:2 ratio, while compound 2 showed *trans*-olefinic hydrogens. Therefore, compounds 1 and 2 are N^1 , N^5 , N^{10} -tri(*E*, *E*, *E*)feruloyl spermidine and N^1 , N^5 , N^{10} -tri(*Z*, *Z*, *Z*)feruloyl spermidine, respectively, which are different in their configuration. The HPLC chromatographic profile agreed with previous works that reported on the separation of this compound by HPLC from *Hippesastrum* × hortorum pollen (Youhnovski, Werner & Hesse, 2001) and *Sambucus nigra* L. (Kite et al., 2013). According to their findings, the HPLC chromatogram of this compound revealed four peaks corresponding to the four isomers: *ZZZ*, *EZZ*, *ZZE*, and *EEE*. These support that the peaks in our chromatogram are N^1 , N^5 , N^{10} -triferuloyl spermidine in different conformations. In fact, the *E*- and *Z*-feruloyl moiety can be interconverted on exposure to UV light, in which the *Z*-isomer is dominant. Although compound 1 contained a small proportion of *Z*-feruloyl moiety, the antioxidation and LOXI activity were comparable to those of compound 2, whose structure had no *Z*-feruloyl residues. These observations preliminarily suggest that the geometry of $\text{C}=\text{C}$ unsaturation contributed to a low or no effect on the biological activity. Thus, the feruloyl spermidines could potentially be applied as a mixture of isomers without the need for further time-consuming separation processes, such as HPLC, to obtain an individual pure isomer. In addition to feruloyl spermidines, other related spermidines acylated by a series of phenolics have been reported. A variety of coumaroyl spermidines isolated from rape BP (Zhang, Liu & Lu, 2020) were also demonstrated to have potent antioxidant activities in the DPPH, ABTS, and FRAP assays.

From the data mentioned above, these results showed that the BP samples harvested in Thailand, especially *M. diplotricha* BP, had potential bioactivities. The findings additionally indicated the relationship between the phytochemicals and functional/pharmacological

application. Also, the obtained data supported the traditional or alternative use of *M. diplotricha* BP in the indigenous medicine of Thailand.

CONCLUSIONS

Mimosa diplotricha BP had the highest antioxidant and LOXI activities, which were correlated to the highest TPC and FC. Although both bioactivities of the DCM-partitioned BP extracts were detected, their fractionation led to an enrichment of triferuloyl spermidines and enhanced specific bioactivities, suggesting these might be the major active compounds in the *Mimosa diplotricha* BP. In addition, this work supported the importance of the floral origin of BPs, since it affected the bioactivity directly. Here, while *Mimosa diplotricha* BP had the best antioxidant and LOXI bioactivities, *Helianthus annuus* L. BP had the best TYRI activity.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was financially supported by the Science Achievement Scholarship of Thailand, the 90th Anniversary of Chulalongkorn University Fund (Ratchadaphiseksomphot Endowment Fund), Toray Science Foundation, and Thailand Science Research and Innovation Fund Chulalongkorn University (CUFRB65_food(6)_114_23_44). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Science Achievement Scholarship of Thailand.

90th Anniversary of Chulalongkorn University Fund (Ratchadaphiseksomphot Endowment Fund).

Toray Science Foundation, and Thailand Science Research and Innovation Fund Chulalongkorn University: CUFRB65_food(6)_114_23_44.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Phanthiwa Khongkarat conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Preecha Phuwapraisirisan conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Chanpen Chanchao conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

The raw data is available in the [Supplemental Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.13506#supplemental-information>.

REFERENCES

- Bittencourt ML, Ribeiro PR, Franco RL, Hilhorst HW, De Castro RD, Fernandez LG. 2015.** Metabolite profiling, antioxidant and antibacterial activities of Brazilian propolis: use of correlation and multivariate analyses to identify potential bioactive compounds. *Food Research International* 76(Pt 3):449–457 DOI 10.1016/j.foodres.2015.07.008.
- Campos MGR, Frigerio C, Lopes J, Bogdanov S. 2010.** What is the future of Bee-Pollen? *JAAS* 2(4):131–144 DOI 10.3896/IBRA.4.02.4.01.
- Chamchumroon V, Suphuntee N, Tetsana N, Poopath M, Tanikkool S. 2017.** *Threatened plants in Thailand*. Bangkok: Omega Printing Co. Ltd.
- Chang TS. 2009.** An updated review of tyrosinase inhibitors. *International Journal of Molecular Sciences* 10(6):2440–2475 DOI 10.3390/ijms10062440.
- Chantarudee A, Phuwapraisirisan P, Kimura K, Okuyama M, Mori H, Kimura A, Chanchao C. 2012.** Chemical constituents and free radical scavenging activity of corn pollen collected from *Apis mellifera* hives compared to floral corn pollen at Nan, Thailand. *BMC Complementary and Alternative Medicine* 12:45 DOI 10.1186/1472-6882-12-45.
- Daudu OM. 2019.** Bee pollen extracts as potential antioxidants and inhibitors of α -amylase and α -glucosidase enzymes *in vitro* assessment. *Journal of Apicultural Science* 63(2):315–325 DOI 10.2478/jas-2019-0020.
- De-Melo AAM, Estevinho LM, Moreira MM, Delerue-Matos C, Freitas ADSD, Barth OM, Almeida-Muradian LBD. 2018.** Phenolic profile by HPLC-MS, biological potential, and nutritional value of a promising food: monofloral bee pollen. *Journal of Food Biochemistry* 42(5):e12536 DOI 10.1111/jfbc.12536.
- Do TML, Duong T-H, Nguyen V-K, Phuwapraisirisan P, Doungwichitrkul T, Ni-amnont N, Jarupinthusophon S, Sichaem J. 2021.** Schomburgkixanthone, a novel bixanthone from the twigs of *Garciniaschomburgkiana*. *Natural Product Research* 35(21):3613–3618 DOI 10.1080/14786419.2020.1716351.
- Durovic S, Sorgic S, Popov S, Pezo L, Maskovic P, Blagojevic S, Zekovic Z. 2022.** Recovery of biologically active compounds from stinging nettle leaves part I: supercritical carbon dioxide extraction. *Food Chemistry* 373:131724 DOI 10.1016/j.foodchem.2021.131724.
- Eshwarappa RS, Ramachandra YL, Subaramaihha SR, Subbaiah SG, Austin RS, Dhananjaya BL. 2016.** Anti-lipoxygenase activity of leaf gall extracts of *Terminalia*

- chebula* (Gaertn.) Retz.(Combretaceae). *Pharmacognosy Research* **8**(1):78–82
DOI 10.4103/0974-8490.171103.
- Khongkarat P, Ramadhan R, Phuwapraisirisan P, Chanchao C. 2020.** Safflospermidines from the bee pollen of *Helianthus annuus* L. exhibit a higher *in vitro* antityrosinase activity than kojic acid. *Heliyon* **6**(3):e03638 DOI 10.1016/j.heliyon.2020.e03638.
- Khongkarat P, Traiyasut P, Phuwapraisirisan P, Chanchao C. 2022.** First report of fatty acids in *Mimosa diplotricha* bee pollen with *in vitro* lipase inhibitory activity. *PeerJ* **10**:e12722 DOI 10.7717/peerj.12722.
- Kieliszek M, Piwowarek K, Kot AM, Blazejak S, Chlebowska-Smigiel A, Wolska I. 2018.** Pollen and bee bread as new health-oriented products: a review. *Trends in Food Science and Technology* **71**:170–180 DOI 10.1016/j.tifs.2017.10.021.
- Kim SB, Jo YH, Liu Q, Ahn JH, Hong IP, Han SM, Hwang BY, Lee MK. 2015.** Optimization of extraction condition of bee pollen using response surface methodology: correlation between anti-melanogenesis, antioxidant activity, and phenolic content. *Molecules* **20**(11):19764–19774 DOI 10.3390/molecules201119656.
- Kite GC, Larsson S, Veitch NC, Porter EA, Ding N, Simmonds MSJ. 2013.** Acyl spermidines in inflorescence extracts of elder (*Sambucus nigra* L., Adoxaceae) and elderflower drinks. *Journal of Agricultural and Food Chemistry* **61**(14):3501–3508 DOI 10.1021/jf304602q.
- Kostic AZ, Pesic MB, Mosic MD, Dojcinovic BP, Natic MM, Trifkovic JD. 2015.** Mineral content of bee pollen from Serbia. *Archives of Industrial Hygiene and Toxicology* **66**(4):251–258 DOI 10.1515/aiht-2015-66-2630.
- Lobo V, Patil A, Phatak A, Chandra N. 2010.** Free radicals, antioxidants, and functional foods: impact on human health. *Pharmacognosy Reviews* **4**(8):118–126 DOI 10.4103/0973-7847.70902.
- Lv H, Wang X, He Y, Wang H, Suo Y. 2015.** Identification and quantification of flavonoid aglycones in rape bee pollen from Qinghai-Tibetan Plateau by HPLC-DADAPCI/MS. *Journal of Food Composition and Analysis* **38**:49–54 DOI 10.1016/j.jfca.2014.10.011.
- Marghitas LA, Stanciu OG, Dezmirean DS, Bobis O, Popescu O, Bogdanov S, Campos MG. 2009.** *In vitro* antioxidant capacity of honeybee-collected pollen of selected floral origin harvested from Romania. *Food Chemistry* **115**(3):878–883 DOI 10.1016/j.foodchem.2009.01.014.
- Mashima R, Okuyama T. 2015.** The role of lipoxygenases in pathophysiology; new insights and future perspectives. *Redox Biology* **6**:297–310 DOI 10.1016/j.redox.2015.08.006.
- Moita E, Sousa C, Andrade PB, Fernandes F, Pinho BR, Silva LR, Valentao P. 2014.** Effects of *Echium plantagineum* L. bee pollen on basophil degranulation: relationship with metabolic profile. *Molecules* **19**(7):10635–10649 DOI 10.3390/molecules190710635.

- Obaid RJ, Mughal EU, Naeem N, Sadiq A, Alsantali RI, Jassas RS, Moussa Z, Ahmed SA. 2021.** Natural and synthetic flavonoid derivatives as new potential tyrosinase inhibitors: a systematic review. *RSC Advances* **11**:22159–22198 DOI [10.1039/d1ra03196a](https://doi.org/10.1039/d1ra03196a).
- Pham-Huy LA, He H, Pham-Huy C. 2010.** Free radicals, antioxidants in disease, and health. *International Journal of Biomedical Science* **4**:89.
- Ratnasari N, Walters M, Tsopmo A. 2017.** Antioxidant and lipoxygenase activities of polyphenol extracts from oat brans treated with polysaccharide degrading enzymes. *Heliyon* **3**(7):e00351 DOI [10.1016/j.heliyon.2017.e00351](https://doi.org/10.1016/j.heliyon.2017.e00351).
- Rebiai A, Lanez T. 2012.** Chemical composition and antioxidant activity of *Apis mellifera* bee pollen from northwest Algeria. *Journal of Fundamental and Applied Sciences* **4**(2):155–163 DOI [10.4314/jfas.v4i2.5](https://doi.org/10.4314/jfas.v4i2.5).
- Rzepecka-Stojko A, Stojko J, Kurek-Gorecka A, Gorecki M, Kabala-Dzik A, Kubina R, Mozdziejz A, Buszman E. 2015.** Polyphenols from bee pollen: structure, absorption, metabolism and biological activity. *Molecules* **20**(12):21732–21749 DOI [10.3390/molecules201219800](https://doi.org/10.3390/molecules201219800).
- Saral O, Sahin H, Saral S, Alkanat M, Akyildiz K, Topcu A, Yilmaz A. 2022.** Bee pollen increases hippocampal brain-derived neurotrophic factor and suppresses neuroinflammation in adult rats with chronic immobilization stress. *Neuroscience Letters* **766**:136342 DOI [10.1016/j.neulet.2021.136342](https://doi.org/10.1016/j.neulet.2021.136342).
- Saric A, Balog T, Sobocanec S, Kusic B, Sverko V, Rusak G, Likic S, Bubalo D, Pinto B, Reali D, Marotti T. 2009.** Antioxidant effects of flavonoid from Croatian *Cystus incanus* L. rich bee pollen. *Food and Chemical Toxicology* **47**(3):547–554 DOI [10.1016/j.fct.2008.12.007](https://doi.org/10.1016/j.fct.2008.12.007).
- Sethiya NK, Mishra SH. 2014.** Investigation of mangiferin, as a promising natural polyphenol xanthone on multiple targets of Alzheimer's disease. *Journal of Biologically Active Products from Nature* **4**:2, 111–119 DOI [10.1080/22311866.2014.921121](https://doi.org/10.1080/22311866.2014.921121).
- Shahidi F, Zhong Y. 2015.** Measurement of antioxidant activity. *Journal of Functional Foods* **18**:757–781 DOI [10.1016/j.jff.2015.01.047](https://doi.org/10.1016/j.jff.2015.01.047).
- Sommano SR, Bhat FM, Wongkeaw M, Sriwichai T, Sunanta P, Chuttong B, Burgett M. 2020.** Amino acid profiling and chemometric relations of black dwarf honey and bee pollen. *Frontiers in Nutrition* **7**:558579 DOI [10.3389/fnut.2020.558579](https://doi.org/10.3389/fnut.2020.558579).
- Sroka Z, Sowa A, Drys A. 2017.** Inhibition of lipoxygenase and peroxidase reaction by some flavonols and flavones: the structure–activity relationship. *Natural Product Communications* **12**(11):1705–1708 DOI [10.1177/1934578X1701201111](https://doi.org/10.1177/1934578X1701201111).
- Sun L, Guo Y, Zhang Y, Zhuang Y. 2017.** Antioxidant and anti-tyrosinase activities of phenolic extracts from rape bee pollen and inhibitory melanogenesis by cAMP/MITF/TYR pathway in B16 mouse melanoma cells. *Frontiers in Pharmacology* **8**:104 DOI [10.3389/fphar.2017.00104](https://doi.org/10.3389/fphar.2017.00104).
- Suriyatem R, Auras RA, Intipunya P, Rachtanapun P. 2017.** Predictive mathematical modeling for EC₅₀ calculation of antioxidant activity and antibacterial ability

- of Thai bee products. *Journal of Applied Pharmaceutical Science* 7(09):122–133
DOI 10.7324/JAPS.2017.70917.
- Teerasripreecha D, Phuwapraisirisan P, Puthong S, Kimura K, Okuyama M, Mori H, Kimura A, Chanchao C. 2012.** *In vitro* antiproliferative/cytotoxic activity on cancer cell lines of a cardanol and a cardol enriched from Thai *Apis mellifera* propolis. *BMC Complementary and Alternative Medicine* 12:27 DOI 10.1186/1472-6882-12-27.
- Umthong S, Phuwapraisirisan P, Puthong S, Chanchao C. 2011.** *In vitro* antiproliferative activity of partially purified *Trigona laeviceps* propolis from Thailand on human cancer cell lines. *BMC Complementary and Alternative Medicine* 11:37 DOI 10.1186/1472-6882-11-37.
- Vegh R, Csoka M, Soros C, Sipos L. 2021.** Food safety hazards of bee pollen—a review. *Trends in Food Science and Technology* 114:490–509 DOI 10.1016/j.tifs.2021.06.016.
- Wang B, Wu L, Chen J, Dong L, Chen C, Wen Z, Hu J, Fleming I, Wang DW. 2021.** Metabolism pathways of arachidonic acids: mechanisms and potential therapeutic targets. *Signal Transduction and Targeted Therapy* 6(1):94 DOI 10.1038/s41392-020-00443-w.
- Yang Y, Wang H, Liu M, Huang W, Wang Y, Wu Y. 2019.** A multiplex real-time PCR method applied to detect eight pollen species in food for the prevention of allergies. *European Food Research and Technology* 245:2195–2207 DOI 10.1007/s00217-019-03327-8.
- Yildiz OK, Karahalil FA, Can Z, Sahin H, Kolayli SE. 2014.** Total monoamine oxidase (MAO) inhibition by chestnut honey, pollen, and propolis. *Journal of Enzyme Inhibition and Medicinal Chemistry* 29(5):690–694 DOI 10.3109/14756366.2013.843171.
- Youhnovski N, Werner C, Hesse M. 2001.** N, N', N'-Triferuloylspermidine, a new UV absorbing polyamine derivative from pollen of *Hippeastrum × hortorum*. *Zeitschrift für Naturforschung C* 56(7–8):526–530 DOI 10.1515/znc-2001-7-809.
- Zamble A, Sahpaz S, Hennebelle T, Carato P, Bailleul F. 2006.** N¹, N⁵, N¹⁰-Tris (4-hydroxycinnamoyl) spermidines from *Microdesmis keayana* roots. *Chemistry & Biodiversity* 3(9):982–989 DOI 10.1002/cbdv.200690107.
- Zhang H, Liu R, Lu Q. 2020.** Separation and characterization of phenolamines and flavonoids from rape bee pollen, and comparison of their antioxidant activities and protective effects against oxidative stress. *Molecules* 25(6):1264 DOI 10.3390/molecules25061264.