

MEETING ABSTRACT

Open Access

Identifying forces that stabilize the oligomeric state of bacterial homologs of neurotransmitter transporters

Kumaresan Jayaraman, Azmat Sohail, Michael Freissmuth, Harald H Sitte*, Thomas Stockner

From 18th Scientific Symposium of the Austrian Pharmacological Society (APHAR). Joint meeting with the Croatian, Serbian and Slovenian Pharmacological Societies. Graz, Austria. 20-21 September 2012

Background

Neurotransmitter transporters of the SLC1 and SLC6 family are found on presynaptic neurons and on glia cells. The function of these transporters is the termination of neurotransmission by the rapid removal of the neurotransmitter molecules from the synaptic cleft. These transporters couple substrate transport to ion gradients of sodium and chloride. Almost all of the eucaryotic transporters have been described to function as oligomers. However, the forces stabilizing the oligomeric state are not well understood. No crystal structures of eucaryotic transporters are available, but recently crystal structures of bacterial homologs thereof have been solved: GltPh (SLC 1 family) was found as a trimer, LeuT (SLC6 family) was crystallized as a dimer. These homologous crystal structures allow rationalizing on the driving forces that stabilize the eucaryotic counterparts.

Methods

The crystal structures of LeuT and GltPh were obtained from the Protein Data Bank (PDB). We identified the interfaces between the protomers and analyzed hydrogen bonding, hydrophobic and hydrophilic interactions as well as size and width of the interface area.

Results

We investigated the protein-protein interfaces between the transporter protomers and identified the dominant forces that stabilize the oligomer. These consist of hydrophobic interactions between the aliphatic side chains

within the interface and of polar interactions by hydrogen bonds between hydroxyl groups.

Conclusions

The contributions of different forces to the stability of oligomer assemblies vary between proteins. While the hydrophobic mismatch is a prominent contributor to the stability of the GltPh transporter, it plays a minor role for LeuT, where helix packing and aromatic interactions seem to dominate.

Acknowledgements

This work was supported by the Austrian Science Fund (FWF, project no. F3506 to H.H.S.).

Published: 17 September 2012

doi:10.1186/2050-6511-13-S1-A59

Cite this article as: Jayaraman et al.: Identifying forces that stabilize the oligomeric state of bacterial homologs of neurotransmitter transporters. *BMC Pharmacology and Toxicology* 2012 **13**(Suppl 1):A59.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

 **BioMed Central**

* Correspondence: harald.sitte@meduniwien.ac.at
Institute of Pharmacology, Center for Physiology and Pharmacology, Medical University Vienna, 1090 Vienna, Austria