

State of the Review



OPEN ACCESS

Received: Mar 26, 2022

Accepted: Apr 13, 2022

Published online: May 16, 2022

Correspondence to

Dong Zhao, MD, PhD

Department of Epidemiology, Beijing Anzhen Hospital, Capital Medical University, Beijing Institute of Heart, Lung and Blood Vessel Diseases, No. 2 Anzhen Road, Chaoyang District, Beijing 100029, China.
Email: deezhao@vip.sina.com

Copyright © 2022. The Korean Society of Cardiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Dong Zhao

<https://orcid.org/0000-0002-3239-3449>

Chen Yao

<https://orcid.org/0000-0003-4224-5535>

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

Conflict of Interest

The authors have no financial conflicts of interest.

Pragmatic Clinical Studies: An Emerging Clinical Research Discipline for Improving Evidence-Based Practice of Cardiovascular Diseases in Asia

Dong Zhao , MD, PhD¹, and Chen Yao , MD²

¹Capital Medical University Beijing Anzhen Hospital-Beijing Institute of Heart, Lung & Blood Vessel Diseases, Beijing, China

²Peking University Clinical Research Institute. Peking University First Hospital, Beijing, China

AUTHOR'S SUMMARY

Pragmatic clinical studies, as an emerging clinical research discipline, include a wide range of studies that are largely embedded with routine clinical practice aiming to evaluate comparative effectiveness and safety of different clinical intervention strategies. In this review, we described the evolution of the conceptual framework of pragmatic clinical studies, shared perspectives on the importance of pragmatic clinical studies for cardiovascular diseases as a complement to conventional randomized controlled trials, as well as highlighted the specific importance of pragmatic clinical research in improving evidence-based practice for cardiovascular disease managements in Asia.

ABSTRACT

Pragmatic clinical studies, an emerging clinical research discipline, include a wide range of studies that are largely embedded with routine clinical practice and aim to evaluate the comparative effectiveness and safety of different clinical intervention strategies. Increased availability and quality of electronic medical/health records drives the development of pragmatic clinical studies. In this review, we describe evolution of the conceptual framework of pragmatic clinical studies and share perspectives on the importance of pragmatic clinical studies in evidence-based practice for cardiovascular diseases, as a complement to conventional randomized controlled trials. We also highlight specific needs of pragmatic clinical studies in improving evidence-based practice for cardiovascular disease in Asian countries. The main challenges of pragmatic clinical studies are discussed briefly in this review.

Keywords: Pragmatic clinical study; Cardiovascular disease; Asia

INTRODUCTION

Cardiovascular disease (CVD) is the leading cause of death in Asia with increasing mortality owing to the combined impact of population aging, increasing epidemics of major risk factors, insufficient implementation of effective strategies in both primary and secondary

Data Sharing Statement

The data generated in this study is available from the corresponding author upon reasonable request.

Author Contributions

Conceptualization: Zhao D, Yao C; Data curation: Zhao D; Formal analysis: Zhao D; Methodology: Zhao D, Yao C; Validation: Zhao D, Yao C; Writing - original draft: Zhao D; Writing - review & editing: Zhao D, Yao C.

prevention and residual risk from emerging risk factors.¹⁾ Dramatic efforts have been made to alleviate the harms to public health caused by CVD with surging activities and substantial progression made in CVD clinical research in many countries and regions of Asia, especially Eastern Asia.²⁾ In recent decade, China, Korea, and Japan have initiated or joined more multicenter randomized controlled trials (RCTs) than in the previous decade. According to the data provided by ClinicalTrials.org of the National Institutes of Health in the United States, among 2305 phase III RCTs conducted in the cardiovascular field from 2011 to 2021, the numbers of RCTs involving China, Korea and Japan were 281, 231 and 201 respectively, approximately two times increase compared with the numbers from 2000 to 2010.³⁾ Pragmatic clinical studies have also increased in these countries, including randomized or nonrandomized pragmatic clinical trials and observational studies (including registry studies) to assess the comparative effectiveness and safety of a wide range of management strategies of major CVDs in clinical practice. In the past 10 years, the importance of pragmatic clinical studies in evidence-based medicine (EBM) practice has been increasingly recognized as a source of evidence that is more applicable to the patients in which the intervention will be applied.⁴⁻⁶⁾ Pragmatic clinical studies for CVD, initiated or conducted jointly by Asian countries have provided important evidence to support decision making for patients in clinical practice.⁷⁻¹⁴⁾

In this review, we aimed to describe the evolution of the conceptual framework of pragmatic clinical studies, share perspectives on the importance of pragmatic clinical studies for CVDs as a complement to conventional RCTs, as well as highlight the specific importance of pragmatic clinical research in improving evidence-based practice for CVD managements in Asia.

EVOLUTION OF THE CONCEPT OF PRAGMATIC CLINICAL STUDIES

The original concept of the pragmatic clinical study in therapeutic trials was proposed by two French statisticians, Schwartz and Lellouch, in a paper published in 1967 entitled “Explanatory and Pragmatic Attitudes in Therapeutical Trials.”¹⁵⁾ This paper was republished in 2009.¹⁶⁾

The authors of this article coined the term “pragmatic” and provided persuasive arguments for the differences between a pragmatic approach and an explanatory approach in therapeutic trials, as well as the importance of a pragmatic approach in clinical decision making for patients in daily practice. The concept was well summarized later by Sackett,¹⁷⁾ as follows: “Pragmatic trials are primarily designed to determine the effects of an intervention under the usual conditions in which it will be applied, whereas explanatory trials are primarily designed to determine the effects of an intervention under ideal circumstances.” A Consolidated Standards of Reporting Trials (CONSORT) document “Improving the reporting of pragmatic trials: an extension of the CONSORT statement.” was published in 2008. This document provided simple comparisons of the characteristics of pragmatic trials and explanatory trials and added eight specific requirements to the reporting of pragmatic trials, based on original CONSORT checklist.¹⁸⁾ Later, the concept of pragmatic study was further developed since differences between pragmatic trials and explanatory trials are multidimensional rather than dichotomy. In 2009, Thorpe et al.¹⁹⁾ developed a tool called PRECIS (Pragmatic–Explanatory Continuum Indicator Summary) to describe multidimensional features of pragmatic trial or explanatory trial. PRECIS was updated to PRECIS II in 2015.²⁰⁾ The PRECIS II provided a more practical tool for the design of pragmatic trial and pragmatic clinical studies. According

Table 1. The features of pragmatic clinical study and factors reduced the extent of pragmatic approach in 9 domains of study design*

Domains	Questions	Very pragmatic approach	Factors reduced the extent of pragmatic approach
Eligibility	To what extent are the participants in this trial similar to those who would receive this intervention if it were part of usual care?	Include anyone with the condition of interest who is likely to be a candidate for the intervention if it was being provided in usual care for this condition.	Eligibility of the patients with the condition of interest is reduced by the extent of inclusion and exclusion criteria.
Recruitment	How much extra effort is made to recruit participants over and above what would be used in the usual care setting to engage with patients?	Recruiting participants through usual appointments or searching EMR or comprehensive registers at a diverse range of clinics or hospitals without overt recruitment effort.	To speed up the recruitment by media advertising or incentives.
Setting	How different are the settings of the trial from the usual care setting?	The setting would be to do the trial in an identical setting to which you intend the results to be applied.	Selecting participating centers.
Organizations	How different are the resources, provider expertise, and the organization of care delivery in the intervention arm of the trial from those available in usual care?	Slot the intervention into the usual organization of care for the condition of interest, making use of no more than the existing healthcare staff and resources in that setting.	Additional requirements on numbers of healthcare providers, training, experience, and specialty certification.
Flexibility in delivery	How different is the flexibility in how the intervention is delivered and the flexibility anticipated in usual care?	To leave the details of how to implement the intervention up to providers like that in usual care.	Specified, protocol driven intervention and monitoring the compliance.
Flexibility in adherence	How different is the flexibility in how participants are monitored and encouraged to adhere to the intervention from flexibility anticipated in usual care?	To allow for full flexibility in how the end user recipients engage with the intervention.	To increase adherence of patients by using strategies of re-screening evaluate, withdraw, and monitor.
Follow-up	How different is the intensity of measurement and follow-up of participants in the trial from the typical follow-up in usual care?	To obtain outcome data from electronic medical records or other usual data to measure mortality or hospital admissions. The trials have no more follow-up than is normal in usual care.	Any additional follow-up than would typically occur under usual care.
Primary outcome	To what extent is the trial's primary outcomes directly relevant to participants?	To select the outcomes that are of obvious importance from the patient's perspective and be relevant to healthcare providers.	To choose outcomes that are less important to patients or participants.
Primary analysis	To what extent are all data included in the analysis of the primary outcome?	To use an intention-to-treat analysis using all available data.	To excluding no-complaint, no-adhered of patients, providers, and sites.

*Extracting from PRECIS II (Pragmatic-Explanatory Continuum Indicator Summary).²⁰⁾

to PRECIS II criteria, trials can be classified into five grades from very explanatory to very pragmatic, using different strategies in nine domains of trial design, including eligibility, recruitment, setting, organizations, flexibility in delivery, flexibility in adherence, follow up, primary outcomes, and primary analysis. **Table 1** provides the features of pragmatic trials in the nine domains.²⁰⁾ Obviously, the conceptual framework of pragmatic trials in PRECIS II covers a wider range of clinical trials designs with a combination of pragmatic and explanatory features, even covering the trials based on observational cohort studies.

One of the underlying drivers of the evolution of pragmatic clinical studies may be the real-world clinical data with extensive coverage for inpatients and outpatients become widely available in electronic medical records (EMR), as well as the increasing feasibility of tracking the long-term outcomes of patients by interlinking medical insurance databases and death registry system databases.²¹⁾²²⁾ More importantly, rapid development of artificial intelligence (AI) in big data science and technology has promoted improvement in clinical data, which would strongly support the advancement of pragmatic clinical studies.²³⁾

With increasing clinical trials using data of EMRs or other electronic health records to identify eligible patients, to deliver an embedded intervention, to identify trial outcomes, or a combination of these purpose, a new checklist in the CONSORT extension for the reporting of RCTs conducted using cohorts and routinely collected data (CONSORT-ROUTINE) was published in 2021.²⁴⁾ This document provides new standards for the reporting of the pragmatic clinical trials conducted using cohorts or routinely collected data.

These documents reflect the evolution of the concept of pragmatic clinical studies, from dichotomous to multidimensional with more comprehensive coverage regarding study types and designs.

THE IMPORTANCE OF PRAGMATIC CLINICAL STUDIES AS A COMPLEMENT TO EXPLANATORY RANDOMIZED CONTROLLED TRIALS IN EVIDENCE-BASED PRACTICE

In the past 30 years, EBM has replaced the old paradigm of empirically based medicine to become the mainstream medical paradigm in health care worldwide.²⁵⁾²⁶⁾ As originally proposed by Dr. Guyatt, the paradigm of EBM referred to practice integrating best evidence, medical expertise, and patients' values.²⁷⁾ The definition of EBM practice was further defined by Sackett et al.²⁸⁾ in 1996 as follows: "The practice of EBM means integrating individual clinical expertise with the best available external clinical evidence from systematic research. By individual clinical expertise we mean the proficiency and judgment that individual clinicians acquire through clinical experience and clinical practice." The original concepts of EBM and EBM practice clearly imply the importance of evidence from pragmatic clinical studies in EBM practice. Indeed, it is not easy to judge where a treatment is suitable for an individual patient without combined evidence from both explanatory RCTs which evaluate the efficacy and safety of an intervention in selected patients under strictly controlled implementation environments, and pragmatic studies, which evaluate effectiveness and safety of an intervention in a largely unselected population of patients that received the intervention in real clinical practice. However, the EBM is often understood as explanatory RCT-EBM. In many countries, including Asian countries, a comprehensive system has been well established to support explanatory RCT-EBM practice. The system includes the following: 1) Regulatory processes for new products for medical treatment: explanatory RCT-based evidence is often a mandatory condition for approving new drugs or new interventional devices; 2) Clinical guidelines: the strength of recommendations for disease managements in clinical practice guidelines is based on the availability of explanatory RCT studies and systematic reviews of these RCTs²⁶⁾²⁹⁾; 3) Quality of care assessment: the utilities of treatment strategies with evidence from RCTs often are used as indicators of performance measures in clinical practice³⁰⁾³¹⁾; 4) Publication: explanatory RCT studies are highly prioritized for acceptance by high quality journals. It is true that high quality randomized double-blind controlled clinical trials can provide an unbiased evaluation of efficacy and safety under experimental conditions. In fact, high-quality explanatory RCT studies are pivotal in the EBM paradigm and are often synonymous with best evidence.

However, there are increasing concerns regarding the limitations of explanatory RCTs.³²⁾³³⁾ The following three core issues may explain the growing demands for pragmatic clinical studies as a complementary source of evidence to guide clinical practice for CVD.

The external validity of explanatory randomized controlled trials for patients with cardiovascular disease in clinical practice

A considerable number of studies and systematic reviews have reported lower representatives and questionable external validity of the evidence from at least some RCTs on interventions for CVD and other diseases.³⁴⁻⁴⁰⁾ One study compared baseline characteristics and outcomes between patients recruited from a clinical registry and patients from 14 RCTs that evaluated

the efficacy and safety of percutaneous coronary intervention (PCI) compared with coronary artery bypass graft. The study found that only 36% of the patients from clinical practice met the inclusion criteria of these RCTs.³⁸⁾ Another study compared the baseline characteristics and clinical outcomes of patients enrolled in a RCT (BIOSCIENCE trial) with patients included in the CARDIOBASE Bern PCI Registry in the same hospital. The study reported that non-participants in the RCT had a higher risk profile, with more women, older peoples, multiple morbidities and adverse prognoses, as compared with RCT participants.³⁹⁾ A large study examined the representativeness of antihypertensive drug trials. Among 110 RCTs on renin-angiotensin aldosterone system (RAAS) drugs, 90% (99 RCTs) of the trials recruited patients under 60-year-old, and 10% (11 RCTs) recruited older patients, defined as those aged ≥ 60 years. The incidence rates of serious adverse events (SAEs) in both the RCTs including older patients, and RCTs without older patients were compared with the incidence rates of SAE in a large number of patients with hypertension using RAAS drugs in real clinical practice by age and sex. Community-dwelling patients with hypertension had a significantly higher risk of SAE than patients in the above two types of RCT. The standardized ratio of SAE, after adjusting for age and sex, was 4.76 times higher than the RCTs including older-patients and 4.23 times higher than the RCTs with younger patients. The authors of that paper stated that the study results suggest “even accounting for age and sex, participants in hypertension trials and people with hypertension in the community are very different populations.”⁴⁰⁾ A recently published systematic review further addressed this issue. That systematic review reported estimated exclusion rates of clinical patients from 305 trials of treatments for 31 physical conditions. The median total exclusion rate was 77.1% of patients. Median exclusion rates for trials of common chronic conditions were high, including for hypertension (83.0%), type 2 diabetes (81.7%), chronic obstructive pulmonary disease (84.3%), and asthma (96.0%). The most frequently applied exclusion criteria were related to age, co-morbidities, and co-prescribing.⁴¹⁾

Evidence gaps for clinical questions in cardiovascular disease management

Several studies have revealed large evidence gaps for major clinical questions in CVD. One study examined the levels of evidence for 6,329 recommendations in 51 current guidelines of the American College of Cardiology/American Heart Association (ACC/AHA) and European Society of Cardiology (ESC). Among 2,930 recommendations in ACC/AHA guidelines, 8.5% was classified as level A, 50% level B and 41.5% level C evidence. In 3,399 recommendations of ESC guidelines, 14.2% was classified as level A, 31.0% level B, and 54.8% level C evidence.⁴⁰⁾ The study compared the current guidelines with prior versions; the proportions with level A evidence showed no increase in either ACC/AHA (median, 9.0% [current] vs. 11.7% [prior]) or ESC guidelines (median, 15.1% [current] vs. 17.6% [prior]). The authors of that study concluded that “among recommendations in major cardiovascular society guidelines, only a small percentage were supported by evidence from multiple RCTs or a single, large RCT. This pattern does not appear to have meaningfully improved from 2008 to 2018.”⁴²⁾ Other studies have presented similar results regarding the quality of evidence in the guidelines for atrial fibrillation.⁴³⁾ The authors of these studies appealed for the development and implementation of inexpensive methods to generate a large volume of evidence. These studies highlight the need for pragmatic clinical studies evaluating the comparative effectiveness of existing treatments, and novel methods of conducting clinical trials with less waste.

The quality of evidence from pragmatic clinical studies

It is a key question whether pragmatic clinical studies can provide reliable evidence or evidence comparable to explanatory RCTs. A good example of a pragmatic randomized clinical

study for CVD intervention is TASTE (The Thrombus Aspiration in ST-Elevation Myocardial Infarction in Scandinavia) trial. This study aimed to evaluate whether thrombus aspiration reduces mortality. The results of that registry-based clinical trial were published in *The New England Journal of Medicine* in 2013.⁴⁴⁾ The TASTE trial provided an example of combining the benefits of randomized treatment assignment with the best features of a large-scale clinical registry. This RCT/pragmatic clinical trial demonstrated some of the most important advantages of pragmatic trials, namely broad inclusion criteria to ensure wide clinical applicability, a simplified enrollment process to maximize the commitment and compliance of participating hospitals, and a substantial reduction in the expense associated with conducting a RCT. The study results have become important evidence in relevant guidelines.⁴⁵⁾

The important RCT DUPLICATE initiative has been launched to emulate RCT with nonrandomized real-world study (RWS). The first report was published in 2021 on the first 10 trial emulations, evaluating CVD outcomes of anti-diabetes or antiplatelet medications.⁴⁶⁾ The researchers used patient-level claims data with the same inclusion and exclusion criteria. The intervention group and control group were matched using propensity scores to control for more than 120 pre-exposure confounders. The study found that the regulatory conclusions were equivalent in 6/10 trials; the effectiveness estimation, indicated using the hazard ratio (HR) was within the 95% confidence intervals of HRs from the corresponding RCT in 8/10 studies.⁴⁶⁾ Whereas the results of that study do not mean that an observational study based on routine clinical data can replace a RCT, the findings at least provide an evaluation of comparability and differences in the effects and safety of certain intervention between pragmatic clinical studies and RCTs.

The importance of pragmatic clinical studies in evidence-based practice for CVD is obvious and may increase with further development of AI technology. It has become possible to evaluate and provide feedback more comprehensively and relatively rapidly using more pragmatic approach for issues of effectiveness and safety in clinical practice for any treatment or diagnostic strategy to support judgments with expertise and the patients' values. Furthermore, the evidence produced using routine clinical data denoted real world evidence has greater extrapolations and lower costs and has been accepted by regulatory agencies of medical products in some countries, including in Asia.⁴⁷⁻⁴⁹⁾ It is critical to emphasize that the evidence from pragmatic clinical studies is not dispensable but essential to guide clinical practice.

SPECIFIC NEEDS FOR PRAGMATIC CLINICAL STUDIES OF CARDIOVASCULAR DISEASE IN ASIA

Among all deaths from CVD, 58% of CVD deaths have occurred in Asia, with specific epidemic characteristics regarding the subtypes of CVD. Asia has a very large number of patients with one or more CVD risk factors, such as hypertension, dyslipidemia, diabetes, and established CVDs, including stroke and coronary heart disease (CHD), heart failure and atrial fibrillation (AF).¹⁾⁵⁰⁾ Most patients with CVDs or risk factors require clinical treatments. Unfortunately, most RCTs in the field of CVD have included no or few patients from Asia.

Table 2 shows a comparison of the numbers of completed RCTs globally classified by the types of CVDs or risk factors and treatment strategies, with the corresponding numbers of RCTs joined by regions of Asia.³⁾ For example, among 177 RCTs for PCI in patients with CHD, 33 (19%) included patients from East Asia, 14 (8%) included patients from South-east Asia

and 9 (5%) included patients from South Asia; Among 69 RCTs for anticoagulant treatments in patients with AF, 24 (35%) included patients from East Asia, 11 (16%) included patients from South-east Asia and 9 (13%) included patients from South Asia.

However, available evidence shows that Asian patients have specific effectiveness and safety issues with certain medical treatments recommended by clinical guidelines of the US or Europe based on evidence of RCTs with few or no Asian patients.

A series of expert consensus, position statements and reviews have highlighted the differences in anti-ischemic effectiveness and bleeding risk of antithrombotic therapies between East Asian patients and White patients.⁵¹⁻⁵⁴⁾ The first expert consensus on this issue was the World Heart Federation (WHF) expert consensus statement on antiplatelet therapy in East Asian patients with ACS or undergoing PCI. This document, published in 2014, clearly indicated that the recommendations on antiplatelet therapy of the ACC/AHA and ESC guidelines were based primarily on large, phase III, randomized, controlled trials of the P2Y₁₂ inhibitors clopidogrel, prasugrel, and ticagrelor. However, few East Asian patients were included in the trials conducted to assess the use of these agents. East Asian patients have differing risk profiles for both thrombophilia and bleeding compared with White patients, and a different 'therapeutic window' of on-treatment platelet reactivity might be appropriate in East Asian patients. The WHF produced this expert consensus statement to determine the antiplatelet treatment strategies that are most appropriate for East Asian patients.⁵¹⁾ An updated expert consensus was published in 2020, in which, the issues regarding East Asian patients having reduced anti-ischemic benefits and increased bleeding risk with antithrombotic therapies as well as relevant strategies for East Asian patients, were further addressed.⁵³⁾

One review by Cho well summarized the challenges regarding Asian patients with nonvalvular atrial fibrillation (NVAf) of using evidence from RCTs on non-vitamin K oral anticoagulants (NOACs) that were conducted among on predominantly non-Asian patients with NVAf.⁵⁵⁾ According to that review, the FDA emphasized the results of the RE-LY study showing that the dabigatran 150-mg dose was superior to warfarin for efficacy and similar for bleeding. But two Asian studies found that a fixed dose of 110 mg might be sufficient based on the

Table 2. Comparison of the numbers of completed RCTs globally with the numbers of RCTs jointed by Asian regions*

Diseases and treatments	Global	East Asia	South Asia	South-East Asia
Clinical trials for coronary heart disease treatment				
PCI	177	33 (19)	9 (5)	14 (8)
Antiplatelet drugs	101	20 (20)	6 (6)	9 (9)
Clinical trials for heart failure treatment				
All	547	57 (10)	23 (4)	24 (4)
Clinical trials of atrial fibrillation treatment				
Anticoagulants	69	24 (35)	9 (13)	11 (16)
Ablation	132	10 (8)	0	0
Clinical trials of hypertension treatment				
All	1,149	156 (14)	31 (3)	51 (4)
Clinical trials of dyslipidemia treatment				
Statins	225	38 (17)	7 (3)	8 (4)
PCSK9 inhibitors	10	3 (30)	0	0
Clinical trials of diabetes treatment				
SGLT-2 inhibitor	209	68 (33)	32 (15)	29 (14)
GLP-1 agonist	27	7 (26)	4 (15)	4 (15)

Values are presented as number (%).

RCT = randomized controlled trial.

*Data are obtained from ClinicalTrial.org of the National Institutes of Health in the United States. <https://clinicaltrial.gov>.³⁾

comparable efficacy and favorable safety to 150-mg dose. Whereas the optimal doses of other agents NOACs in Asian population are uncertain, several real-world studies have provided pragmatic evidence that a reduced dose is associated with a significantly lower risk of stroke, major bleeding, intracranial hemorrhage, and mortality. Pharmacokinetic studies in Japan estimated that a 15 mg dose of rivaroxaban in Japanese patients was pharmacokinetically equivalent to a 20-mg dose in White patients.⁵⁶⁾ Another review reported that intracranial hemorrhage is a major concern in selection and dosing of NOACs among Asians because the risk of intracranial hemorrhage, the most devastating complication, was higher in Asians than in non-Asians and greater in patients with prior stroke/transient ischemic attack than in those without.⁵⁷⁾

Some studies have also found high effectiveness and a high risk of adverse effects with statins in Asian patients.⁵⁸⁻⁵⁹⁾ Therefore, only medium dose statins are recommended for CVD patients or patients with high CVD risk in the clinical guidelines issued by some Asian countries, such as China.⁶⁰⁾

The cost-effectiveness of a treatment is critically important for many low- and middle-income countries. Asian countries need to conduct more comparative effectiveness studies and cost effectiveness studies based on EMR to help in decision making regarding clinical applications of old medications or new treatment strategies.⁶¹⁾

Pragmatic clinical studies can be an affordable and useful way for Asian countries to re-evaluate the evidence from RCTs with no or few Asian patients. In recent years, many pragmatic clinical studies on important issues in CVDs conducted in Asian countries have been published.⁹⁻¹⁴⁾

CHALLENGES OF PRAGMATIC CLINICAL STUDIES IN ASIA

Compared with the well-established theoretical and practice system of explanatory RCTs over the past nearly 70 years, pragmatic clinical studies remain a developing discipline with an incomplete theoretical and practice system.⁶²⁾

Unclear scope and terminology

Just like at the beginning of any new sub-discipline in clinical research, the scope of the concept of pragmatic clinical studies is not well defined. Some researchers have considered pragmatic clinical studies to only include pragmatic RCTs, whereas others consider that pragmatic clinical studies also cover registry studies, as well as prospective or retrospective patient cohort studies in a clinical setting.⁶⁾²⁴⁾⁶³⁾ Some reports have used the term RWS as an umbrella term covering pragmatic clinical studies.⁶⁾ There is also some overlap in the scope of the conceptual framework between pragmatic clinical study and outcome studies, and comparative effectiveness research. It is important to conduct more systematic carding to clarify the current knowledge and rationality regarding the scope of pragmatic clinical studies, and develop unified definition, classification, and terminology for this emerging subdiscipline of clinical research.

Accessibility and quality of electronic medical records

Some common problematic issues and barriers exist to the use of EMR in pragmatic clinical research. One paper summarized all large EMR databases in China. However, the accessibility

of these EMR databases was very low. Additionally, the availability of information necessary for clinical research is much lower than that in the United States and Europe.⁶⁴⁾ Other barriers to the use of EMR in pragmatic clinical studies include funding, ethical issues, and data quality including accuracy, completeness, and consistency.⁶³⁾⁶⁴⁾

Residual confounding effects in observational cohort studies

Whereas some pragmatic clinical studies use randomization methods to assign participants to an interventional group and control group, many pragmatic clinical studies used non-randomization or observational approaches to assess the comparative effects of an intervention strategy. When patients in the intervention and control groups are not assigned randomly, one main issue is how to ensure comparability between those patients that receive an intervention and those who did not. The risk of bias is sometime high for non-randomized pragmatic studies because the observed effects or the safety of an intervention may be affected by confounding factors.⁶⁵⁾ Certain methods are recommended to increase the comparability in observational cohort trials, including propensity score matching, inverse probability weighting and using instrumental variables et al.⁶⁶⁻⁶⁸⁾

CONCLUSIONS

Pragmatic clinical studies, as an emerging clinical research discipline, include a wide range of studies that are largely embedded with routine clinical practice aiming to evaluate comparative effectiveness and safety of different clinical intervention strategies. To increase awareness of the importance of pragmatic clinical studies in EBM practice for CVD in Asia, we described evolution of the conceptual framework of pragmatic clinical studies, from a dichotomous “pragmatic or explanatory” concept to an emerging clinical research discipline with multidimensional features and a comprehensive scope. A considerable number of studies support pragmatic clinical studies as an indispensable source of evidence in EBM practice because the evidence is more applicable to the patients in which the intervention will be applied. It is particularly important to increase the number of pragmatic clinical studies on CVD conducted in Asia because most of available evidence from explanatory RCTs has included no or few Asian patients. However, the available evidence shows there are considerable racial differences in the effects and safety of certain interventions between Asian and non-Asian patients. Clearly, it is not feasible to identify potential racial differences by repeating all explanatory RCTs that included no or few Asian patients in re-conducted RCTs among Asian population. Furthermore, as a continent mainly comprising low- and middle-income countries, Asia needs more cost-effective studies based on pragmatic clinical studies. Despite the many challenges and barriers, we expect that pragmatic clinical studies will continuously develop and become a mature sub-discipline of clinical research in Asia and worldwide.

ACKNOWLEDGMENTS

We thank Liwen Bianji (Edanz) (www.liwenbianji.cn) for editing the language of a draft of this manuscript.

REFERENCES

1. Zhao D. Epidemiology features of cardiovascular disease in Asia. *JACC-Asia*. 2021;1:1-13.
[CROSSREF](#)
2. Liu J, Ma C. Current state of cardiovascular research in China. *Nat Rev Cardiol* 2019;16:575-6.
[PUBMED](#) | [CROSSREF](#)
3. U.S. National Library of Medicine. ClinicalTrials.gov [Internet]. Bethesda (MD): U.S. National Library of Medicine; 2022 [cited 2022 February 15]. Available from: <https://www.clinicaltrials.gov/>.
4. Ford I, Norrie J. Pragmatic trials. *N Engl J Med* 2016;375:454-63.
[PUBMED](#) | [CROSSREF](#)
5. Jones WS, Roe MT, Antman EM, et al. The changing landscape of randomized clinical trials in cardiovascular disease. *J Am Coll Cardiol* 2016;68:1898-907.
[PUBMED](#) | [CROSSREF](#)
6. Zhou Q, Dong CY, Wang YM, Zhang BH, Cheng YL, Yao C. The role of evidence-based medicine in guiding clinical research and practice expert opinion based on studies on antiviral drugs for the treatment of COVID-19. *Chin J Evid Based Med* 2022;22:373-379.
7. You SC, Rho Y, Bikdeli B, et al. Association of ticagrelor vs clopidogrel with net adverse clinical events in patients with acute coronary syndrome undergoing percutaneous coronary intervention. *JAMA* 2020;324:1640-50.
[PUBMED](#) | [CROSSREF](#)
8. Anderson CS, Arima H, Lavados P, et al. Cluster-randomized, crossover trial of head positioning in acute stroke. *N Engl J Med* 2017;376:2437-47.
[PUBMED](#) | [CROSSREF](#)
9. Zhou M, Zhang J, Liu J, et al. Proton pump inhibitors and in-hospital gastrointestinal bleeding in patients with acute coronary syndrome receiving dual antiplatelet therapy. *Mayo Clin Proc* 2022;97:682-92.
[PUBMED](#) | [CROSSREF](#)
10. Shoji S, Sawano M, Sandhu AT, et al. Ischemic and bleeding events among patients with acute coronary syndrome associated with low-dose prasugrel vs standard-dose clopidogrel treatment. *JAMA Netw Open* 2020;3:e202004.
[PUBMED](#) | [CROSSREF](#)
11. Nakamura M, Kadota K, Takahashi A, et al. Relationship between platelet reactivity and ischemic and bleeding events after percutaneous coronary intervention in East Asian patients: 1-year results of the PENDULUM registry. *J Am Heart Assoc* 2020;9:e015439.
[PUBMED](#) | [CROSSREF](#)
12. Zheng B, Huo Y, Lee SW, et al. Long-term antithrombotic management patterns in Asian patients with acute coronary syndrome: 2-year observations from the EPICOR Asia study. *Clin Cardiol* 2020;43:999-1008.
[PUBMED](#) | [CROSSREF](#)
13. Yoon YH, Ahn JM, Kang DY, et al. Pragmatic trial comparing routine versus no routine functional testing in high-risk patients who underwent percutaneous coronary intervention: rationale and design of POST-PCI trial. *Am Heart J* 2020;224:156-65.
[PUBMED](#) | [CROSSREF](#)
14. Zheng Y, Reinhardt JD, Li J, et al. Can clinical and functional outcomes be improved with an intelligent "internet plus"-based full disease cycle remote ischemic conditioning program in acute ST-elevation myocardial infarction patients undergoing percutaneous coronary intervention? Rationale and design of the i-RIC trial. *Cardiovasc Drugs Ther* 2022;36:45-57.
[PUBMED](#) | [CROSSREF](#)
15. Schwartz D, Lellouch J. Explanatory and pragmatic attitudes in therapeutical trials. *J Chronic Dis* 1967;20:637-48.
[PUBMED](#) | [CROSSREF](#)
16. Schwartz D, Lellouch J. Explanatory and pragmatic attitudes in therapeutical trials. *J Clin Epidemiol* 2009;62:499-505.
[PUBMED](#) | [CROSSREF](#)
17. Sackett DL. *Clinical epidemiology: how to do clinical practice research*. Philadelphia (PA): Lippincott, Williams and Wilkins; 2006.
18. Zwarenstein M, Treweek S, Gagnier JJ, et al. Improving the reporting of pragmatic trials: an extension of the CONSORT statement. *BMJ* 2008;337:a2390.
[PUBMED](#) | [CROSSREF](#)
19. Thorpe KE, Zwarenstein M, Oxman AD, et al. A pragmatic-explanatory continuum indicator summary (PRECIS): a tool to help trial designers. *J Clin Epidemiol* 2009;62:464-75.
[PUBMED](#) | [CROSSREF](#)

20. Loudon K, Treweek S, Sullivan F, Donnan P, Thorpe KE, Zwarenstein M. The PRECIS-2 tool: designing trials that are fit for purpose. *BMJ* 2015;350:h2147.
[PUBMED](#) | [CROSSREF](#)
21. Kwakkenbos L, Imran M, McCall SJ, et al. CONSORT extension for the reporting of randomised controlled trials conducted using cohorts and routinely collected data (CONSORT-ROUTINE): checklist with explanation and elaboration. *BMJ* 2021;373:n857.
[PUBMED](#) | [CROSSREF](#)
22. Evans RS. Electronic health records: then, now, and in the future. *Yearb Med Inform* 2016;25 Suppl 1:S48-61.
[PUBMED](#) | [CROSSREF](#)
23. Hemingway H, Asselbergs FW, Danesh J, et al. Big data from electronic health records for early and late translational cardiovascular research: challenges and potential. *Eur Heart J* 2018;39:1481-95.
[PUBMED](#) | [CROSSREF](#)
24. Antman EM, Bierer BE. Standards for clinical research: keeping pace with the technology of the future. *Circulation* 2016;133:823-5.
[PUBMED](#) | [CROSSREF](#)
25. Djulbegovic B, Guyatt GH. Progress in evidence-based medicine: a quarter century on. *Lancet* 2017;390:415-23.
[PUBMED](#) | [CROSSREF](#)
26. Tricoci P, Allen JM, Kramer JM, Califf RM, Smith SC Jr. Scientific evidence underlying the ACC/AHA clinical practice guidelines. *JAMA* 2009;301:831-41.
[PUBMED](#) | [CROSSREF](#)
27. Guyatt G. Evidence-based medicine. *ACP J Club* 1991;114:A16.
28. Sackett DL, Rosenberg WM, Gray JA, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. *BMJ* 1996;312:71-2.
[PUBMED](#) | [CROSSREF](#)
29. Fanaroff AC, Fudim M, Califf RM, Windecker S, Smith SC Jr, Lopes RD. Levels of evidence supporting drug, device, and other recommendations in the American Heart Association/American College of Cardiology guidelines. *Am Heart J* 2020;226:4-12.
[PUBMED](#) | [CROSSREF](#)
30. Jneid H, Addison D, Bhatt DL, et al. 2017 AHA/ACC clinical performance and quality measures for adults with ST-elevation and non-ST-elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on performance measures. *J Am Coll Cardiol* 2017;70:2048-90.
[PUBMED](#) | [CROSSREF](#)
31. Schiele F, Aktaa S, Rossello X, et al. 2020 Update of the quality indicators for acute myocardial infarction: a position paper of the Association for Acute Cardiovascular Care: the study group for quality indicators from the ACVC and the NSTEMI-ACS guideline group. *Eur Heart J Acute Cardiovasc Care* 2021;10:224-33.
[PUBMED](#) | [CROSSREF](#)
32. Lauer MS, Gordon D, Wei G, Pearson G. Efficient design of clinical trials and epidemiological research: is it possible? *Nat Rev Cardiol* 2017;14:493-501.
[PUBMED](#) | [CROSSREF](#)
33. Kostis JB, Dobrzynski JM. Limitations of randomized clinical trials. *Am J Cardiol* 2020;129:109-15.
[PUBMED](#) | [CROSSREF](#)
34. Laursen PN, Holmvang L, Lønborg J, et al. Comparison between patients included in randomized controlled trials of ischemic heart disease and real-world data. A nationwide study. *Am Heart J* 2018;204:128-38.
[PUBMED](#) | [CROSSREF](#)
35. Smyth B, Haber A, Trongtrakul K, et al. Representativeness of randomized clinical trial cohorts in end-stage kidney disease: a meta-analysis. *JAMA Intern Med* 2019;179:1316-24.
[PUBMED](#) | [CROSSREF](#)
36. Jackson MW, Austin D, Swanson N, et al. Outcomes after culprit-only percutaneous coronary intervention for multivessel disease during ST-segment elevation myocardial infarction: a comparison of registry and clinical trial outcomes. *Coron Artery Dis* 2018;29:564-72.
[PUBMED](#) | [CROSSREF](#)
37. Megaly M, Buda K, Alaswad K, et al. Comparative analysis of patient characteristics in cardiogenic shock studies: differences between trials and registries. *JACC Cardiovasc Interv* 2022;15:297-304.
[PUBMED](#) | [CROSSREF](#)
38. Hordijk-Trion M, Lenzen M, Wijns W, et al. Patients enrolled in coronary intervention trials are not representative of patients in clinical practice: results from the Euro Heart Survey on Coronary Revascularization. *Eur Heart J* 2006;27:671-8.
[PUBMED](#) | [CROSSREF](#)

39. Franzone A, Heg D, Räber L, et al. External validity of the “all-comers” design: insights from the BIOSCIENCE trial. *Clin Res Cardiol* 2016;105:744-54.
[PUBMED](#) | [CROSSREF](#)
40. Hanlon P, Corcoran N, Rughani G, et al. Observed and expected serious adverse event rates in randomised clinical trials for hypertension: an observational study comparing trials that do and do not focus on older people. *Lancet Healthy Longev* 2021;2:e398-406.
[PUBMED](#) | [CROSSREF](#)
41. He J, Morales DR, Guthrie B. Exclusion rates in randomized controlled trials of treatments for physical conditions: a systematic review. *Trials* 2020;21:228.
[PUBMED](#) | [CROSSREF](#)
42. Fanaroff AC, Califf RM, Windecker S, Smith SC Jr, Lopes RD. Levels of evidence supporting American College of Cardiology/American Heart Association and European Society of Cardiology Guidelines, 2008–2018. *JAMA* 2019;321:1069-80.
[PUBMED](#) | [CROSSREF](#)
43. Barnett AS, Lewis WR, Field ME, et al. Quality of evidence underlying the American Heart Association/American College of Cardiology/Heart Rhythm Society Guidelines on the management of atrial fibrillation. *JAMA Cardiol* 2017;2:319-23.
[PUBMED](#) | [CROSSREF](#)
44. Fröbert O, Lagerqvist B, Olivecrona GK, et al. Thrombus aspiration during ST-segment elevation myocardial infarction. *N Engl J Med* 2013;369:1587-97.
[PUBMED](#) | [CROSSREF](#)
45. Windecker S, Kolh P, Alfonso F, et al. 2014 ESC/EACTS Guidelines on myocardial revascularization: The Task Force on Myocardial Revascularization of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS) developed with the special contribution of the European Association of Percutaneous Cardiovascular Interventions (EAPCI). *Eur Heart J* 2014;35:2541-619.
[PUBMED](#) | [CROSSREF](#)
46. Franklin JM, Patorno E, Desai RJ, et al. Emulating randomized clinical trials with nonrandomized real-world evidence studies: first results from the RCT DUPLICATE initiative. *Circulation* 2021;143:1002-13.
[PUBMED](#) | [CROSSREF](#)
47. Baumfeld Andre E, Reynolds R, Caubel P, Azoulay L, Dreyer NA. Trial designs using real-world data: the changing landscape of the regulatory approval process. *Pharmacoepidemiol Drug Saf* 2020;29:1201-12.
[PUBMED](#) | [CROSSREF](#)
48. Li M, Chen S, Lai Y, et al. Integrating real-world evidence in the regulatory decision-making process: a systematic analysis of experiences in the US, EU, and China using a logic model. *Front Med (Lausanne)* 2021;8:669509.
[PUBMED](#) | [CROSSREF](#)
49. Nishioka K, Makimura T, Ishiguro A, Nonaka T, Yamaguchi M, Uyama Y. Evolving acceptance and use of RWE for regulatory decision making on the benefit/risk assessment of a drug in Japan. *Clin Pharmacol Ther* 2022;111:35-43.
[PUBMED](#) | [CROSSREF](#)
50. Chiang CE, Wang KL, Lip GY. Stroke prevention in atrial fibrillation: an Asian perspective. *Thromb Haemost* 2014;111:789-97.
[PUBMED](#) | [CROSSREF](#)
51. Levine GN, Jeong YH, Goto S, et al. Expert consensus document: World Heart Federation expert consensus statement on antiplatelet therapy in East Asian patients with ACS or undergoing PCI. *Nat Rev Cardiol* 2014;11:597-606.
[PUBMED](#) | [CROSSREF](#)
52. Huo Y, Jeong YH, Gong Y, et al. 2018 update of expert consensus statement on antiplatelet therapy in East Asian patients with ACS or undergoing PCI. *Sci Bull (Beijing)* 2019;64:166-79.
[CROSSREF](#)
53. Kim HK, Tantry US, Smith SC Jr, et al. The East Asian paradox: an updated position statement on the challenges to the current antithrombotic strategy in patients with cardiovascular disease. *Thromb Haemost* 2021;121:422-32.
[PUBMED](#) | [CROSSREF](#)
54. Kang J, Kim HS. The evolving concept of dual antiplatelet therapy after percutaneous coronary intervention: focus on unique feature of East Asian and “Asian Paradox”. *Korean Circ J* 2018;48:537-51.
[PUBMED](#) | [CROSSREF](#)
55. Cho IY. Using non-vitamin K oral anticoagulants in specific patient populations: a study of Korean cases. *Ther Clin Risk Manag* 2019;15:1183-206.
[PUBMED](#) | [CROSSREF](#)

56. Cho IY, Choi KH, Sheen YY. How does “Regulatory Practice” create discrepancies in drug label information between Asian and Western countries? Different label information for direct oral anticoagulants approved in the United States, Europe, Korea, and Japan. *Ther Innov Regul Sci* 2019;53:233-42.
[PUBMED](#) | [CROSSREF](#)
57. Bang OY, Hong KS, Heo JH. Asian patients with stroke plus atrial fibrillation and the dose of non-vitamin K oral anticoagulants. *J Stroke* 2016;18:169-78.
[PUBMED](#) | [CROSSREF](#)
58. HPS2-THRIVE Collaborative Group. HPS2-THRIVE randomized placebo-controlled trial in 25 673 high-risk patients of ER niacin/laropiprant: trial design, pre-specified muscle and liver outcomes, and reasons for stopping study treatment. *Eur Heart J* 2013;34:1279-91.
[PUBMED](#) | [CROSSREF](#)
59. Lin JL, Chen PS, Lin HW, Tsai LM, Lin SH, Li YH. Real-world analyses of the safety outcome among a general population treated with statins: an Asian population-based study. *J Atheroscler Thromb* 2021;28:63076.
[PUBMED](#) | [CROSSREF](#)
60. Joint Committee for Guideline Revision. 2016 Chinese guidelines for the management of dyslipidemia in adults. *J Geriatr Cardiol* 2018;15:1-29.
[PUBMED](#)
61. Daccache C, Rizk R, Dahham J, Evers SM, Hiligsmann M, Karam R. Economic evaluation guidelines in low- and middle-income countries: a systematic review. *Int J Technol Assess Health Care* 2021;38:e1.
[PUBMED](#) | [CROSSREF](#)
62. French C, Pinnock H, Forbes G, Skene I, Taylor SJ. Process evaluation within pragmatic randomised controlled trials: what is it, why is it done, and can we find it? - a systematic review. *Trials* 2020;21:916.
[PUBMED](#) | [CROSSREF](#)
63. Sun X, Tan J, Tang L, Guo JJ, Li X. Real world evidence: experience and lessons from China. *BMJ* 2018;360:j5262.
[PUBMED](#) | [CROSSREF](#)
64. Xie J, Wu EQ, Wang S, et al. Real-world data for healthcare research in China: call for actions. *Value Health Reg Issues* 2022;27:72-81.
[PUBMED](#) | [CROSSREF](#)
65. Bosdriesz JR, Stel VS, van Diepen M, et al. Evidence-based medicine-When observational studies are better than randomized controlled trials. *Nephrology (Carlton)* 2020;25:737-43.
[PUBMED](#) | [CROSSREF](#)
66. Benedetto U, Head SJ, Angelini GD, Blackstone EH. Statistical primer: propensity score matching and its alternatives. *Eur J Cardiothorac Surg* 2018;53:1112-7.
[PUBMED](#) | [CROSSREF](#)
67. Mansournia MA, Altman DG. Inverse probability weighting. *BMJ* 2016;352:i189.
[PUBMED](#) | [CROSSREF](#)
68. Lousdal ML. An introduction to instrumental variable assumptions, validation and estimation. *Emerg Themes Epidemiol* 2018;15:1.
[PUBMED](#) | [CROSSREF](#)