



DRPLA: understanding the natural history and developing biomarkers to accelerate therapeutic trials in a globally rare repeat expansion disorder

Aiysha Chaudhry¹ · Alkyoni Anthanasiou-Fragkouli¹ · Henry Houlden¹

Received: 14 August 2020 / Revised: 2 September 2020 / Accepted: 3 September 2020 / Published online: 26 October 2020
© The Author(s) 2020

Abstract

Dentatorubral–pallidoluysian atrophy (DRPLA) is a rare neurodegenerative disorder caused by CAG repeat expansions in the atrophin-1 gene and is inherited in an autosomal dominant fashion. There are currently no disease-modifying treatments available. The broad development of therapies for DRPLA, as well as other similar rare diseases, has hit a roadblock due to the rarity of the condition and the wide global distribution of patients and families, consequently inhibiting biomarker development and therapeutic research. Considering the shifting focus towards diverse populations, widespread genetic testing, rapid advancements in the development of clinical and wet biomarkers for Huntington’s disease (HD), and the ongoing clinical trials for antisense oligonucleotide (ASO) therapies, the prospect of developing effective treatments in rare disorders has completely changed. The awareness of the HD ASO program has prompted global collaboration for rare disorders in natural history studies and the development of biomarkers, with the eventual goal of undergoing treatment trials. Here, we discuss DRPLA, which shares similarities with HD, and how in this and other repeat expansion disorders, neurogenetics groups like ours at UCL are gearing up for forthcoming natural history studies to accelerate future ASO treatment trials to hopefully emulate the progress seen in HD.

Current understanding of DRPLA

Dentatorubral–pallidoluysian atrophy (DRPLA) is a rare autosomal dominant neurodegenerative disorder, characterized by progressive cerebellar ataxia, myoclonus, epilepsy, dementia, choreoathetosis, and psychiatric symptoms [90]. The condition was first described by Titica and van Boegard in 1946, whereby two cases in a family with progressive choreoathetosis, ataxia, and dementia were reported [88]. The term “hereditary DRPLA” was later coined by Naito and Oyanagi in 1982 [61]. DRPLA is classified within the spinocerebellar ataxia (SCA) group, which represents a heterogeneous group of > 40 autosomal dominantly inherited diseases [44]. DRPLA is caused by a CAG-polyglutamine (polyQ) repeat expansion. Nine such polyQ diseases have currently been identified in humans, including Huntington’s

disease (HD), spinal and bulbar muscular atrophy (SBMA), SCA 1, 2, 3, 6, 7, and 17 [82].

DRPLA is the result of an unstable CAG repeat expansion in exon 5 of the atrophin-1 (*ATN1*) gene [46, 60]. The number of repeats in normal individual chromosomes ranges typically between 6 and 35. Full penetrance occurs at ≥ 48 CAG repeats, whilst alleles of 35–47 repeats are incompletely penetrant and are usually associated with a milder clinical phenotype [13, 37, 38, 46, 55, 60]. Characterized by genetic anticipation, with paternal transmission resulting in more prominent anticipation than maternal transmission, DRPLA symptoms present more severely and earlier in each subsequent generation [56, 90]. The CAG repeat load is also associated with the phenotype, whereby the longer the size of expanded CAG repeats, the earlier the age of onset and death, the more severe the symptoms and long-term disability, and the poorer the prognosis [34, 37, 55]. Figure 1 illustrates the currently known features of DRPLA.

Due to the heterogeneity in clinical presentation, based on the prominent genetic anticipation and age of onset, diagnosing DRPLA can often be challenging, with symptoms associated with a broad differential diagnosis. Whilst epileptic seizures are common in juvenile-onset patients (onset prior

✉ Henry Houlden
h.houlden@ucl.ac.uk

¹ Department of Neuromuscular Disorders, UCL Institute of Neurology, Queen Square, London WC1N 3BG, UK




Fig. 1 Main genetic, clinical, neuroimaging and neuropathological features of DRPLA. **a** Illustration of location of the trinucleotide repeat within the gene—In DRPLA, the CAG trinucleotide repeat expansion occurs in coding exon 5 of the atrophin-1 (*ATN1*) gene, which is located on chromosome 12p13.31. Normal alleles in the *ATN1* gene have 6–35 CAG repeats. ≥ 20 repeats are considered normal mutable alleles that expand on transmission and result in symptoms in the next generation, and ≥ 48 repeats demonstrate fully pen-

etrant clinical phenotype. The unstable CAG repeat sequence causes a polyglutamine (polyQ) expansion in the atrophin-1 protein [13, 25, 44, 55, 60, 68, 94]. **b** Clinical features—The primary clinical features of DRPLA are ataxia and cognitive impairment, however, the age of onset affects the clinical presentation, with different symptoms observed between adult-onset and juvenile-onset DRPLA (Wardle et al. [98]; [34, 48, 55]. **c** Neuroimaging findings—[47, 106]. **d** Neuropathological findings—[36, 84, 106, 107]

to the age of 20), the frequency of seizures is reduced after the age of 20, and rare in patients with an onset after the age of 40. Patients with an onset after the age of 20 tend to present with cerebellar ataxia, choreoathetosis and dementia, often making the disease difficult to differentiate from clinical mimics including HD and other hereditary SCAs [13, 61]. Further, brain MRI findings in DRPLA are variable, with case reports of early-stage patients often presenting with only mild changes, whilst late stages of the disease are associated with non-specific changes such as atrophy of the cerebellum and brainstem, complicating the differentiation of the disease from other neurological disorders [35, 45, 77, 83, 92].

Global burden

Defining global burden through natural history studies is important to understand the impact of condition and to identify disease biomarkers in the preparation for therapeutic trials. DRPLA is most commonly recognised in populations of Japanese ancestry and has an estimated incidence in Japan of 2–7 per million [26, 71]. DRPLA is considered to be the third most common autosomal dominant ataxia in the Japanese population, accounting for approximately 7.3–20% of autosomal dominant SCA [54, 87, 91]. Whilst it is believed to be rare in non-Asian populations, there are no accurate reports on the worldwide prevalence of DRPLA,

with current estimates based on the evaluation of cohorts diagnosed with SCA, suggesting that the prevalence of DRPLA is likely to be underestimated [6, 91, 96]. In Singapore, Korea and China, the frequencies of DRPLA have been found to be 3.4%, 3.4%, and 1%, respectively [42, 50, 111]. In South America, the DRPLA frequency has found to be 0.14% and 3.1% in SCA cohorts from Brazil and Venezuela, respectively [8, 65]. In Europe, findings have been variable, with reports of the frequency in Portuguese cohorts with autosomal dominant ataxias of DRPLA ranging from 4 to 11.2%, whilst in Spain, the frequency was reported as 3.3% [18, 39, 93]. In South Wales, France, and Italy, the frequency amongst cohorts with SCA has been reported as 5%, 0.25%, and 0.45–1%, respectively [11, 27, 49, 97]. Figure 2 illustrates the estimated number of cases around the world.


Developing a treatment approach for DRPLA and other repeat expansion disorders: drawing upon insights observed for HD

There are currently no treatments to prevent or stop the disease progression in DRPLA [90]. Whilst the exact pathophysiology of DRPLA is unclear, the literature overall points towards the idea that the expanded polyQ stretch leads to a “gain-of-toxic” function of the mutant protein on neuronal cells [89]. To downregulate the levels of the pathological polyQ proteins, RNA-targeting therapies may hold promise in the treatment of DRPLA, in particular, antisense oligonucleotides (ASO) therapy [16]. Therapeutic ASOs are single-stranded synthetic DNA molecules that work by binding to complementary target mRNA through Watson and Crick hybridization to interfere with normal gene expression and protein synthesis. ASOs affect gene expression through three mechanisms: RNase H-mediated degradation of mRNA, blocking ribosomes from binding to mRNA and preventing protein translation, or by modulating splicing of pre-mRNA [70, 102]. Figure 3 shows the normal steps of gene expression and the mechanisms by which therapeutic ASOs can influence this process. The scope of ASO therapeutics has expanded considerably in recent years, with an emphasis particularly placed on rare untreatable conditions, which cannot be easily addressed with small molecule drugs. ASO therapeutics have shown promise in several neurological disorders. For example, Nusinersen and Eteplisren are FDA-approved ASO treatment options for spinal muscular atrophy (SMA) and Duchenne muscular dystrophy (DMD), respectively, whilst clinical trials are ongoing for ASO treatments

for amyotrophic lateral sclerosis (ALS), Alzheimer’s disease (AD), and HD [75, 104]. Figure 4 highlights the progress of therapeutic ASO development for repeat expansion neurological disorders.

HD is the most widely studied CAG repeat expansion disorder and has gained significant attention for ASO therapeutics. Based on the success seen in preclinical studies of non-human animals, ASO clinical trials are ongoing for HD [78]. HTT_{RX} is an ASO that targets the mutant and wild-type alleles with the purpose of reducing levels of the mutant Huntingtin protein (mHTT). Through phase 1–2a clinical trial in early-stage HD patients, it was found that CSF mHTT levels showed dose-dependent decrease by up to 40%. No significant safety concerns were reported, though levels of CSF neurofilament light chain (NfL), a marker of neuroaxonal damage, were shown to be increased in the final study visit [86]. Preclinical studies have also been conducted for SCA. In early manifest transgenic SCA3 mice, *ATXN3*-targeting ASO resulted in sustained reduction of polyQ-expanded *ATXN3*, accompanied by rescued motor impairment [57]. Further, in SCA2 mouse models, the delivery of *ATXN2*-targeting ASO led to the downregulation of the *ATXN2* mRNA and protein, delayed onset of the SCA2 phenotype, with improved motor performance [74]. These findings indicate a promising proof-in-concept for ASO therapy as an approach for polyQ conditions.

The advances in HD are suggestive of the possibility of adopting similar methods to define biomarkers and treat DRPLA. Before clinically meaningful interventions can be discovered, a greater understanding of DRPLA disease progression and the identification of wet biomarkers must be pursued. Despite the significant advancements made for other neurodegenerative diseases, such as HD and AD, biomarkers in biological fluid, such as blood and CSF, have not been found for DRPLA [16]. Potential biomarkers including glial fibrillary acidic protein, DJ-1, and tau have been studied in SCA1, SCA2, and SCA6 patients, where only CSF tau was significantly higher in patients than controls, though levels did not correlate with CAG repeat size and disease severity [9]. CSF and plasma/serum NfL have been shown to be a notable biomarker in many neurodegenerative conditions, including HD, AD, ALS, and multiple sclerosis [12, 23, 30, 101]. In a small cohort of repeat-expansion SCA patients, serum NfL was found to be higher in patients than controls; however, the correlation with disease severity was not analysed [103]. Another potentially important biomarker for DRPLA disease progression and severity, which may



B

	Country	Number of cases	Location	Source
Europe	Portugal	5 families, 25 cases 8 families, 34 cases	Lisbon Population-based (1994-2004)	Vale, 2010 [93] Coutinho, 2013 [18]
	Hungary	1 case	Szeged	Zádori, 2015 [110]
	Spain	1 family, 3 cases 1 family, 12 cases 1 case	Cantabria Barcelona Barcelona	Infante, 2005 [39] Muñoz, 1999 [59] Pujana, 1999 [67]
	Turkey	1 family, 6 cases	Gaziantep	Yis, 2009 [109]
	France	1 family, 2 cases 1 family, 4 cases 1 family, 2 cases (+1 case)	Marseille Lille Paris	Casseron, 2004 [14] Destée, 2000 [21] Le Ber, 2003 [49]
	Wales	4 families, 9 cases	Cardiff	Wardle, 2009 [100]
	England	1 family, 6 cases 2 families, 9 cases 2 families, 4 cases 1 family, 2 cases	Birmingham London Region unclear Salisbury	Cox, 2000 [19] Wamer, 1995 [101] Becher, 1997 [4] Connarty, 1996 [17]
	Belgium	2 cases	Brussels	Titica, 1946 [88]
	Denmark	1 family, 6 cases	Copenhagen	Nøremølle, 1995 [63]
	Malta	1 family, 11 cases	Region unclear	Wamer, 1995 [101]
	(former) Yugoslavia	1 case	Region unclear	Smith, 1958 [81]
	Germany	3 cases	Luebek and Tübingen	Personal communication
	Italy	2 families, 3 cases 6 families, 51 cases 1 family 1 family, 1 case 1 case	Rome/Sicily Sicily Naples/Sicily Torino Foggia	Veneziano, 2014 [95] Gimaldi, 2019 [32] Filla, 2000 [27] Brusco, 2004 [11] Bidollari, 2019 [7]
Asia	India	3 cases 1 family, 9 cases	Mumbai New Delhi	Bhanushali, 2020 [5] Sharma, 2020 [76]
	Singapore	2 families, 2 cases	Singapore	Zhao, 2002 [112]
	China	2 families, 4 cases 1 family, 2 cases 1 family, 4 cases 1 family, 5 cases 3 cases 1 case	Hong Kong Taiwan Guangzhou Shenzhen Beijing Beijing	Yam, 2004 [106] Lee, 2001 [50] Liu, 2017 [52] Chen, 2013 [15] Zhang, 2013 [111] Hao, 2010 [33]
	Thailand	6 cases	Bangkok	Personal communication
	Korea	1 family, 5 cases 3 cases	Incheon Seoul	Kim, 2018 [43] Jin, 1999 [42]
	South America	Brazil	1 case	Fortaleza
Venezuela		2 families, 2 patients	Caracas	Paradisi, 2016 [65]
North America	USA	2 families, 6 cases 1 case 1 family, 6 cases 1 case 1 family, 14 cases	North Carolina and Maryland Texas Pennsylvania Georgia Tennessee	Becher, 1997 [4] Brunetti-Pierri, 2006 [10] Licht, 2002 [51] Silver, 2015 [79] Potter, 1995 [66]
	Canada	1 case	Toronto	Munhoz, 2004 [58]
Oceania	Australia	1 family, 3 cases	Melbourne	Vinton, 2005 [96]
	New Zealand	1 family, 3 cases	Auckland	Simpson, 2012 [80]

Fig. 2 Estimated number of reported cases of DRPLA based on a literature search. **a** World map with dots corresponding to the number of reported cases of DRPLA, as per studies shown in **(b)**. **b** Table of DRPLA families/cases found outside of Japan as reported in literature. DRPLA literature search for studies written in English published between January 1945 and July 2020 was performed on PubMed and Scopus databases, using the following key words: DRPLA; dentatorubral–pallidoluysian atrophy; Naito-Oyanagi disease; Haw-River syndrome; or ataxia. Cases for Thailand and Germany based on personal communication

also act as a potential therapeutic target, is repeat expansion somatic instability. This has been observed in HD mouse models and human brain tissue; it is worth exploring as a prominent biomarker for DRPLA and other repeat expansion disorders [3, 24, 85]. In addition to being clinically beneficial in regard to improving diagnostic accuracy and monitoring disease progression, biological biomarkers for DRPLA would also be influential in research. For example, they would provide greater accuracy in clinical trial recruitment, objective monitoring of disease-related biological changes, tracking adverse effects and response to treatment interventions [1]. Studies in larger cohorts are needed to gather data on the role of biological biomarkers DRPLA.

Insight into the natural progression of rare diseases is an essential step in facilitating the process of drug development [64]. To garner progress towards the discovery of disease-modifying treatments for DRPLA, emphasis must be placed on natural history studies to enhance our understanding of disease progression and to identify reproducible, validated biomarkers (Fig. 5). For rare diseases, this entails international collaboration to understand the longitudinal clinical progression in statistically large numbers of cases, ideally with $n > 20$ patients from multiple geographical areas, alongside matched controls. Initial investigative markers of disease progression would include clinical rating scales, imaging techniques and EEG, and objective, fluid-based biomarkers. In DRPLA, and other rare repeat expansion disorders, prominent analysis will comprise of DNA extracted from multiple fluids to investigate somatic instability, RNA extracted from blood and fibroblast cell lines, and extraction of serum/plasma and/or CSF for the examination of biomarkers such as NfL. The goal over the next 3 years will be to enhance our insight of DRPLA clinical features, imaging and fluid biomarkers, disease progression, and to uncover methods to monitor response to therapeutic intervention.




Fig. 3 Normal process of protein synthesis and common mechanisms of ASOs on target RNA. *Normal process of protein synthesis* (left): DNA is transcribed to pre-messenger RNA (pre-mRNA), which contains coding (exon) and non-coding (intron) regions between 5' and 3' untranslated regions (UTR). Pre-mRNA undergoes post-transcriptional modifications into mature mRNA, including 5' capping, removal of introns (splicing), and polyadenylation (poly-A tail). The mature mRNA undergoes ribosome-dependent protein synthesis. *Mechanisms of ASOs on target RNA* (right): Splicing modulation—ASO binds to pre-mRNA intron/exon junctions and modulate splic-

ing to include or skip the target exon, resulting in the synthesis of a modified protein. Translation inhibition—ASO binds to the mRNA and sterically blocks and prevents the binding of ribosomes to the mRNA, inhibiting translation and resulting in reduced protein synthesis. RNase H-mediated degradation of mRNA—ASO binds to the mRNA to form an RNA–DNA hybrid, allowing the recruitment of RNase H nuclease, inducing degradation of the target mRNA, resulting in reduced protein synthesis [22, 70, 73, 78, 102]. Diagram created on *biorender.com*

Disease Chromosome/gene	Repeat expansion	Progress of therapeutic ASO development	References
HD Chr4/Huntingtin	(CAG) _n	<i>Clinical trial</i> IONIS-HTT _{rx} phase III (NCT03761849).	Scoles <i>et al</i> , 2019 Tabrizi <i>et al</i> , 2019
ALS Chr9/C9orf72	(GGGGCC) _n	<i>Clinical trial</i> BIIB078 phase I (NCT04288856).	Scoles <i>et al</i> , 2019
SCA1 Chr6/ATXN1	(CAG) _n	<i>Preclinical research</i> Improved motor deficits and survival, reduced ATXN1 mRNA and protein in mice.	Freidrich <i>et al</i> , 2018
SCA2 Chr12/ATXN2	(CAG) _n	<i>Preclinical research</i> Improved motor phenotype, reduced ATXN2 mRNA and protein levels in mice.	Scoles <i>et al</i> , 2017
SCA3 Chr14/ATXN3	(CAG) _n	<i>Preclinical research</i> Rescued motor impairment, reduced polyQ-expanded ATXN3 in mice.	McLoughlin <i>et al</i> , 2018
SCA7 Chr3/ATXN7	(CAG) _n	<i>Preclinical research</i> Improved retinal degeneration and visual function, silencing of ATXN7 and reduction of protein aggregation in mice.	Niu <i>et al</i> , 2018
SCA36 Chr20/NOP35	(GGCCTG) _n	<i>Preclinical research</i> Reduced NOP56 mRNA expression SCA36 patient iPSCs and iPSC-derived neurons.	Matsuzono <i>et al</i> , 2017
DM1 Chr19/DMPK	(CTG) _n	<i>Preclinical research</i> Improved muscle strength, muscle histology and body weight, and reduced mutant transcripts in mice. <i>Clinical trial</i> IONIS-DMPK _{rx} phase I-II completed and significant biological changes not found (NCT02312011).	Jauvin <i>et al</i> , 2017
Fragile X syndrome X-linked/FMR1	(CGG) _n	<i>Preclinical research</i> Improved survival and suppressed repeat toxicity in mice and human neurons.	Rodriguez <i>et al</i> , 2020
SBMA X-linked/AR	(CAG) _n	<i>Preclinical research</i> Delayed onset and progression of motor dysfunction, improved survival, and suppressed mutant gene expression in mice.	Sahashi <i>et al</i> , 2015

Fig. 4 Progress in the development of ASO therapies for repeat expansions associated with neurological disorders. Table shows stages in the research towards developing ASO therapies for neurological repeat expansion disorders. Information. Chr: chromosome;

HD; Huntington's disease; ALS: amyotrophic lateral sclerosis; SCA: spinocerebellar ataxia; DM1: myotonic dystrophy; SBMA: spinobulbar muscular atrophy

The next few years for DRPLA

The future for DRPLA and other rare disorders is one of momentous opportunity. The knowledge gained in the scientific community from previous successful (and many unsuccessful) trials for SMA, HD, and other similar diseases have defined the foundations required to understand disease progression and how to see the reversal. The current development of collaborative natural history and biomarker studies for DRPLA by our group at UCL alongside other institutions gives hope to DRPLA patients and families for advancements over the next few years. Whilst it is expected

that many clinical, fluid or imaging markers of disease will overlap with other conditions, namely fluid NfL and MRI sequences, it can be postulated that DRPLA-specific markers may be discovered, for example, disease-associated protein levels such as ATN1, or somatic instability in the repeat expansion in biosamples. Though funding is challenging in rare disorders, natural history studies, in tandem with open-access data, imaging, wet biomarker and fibroblast repositories are essential. Ascertaining a wide resource for use by future researchers is crucial in the drive towards discoveries that may potentially benefit patient care.




Fig. 5 Facilitating DRPLA therapeutic development through understanding of natural history and discovery of biomarkers. Natural history studies follow the course of a disease from prior to inception, through the presymptomatic and clinical stages, to the point it ends (the patient is either cured, chronically disabled or dead, without external intervention) [20]. International, collaborative clinical studies are paramount to the DRPLA drug discovery process by identifying milestones of the disease progression and facilitating the discovery of longitudinal or cross-sectional biomarkers to objectively track disease-related biological changes. The discovery of biomarkers is, in turn, essential for clinical trials. The figure showcases the process by which a clinical and biobank resource for DRPLA can be uncovered, through collaborative efforts. Several methods are used for biomarker

discovery; in particular, ‘omics’ technologies contribute towards the rapid discovery and validation of biomarkers. Genomics allows the identification of gene mutations or polymorphisms; transcriptomics can identify changes in RNA; epigenetics can identify modified epigenetic mechanisms; metabolomics and proteomics can identify small molecule metabolites and protein biomarkers in human biological fluid, respectively [1, 2, 28, 31, 41]. MS: multiple spectrometry; NfL: neurofilament light chain; NMR: nuclear magnetic resonance; MALDI-TOF MS: matrix-assisted laser desorption/ionization time of flight mass spectrometry; MRI: magnetic resonance imaging; EEG: electroencephalogram; EDTA: Ethylenediamine tetraacetic acid; CSF: cerebrospinal fluid

Acknowledgements We thank all patients and families for their support, as well as clinical and laboratory collaborators who are essential to our work. We are grateful to Ataxia UK, CureDRPLA and The National Institute for Health Research University College London Hospitals Biomedical Research Centre for funding and supporting the current recruitment and forthcoming natural history study into DRPLA; for which we are enthusiastic to receive email contact (h.houlden@ucl.ac.uk) and enroll patients from around the world.

Compliance with ethical standards

Conflicts of interest The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Ashizawa T, Öz G, Paulson HL (2018) Spinocerebellar ataxias: prospects and challenges for therapy development. *Nat Rev Neurol* 14(10):590–605. <https://doi.org/10.1038/s41582-018-0051-6>
- Augustine EF, Adams HR, Mink JW (2013) Clinical trials in rare disease: challenges and opportunities. *J Child Neurol* 28(9):1142–1150. <https://doi.org/10.1177/0883073813495959>
- Aviolat H, Pinto RM, Godschall E et al (2019) Assessing average somatic CAG repeat instability at the protein level. *Sci Rep* 9:19152. <https://doi.org/10.1038/s41598-019-55202-x>
- Becher MW, Rubinsztein DC, Leggo J, Wagster MV, Stine OC, Ranen NG, Barron L (1997) Dentatorubral and pallidolusian atrophy (DRPLA) Clinical and neuropathological findings in genetically confirmed north American and European pedigrees. *Movement Disord* 12(4):519–530. <https://doi.org/10.1002/mds.870120408>
- Bhanushali AA, Venkatesan R, Das BR (2020) Spinocerebellar Ataxias in India: three-year molecular data from a central reference laboratory. *Neurol India* 68:86–91. <https://www.neurologyindia.com/text.asp?2020/68/1/86/279666>
- Bhowmik A, Rangaswami S et al (2015) Molecular genetic analysis of trinucleotide repeat disorders (TRDs) in Indian population and application of repeat primed PCR. *Eur J Med Genet* 58:160–167. <https://doi.org/10.1016/j.ejmg.2014.12.010>
- Bidollari E, Rotundo G, Altieri F, Amicucci M, Wiquel D, Ferrari D, Goldoni M, Bernardini L, Consoli F, De Luca A, Fanelli S, Lamorte G, D'Agruma L, Vescovi AL, Squitieri F, Rosati J (2019) Generation of induced pluripotent stem cell line CSSi008-A (4698) from a patient affected by advanced stage of Dentato-Rubral-Pallidolusian atrophy (DRPLA). *Stem Cell Res* 40:101551. <https://doi.org/10.1016/j.scr.2019.101551>
- Braga-Neto P, Pedroso JL, Furtado GV et al (2017) Dentatorubro-Pallidolusian Atrophy (DRPLA) among 700 Families with Ataxia in Brazil. *Cerebellum* 16:812–816. <https://doi.org/10.1007/s12311-017-0862-9>
- Brouillette A, Gulin O, Gomez C (2015) Cerebrospinal fluid biomarkers in spinocerebellar ataxia: a pilot study. *Dis Markers*. <https://doi.org/10.1155/2015/413098>
- Brunetti-Pierri N, Wilfong AA, Hunter JV, Craigen WJ (2006) A severe case of dentatorubro-pallidolusian atrophy (DRPLA) with microcephaly, very early onset of seizures, and cerebral white matter involvement. *Neuropediatrics* 37(5):308–311. <https://doi.org/10.1055/s-2006-955967>
- Brusco A, Gellera C, Cagnoli C et al (2004) Molecular genetics of hereditary spinocerebellar ataxia: mutation analysis of spinocerebellar ataxia genes and CAG/CTG repeat expansion detection in 225 Italian Families. *Arch Neurol* 61(5):727–733. <https://doi.org/10.1001/archneur.61.5.727>
- Byrne LM, Rodrigues FB, Blennow K, Durr A, Leavitt BR, Roos R, Scahill RI, Tabrizi SJ, Zetterberg H, Langbehn D, Wild EJ (2017) Neurofilament light protein in blood as a potential biomarker of neurodegeneration in Huntington's disease: a retrospective cohort analysis. *Lancet Neurol* 16(8):601–609. [https://doi.org/10.1016/S1474-4422\(17\)30124-2](https://doi.org/10.1016/S1474-4422(17)30124-2)
- Carroll LS, Massey TH, Wardle M, Peall KJ (2018) Dentatorubral-pallidolusian atrophy: an update. *Tremor Other Hyperkinetic Movements* 8:577. <https://doi.org/10.7916/D81N9HST>
- Casseron W, Azulay JP, Broglin D, Kaphan E, Genton P, Le Ber I, Gastaut JL (2004) Phenotype variability in a caucasian family with dentatorubral-pallidolusian atrophy. *Eur Neurol* 52(3):175–176. <https://doi.org/10.1159/000081859>
- Chen J, Zeng Z, Wu J et al (2013) The clinical and genetical characteristic of one dentatorubral-pallidolusian atrophy pedigree with an onset of cognitive impairment. *Chin J Neurol* 46:962–969. <https://doi.org/10.3760/cma.j.issn.1006-7876.2013.10.011>
- Coarelli G, Brice A, Durr A (2018) Recent advances in understanding dominant spinocerebellar ataxias from clinical and genetic points of view. *F1000Research*. 7: F1000. <https://doi.org/10.12688/f1000research.15788.1>
- Connarty M, Dennis NR, Patch C et al (1996) Molecular re-investigation of patients with Huntington's disease in Wessex reveals a family with dentatorubral and pallidolusian atrophy. *Hum Genet* 97:76–78. <https://doi.org/10.1007/BF00218837>
- Coutinho P, Ruano L, Loureiro JL et al (2013) Hereditary ataxia and spastic paraplegia in Portugal: a population-based prevalence study. *JAMA Neurol* 70(6):746–755. <https://doi.org/10.1001/jamaneurol.2013.1707>
- Cox H, Costin-Kelly NM, Ramani P, Whitehouse WP (2000) An established case of dentatorubral pallidolusian atrophy (DRPLA) with unusual features on muscle biopsy. *Eur J Paediat Neurol* 4(3):119–123. <https://doi.org/10.1053/ejpn.2000.0279>
- de Paz MP et al (2010) Rare diseases epidemiology research. In: Posada M, Groft S (eds) *Rare diseases epidemiology. Advances in experimental medicine and biology*, vol 686. Springer, Dordrecht. https://doi.org/10.1007/978-90-481-9485-8_2
- Destée A, Delalande I, Vuillaume I, Schraen-Maschke S, Defebvre L, Sablonnière B (2000) The first identified French family with dentatorubral-pallidolusian atrophy. *Movement Disord* 15(5):996–999. [https://doi.org/10.1002/1531-8257\(200009\)15:5<996::aid-mds1036>3.0.co;2-9](https://doi.org/10.1002/1531-8257(200009)15:5<996::aid-mds1036>3.0.co;2-9)
- Dhuri K, Bechtold C, Quijano E, Pham H, Gupta A, Vikram A, Bahal R (2020) Antisense oligonucleotides: an emerging area in drug discovery and development. *J Clin Med* 9(6):2004. <https://doi.org/10.3390/jcm9062004>
- Disanto G, Barro C, Benkert P, Naegelin Y, Schädelin S, Giardiello A, Zecca C, Blennow K, Zetterberg H, Leppert D, Kappos L, Gobbi C, Kuhle J (2017) Serum neurofilament light: a biomarker of neuronal damage in multiple sclerosis. *Ann Neurol* 81(6):857–870. <https://doi.org/10.1002/ana.24954>
- Dragivela E, Henricks A, Teed A et al (2009) Intergenerational and striatal CAG repeat instability in Huntington's disease

- knock-in mice involve different DNA repair genes. *Neurobiol Dis* 33:37–47. <https://doi.org/10.1016/j.nbd.2008.09.014>
25. Ellerby L (2019) Repeat expansion disorders: mechanisms and therapeutics. *Neurotherapeutics* 16:924–927. <https://doi.org/10.1007/s13311-019-00823-3>
 26. Fanto M, Charroux B (2010) Atrophin-1. *Encyclopedia Movement Disord* 2010:100–103. <https://doi.org/10.1016/B978-0-12-374105-9.00303-8>
 27. Filla A, Mariotti C, Caruso G, Coppola G, Coccozza S, Castaldo I, Calabrese O, Salvatore E, De Michele G, Riggio MC, Parayson D, Gellera C, Di Donato S (2000) Relative frequencies of CAG expansions in spinocerebellar ataxia and dentatorubropallidolusian atrophy in 116 Italian families. *Eur Neurol* 44(1):31–36. <https://doi.org/10.1159/000008189>
 28. Fitzgerald G (2016) Measure for measure: biomarker standards and transparency. *Sci Transl Med*. 8:34310. <https://doi.org/10.1126/scitranslmed.aaf8590>
 29. Friedrich J, Kordasiewicz HB, O’Callaghan B, Handler HP, Wagener C, Duvick L, Swayze EE, Rainwater O, Hofstra B, Benneyworth M, Nichols-Meade T, Yang P, Chen Z, Ortiz JP, Clark HB, Öz G, Larson S, Zoghbi HY, Henzler C, Orr HT (2018) Antisense oligonucleotide-mediated ataxin-1 reduction prolongs survival in SCA1 mice and reveals disease-associated transcriptome profiles. *JCI insight* 3(21):e123193. <https://doi.org/10.1172/jci.insight.123193>
 30. Gaiani A, Martinelli I, Bello L, Querin G, Puthenparampil M, Ruggero S, Toffanin E, Cagnin A, Briani C, Pegoraro E, Sorarù G (2017) Diagnostic and prognostic biomarkers in amyotrophic lateral sclerosis: neurofilament light chain levels in definite subtypes of disease. *JAMA Neurol* 74(5):525–532. <https://doi.org/10.1001/jamaneurol.2016.5398>
 31. Garcia-Gutierrez MS, Navarrete F, Sala F, Gasparian A, Austrich-Olivares A, Manzanares J (2020) Biomarkers in psychiatry: concept, definition, types and relevance to the clinical reality. *Front Psychiatry* 11:432. <https://doi.org/10.3389/fpsyg.2020.00432>
 32. Grimaldi S, Cupidi C, Smirne N, Bernardi L, Giacalone F, Piccione G, Basiricò S, Mangano GD, Nardello R, Orsi L, Grosso E, Laganà V, Mitolo M, Maletta RG, Bruni AC (2019) The largest caucasian kindred with dentatorubral-pallidolusian atrophy: a founder mutation in Italy. *Movement Disord* 34(12):1919–1924. <https://doi.org/10.1002/mds.27879>
 33. Hao Y, Gu W, Wang G et al (2010) Clinical and genetic study of one DRPLA case. *Chin J Contemporary Neurol Neurosurg* 10:637–641. <https://doi.org/10.3969/j.issn.1672-6731.2010.06.012>
 34. Hasegawa A, Ikeuchi T, Koike R, Matsubara N, Tsuchiya M, Nozaki H, Homma A, Idezuka J, Nishizawa M, Onodera O (2010) Long-term disability and prognosis in dentatorubral-pallidolusian atrophy: a correlation with CAG repeat length. *Movement Disord* 25(11):1694–1700. <https://doi.org/10.1002/mds.23167>
 35. Hatano T, Okuma Y, Iijima M, Fujishima K, Goto K, Mizuno Y (2003) Cervical dystonia in dentatorubral-pallidolusian atrophy. *Acta Neurol Scand* 108(4):287–289. <https://doi.org/10.1034/j.1600-0404.2003.00150.x>
 36. Hayashi Y, Kakita A, Yamada M, Egawa S, Oyanagi S, Naito H, Takahashi H (1998) Hereditary dentatorubral-pallidolusian atrophy: ubiquitinated filamentous inclusions in the cerebellar dentate nucleus neurons. *Acta Neuropathol* 95(5):479–482. <https://doi.org/10.1007/s004010050828>
 37. Ikeuchi T, Koide R, Onodera O et al (1995) Dentatorubral-pallidolusian atrophy (DRPLA). Molecular basis for wide clinical features of DRPLA. *Clin Neurosci* 3:23–27
 38. Ikeuchi T, Koide R, Tanaka H, Onodera O, Igarashi S, Takahashi H, Sato K (1995) Dentatorubral-pallidolusian atrophy: clinical features are closely related to unstable expansions of trinucleotide (CAG) repeat. *Ann Neurol* 37(6):769–775. <https://doi.org/10.1002/ana.410370610>
 39. Infante J, Combarros O, Volpini V, Corral J, Llorca J, Berciano J (2005) Autosomal dominant cerebellar ataxias in Spain: molecular and clinical correlations, prevalence estimation and survival analysis. *Acta Neurol Scand* 111(6):391–399. <https://doi.org/10.1111/j.1600-0404.2005.00400.x>
 40. Jauvin D, Chrétien J, Pandey SK, Martineau L, Revillod L, Bassez G, Lachon A, MacLeod AR, Gourdon G, Wheeler TM, Thornton CA, Bennett CF, Puymirat J (2017) Targeting DMPK with antisense oligonucleotide improves muscle strength in myotonic dystrophy type 1 mice. *Mol Ther Nucleic Acids* 7:465–474. <https://doi.org/10.1016/j.omtn.2017.05.007>
 41. Jewell NP (2016) Natural history of diseases: statistical designs and issues. *Clin Pharmacol Ther* 100(4):353–361. <https://doi.org/10.1002/cpt.423>
 42. Jin D, Oh M, Song S et al (1999) Frequency of spinocerebellar ataxia types 1, 2, 3, 6, 7 and dentatorubral pallidolusian atrophy mutations in Korean patients with spinocerebellar ataxia. *J Neurol* 246:207–210. <https://doi.org/10.1007/s004150050335>
 43. Kim H, Yun JY, Choi KG, Koo H, Han HJ (2018) Sleep related problems as a nonmotor symptom of dentatorubropallidolusian atrophy. *J Korean Med Sci* 33(17):e130. <https://doi.org/10.3346/jkms.2018.33.e130>
 44. Klockgether T, Mariotti C, Paulson HL (2019) Spinocerebellar ataxia. *Nat Rev Dis Primers* 5:24. <https://doi.org/10.1038/s41572-019-0074-3>
 45. Kobayashi J, Nagao M, Kawata A, Matsubara S (2009) A case of late adult-onset dentatorubral-pallidolusian atrophy mimicking central pontine myelinolysis. *J Neurol* 256(8):1369–1371. <https://doi.org/10.1007/s00415-009-5111-9>
 46. Koide R, Ikeuchi T, Onodera O, Tanaka H, Igarashi S, Endo K, Saito M (1994) Unstable expansion of CAG repeat in hereditary dentatorubral-pallidolusian atrophy (DRPLA). *Nat Genet* 6(1):9–13. <https://doi.org/10.1038/ng0194-9>
 47. Koide R, Onodera O, Ikeuchi T, Kondo R, Tanaka H, Tokiguchi S, Shimizu N (1997) Atrophy of the cerebellum and brainstem in dentatorubral pallidolusian atrophy: influence of CAG repeat size on MRI findings. *Neurology* 49(6):1605–1612. <https://doi.org/10.1212/WNL.49.6.1605>
 48. Latimer CS, Flanagan ME, Cimino PJ, Jayadev S, Davis M, Hoffer ZS, Montine TJ, Gonzalez-Cuyar LF, Bird TD, Keene CD (2017) Neuropathological comparison of adult onset and juvenile Huntington’s disease with cerebellar atrophy: a report of a father and son. *J Huntington’s Dis* 6(4):337–348. <https://doi.org/10.3233/JHD-170261>
 49. Le Ber I, Camuzat A, Castelnovo G et al (2003) Prevalence of dentatorubral-pallidolusian atrophy in a large series of white patients with cerebellar ataxia. *Arch Neurol* 60(8):1097–1099. <https://doi.org/10.1001/archneur.60.8.1097>
 50. Lee IH, Soong BW, Lu YC, Chang YC (2001) Dentatorubropallidolusian atrophy in Chinese. *Arch Neurol* 58(11):1905–1908. <https://doi.org/10.1001/archneur.58.11.1905>
 51. Licht DJ, Lynch DR (2002) Juvenile dentatorubral-pallidolusian atrophy: new clinical features. *Pediatr Neurol* 26(1):51–54. [https://doi.org/10.1016/s0887-8994\(01\)00346-0](https://doi.org/10.1016/s0887-8994(01)00346-0)
 52. Liu H, Xie L, Su C et al (2017) Clinical features, electrophysiology, neuroimaging and gene analysis of one juvenile dentatorubral-pallidolusian atrophy pedigree. *Chin J Neurol* 50:506–510. <https://doi.org/10.3760/cma.j.issn.1006-7876.2017.07.006>
 53. Matsuzono K, Imamura K, Murakami N, Tsukita K, Yamamoto T, Izumi Y, Kaji R, Ohta Y, Yamashita T, Abe K, Inoue H (2017) Antisense oligonucleotides reduce RNA foci in spinocerebellar

- ataxia 36 patient iPSCs. *Mol Ther Nucleic ACIDS* 8:211–219. <https://doi.org/10.1016/j.omtn.2017.06.017>
54. Maruyama H, Izumi Y, Morino H et al (2002) Difference in disease-free survival curve and regional distribution according to subtype of spinocerebellar ataxia: A study of 1,286 Japanese patients. *Int J Med Genetics* 114:578–583. <https://doi.org/10.1002/ajmg.10514>
 55. Maruyama S, Saito Y, Nakagawa E, Saito T, Komaki H, Sugai K, Sasaki M, Kumada S, Saito Y, Tanaka H, Minami N, Goto Y (2012) Importance of CAG repeat length in childhood-onset dentatorubral-pallidoluysian atrophy. *J Neurol* 259(11):2329–2334. <https://doi.org/10.1007/s00415-012-6493-7>
 56. Mazzoni P, Jain S (2007) Myoclonus. *Neurobiol Dis* 30:305–317. <https://doi.org/10.1016/B978-012088592-3/50032-3>
 57. McLoughlin H, Moore L, Chopra R et al (2018) Oligonucleotide therapy mitigates disease in spinocerebellar ataxia type 3. *Ann Neurol* 84:64–77. <https://doi.org/10.1002/ana.25264>
 58. Munhoz RP, Bergeron C, Lang AE (2004) Sporadic case of dentatorubral pallidoluysian atrophy with no CAG repeat expansion and no intranuclear inclusions. *Movement Disord* 19(5):580–583. <https://doi.org/10.1002/mds.10674>
 59. Muñoz E, Milà M, Sánchez A et al (1999) Dentatorubro-pallidoluysian atrophy in a Spanish family: a clinical, radiological, pathological, and genetic study. *J Neurol Neurosurg Psychiatry* 67:811–814
 60. Nagafuchi S, Yanagisawa H, Sato K et al (1994) Expansion of an unstable CAG trinucleotide on chromosome 12p in dentatorubral and pallidoluysian atrophy. *Nat Genet* 6:14–18
 61. Naito H, Oyanagi S (1982) Familial myoclonus epilepsy and choreoathetosis: hereditary dentatorubral-pallidoluysian atrophy. *Neurology* 32(8):798–798. <https://doi.org/10.1212/WNL.32.8.798>
 62. Niu C, Prakash TP, Kim A, Quach JL, Hury LA, Yang Y, Lopez E, Jazayeri A, Hung G, Sopher BL, Brooks BP, Swayze EE, Bennett CF, La Spada AR (2018) Antisense oligonucleotides targeting mutant Ataxin-7 restore visual function in a mouse model of spinocerebellar ataxia type 7. *Sci Transl Med* 10(465):8677. <https://doi.org/10.1126/scitranslmed.aap8677>
 63. Nørremølle A, Nielsen JE, Sørensen SA, Hasholt L (1995) Elongated CAG repeats of the B37 gene in a Danish family with dentato-rubro-pallido-luysian atrophy. *Hum Genet* 95(3):313–318
 64. Gavin P (2015) The importance of natural histories for rare diseases. *Expert Opin Orphan Drugs* 3(8):855–857. <https://doi.org/10.1517/21678707.2015.1063415>
 65. Paradisi I, Ikonomu V, Arias S (2016) Spinocerebellar ataxias in Venezuela: genetic epidemiology and their most likely ethnic descent. *J Hum Genet* 61:215–222. <https://doi.org/10.1038/jhg.2015.131>
 66. Potter NT, Meyer MA, Zimmerman AW, Eisenstadt ML, Anderson IJ (1995) Molecular and clinical findings in a family with dentatorubral-pallidoluysian atrophy. *Ann Neurol* 37(2):273–277. <https://doi.org/10.1002/ana.410370220>
 67. Pujana MA, Corral J, Gratacòs M, Combarros O, Berciano J, Genís D (1999) Spinocerebellar ataxias in Spanish patients: genetic analysis of familial and sporadic cases. *Hum Genet* 104(6):516–522
 68. Rodriguez C, Todd P (2019) New pathologic mechanisms in nucleotide repeat expansion disorders. *Neurobiol Dis* 130:104515. <https://doi.org/10.1016/j.nbd.2019.104515>
 69. Rodriguez CM, Wright SE, Kearse MG et al (2020) A native function for RAN translation and CGG repeats in regulating fragile X protein synthesis. *Nat Neurosci* 23:386–397. <https://doi.org/10.1038/s41593-020-0590-1>
 70. Rossor A, Reilly M, Sleight J (2018) Antisense oligonucleotides and other genetic therapies made simple. *Pract Neurol* 18:126–131. <https://doi.org/10.1136/practneurol-2017-001764>
 71. Ruano L, Melo C, Silva MC, Coutinho P (2014) The global epidemiology of hereditary ataxia and spastic paraplegia: a systematic review of prevalence studies. *Neuroepidemiology* 42(3):174–183. <https://doi.org/10.1159/000358801>
 72. Sahashi K, Katsuno M, Hung G et al (2015) Silencing neuronal mutant androgen receptor in a mouse model of spinal and bulbar muscular atrophy. *Hum Mol Genet* 24:5985–5994. <https://doi.org/10.1093/hmg/ddv300>
 73. Schoch K, Miller T (2017) Antisense oligonucleotides: translation from mouse models to human neurodegenerative diseases. *Neuron* 94:1056–1070. <https://doi.org/10.1016/j.neuron.2017.04.010>
 74. Scoles D, Meera P, Schneider M et al (2017) Antisense oligonucleotide therapy for spinocerebellar ataxia type 2. *Nature* 544:362–366. <https://doi.org/10.1038/nature22044>
 75. Scoles DR, Minikel EV, Pulst SM (2019) Antisense oligonucleotides: a primer. *Neurol Genetics* 5(2):e323. <https://doi.org/10.1212/NXG.0000000000000323>
 76. Sharma P, Shaikh R, Shamim U et al (2020) Genetically confirmed first Indian dentato-rubral-pallidoluysian atrophy kindred: a case report. *Ann Mov Disord* 3:60–64
 77. Shiraishi H, Egawa K, Ito T, Kawano O, Asahina N, Kohsaka S (2017) Efficacy of perampamilol for controlling seizures and improving neurological dysfunction in a patient with dentatorubral-pallidoluysian atrophy (DRPLA). *Epilepsy Behav Case Rep* 8:44–46. <https://doi.org/10.1016/j.ebcrc.2017.05.004>
 78. Silva A, Lobo D, Martins I et al (2020) Antisense oligonucleotide therapeutics in neurodegenerative diseases: the case of polyglutamine disorders. *Brain* 143:407–429. <https://doi.org/10.1093/brain/awz328>
 79. Silver MR, Sethi KD, Mehta SH et al (2015) Case report of optic atrophy in Dentatorubropallidoluysian Atrophy (DRPLA). *BMC Neurol* 15:260. <https://doi.org/10.1186/s12883-015-0520-0>
 80. Simpson M, Smith A, Kent H, Roxburgh R (2012) Neurological picture. Distinctive MRI abnormalities in a man with dentatorubral-pallidoluysian atrophy. *J Neurol Neurosurg Psychiatry* 83(5):529–530. <https://doi.org/10.1136/jnnp-2011-301612>
 81. Smith J, Gonda V, Malamud N (1958) Unusual form of cerebellar ataxia: combined dentato-rubral and pallido-luysian degeneration. *Neurology* 8:205
 82. Stoyas C, Spada A (2018) The CAG–polyglutamine repeat diseases: a clinical, molecular, genetic, and pathophysiologic nosology. *Handbook Clin Neurol* 147:143–170. <https://doi.org/10.1016/B978-0-444-63233-3.00011-7>
 83. Sugiyama A, Sato N, Nakata Y et al (2018) Clinical and magnetic resonance imaging features of elderly onset dentatorubral-pallidoluysian atrophy. *J Neurol* 265:322–329. <https://doi.org/10.1007/s00415-017-8705-7>
 84. Sunami Y, Koide R, Arai N, Yamada M, Mizutani T, Oyanagi K (2011) Radiologic and neuropathologic findings in patients in a family with dentatorubral-pallidoluysian atrophy. *AJNR Am J Neuroradiol* 32(1):109–114. <https://doi.org/10.3174/ajnr.A2252>
 85. Swami M, Hendricks AE, Gillis T, Massood T, Mysore J, Myers RH, Wheeler VC (2009) Somatic expansion of the Huntington's disease CAG repeat in the brain is associated with an earlier age of disease onset. *Hum Mol Genet* 18(16):3039–3047. <https://doi.org/10.1093/hmg/ddp242>
 86. Tabrizi S, Leavitt B, Landwehrmeyer B et al (2019) Targeting Huntingtin expression in patients with Huntington's disease. *NEJM* 380:2307–2316. <https://doi.org/10.1056/NEJMoa190090>
 87. Takano H, Cancel G, Ikeuchi T, Lorenzetti D, Mawad R, Stevanin G, Sasaki R (1998) Close associations between prevalences of dominantly inherited spinocerebellar ataxias with CAG-repeat expansions and frequencies of large normal CAG

- alleles in Japanese and Caucasian populations. *Am J Hum Genet* 63(4):1060–1066. <https://doi.org/10.1086/302067>
88. Titica J, van Bogaert LUDO (1946) Heredo-degenerative hemiballismus: a contribution to the question of primary atrophy of the corpus Luysii. *Brain* 69(4):251–263
 89. Tsuji S (2000) [Dentatorubral-pallidolusian atrophy (DRPLA)—discovery of the disease, DRPLA gene and the pathophysiology] *Rinsho Shinkeigaku. Clin Neurol* 40(12):1287–1289
 90. Tsuji S (2012) Dentatorubral-pallidolusian atrophy. *Handbook Clin Neurol* 103:587–594. <https://doi.org/10.1016/B978-0-444-51892-7.00041-3>
 91. Tsuji S, Onodera O, Goto J et al (2008) Sporadic ataxias in Japan—a population-based epidemiological study. *Cerebellum* 7:189. <https://doi.org/10.1007/s12311-008-0028-x>
 92. Tunc S, Tadic V, Zühlke C, Hellenbroich Y, Brüggemann N (2018) Pearls & Oy-sters: Family history of Huntington disease disguised a case of dentatorubral-pallidolusian atrophy. *Neurology* 90(3):142–143. <https://doi.org/10.1212/WNL.0000000000004833>
 93. Vale J, Bugalho P, Silveira I, Sequeiros J, Guimarães J, Coutinho P (2010) Autosomal dominant cerebellar ataxia: frequency analysis and clinical characterization of 45 families from Portugal. *Eur J Neurol* 17(1):124–128. <https://doi.org/10.1111/j.1468-1331.2009.02757.x>
 94. Veneziano L, Mantuano E, Catalli C, Gellera C, Durr A, Romano S, Spadaro M, Frontali M, Novelletto A (2014) A shared haplotype for dentatorubropallidolusian atrophy (DRPLA) in Italian families testifies of the recent introduction of the mutation. *J Hum Genet* 59(3):153–157. <https://doi.org/10.1038/jhg.2013.137>
 95. Vinton A, Fahey M, O'Brien T et al (2005) Dentatorubral-pallidolusian atrophy in three generations, with clinical courses from nearly asymptomatic elderly to severe juvenile, in an Australian family of Macedonian descent. *AJMG* 136A:201–204
 96. Wardle M, Majounie E, Williams NM et al (2008) Dentatorubral pallidolusian atrophy in South Wales. *J Neurol Neurosurg Psychiatry* 79:804–807. <https://doi.org/10.1136/jnnp.2007.128074>
 97. Wardle M, Majounie E, Muzaimi MB, Williams NM, Morris HR, Robertson NP (2009) The genetic aetiology of late-onset chronic progressive cerebellar ataxia. A population-based study. *J Neurol* 256(3):343–348. <https://doi.org/10.1007/s00415-009-0015-2>
 98. Wardle M, Majounie E, Muzaimi MB, Williams NM, Morris HR, Robertson NP (2009) The genetic aetiology of late-onset chronic progressive cerebellar ataxia A population-based study. *J Neurol* 256(3):343–348. <https://doi.org/10.1007/s00415-009-0015-2>
 99. Wardle M, Morris HR, Robertson NP (2009) Clinical and genetic characteristics of non-Asian dentatorubral-pallidolusian atrophy: a systematic review. *Movement Disord* 24(11):1636–1640. <https://doi.org/10.1002/mds.22642>
 100. Warner TT, Williams LD, Walker RWH, Flinter F, Robb SA, Bunday SE, Harding AE (1995) A clinical and molecular genetic study of dentatorubropallidolusian atrophy in four European families. *Ann Neurol* 37(4):452–459. <https://doi.org/10.1002/ana.410370407>
 101. Weston P, Poole T, Ryan NS, Nair A, Liang Y, Macpherson K, Drueh R, Malone IB, Ahsan RL, Pemberton H, Klimova J, Mead S, Blennow K, Rossor MN, Schott JM, Zetterberg H, Fox NC (2017) Serum neurofilament light in familial Alzheimer disease: a marker of early neurodegeneration. *Neurology* 89(21):2167–2175. <https://doi.org/10.1212/WNL.0000000000004667>
 102. Wild EJ, Tabrizi SJ (2017) Therapies targeting DNA and RNA in Huntington's disease. *Lancet Neurol* 16(10):837–847. [https://doi.org/10.1016/S1474-4422\(17\)30280-6](https://doi.org/10.1016/S1474-4422(17)30280-6)
 103. Wilke C, Bender F, Hayer SN, Brockmann K, Schöls L, Kuhle J, Synofzik M (2018) Serum neurofilament light is increased in multiple system atrophy of cerebellar type and in repeat-expansion spinocerebellar ataxias: a pilot study. *J Neurol* 265(7):1618–1624. <https://doi.org/10.1007/s00415-018-8893-9>
 104. Wurster CD, Ludolph AC (2018) Antisense oligonucleotides in neurological disorders. *Thera Adv Neurol Disord* 11:1756286418776932. <https://doi.org/10.1177/1756286418776932>
 105. Yam WK, Wu NS, Lo IF, Ko CH, Yeung WL, Lam ST (2004) Dentatorubral-pallidolusian atrophy in two Chinese families in Hong Kong. *Hong Kong Med J* 10(1):53–56
 106. Yamada M (2010) Dentatorubral-pallidolusian atrophy (DRPLA): The 50th Anniversary of Japanese Society of Neuropathology. *Neuropathology* 30:453–457. <https://doi.org/10.1111/j.1440-1789.2010.01120.x>
 107. Yamada M, Wood JD, Shimohata T, Hayashi S, Tsuji S, Ross CA, Takahashi H (2001) Widespread occurrence of intranuclear atrophin-1 accumulation in the central nervous system neurons of patients with dentatorubral-pallidolusian atrophy. *Ann Neurol* 49(1):14–23. [https://doi.org/10.1002/1531-8249\(200101\)49:1%3C14:AID-ANA5%3E3.0.CO;2-X](https://doi.org/10.1002/1531-8249(200101)49:1%3C14:AID-ANA5%3E3.0.CO;2-X)
 108. Yiş U, Dirik E, Gündoğdu-Eken A, Başak AN (2009) Dentatorubral pallidolusian atrophy in a Turkish family. *Turk J Pediatr* 51(6):610–612
 109. Zádori D, Tánzos T, Jakab K, Vécsei L, Klivényi P (2015) The first identified Central-Eastern European patient with genetically confirmed dentatorubral-pallidolusian atrophy. *Ideggyogyaszati szemle* 68(1–2):68–71
 110. Zhang Z, Hao Y (2013) Genetics and clinical study of Chinese kindreds with dentatorubral pallidolusian atrophy. *Chin J Med Genetics* 30:31–35. <https://doi.org/10.3760/cma.j.issn.1003-9406.2013.01.008>
 111. Zhao Y, Tan EK, Law HY, Yoon CS, Wong MC, Ng I (2002) Prevalence and ethnic differences of autosomal-dominant cerebellar ataxia in Singapore. *Clin Genet* 62:478–481. <https://doi.org/10.1034/j.1399-0004.2002.620610.x>