

Diagnostic value of the posterior talofibular ligament area for chronic lateral ankle instability

Young Joo, MD, PhD^a, JeeYoun Moon, MD, PhD^b, Billy Huh, MD, PhD^c, Geung Kyu Lee, MD^d, Hyung Rae Cho, MD^d, Keum Nae Kang, MD^e, Soho Lee, MD^f, Young Uk Kim, MD, PhD^{f,*}

Abstract

An injured posterior talofibular ligament (PTFL) is one of the reasons for chronic lateral ankle instability (CLAI). Previous researches have demonstrated that the PTFL thickness (PTFLT) is associated with chronic ligament injuries. However, ligament hypertrophy is different from ligament thickness. Thus, we created the PTFL cross-sectional area (PTFLCSA) as a diagnostic image parameter to assess the hypertrophy of the whole PTFL. We assumed that the PTFLCSA is a key morphological diagnostic parameter in CLAI. PTFL data were obtained from 15 subjects with CLAI and from 16 normal individuals. The T1-weighted axial ankle-MR (A-MR) images were acquired at the level of PTFL. We measured the PTFLT and PTFLCSA at the posterior aspect of the ankle using our imaging analysis program. The PTFLT was measured as the thickness between point of anterior and posterior fiber of PTFL. The PTFLCSA was calculated as the whole cross-sectional PTFL area. The average PTFLT was 3.43 ± 0.52 mm in the healthy group and 4.89 ± 0.80 mm in the CLAI group. The mean PTFLCSA was 41.06 ± 12.18 mm² in the healthy group and 80.41 ± 19.14 mm² in the CLAI group. CLAI patients had significantly greater PTFLT ($P < .001$) and PTFLCSA ($P < .001$) than the healthy group. A receiver operating characteristic curve analysis demonstrated that the optimal cutoff score of the PTFLT was 4.19 mm, with 93.3% sensitivity, 93.7% specificity, and an area under the curve of 0.97. The most suitable cutoff value of the PTFLCSA was 61.15 mm², with 93.3% sensitivity, 100% specificity, and area under the curve of 0.99. Even though the PTFLT and PTFLCSA were both significantly associated with CLAI, the PTFLCSA was a more exact morphological measurement parameter.

Abbreviations: A-MR = ankle-magnetic resonance, ATFL = anterior talofibular ligament, CFL = calcaneofibular ligament, CLAI = chronic lateral ankle instability, MRI = MR imaging, PTFL = posterior talofibular ligament, PTFLCSA = PTFL cross-sectional area, PTFLT = PTFL thickness, ROC = receiver operating characteristic.

Keywords: chronic lateral ankle instability, cross-sectional area, posterior talofibular ligament, thickness

1. Introduction

Lateral ankle ligamentous sprain is considered as one of the most common injuries affecting the lateral ligaments of the ankle. According to previous researches, 20 to 30% of all sports injuries are ankle injuries. Lateral ankle ligamentous sprain are frequently partially treated. The rate of repeated ankle sprain is >40%, and recurrent ankle injury can lead to chronic lateral ankle instability (CLAI) and ankle osteoarthritis.^[1-4] The lateral ankle ligament is very important to maintain static and dynamic ankle stability and intact ligaments for movement and support functions of ankle and foot.^[5,6] As we already know, frequent injuries of the ligaments such as calcaneofibular ligament (CFL), deltoid ligament and anterior talofibular ligament (ATFL), lead to CLAI.^[5-8] However, there

is no attentions about the relationship between CLAI and posterolateral ankle ligaments such as posterior talofibular ligament (PTFL). PTFL on the ankle-magnetic resonance (A-MR) image is visualized between the medial surface of the lateral malleolus and talus.^[9-11] The original insertions of the posterior and anterior fibers of the PTFL are the medial aspect of the lateral malleolus of the fibula. The posterior fibers of the PTFL were attached into the lateral tubercle of the posterior process of the talus. The anterior fibers were attached into the surface of the lateral talus posterior to the lateral malleolar facet.^[11] An injured PTFL has been thought to be one of an important finding of CLAI.^[11,12] A-MR images facilitate the analysis of the pathologic disorders of the PTFL.^[13] However, many treating doctors rarely consider the A-MR findings when assessing chronic morphologic changes in the PTFL in the CLAI patients.

The authors have no funding to disclose.

The authors have no conflicts of interest to disclose.

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

^a Department of Anesthesiology and Pain Medicine, CHA Ilsan Medical Center, CHA University, Goyang, Republic of Korea, ^b Department of Anesthesiology and Pain Medicine, Seoul National University Hospital, Seoul National University School of Medicine, Seoul, Republic of Korea, ^c Department of Pain Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX, ^d Department of Anesthesiology and Pain Medicine, Myongji Hospital, Hanyang University College of Medicine, Goyang, Republic of Korea, ^e Department of Anesthesiology and Pain Medicine, National Police Hospital, Seoul, Republic of Korea, ^f Department of Anesthesiology and Pain Medicine, Catholic Kwandong University of Korea College of Medicine, International ST. Mary's Hospital, Incheon, Republic of Korea.

* Correspondence: Young Uk Kim, Department of Anesthesiology and Pain Medicine, Catholic Kwandong University of Korea College of Medicine, International ST. Mary's Hospital, Simgokro, 100 Gil 25, Seo-Gu, Incheon City 22711, Republic of Korea (e-mail: uk201@hanmail.net).

Copyright © 2023 the Author(s). Published by Wolters Kluwer Health, Inc. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial License 4.0 (CCBY-NC), where it is permissible to download, share, remix, transform, and buildup the work provided it is properly cited. The work cannot be used commercially without permission from the journal.

How to cite this article: Joo Y, Moon J, Huh B, Lee GK, Cho HR, Kang KN, Lee S, Kim YU. Diagnostic value of the posterior talofibular ligament area for chronic lateral ankle instability. *Medicine* 2023;102:5(e32827).

Received: 22 October 2021 / Received in final form: 10 January 2023 / Accepted: 11 January 2023

<http://dx.doi.org/10.1097/MD.00000000000032827>

Moreover, previous researches only analyzed the PTFL using a single measurement at the approximate “halfway” of the PTFL. However, an asymmetrical thickening and partial tear of the PTFL can occur anywhere.^[14] Therefore, measurement mistakes could occur in some cases. In contrast to the PTFL thickness (PTFLT) between anterior and posterior fiber, the cross-sectional area of PTFL does not worry about these measurement mistakes because the PTFL cross-sectional area (PTFLCSA) measures the whole PTFLCSA. Thus, to assess the hypertrophy of the PTFL, we created the PTFLCSA as an adjuvant new image diagnostic parameter. We assumed that the PTFLCSA is an important morphologic parameter in CLAI diagnosis. Thus, we used A-MR to compare the PTFLT and PTFLCSA between CLAI patients and healthy groups.

2. Materials and methods

2.1. Patients

This retrospective research has been examined by the Catholic Kwandong University Institutional Review Board. (IRB no: IS19RISI0049). We reviewed CLAI patients who diagnosed our orthopedic clinic with ankle discomfort from May 2016 to January 2019 and who had taken A-MR. The criteria for inclusion of CLAI were as follows: past history of recurrent ankle injuries; chronic lateral ankle discomfort and persistent pain on anterior drawer test; no feel confident exercise; had not responded to conservative treatment; and at least 1 year has elapsed since the first ligament injury, but ankle pain persists. Patients were excluded if they had any following issues: past

ankle surgical history such as the ankle arthroscopy, modified Broström procedure, peroneal nerve disorder, and hindfoot varus.

There were 4 (26.6%) males and 11 (73.4%) females with a mean age of 38.60 ± 14.36 years (range, 16 to 59 years) (Table 1). To compare the PTFLT and PTFLCSA between the CLAI group and normal group, we also enrolled healthy individuals. In the normal group, 16 subjects (8 males and 8 females) were enrolled with a mean age of 38.13 ± 17.51 years (range, 16–65 years).

2.2. Imaging parameters

A-MR scans were performed on a 3.0-T Siemens MAGNETOM Avanto (Siemens Healthcare, Munich, Germany) and 3T Ingina scanners (Philips, Eindhoven, Netherlands). The A-MR imaging (MRI) protocols included the following sequences: repetition time 603-ms/echo time 19-ms, high resolution axial T1-weighted images, a slice thickness of 3.0mm, intersection gap of 0.9mm, 484×471 matrix, 140×140 cm field of view, and >3.0 ETL.

2.3. Image analysis

PTFLT and PTFLCSA measurements were analyzed by the one specialist, who was blinded to the ankle classification. We acquired the T1 weighted axial MR cut at the thickest view of the PTFL. We measured the PTFLT and PTFLCSA on MR images using an image analysis software (INFINITT PACS system, Seoul, South Korea). (Fig. 1A and B). The PTFLT was measured as the thickness between point of anterior and posterior fiber of PTFL. The PTFLCSA was measured as the cross-sectional area of the PTFL that was extremely hypertrophied on T1-weighted MR images.

2.4. Statistical analysis

We compared the PTFLT and PTFLCSA between the CLAI and the normal healthy groups using unpaired *t* tests. Data are presented the mean standard deviations. A receiver operating characteristic (ROC) curve analysis was generated for diagnostic method. *P* values $<.05$ were considered statistically significantly different. We used options in the SPSS package (IBM/SPSS for Windows ver 22.0, Chicago, IL) for the presentation and area under the curve (AUC) calculation of the ROC curve.

Table 1

Comparison of the characteristics of the control and CLAI groups.

Variable	Control group n = 16	CLAI group n = 15	Statistical significance
Gender (male/female)	8/8	4/11	NS
Ankle image (Rt/Lt)	10/6	8/7	NS
Age (yr)	38.13 ± 17.51	38.60 ± 14.36	NS
PTFLT (mm)	3.43 ± 0.52	4.89 ± 0.80	<i>P</i> < .001
PTFLCSA (mm ²)	41.06 ± 12.18	80.41 ± 19.14	<i>P</i> < .001

Data represent the mean \pm standard deviation (SD) or the numbers of patients. CLAI = chronic lateral ankle instability, NS = not statistically significant (*P* > .05), PTFLCSA = posterior talofibular ligament cross-sectional area, PTFLT = posterior talofibular ligament thickness.




Figure 1. (A) Measurement of both posterior talofibular ligament thickness (PTFLT) (white arrow) and (B) posterior talofibular ligament cross-sectional area (PTFLCSA) (white arrow) in the chronic lateral ankle instability group was carried out on MR T1 weighted images. F = fibula, MR = magnetic resonance, T = tibia.

There are several important limitations of this research. First, small sample may be weakness of this research. Second, the ankle lateral ligament complex comprises the CFL ATFL, and the PTFL. In this study, we only focused on PTFL. Third, there are several different methods to assess CLAI, such as manual anterior drawer test, stress radiography, ultrasound examination or arthroscopy have been proved to be effective at discriminating CLAI.^[24–33] However, we analyzed the PTFLCSA and PTFLT on MRI only.

Despite these limitations, this is the first research to document the association of PTFLCSA with CLAI. The PTFLCSA is a simple, reliable measurement tool with high sensitive value to evaluate CLAI.

5. Conclusion

The PTFLCSA was a more reliable measurement parameter for CLAI than PTFLT. We demonstrated that the best cutoff point of the PTFLCSA as 61.15 mm², with 93.3% sensitivity, 100% specificity. When assessing patients with CLAI, physicians should evaluate carefully the PTFLCSA rather than the PTFLT.

Acknowledgments

The authors thank the International ST. Mary's Hospital.

Author contributions

Supervision: Young Uk Kim.

Validation: Young Uk Kim.

Visualization: Young Uk Kim.

Writing – original draft: Young Uk Kim.

Writing – review & editing: Young Joo, JeeYoun Moon, Billy Huh, Geung Kyu Lee, Hyung Rae Cho, Keum Nae Kang, Soho Lee, Young Uk Kim.

References

- Brambilla L, Bianchi A, Malerba F, et al. Lateral ankle ligament anatomic reconstruction for chronic ankle instability: Allograft or auto-graft? A systematic review. *Foot Ankle Surg.* 2020;26:85–93.
- Chew CP, Koo KOT, Lie DTT. Periosteal flap augmentation of the Modified Brostrom-Gould procedure for chronic lateral ankle instability. *J Orthop Surg (Hong Kong).* 2018;26:2309499018757532309499018757530.
- Hu Y, Tao H, Qiao Y, et al. Evaluation of the Talar cartilage in chronic lateral ankle instability with lateral ligament injury using biochemical T2* mapping: correlation with clinical symptoms. *Acad Radiol.* 2018;25:1415–21.
- Lee JH, Yoon TL. Effective treatment for chronic ankle instability during lateral step-down - kinesiology tape, resistance exercise, or both accompanied with heel raise-lower exercise? *J Sport Rehabil.* 2019;28:809–16.
- Brown AJ, Shimozone Y, Hurley ET, et al. Arthroscopic repair of lateral ankle ligament for chronic lateral ankle instability: a systematic review. *Arthroscopy.* 2018;34:2497–503.
- Choi WJ, Lee JW, Han SH, et al. Chronic lateral ankle instability: the effect of intra-articular lesions on clinical outcome. *Am J Sports Med.* 2008;36:2167–72.
- Noailles T, Lopes R, Padiolleau G, et al. Non-anatomical or direct anatomical repair of chronic lateral instability of the ankle: a systematic review of the literature after at least 10 years of follow-up. *Foot Ankle Surg.* 2018;24:80–5.
- Zhan Y, Yan X, Xia R, et al. Anterior-inferior tibiofibular ligament anatomical repair and augmentation versus trans-syndesmosis screw fixation for the syndesmosis instability in external-rotation type ankle fracture with posterior malleolus involvement: a prospective and comparative study. *Injury.* 2016;47:1574–80.
- Chandnani VP, Harper MT, Ficke JR, et al. Chronic ankle instability: evaluation with MR arthrography, MR imaging, and stress radiography. *Radiology.* 1994;192:189–94.
- Chun KY, Choi YS, Lee SH, et al. Deltoid ligament and tibiofibular syndesmosis injury in chronic lateral ankle instability: magnetic resonance imaging evaluation at 3t and comparison with arthroscopy. *Korean J Radiol.* 2015;16:1096–103.
- Gursoy M, Dag F, Mete BD, et al. The anatomic variations of the posterior talofibular ligament associated with os trigonum and pathologies of related structures. *Surg Radiol Anat.* 2015;37:955–62.
- Doring S, Provyn S, Marcelis S, et al. Ankle and midfoot ligaments: ultrasound with anatomical correlation: a review. *Eur J Radiol.* 2018;107:216–26.
- Schneck CD, Mesgarzadeh M, Bonakdarpour A, et al. MR imaging of the most commonly injured ankle ligaments. Part I. Normal anatomy. *Radiology.* 1992;184:499–506.
- Taser F, Shafiq Q, Ebraheim NA. Anatomy of lateral ankle ligaments and their relationship to bony landmarks. *Surg Radiol Anat.* 2006;28:391–7.
- D'Hooghe P, Alkhelaifi K, Almusa E, et al. Chronic lateral ankle instability increases the likelihood for surgery in athletes with os trigonum syndrome. *Knee Surg Sports Traumatol Arthrosc.* 2019;27:2813–7.
- Trichine F, Friha T, Boukabou A, et al. Surgical treatment of chronic lateral ankle instability using an inferior extensor retinaculum flap: a retrospective study. *J Foot Ankle Surg.* 2018;57:226–31.
- Porter DA, Kamman KA. Chronic lateral ankle instability: open surgical management. *Foot Ankle Clin.* 2018;23:539–54.
- Miklovic TM, Donovan L, Protzuk OA, et al. Acute lateral ankle sprain to chronic ankle instability: a pathway of dysfunction. *Phys Sportsmed.* 2018;46:116–22.
- Cho JH, Lee DH, Song HK, et al. Value of stress ultrasound for the diagnosis of chronic ankle instability compared to manual anterior drawer test, stress radiography, magnetic resonance imaging, and arthroscopy. *Knee Surg Sports Traumatol Arthrosc.* 2016;24:1022–8.
- Cha SD, Kim HS, Chung ST, et al. Intra-articular lesions in chronic lateral ankle instability: comparison of arthroscopy with magnetic resonance imaging findings. *Clin Orthop Surg.* 2012;4:293–9.
- Zhu ZJ, Zhu Y, Liu JF, et al. Posterolateral ankle ligament injuries affect ankle stability: a finite element study. *BMC Musculoskelet Disord.* 2016;17:96.
- Liu K, Gustavsen G, Royer T, et al. Increased ligament thickness in previously sprained ankles as measured by musculoskeletal ultrasound. *J Athl Train.* 2015;50:193193e198.
- Ahmad MA, Pandey UC, Crerand JJ, et al. Magnetic resonance imaging of the normal and injured lateral collateral ligaments of the ankle. *Ann Chir Gynaecol.* 1998;87:311–6.
- Baytar MS, Yilmaz C, Karasu D, et al. Comparison of ultrasonography guided serratus anterior plane block and thoracic paravertebral block in video-assisted thoracoscopic surgery: a prospective randomized double-blind study. *Korean J Pain.* 2021;34:234–40.
- Cankurtaran D, Yigman ZA, Umay E. Factors associated with paravertebral muscle cross-sectional area in patients with chronic low back pain. *Korean J Pain.* 2021;34:454–62.
- Doo AR, Kim JW, Lee JH, et al. A comparison of two techniques for ultrasound-guided caudal injection: the influence of the depth of the inserted needle on caudal block. *Korean J Pain.* 2015;28:122–8.
- Ghai B, Kumar M, Makkar JK, et al. Comparison of ultrasound guided pulsed radiofrequency of genicular nerve with local anesthetic and steroid block for management of osteoarthritis knee pain. *Korean J Pain.* 2022;35:183–90.
- Jain A, Jain S, Barasker SK, et al. Predictors of discogenic pain in magnetic resonance imaging: a retrospective study of provocative discography performed by posterolateral approach. *Korean J Pain.* 2021;34:447–53.
- Jeong H, Choi JW, Sim WS, et al. Ultrasound-guided erector spinae plane block for pain management after gastrectomy: a randomized, single-blinded, controlled trial. *Korean J Pain.* 2022;35:303–10.
- John H, Sohn K, Kim JH. Relationship between needle depth for lumbar transforaminal epidural injection and patients' height and weight using magnetic resonance imaging. *Korean J Pain.* 2022;35:345–52.
- Kose SG, Kose HC, Celikel F, et al. Predictive factors associated with successful response to ultrasound guided genicular radiofrequency ablation. *Korean J Pain.* 2022;35:447–57.
- Mendonca CR, Noll M, de Carvalho Santos A, et al. High prevalence of musculoskeletal pain in individuals with severe obesity: sites, intensity, and associated factors. *Korean J Pain.* 2020;33:245–57.
- Yoo SW, Ki MJ, Doo AR, et al. Prediction of successful caudal epidural injection using color Doppler ultrasonography in the paramedian sagittal oblique view of the lumbosacral spine. *Korean J Pain.* 2021;34:339–45.