



## Molecular epidemiology of DNA repair gene polymorphisms and head and neck cancer

Meilin Wang<sup>a,b</sup>, Haiyan Chu<sup>a,b</sup>, Zhengdong Zhang<sup>a,b,✉</sup>, Qingyi Wei<sup>c,✉</sup>

<sup>a</sup>Department of Genetic Toxicology, the Key Laboratory of Modern Toxicology of Ministry of Education, School of Public Health, Nanjing Medical University, Nanjing, Jiangsu 211166, China;

<sup>b</sup>Department of Environmental Genomics, Jiangsu Key Laboratory of Cancer Biomarkers, Prevention and Treatment, Cancer Center, Nanjing Medical University, Nanjing, Jiangsu 211166, China;

<sup>c</sup>Department of Epidemiology, the University of Texas M. D. Anderson Cancer Center, Houston, Texas, USA.

Received 20 March 2013, Accepted 27 March 2013, Epub 16 April 2013

### Abstract

Although tobacco and alcohol consumption are two common risk factors of head and neck cancer (HNC), other specific etiologic causes, such as viral infection and genetic susceptibility factors, remain to be understood. Human DNA is often damaged by numerous endogenous and exogenous mutagens or carcinogens, and genetic variants in interaction with environmental exposure to these agents may explain interindividual differences in HNC risk. Single nucleotide polymorphisms (SNPs) in genes involved in the DNA damage-repair response are reported to be risk factors for various cancer types, including HNC. Here, we reviewed epidemiological studies that have assessed the associations between HNC risk and SNPs in DNA repair genes involved in base-excision repair, nucleotide-excision repair, mismatch repair, double-strand break repair and direct reversion repair pathways. We found, however, that only a few SNPs in DNA repair genes were found to be associated with significantly increased or decreased risk of HNC, and, in most cases, the effects were moderate, depending upon locus-locus interactions among the risk SNPs in the pathways. We believe that, in the presence of exposure, additional pathway-based analyses of DNA repair genes derived from genome-wide association studies (GWASs) in HNC are needed.

**Keywords:** DNA repair, molecular epidemiology, head and neck cancer, genetic variation, genome-wide association study

### INTRODUCTION

Head and neck cancer (HNC), of which the vast majority are squamous cell carcinomas of the head and neck (SCCHN), is the sixth most common cancer and the seventh leading cause of cancer-related deaths

worldwide<sup>[1]</sup>. In the United States, it has been estimated that 52,610 new cases and 11,500 deaths from HNC may have occurred in 2012, which accounts for 3% of all malignancies<sup>[2]</sup>. Anatomically, HNC includes cancers of the oral cavity, oropharynx, hypopharynx, and larynx. HNC is considered a complex disease because

✉ Corresponding authors: Qingyi Wei, MD, PhD, Department of Epidemiology, The University of Texas M.D. Anderson Cancer Center, 1515 Holcombe Blvd, Houston, Texas 77030, USA. Tel/Fax: 1-713-792-3020/1-713-563-0999, E-mail: [qwei@mdanderson.org](mailto:qwei@mdanderson.org); and Zhengdong Zhang, MD, PhD, Department of Environmental

Genomics, School of Public Health, Nanjing Medical University, 818 East Tianyuan Road, Nanjing, Jiangsu 211166, China. Tel/Fax: +86-25-8686-8423/+86-25-8686-8499, E-mail: [drzdzhang@gmail.com](mailto:drzdzhang@gmail.com).

The authors reported no conflict of interests.

both genetic and environmental risk factors contribute to its etiology. Epidemiological studies have demonstrated that tobacco use and alcohol consumption are the most known risk factors for HNC<sup>[3,4]</sup>. Furthermore, infection with high-risk types of human papilloma viruses (HPVs) has also been recognized as an increasingly important risk factor for HNC, especially for oropharyngeal squamous cell carcinoma<sup>[5,6]</sup>. Family history of cancer is another important risk factor for HNC, suggesting that genetic factors may contribute to HNC susceptibility<sup>[7]</sup>.

DNA is constantly damaged by endogenous oxygen free radicals from metabolism and exogenous (both chemical and physical) mutagens. The DNA damage repair process is a crucial mechanism to maintain the stability of genetic materials in mammalian cells, and different repair pathways are available to repair different types of DNA damage<sup>[8]</sup>. If this mechanism fails, however, unrepaired damage can result in apoptosis or otherwise may lead to mutation fixation, unregulated cell growth and cancer<sup>[9]</sup>. There are several DNA repair pathways in human cells, including base excision repair (BER), nucleotide excision repair (NER), mismatch repair (MMR), double-strand break repair (DSBR), and direct reversion repair (DRR)<sup>[8]</sup>.

Depending upon the nature of DNA damaging agents, different repair pathways involving distinct genes could be activated. For example, the BER pathway is one of the important mechanisms responsible for the removal of the oxidative and hydrolytic decay of DNA damage. The damaged base is recognized and cleaved from the DNA backbone to generate an apurinic/apyrimidinic (AP) site, and the abasic site is then restored by apurinic/apyrimidinic endonuclease 1 (APE1); the oxidized base 8-oxoguanine is excised by 8-oxoguanine DNA glycosylase (OGG1), and then the DNA synthesis and ligation are excised by polymerase. These processes require the participation of XRCC1, ADPRT and other proteins, such as DNA ligase III (LIG3)<sup>[10]</sup>. The NER pathway identifies and repairs bulky DNA lesions caused by environmental agents, such as ultraviolet light (UV) radiation and mutagenic chemicals<sup>[11]</sup>. In general, NER has four steps in the repair activity as follows: (a) damage recognition by a complex of bound proteins including XPC; (b) unwinding of the DNA by the TFIIH complex that includes XPD; (c) removal of the damaged single-stranded fragment by molecules including an ERCC1/XPF complex; and (d) synthesis by DNA polymerases<sup>[12]</sup>. The DSBR pathway involves two principal pathways, namely homologous recombination (HR) and non-homologous end joining (NHEJ). HR repairs DSBs with a template, such

as a sister chromatid or homologous chromosome, whereas NHEJ (including XRCC4, XRCC5, XRCC6, and LIG4) does not require any repair template at all<sup>[13]</sup>. HR can result in deletions and rearrangements, which involves many molecules, including XRCC2, XRCC3, and RAD51. The MMR pathway corrects the errors that occur during DNA replication and recombination and maintains DNA homeostasis. Loss of MMR function results in the accumulation of potential mutations and genetic instability<sup>[14]</sup>. The major components involved in this pathway include MLH1, MSH2, MSH6, and PMS2. The direct reversal of DNA damage can activate O<sup>6</sup>-methylguanine methyltransferase (MGMT), which removes a methyl group from the O<sup>6</sup>-position in guanine and transfers it to its own cysteine residue at codon 145 in the protein<sup>[15]</sup>.

It has been shown that a DNA repair phenotype is genetically determined and thus varies in the general population. For example, our studies have shown inter-individual variations of DNA repair capacity (DRC) in the general population and the effect of a suboptimal DRC on the risk of HNC<sup>[16,17]</sup>. One of the genetic variations is single nucleotide polymorphism (SNP), which may be located in coding regions of genes, non-coding sequences of genes, or in the intergenic regions between genes. SNPs in the coding region can be synonymous, where no amino acid substitution is observed, or non-synonymous, where the base substitution results in an alteration in the amino acid codes that could affect the structure and function of the proteins. For example, SNPs in the 3'- and 5'- untranslated regions (UTRs) of genes can alter the binding of transcriptional factors, microRNA, and mRNA stability<sup>[18]</sup>. In the past few years, DNA repair gene SNPs have been extensively studied in the etiology of HNC and found to be associated with HNC risk<sup>[19]</sup>. In this review, we summarize most of the reported SNPs in DNA repair genes (**Table 1**), which have been investigated in published epidemiologic studies of HNC.

## THE BER PATHWAY

### APE1

APE1, also known as APE, APEX, HAP1, and REF-1, plays a central role in the BER pathway by hydrolyzing the phosphodiester backbone immediately at 5' to the AP site. APE1 can also act as a 3'-phosphodiesterase to initiate repair of DNA single strand breaks that are produced either directly by reactive oxygen species or indirectly through the enzymatic removal of damaged bases<sup>[20,21]</sup>. A total of 123 SNPs in *APE1* have been reported (<http://www.ncbi.nlm.nih.gov/projects/SNP/>, as of Mar 1, 2013), 18 of which

are common and potentially functional<sup>[22]</sup>, but only -656T>G (rs1760944) and Asp148Glu (rs3136820) are the most extensively studied SNPs. Studies on the functional impact of the -656T>G SNP showed that this SNP influenced the transcriptional activity

of APE1 and contributed to lung cancer susceptibility<sup>[23,24]</sup>. As for the Asp148Glu SNP, functional studies suggested that the G variant allele was associated with increased mitotic delay after exposure to ionizing irradiation<sup>[25]</sup>. Additionally, Li et al.<sup>[26]</sup> studied the effects

**Table 1** Commonly reported SNPs in DNA repair genes implicated in head and neck cancer

Pathway	Reported SNPs <sup>a</sup>	Most studied SNPs	Ref. SNP IDs	Base changes
<b>Base excision repair</b>				
<i>APE1</i>	123	-656T>G	rs1760944	T>G
		Asp148Glu	rs3136820	T>G
<i>ADPRT</i>	1063	Val762Ala	rs1136410	T>C
<i>OGG1</i>	439	Ser326Gys	rs1052133	C>G
<i>XRCC1</i>	200	Arg194Trp	rs1799782	C>T
		Arg280His	rs25489	G>A
		Arg399Gln	rs25487	G>A
<i>LIG3</i>	449	-39G>A	rs2074522	G>A
		Arg780His	rs3136025	G>A
<i>PCNA</i>	269	Arg39Ser	rs1050525	A>C
<b>Nucleotide excision repair</b>				
<i>XPA</i>	461	23A>G	rs1800975	A>G
<i>XPC</i>	692	PAT	-	-/+
		Ala499Val	rs2228000	C>T
		Lys939Gln	rs2228001	A>C
<i>ERCC1</i>	437	Asn118Asn	rs11615	T>C
		IVS5 +33C>A	rs3212961	C>A
		8092C>A	rs3212986	C>A
<i>ERCC2/XPD</i>	560	Asp312Asn	rs1799793	G>A
		Lys751Gln	rs1052559	A>C
<i>ERCC4/XPF</i>	766	Arg415Gln	rs1800067	G>A
		Ser835Ser	rs1799801	C>T
<i>ERCC5/XPG</i>	742	Asp1104His	rs17655	G>C
<b>Mismatch repair</b>				
<i>MLH1</i>	1677	-93G>A	rs1800734	G>A
<i>MSH2</i>	2635	Gly322Asp	rs4987188	G>A
<i>MSH6</i>	1013	Gly39Glu	rs1042821	G>A
<i>PMS2</i>	945	Ser775Asn	rs1059060	G>A
<b>Double-strand break repair</b>				
<b>Homologous recombination</b>				
<i>XRCC2</i>	661	Arg188His	rs3218536	G>A
<i>XRCC3</i>	438	Thr241Met	rs861539	C>T
<i>RAD51</i>	737	135G>C	rs1801320	G>C
		172G>T	rs1801321	G>T
<i>NBS1</i>	1040	Glu185Gln	rs1805794	G>C
<b>Non-homologous end-joining</b>				
<i>XRCC4</i>	5014	Ser298Ala	rs3734091	A>C
<i>XRCC5</i>	1092	69506G>A	rs3835	G>A
<i>XRCC6</i>	843	-991T>C	rs5751129	T>C
		-61C>G	rs2267437	C>G
<i>LIG4</i>	323	Ile658Val	rs2232641	A>G
		Thr9Ile	rs1805388	G>C
		Asp568Asp	rs1805386	T>C
<b>Direct reversion repair</b>				
<i>MGMT</i>	5924	Leu84Phe	rs12917	C>T
		Ile143Val	rs2308321	C>T

<sup>a</sup><http://www.ncbi.nlm.nih.gov/projects/SNP/>, as of Mar 1, 2013.

of SNPs in *APE1*, *ADPRT*, and *XRCC1* on *SCCHN* risk in a hospital-based case-control study and found that no altered risk was associated with the *APE1* Asp148Glu SNP in *SCCHN*. In contrast, Matullo et al. found that the Asp148Glu variant allele was associated with a significantly reduced incidence of upper aero-digestive cancer (UADC), including HNC<sup>[9]</sup>. A meta-analysis of 28 case-control studies indicated that the Asp148Glu SNP may be a low-penetrance risk factor for cancer development<sup>[27]</sup>.

### ADPRT

ADPRT, also known as PARP1, is involved in DNA damage signaling, genomic stability of damaged cells, BER, recombination, and transcriptional regulation of tumor suppressor genes<sup>[28]</sup>. Accordingly, activation of ADPRT has been shown in a variety of diseases<sup>[29]</sup>. It has been reported that 1063 SNPs have been identified in this gene (<http://www.ncbi.nlm.nih.gov/projects/SNP/>), but a few are common and potentially functional. A Val762Ala SNP (rs1136410) in *ADPRT* was shown to contribute to significantly lower poly(ADP-ribosyl)ation activities in a dose-dependent manner in response to oxidative damage<sup>[30]</sup>. In an association study, Li et al. found some evidence of an interaction between the *ADPRT* 762Ala SNP and variants of other BER genes in the etiology of *SCCHN*, although this SNP was not found to be associated with risk of other cancers<sup>[26]</sup>. These discrepant findings across studies may reflect a cancer-specific effect in the etiology of cancers. In addition, previous studies have also indicated that ADPRT expression levels appear to be low in breast cancer<sup>[31]</sup>, but high in endometrial cancer<sup>[32]</sup>, which appears to be validated by recently pooled results<sup>[33]</sup>.

### OGG1

The human OGG1 (hOGG1) catalyzes the cleavage of the glycosylic bonds between the modified base and the sugar moiety, leaving an abasic AP site in DNA; the resultant AP site is then incised, and the repair is completed by successive actions of a phosphodiesterase, a DNA polymerase, and a DNA ligase<sup>[34]</sup>. Among the 439 SNPs identified to date (<http://www.ncbi.nlm.nih.gov/projects/SNP/>), the most extensively studied Ser326Cys SNP (rs1052133) is located in exon 7 of *hOGG1*, resulting in an amino acid substitution of serine with cysteine at codon 326. Studies on the functional impact of the *hOGG1* Ser326Cys SNP yielded inconsistent results. Many studies reported that no difference was observed in catalytic activities by the Ser326Cys genotype<sup>[35-37]</sup>. However, other studies revealed that the Cys allele

exhibited reduced DNA repair activity and was associated with risk of cancers<sup>[38]</sup>. The role of the Ser326Cys SNP and HNC risk was investigated in six studies<sup>[39-44]</sup>. The first case-control study, with a small sample size (167 cases and 331 controls), reported a positive association with HNC risk<sup>[44]</sup> which was confirmed by Pawlowska et al.<sup>[40]</sup>. Moreover, Sliwinski et al. showed that the Ser326Cys SNP might modify the risk of *SCCHN* associated with smoking<sup>[43]</sup>. Nevertheless, Zhang et al. conducted a hospital-based study and found no association between the Ser326Cys SNP and HNC risk, which may be due to the possibility of selection bias of subjects<sup>[41]</sup>. The discrepant findings in these studies may reflect the differences in genetic background among different ethnic groups under the investigation, carcinogen exposure in different populations, and limited study power as indicated by study sample sizes.

### XRCC1

*XRCC1*, as a scaffold protein for other BER proteins, recognizes single-strand breaks. Although more than 200 SNPs in *XRCC1* have been identified (<http://www.ncbi.nlm.nih.gov/projects/SNP/>), only three common SNPs have been widely investigated in cancer risk, which are Arg194Trp (rs1799782), Arg280His (rs25489), and Arg399Gln (rs25487), located in exon 6, 9, and 10, respectively. The Arg399Gln SNP is the most frequently investigated SNP of the BER genes. Several studies reported that the Arg399Gln SNP was associated with risk of various cancers<sup>[45-53]</sup>. A meta-analysis conducted by Hu et al. found that the Arg194Trp SNP was associated with a decreased risk of cancer, whereas the Arg280His SNP was associated with an increased risk of cancer<sup>[54]</sup>. Huang et al. performed a pooled analysis for an association between the Arg399Gln SNP and HNC risk. They found that the *XRCC1* 399Gln/Gln genotype was associated with a decreased risk among Caucasians<sup>[50]</sup>, which was supported by other reports that this SNP was associated with measurably reduced DNA repair capacity, as detected by the persistent DNA adducts and elevated levels of sister chromatid exchanges<sup>[55,56]</sup>. However, Li et al.<sup>[26]</sup> failed to identify any association between the Arg399Gln SNP on the risk of *SCCHN* in a moderately large sample size. They further performed a meta-analysis and found that the Arg399Gln SNP was significantly associated with HNC risk among Asians but not among Caucasians, suggesting that this SNP has an effect or a higher penetrance in Asian populations. As for the Arg194Trp SNP, many studies reported detectable effects on DNA-adduct levels, frequency of mutations, and sensitiv-

ity to ionizing radiation<sup>[25,57-59]</sup>. However, the function of the Arg280His SNP is still not fully understood. This SNP could produce an amino acid change and decrease genomic stability<sup>[60]</sup>. A recent meta-analysis suggested that the Arg280His SNP may play a role in cancer development, though only one study on HNC was included in the analysis<sup>[61]</sup>. Inconsistent with the previous findings, Applebaum et al. reported that the *XRCC1* polymorphisms might confer susceptibility to SCCHN in the context of smoking, among those who are seronegative for HPV16<sup>[62]</sup>.

### LIG3 and PCNA

*LIG3* encodes ligase III, which is DNA- and ATP-dependent, an essential component in the BER pathway in the repair of damages caused by free radicals<sup>[63]</sup> and, the levels of damage was found to be also directly associated with the expression levels of *XRCC1*<sup>[64]</sup>. Totally, 449 SNPs in *LIG3* and 269 SNPs in *PCNA* have been reported (<http://www.ncbi.nlm.nih.gov/projects/SNP/>). Although many SNPs in *LIG3* and *PCNA* genes were reported to have been studied in many cancers, such as rs3136025 G>A and rs2074522 G>A in *LIG3*, and rs1050525 in *PCNA*, no significant associations were found<sup>[65-70]</sup>. Therefore, the associations of *LIG3* and *PCNA* SNPs with HNC risk are still not fully understood.

## THE NER PATHWAY

### XPA

The *XPA* gene encodes a zinc-finger DNA-binding protein that participates in DNA excision repair to maintain genomic integrity. The XPA protein is also involved in both global genome and transcription-coupled repair pathways<sup>[71]</sup>. Of 461 SNPs reported to date, a common 23A>G SNP (rs1800975) in the 5'-UTR has been widely studied, although the functional significance of this SNP is still unknown. It has been demonstrated that the 5'-UTR may regulate gene expression through transcriptional and post-transcriptional control mechanisms, that this SNP may affect the binding between translational factors and the promoter, and that individuals with the G variant had a reduced risk of lung cancer in Caucasian<sup>[72]</sup> and Korean populations<sup>[73]</sup>. Previously, five case-control studies with small sample sizes had reported the association between the 23A>G SNP and HNC risk<sup>[74-77]</sup>, and the results were inconclusive, partially because of the possible weak effect of the SNP on HNC and the relatively small size in each of the published studies. For example, Sugimura et al. conducted a case-control study with 122 oral squamous cell carcinoma (OSCC)

cases and 241 controls and found that the 23A>G SNP was associated with OSCC risk<sup>[74]</sup>. Another study found similar results, but also the synergistic effects of *XPA* and *XPB* SNPs, as well as smoking status, on oral cancer risk<sup>[75]</sup>. In contrast, Jelonek et al. did not find any association between this SNP and risk of HNC<sup>[77]</sup>. Later, a meta-analysis on all eligible case-control studies between the 23A>G SNP and HNC risk demonstrated that the G allele was a low-penetrant risk factor for cancer development<sup>[78]</sup>.

### XPC

The *XPC* gene encodes part of the XPC-HR23B complex that plays a role in the early step of NER by initially detecting the DNA damage. The complex can bind to damaged DNA and change the DNA conformation around the lesion<sup>[79]</sup>. Among all identified 692 SNPs of *XPC* (<http://www.ncbi.nlm.nih.gov/projects/SNP/>), three common biallelic poly (AT) insertion/deletion (PAT, -/+), Ala499Val (rs2228000), and Lys939Gln (rs2228001) have been extensively studied. The *XPC*-PAT polymorphism, located in intron 9, was found to be in linkage disequilibrium with the Ala499Val SNP<sup>[80]</sup>, which has been reported to be associated with a decreased DNA repair capacity as measured by chromatid aberrations [81]. However, the functionality of the Lys939Gln SNP is still unknown. Shen et al. firstly reported that the *XPC*-PAT + allele was associated with the risk of SCCHN in a small study (287 cases and 311 controls)<sup>[82]</sup>. However, in their subsequent large case-control study, they found that only the XPC Ala499Val SNP was associated with a significantly increased risk of SCCHN<sup>[83]</sup>. Although Yang et al. found that XPC expression may influence SCCHN risk, no effect of *XPC*-PAT genotypes on SCCHN risk was observed in a Korean population<sup>[84]</sup>. These findings should be validated in larger studies with the same ethnic background.

### ERCC1

ERCC1 is essential for the NER pathway and plays a critical role in genomic stability. SNPs in *ERCC1* have been reported to contribute to susceptibility to carcinogenesis<sup>[85]</sup>. Among the 437 SNPs reported (<http://www.ncbi.nlm.nih.gov/projects/SNP/>), at least three common *ERCC1* SNPs, namely Asn118Asn (rs11615), IVS5 + 33C>A (rs3212961), and 8092>A (rs321986), are probably functional. Two meta-analyses revealed that Asn118Asn and IVS5 +33C>A SNPs, but not 8092C>A, are low-penetrance risk factors for cancer risk<sup>[86,87]</sup>. As for HNC, most studies focused on 8092C>A SNP and HNC risk<sup>[74,76,83,88-90]</sup>. In a small study with 67 SCCHN cases and 73 controls,

Yang et al. studied the association between mRNA expression and the 8092C>A SNP of *ERCC1*, and found lower *ERCC1* mRNA expression in SCCHN patients than in controls, but no association of the 8092C>A SNP with the risk of SCCHN or expression of *ERCC1* mRNA<sup>[88]</sup>. In a larger study, Sturgis et al. found that the 8092C>A SNP combined with the *XPD* Asp312Asn SNP may contribute to the risk of SCCHN<sup>[89]</sup>. However, the results have not been replicated by An et al. in a much large hospital-based case-control study<sup>[83]</sup>. In the stratification of tumor site in the meta-analysis by Zhang et al., no significant risk associated with the 8092C>A SNP was found for SCCHN in different genetic models<sup>[86]</sup>.

### ERCC2/XPD

XPD is an enzyme in the NER pathway that removes certain DNA cross-links, UV photo-lesions, and bulky chemical adducts<sup>[91]</sup>. Of 560 SNPs reported to date (<http://www.ncbi.nlm.nih.gov/projects/SNP/>), two *XPD* polymorphic loci that are of most interest in molecular epidemiology studies are Asp312Asn (rs1799793) and Lys751Gln (rs13181). It has been suggested that the *XPD* 312 Asn variant allele may be associated with reduced repair of aromatic DNA adducts<sup>[92]</sup>. In Lys751Gln SNP, glutamine is changed to lysine at codon 751, which has been reported to have a reduced DNA repair proficiency<sup>[93]</sup>. Several studies have investigated the associations of *XPD* Asp312Asn and Lys751Gln with HNC risk but generated inconsistent results<sup>[9,76,77,83,89,94-97]</sup>. Sturgis et al. firstly studied the association between the Asp312Asn SNP and HNC risk and found that this SNP had a significant association with HNC risk when combined with the *ERCC1* 8092C>A SNP<sup>[89]</sup>. Other studies also showed that HNC risk was not significantly associated with the Asp312Asn SNP<sup>[9,76,77,95,96]</sup>. A recent comprehensive meta-analysis of nine case-control studies on *XPD* Asp312Asn SNP and HNC risk reported that the Asp312Asn SNP might not be associated with HNC risk<sup>[98]</sup>. As for the Lys751Gln SNP, Yuan et al. performed a pooled analysis of 15 case-control studies to assess the overall association between this SNP and HNC risk and found that a significantly elevated HNC risk was associated with Lys751Gln SNP<sup>[99]</sup>, and this increased risk was more pronounced among Europeans but not among Asians.

### ERCC4/XPF

ERCC4 is a key enzyme responsible for excising bulky adducts from damaged DNA because of its unique function in damage site recognition. ERCC4 contains the catalytic domain of the nuclease, and

ERCC1 is required for DNA binding and stabilization of ERCC4<sup>[8]</sup>. To date, 766 SNPs in *ERCC4* have been reported (<http://www.ncbi.nlm.nih.gov/projects/SNP/>) and subsequent association studies found that some of them are likely to contribute to susceptibility to several kinds of cancers, including HNC, but only two common SNPs Arg415Gln (rs1800067) and Ser835Ser (rs1799801) in the exons of *ERCC4* have been extensively studied in the epidemiologic investigations<sup>[76,83,100]</sup>. Although the results from these three reported studies are not consistent, a recent meta-analysis indicated that these two SNPs were not associated with HNC risk. Some publication bias was observed in the meta-analysis, because the results were not adjusted for other covariants, including age, smoking status, drinking status, and other environmental factors<sup>[101]</sup>.

### ERCC5/XPG

ERCC5 has structure-specific endonuclease activity that is essential for two incision steps in NER<sup>[102]</sup>. Of 742 SNPs reported to date (<http://www.ncbi.nlm.nih.gov/projects/SNP/>), the Asp1104 His SNP (rs17655), located in exon 15 of *XPG*, has been largely studied for susceptibility to HNC. Although the His1104Asp SNP together with SNPs of several other NER genes may jointly contribute to the variability of the DRC phenotype, all published data did not show any significant association with risk of HNC<sup>[76,103-105]</sup>.

## THE MMR PATHWAY

### MLH1 and MSH2

MLH1 and MSH2 are the key components of the MMR system that participates in the recognition of nucleotide mismatches occurring during DNA replication and in the recruitment of additional repair proteins to correct replication errors. Numerous SNPs have been identified in *MLH1* and *MSH2* (1677 and 2635, respectively, <http://www.ncbi.nlm.nih.gov/projects/SNP/>), but few of which have been investigated for their associations with cancer. For example, the *MLH1* -93G>A SNP (rs1800734) is located in the promoter region that is responsible for maximal transcriptional activity of this gene<sup>[106,107]</sup>. The association between the -93G>A SNP and cancer risk was investigated most in colorectal cancer<sup>[108,109]</sup>. Reduced expression levels of MSH2 have been reported in hereditary nonpolyposis colon cancer (HNPCC) and some other human cancers<sup>[110-113]</sup>. The Gly322Asp SNP is located in the coding region of the *MSH2* gene, resulting in a modest decrease in mismatch repair efficiency<sup>[114,115]</sup>. However, no study on the association of both the -93G>A and Gly322Asp SNPs with HNC risk has been reported.

## MSH6 and PMS2

The MMR proteins MSH6 and PMS2 have been shown to interact with MSH2 and MLH1, respectively<sup>[116]</sup>, and the alteration in the expression of MSH6 can perturb MMR, while the protein encoded by *PMS2* plays an essential role in repairing DNA by forming an active protein complex with MLH1 that interacts with MSH2 bound to mismatched bases. Although many SNPs have been identified in these genes (1013 for MSH6 and 945 for PMS2, <http://www.ncbi.nlm.nih.gov/projects/SNP/>), inherited mutations are the most studied topics for colorectal cancer. The mutations of *PMS2* were firstly found in a HNPCC patient<sup>[117]</sup>, and inherited mutations in *MSH2* and *MLH1* are the most prevalent cause of colorectal tumors or HNPCC, as these tumors typically have microsatellites, a marker for malfunction of MMR<sup>[118]</sup>, which were not common in HNC.

## THE DSB REPAIR PATHWAY

### Homologous recombination (HR)

#### *XRCC2*

The *XRCC2* gene, located at 7q36.1, is an essential part of the HR repair pathway and also a functional candidate for its involvement in cancer progression, including HNC. Among the 661 SNPs reported to date (<http://www.ncbi.nlm.nih.gov/projects/SNP/>), a common SNP Arg188His (rs3218536) in *XRCC2* has been reported in epidemiological studies. Benhamou et al. found a significant association between the Arg188His SNP and risk of upper aerodigestive tract (UADT) cancers<sup>[119]</sup>; in particular, the 188His allele was associated with a significantly increased risk of pharyngeal cancer. Furthermore, this SNP was reported to be associated with an increased risk of oral cancer risk<sup>[120]</sup>. In contrast, the Arg188His and three other SNPs in the promoter region have been associated with reduced risk in other cancers, such as bladder cancer<sup>[121]</sup>.

#### *XRCC3*

*XRCC3*, involved in the HR of DNA DSB and cross-links, is a member of an emerging family of RAD51-related proteins that likely participate in HR to maintain genomic stability and repair DNA damage<sup>[122]</sup>. *XRCC3* deficient cell lines display reduced HR. Of the 438 SNPs reported to date (<http://www.ncbi.nlm.nih.gov/projects/SNP/>), the main Thr241Met (rs861539) SNP in *XRCC3* leads to an amino acid substitution at codon 241, which has been reported to be associated with slightly but not significantly de-

creased DNA repair capacity<sup>[123]</sup>. Recently, a meta-analysis of 15 eligible studies with 3,191 cases and 5,090 controls found some evidence for an association between the Thr241Met SNP and HNC risk<sup>[124]</sup>. However, another meta-analysis by Flores-Obando et al. revealed that the SNP did not contribute to the risk of HNC<sup>[125]</sup>. These two different results may be due to the number of studies included in the analyses (15 vs. 10) with various study powers.

#### *RAD51*

RAD51 is involved in HR and DSB in DNA and DNA cross-links and for the maintenance of chromosome stability<sup>[126]</sup>. RAD51 is associated with *BRCA1* and *BRCA2* tumor suppressor gene products, suggesting that a defect in recombination may lead to tumor development<sup>[127]</sup>. Of the 737 SNPs reported to date (<http://www.ncbi.nlm.nih.gov/projects/SNP/>), two SNPs in the 5'-UTR of *RAD51*, 135G>C (rs1801320) and 172G>T rs1801321, have been investigated for their associations with the risk of sporadic breast cancer or breast cancer in non-*BRCA1/2* carriers, though the results are yet controversial and inconclusive<sup>[128]</sup>. There are only few studies investigating the association of the 135G>C SNP as a risk factor for HNC so far, but they reported inconsistent findings of either increased, decreased or lack of any HNC risk associated with the 135C allele<sup>[129,130]</sup>. A small study of 81 HNC cases and 111 controls also suggested that no significant effect of this SNP and HNC risk was observed, which was modulated by smoking status and gene-gene interactions<sup>[131]</sup>. Moreover, Lu et al. observed that the 172G>T SNP was associated with significantly reduced risk of SCCHN<sup>[132]</sup>, which was confirmed by Gresner et al.<sup>[131]</sup>. A possible mechanism underlying the protective effect of the 172G>T SNP in HNC may be due to increased cellular apoptotic potential, or the capacity of the DNA DSB repair system.

#### *NBS1*

NBS1 is part of a protein complex that forms in response to DNA damage to maintain chromosomal integrity. The exact role of NBS1 in DNA repair is not fully understood, because NBS1 does not have a DNA binding site or kinase activity. As one of the most commonly studied SNPs, among the 1040 reported SNPs (<http://www.ncbi.nlm.nih.gov/projects/SNP/>), the Glu185Gln SNP is non-synonymous, with an amino acid change that may affect the function of NBS1 and the protein-protein interaction<sup>[133]</sup>. Zheng et al. observed a significant difference in genotype frequencies of the Glu185Gln SNP between nasophar-

ryngeal carcinoma cases and controls and found that the C allele was associated with the risk of invasive or metastatic disease, compared with the G allele<sup>[134]</sup>.

## Non-homologous end-joining (NHEJ)

### *XRCC4*

*XRCC4*, a key component of the NHEJ repair pathway, is found to restore DNA DSB repair. It is basically required for precise end-joining of blunt DNA DSBs. There are 5014 SNPs that have been identified to date (<http://www.ncbi.nlm.nih.gov/projects/SNP/>). Of these SNPs, Tseng et al. reported that the Ser298Ala (rs3734091) SNP in exon 3 of *XRCC4* was associated with the risk of oral cancer in Taiwanese populations<sup>[135]</sup> and found that individuals with 247A allele had a 2.04-fold higher risk of cancer. In addition, Chiu et al. firstly observed that an intron 3 del genotype (rs28360071) may be associated with oral cancer in Taiwanese patients<sup>[136]</sup>.

### *XRCC5 and XRCC6*

*XRCC5* (also named Ku80 or Ku86) and *XRCC6* (also known Ku70), are the components of the major repair pathway of DSBs in eukaryotic cells during most phases of the cell cycle, particularly the G0/G1 phases<sup>[137]</sup>. These proteins are essential for the maintenance of chromosomal integrity and the viability of cell and organism, as NHEJ plays a more predominant role in adult mammals than the alternative DSB repair mechanism, homologous recombination<sup>[138,139]</sup>. There are 1092 and 843 SNPs that have been identified to date for *XRCC5* and *XRCC6*, respectively (<http://www.ncbi.nlm.nih.gov/projects/SNP/>). A small study (152 SCCHN patients and 157 controls) found that the genotypes of rs3835 in *XRCC5* and rs2267437 in *XRCC6* were neither associated with SCCHN risk, nor was gene-smoking or gene-alcohol interaction observed<sup>[130]</sup>. Bau et al. found that the *XRCC6* promoter SNP rs5751129 was associated with oral cancer risk, but not rs2267437, rs132770, or rs132774<sup>[140]</sup>. The same group reported that the *XRCC5* rs828907, but not rs11685387 or rs9288518, was associated with oral cancer susceptibility<sup>[141]</sup>. In addition, Yang et al. investigated subjects with oral premalignant lesions in a case-control study and found no association between the *XRCC5* rs1051685 SNP and the risk of oral cancer<sup>[142]</sup>.

### *LIG4*

*LIG4* jointed the final rejoining step of NHEJ with *XRCC4*, and its deficiency leads to significant genomic instability<sup>[143]</sup>. Among the 323 SNPs reported

to date (<http://www.ncbi.nlm.nih.gov/projects/SNP/>), an Ile658Val (rs2232641) SNP in the exon of *LIG4*, affecting *LIG4* function, has been found to be slightly associated with the developing of cervical carcinoma in a Chinese population<sup>[144]</sup>. However, no published studies have investigated the association between the Ile658Val SNP and HNC risk. Werbrouck et al. investigated the role of another two SNPs, Thr9Ile (rs1805388) and Asp568Asp (rs1805386), in SCCHN development and found that the Thr9Ile SNP was associated with a significantly reduced risk for SCCHN<sup>[130]</sup>.

## THE DR PATHWAY

### MGMT

The only known enzyme in the DR pathway is *MGMT* that removes a methyl group from the O6-position in guanine and transfers it to its own cysteine residue at codon 145 in the protein, inactivating the *MGMT* protein itself while repairing guanine. It is reported that the *MGMT* knockout mice had a higher incidence of nitrosamine-induced tumorigenesis<sup>[145]</sup>. Of the SNPs reported to date (<http://www.ncbi.nlm.nih.gov/projects/SNP/>), two functional common SNPs, Leu84Phe (rs12917) and Ile143Val (rs2308321), have been extensively investigated for their roles in HNC. Recently, Zhang et al. extended the SNPs in *MGMT* including the SNPs in the promoter region and found that none of SNPs in *MGMT* had a substantial effect on SCCHN risk, but a joint effect of several *MGMT* variants may contribute to risk and progression of SCCHN<sup>[146]</sup>. Furthermore, they did a mini-pooled analysis of previously published studies on the Leu84Phe SNP and HNC risk<sup>[50,147,148]</sup> and found that this single variant was not associated with risk of SCCHN.

### GWAS IN HNC

During the past few years, GWASs have identified a large number of robust associations between specific chromosomal loci and complex human diseases<sup>[149]</sup>. For HNC, a recent GWAS identified two novel variants (rs1494961 at 4q21 and rs4767364 at 12q24) and three remaining variants (rs1573496, rs1229984, and rs698) located in the *ADH* gene cluster that is associated with risk of UADT cancers in Caucasians<sup>[150]</sup>. Yuan et al. performed a validation study of GWAS findings for their associations with HNC risk in a hospital-based case-control study and found that rs1229984 at 4q23 was associated with significantly increased risk of HNC in a Chinese population<sup>[151]</sup>. However, the biological relevancy of



GWAS discoveries is still largely unknown. Many of the disease associations identified by GWAS do not involve previously known candidate genes, such as DNA repair pathway genes, and many associated markers are in genomic regions harboring unknown genes<sup>[152]</sup>. Therefore, the functions of those variants identified in GWAS should be validated via biological assays, pathway analysis or bioinformatics in the post-GWAS era.

## CONCLUSION

DNA repair mechanisms play an important role in correcting DNA damage and preserving genomic integrity, and it is obvious that abnormalities of these mechanisms are strongly associated with cancer risk, including HNC. Specifically, genetic variants in DNA repair genes (**Table 1**) are associated with alteration of an individual's susceptibility to malignancy. Here, we have summarized most of previously published studies that have investigated associations between SNPs in DNA repair genes and HNC risk. It is interesting that the only GWAS performed in Caucasian populations did not identify any DNA repair gene SNPs as the top-hits. More GWAS of HNC in different ethnic groups are needed to provide the whole picture of the importance of the SNPs in these genes. Many association studies have focused on the roles of those common and functional SNPs in the BER, NER, MMR, DSBR, and DR pathways. Among the above-reviewed genes, little is known about the prognosis and therapeutic values in HNC. Therefore, further investigations of alterations in DNA repair pathways are needed, which are essential to identify their clinical application in HNC.

## References

- [1] Parkin DM, Pisani P, Ferlay J. Global cancer statistics. *CA Cancer J Clin* 1999; 49: 33-64, 1.
- [2] Siegel R, Naishadham D, Jemal A. Cancer statistics, 2012. *CA Cancer J Clin* 2012; 62: 10-29.
- [3] Hashibe M, Boffetta P, Zaridze D, Shangina O, Szeszenia-Dabrowska N, Mates D, et al. Evidence for an important role of alcohol- and aldehyde-metabolizing genes in cancers of the upper aerodigestive tract. *Cancer Epidemiol Biomarkers Prev* 2006; 15: 696-703.
- [4] Boffetta P, Hecht S, Gray N, Gupta P, Straif K. Smokeless tobacco and cancer. *Lancet Oncol* 2008; 9: 667-75.
- [5] Fakhry C, Westra WH, Li S, Cmelak A, Ridge JA, Pinto H, et al. Improved survival of patients with human papillomavirus-positive head and neck squamous cell carcinoma in a prospective clinical trial. *J Natl Cancer Inst* 2008; 100: 261-9.
- [6] Chaturvedi AK, Engels EA, Pfeiffer RM, Hernandez BY, Xiao W, Kim E, et al. Human papillomavirus and rising oropharyngeal cancer incidence in the United States. *J Clin Oncol* 2011; 29: 4294-301.
- [7] Negri E, Boffetta P, Berthiller J, Castellsague X, Curado MP, Dal Maso L, et al. Family history of cancer: pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *Int J Cancer* 2009; 124: 394-401.
- [8] Wood RD, Mitchell M, Sgouros J, Lindahl T. Human DNA repair genes. *Science* 2001; 291: 1284-9.
- [9] Matullo G, Dunning AM, Guarrera S, Baynes C, Polidoro S, Garte S, et al. DNA repair polymorphisms and cancer risk in non-smokers in a cohort study. *Carcinogenesis* 2006; 27: 997-1007.
- [10] Lu AL, Li X, Gu Y, Wright PM, Chang DY. Repair of oxidative DNA damage: mechanisms and functions. *Cell Biochem Biophys* 2001; 35: 141-70.
- [11] Leibel D, Laspe P, Emmert S. Nucleotide excision repair and cancer. *J Mol Histol* 2006; 37: 225-38.
- [12] Friedberg EC. How nucleotide excision repair protects against cancer. *Nat Rev Cancer* 2001; 1: 22-33.
- [13] Khanna KK, Jackson SP. DNA double-strand breaks: signaling, repair and the cancer connection. *Nat Genet* 2001; 27: 247-54.
- [14] Mohrenweiser HW, Wilson DM, 3rd, Jones IM. Challenges and complexities in estimating both the functional impact and the disease risk associated with the extensive genetic variation in human DNA repair genes. *Mutat Res* 2003; 526: 93-125.
- [15] Pegg AE, Byers TL. Repair of DNA containing O6-alkylguanine. *FASEB J* 1992; 6: 2302-10.
- [16] Cheng L, Eicher SA, Guo Z, Hong WK, Spitz MR, Wei Q. Reduced DNA repair capacity in head and neck cancer patients. *Cancer Epidemiol Biomarkers Prev* 1998; 7: 465-8.
- [17] Wang LE, Hu Z, Sturgis EM, Spitz MR, Strom SS, Amos CI, et al. Reduced DNA repair capacity for removing tobacco carcinogen-induced DNA adducts contributes to risk of head and neck cancer but not tumor characteristics. *Clin Cancer Res* 2010; 16: 764-74.
- [18] Sachidanandam R, Weissman D, Schmidt SC, Kakol JM, Stein LD, Marth G, et al. A map of human genome sequence variation containing 1.42 million single nucleotide polymorphisms. *Nature* 2001; 409: 928-33.
- [19] Ford BN, Ruttan CC, Kyle VL, Brackley ME, Glickman BW. Identification of single nucleotide polymorphisms in human DNA repair genes. *Carcinogenesis* 2000; 21: 1977-81.
- [20] Izumi T, Hazra TK, Boldogh I, Tomkinson AE, Park MS, Ikeda S, et al. Requirement for human AP endonuclease 1 for repair of 3'-blocking damage at DNA single-strand breaks induced by reactive oxygen species. *Carcinogenesis* 2000; 21: 1329-34.
- [21] Fortini P, Pascucci B, Parlanti E, D'Errico M, Simonelli V, Dogliotti E. The base excision repair: mechanisms and its relevance for cancer susceptibility. *Biochimie* 2003; 85: 1053-71.
- [22] Xi T, Jones IM, Mohrenweiser HW. Many amino acid

- substitution variants identified in DNA repair genes during human population screenings are predicted to impact protein function. *Genomics* 2004; 83: 970-9.
- [23] Lu J, Zhang S, Chen D, Wang H, Wu W, Wang X, et al. Functional characterization of a promoter polymorphism in APE1/Ref-1 that contributes to reduced lung cancer susceptibility. *FASEB J* 2009; 23: 3459-69.
- [24] Lo YL, Jou YS, Hsiao CF, Chang GC, Tsai YH, Su WC, et al. A polymorphism in the APE1 gene promoter is associated with lung cancer risk. *Cancer Epidemiol Biomarkers Prev* 2009; 18: 223-9.
- [25] Hu JJ, Smith TR, Miller MS, Mohrenweiser HW, Golden A, Case LD. Amino acid substitution variants of APE1 and XRCC1 genes associated with ionizing radiation sensitivity. *Carcinogenesis* 2001; 22: 917-22.
- [26] Li C, Hu Z, Lu J, Liu Z, Wang LE, El-Naggar AK, et al. Genetic polymorphisms in DNA base-excision repair genes ADPRT, XRCC1, and APE1 and the risk of squamous cell carcinoma of the head and neck. *Cancer* 2007; 110: 867-75.
- [27] Gu D, Wang M, Zhang Z, Chen J. The DNA repair gene APE1 T1349G polymorphism and cancer risk: a meta-analysis of 27 case-control studies. *Mutagenesis* 2009; 24: 507-12.
- [28] Dantzer F, Schreiber V, Niedergang C, Trucco C, Flatte E, De La Rubia G, et al. Involvement of poly(ADP-ribose) polymerase in base excision repair. *Biochimie* 1999; 81: 69-75.
- [29] Masutani M, Nakagama H, Sugimura T. Poly(ADP-ribosylation) in relation to cancer and autoimmune disease. *Cell Mol Life Sci* 2005; 62: 769-83.
- [30] Lockett KL, Hall MC, Xu J, Zheng SL, Berwick M, Chuang SC, et al. The ADPRT V762A genetic variant contributes to prostate cancer susceptibility and deficient enzyme function. *Cancer Res* 2004; 64: 6344-8.
- [31] Bieche I, de Murcia G, Lidereau R. Poly(ADP-ribose) polymerase gene expression status and genomic instability in human breast cancer. *Clin Cancer Res* 1996; 2: 1163-7.
- [32] Ghabreau L, Roux JP, Frappart PO, Mathevet P, Patricot LM, Mokni M, et al. Poly(ADP-ribose) polymerase-1, a novel partner of progesterone receptors in endometrial cancer and its precursors. *Int J Cancer* 2004; 109: 317-21.
- [33] Pabalan N, Francisco-Pabalan O, Jarjanazi H, Li H, Sung L, Ozcelik H. Racial and tissue-specific cancer risk associated with PARP1 (ADPRT) Val762Ala polymorphism: a meta-analysis. *Mol Biol Rep* 2012; 39: 11061-72.
- [34] Dianov GL, Souza-Pinto N, Nyaga SG, Thybo T, Stevnsner T, Bohr VA. Base excision repair in nuclear and mitochondrial DNA. *Prog Nucleic Acid Res Mol Biol* 2001; 68: 285-97.
- [35] Dherin C, Radicella JP, Dizdaroglu M, Boiteux S. Excision of oxidatively damaged DNA bases by the human alpha-hOgg1 protein and the polymorphic alpha-hOgg1(Ser326Cys) protein which is frequently found in human populations. *Nucleic Acids Res* 1999; 27: 4001-7.
- [36] Park YJ, Choi EY, Choi JY, Park JG, You HJ, Chung MH. Genetic changes of hOGG1 and the activity of oh-8Gua glycosylase in colon cancer. *Eur J Cancer* 2001; 37: 340-6.
- [37] Blons H, Radicella JP, Laccourreye O, Brasnu D, Beaune P, Boiteux S, et al. Frequent allelic loss at chromosome 3p distinct from genetic alterations of the 8-oxoguanine DNA glycosylase 1 gene in head and neck cancer. *Mol Carcinog* 1999; 26: 254-60.
- [38] Weiss JM, Goode EL, Ladiges WC, Ulrich CM. Polymorphic variation in hOGG1 and risk of cancer: a review of the functional and epidemiologic literature. *Mol Carcinog* 2005; 42: 127-41.
- [39] Cho EY, Hildesheim A, Chen CJ, Hsu MM, Chen IH, Mittl BF, et al. Nasopharyngeal carcinoma and genetic polymorphisms of DNA repair enzymes XRCC1 and hOGG1. *Cancer Epidemiol Biomarkers Prev* 2003; 12: 1100-4.
- [40] Pawlowska E, Janik-Papis K, Rydzanicz M, Zuk K, Kaczmarczyk D, Olszewski J, et al. The Cys326 allele of the 8-oxoguanine DNA N-glycosylase 1 gene as a risk factor in smoking- and drinking-associated larynx cancer. *Tohoku J Exp Med* 2009; 219: 269-75.
- [41] Zhang Z, Shi Q, Wang LE, Sturgis EM, Spitz MR, El-Naggar AK, et al. No Association between hOGG1 Ser326Cys polymorphism and risk of squamous cell carcinoma of the head and neck. *Cancer Epidemiol Biomarkers Prev* 2004; 13: 1081-3.
- [42] Wikman H, Risch A, Klimek F, Schmezer P, Spiegelhalter B, Dienemann H, et al. hOGG1 polymorphism and loss of heterozygosity (LOH): significance for lung cancer susceptibility in a caucasian population. *Int J Cancer* 2000; 88: 932-7.
- [43] Sliwinski T, Przybylowska K, Markiewicz L, Rusin P, Pietruszewska W, Zelinska-Blizniewska H, et al. MUYH Tyr165Cys, OGG1 Ser326Cys and XPD Lys751Gln polymorphisms and head neck cancer susceptibility: a case control study. *Mol Biol Rep* 2011; 38: 1251-61.
- [44] Elahi A, Zheng Z, Park J, Eyring K, McCaffrey T, Lazarus P. The human OGG1 DNA repair enzyme and its association with orolaryngeal cancer risk. *Carcinogenesis* 2002; 23: 1229-34.
- [45] Kowalski M, Przybylowska K, Rusin P, Olszewski J, Morawiec-Sztandera A, Bielecka-Kowalska A, et al. Genetic polymorphisms in DNA base excision repair gene XRCC1 and the risk of squamous cell carcinoma of the head and neck. *J Exp Clin Cancer Res* 2009; 28: 37.
- [46] Kang SY, Lee KG, Lee W, Shim JY, Ji SI, Chung KW, et al. Polymorphisms in the DNA repair gene XRCC1 associated with basal cell carcinoma and squamous cell carcinoma of the skin in a Korean population. *Cancer Sci* 2007; 98: 716-20.
- [47] Sturgis EM, Castillo EJ, Li L, Zheng R, Eicher SA, Clayman GL, et al. Polymorphisms of DNA repair gene XRCC1 in squamous cell carcinoma of the head and neck. *Carcinogenesis* 1999; 20: 2125-9.

- [48] Pratesi N, Mangoni M, Mancini I, Paiar F, Simi L, Livi L, et al. Association between single nucleotide polymorphisms in the XRCC1 and RAD51 genes and clinical radiosensitivity in head and neck cancer. *Radiother Oncol* 2011; 99: 356-61.
- [49] Csejtei A, Tibold A, Koltai K, Varga Z, Szanyi I, Gobel G, et al. Association between XRCC1 polymorphisms and head and neck cancer in a Hungarian population. *Anticancer Res* 2009; 29: 4169-73.
- [50] Huang WY, Olshan AF, Schwartz SM, Berndt SI, Chen C, Llaca V, et al. Selected genetic polymorphisms in MGMT, XRCC1, XPD, and XRCC3 and risk of head and neck cancer: a pooled analysis. *Cancer Epidemiol Biomarkers Prev* 2005; 14: 1747-53.
- [51] Demokan S, Demir D, Suoglu Y, Kiyak E, Akar U, Dalay N. Polymorphisms of the XRCC1 DNA repair gene in head and neck cancer. *Pathol Oncol Res* 2005; 11: 22-5.
- [52] Tae K, Lee HS, Park BJ, Park CW, Kim KR, Cho HY, et al. Association of DNA repair gene XRCC1 polymorphisms with head and neck cancer in Korean population. *Int J Cancer* 2004; 111: 805-8.
- [53] Olshan AF, Watson MA, Weissler MC, Bell DA. XRCC1 polymorphisms and head and neck cancer. *Cancer Lett* 2002; 178: 181-6.
- [54] Hu Z, Ma H, Chen F, Wei Q, Shen H. XRCC1 polymorphisms and cancer risk: a meta-analysis of 38 case-control studies. *Cancer Epidemiol Biomarkers Prev* 2005; 14: 1810-8.
- [55] Duell EJ, Wiencke JK, Cheng TJ, Varkonyi A, Zuo ZF, Ashok TD, et al. Polymorphisms in the DNA repair genes XRCC1 and ERCC2 and biomarkers of DNA damage in human blood mononuclear cells. *Carcinogenesis* 2000; 21: 965-71.
- [56] Lei YC, Hwang SJ, Chang CC, Kuo HW, Luo JC, Chang MJ, et al. Effects on sister chromatid exchange frequency of polymorphisms in DNA repair gene XRCC1 in smokers. *Mutat Res* 2002; 519: 93-101.
- [57] Hu JJ, Smith TR, Miller MS, Lohman K, Case LD. Genetic regulation of ionizing radiation sensitivity and breast cancer risk. *Environ Mol Mutagen* 2002; 39: 208-15.
- [58] Lunn RM, Langlois RG, Hsieh LL, Thompson CL, Bell DA. XRCC1 polymorphisms: effects on aflatoxin B1-DNA adducts and glycophorin A variant frequency. *Cancer Res* 1999; 59: 2557-61.
- [59] Matullo G, Palli D, Peluso M, Guarrera S, Carturan S, Celentano E, et al. XRCC1, XRCC3, XPD gene polymorphisms, smoking and (32)P-DNA adducts in a sample of healthy subjects. *Carcinogenesis* 2001; 22: 1437-45.
- [60] Kiuru A, Lindholm C, Heilimo I, Ceppi M, Koivistoinen A, Ilus T, et al. Influence of DNA repair gene polymorphisms on the yield of chromosomal aberrations. *Environ Mol Mutagen* 2005; 46: 198-205.
- [61] Zhang K, Zhou B, Wang Y, Rao L, Zhang L. The XRCC1 Arg280His polymorphism contributes to cancer susceptibility: an update by meta-analysis of 53 individual studies. *Gene* 2012; 510: 93-101.
- [62] Applebaum KM, McClean MD, Nelson HH, Marsit CJ, Christensen BC, Kelsey KT. Smoking modifies the relationship between XRCC1 haplotypes and HPV16-negative head and neck squamous cell carcinoma. *Int J Cancer* 2009; 124: 2690-6.
- [63] Dianov GL, O'Neill P, Goodhead DT. Securing genome stability by orchestrating DNA repair: removal of radiation-induced clustered lesions in DNA. *Bioessays* 2001; 23: 745-9.
- [64] Caldecott KW, McKeown CK, Tucker JD, Ljungquist S, Thompson LH. An interaction between the mammalian DNA repair protein XRCC1 and DNA ligase III. *Mol Cell Biol* 1994; 14: 68-76.
- [65] Zhang Y, Newcomb PA, Egan KM, Titus-Ernstoff L, Chanock S, Welch R, et al. Genetic polymorphisms in base-excision repair pathway genes and risk of breast cancer. *Cancer Epidemiol Biomarkers Prev* 2006; 15: 353-8.
- [66] Figueroa JD, Malats N, Real FX, Silverman D, Kogevinas M, Chanock S, et al. Genetic variation in the base excision repair pathway and bladder cancer risk. *Hum Genet* 2007; 121: 233-42.
- [67] Li D, Suzuki H, Liu B, Morris J, Liu J, Okazaki T, et al. DNA repair gene polymorphisms and risk of pancreatic cancer. *Clin Cancer Res* 2009; 15: 740-6.
- [68] Qian B, Zhang H, Zhang L, Zhou X, Yu H, Chen K. Association of genetic polymorphisms in DNA repair pathway genes with non-small cell lung cancer risk. *Lung Cancer* 2011; 73: 138-46.
- [69] Zienolddiny S, Campa D, Lind H, Ryberg D, Skaug V, Stangeland L, et al. Polymorphisms of DNA repair genes and risk of non-small cell lung cancer. *Carcinogenesis* 2006; 27: 560-7.
- [70] Rajaraman P, Hutchinson A, Wichner S, Black PM, Fine HA, Loeffler JS, et al. DNA repair gene polymorphisms and risk of adult meningioma, glioma, and acoustic neuroma. *Neuro Oncol* 2010; 12: 37-48.
- [71] Volker M, Mone MJ, Karmakar P, van Hoffen A, Schul W, Vermeulen W, et al. Sequential assembly of the nucleotide excision repair factors in vivo. *Mol Cell* 2001; 8: 213-24.
- [72] Wu X, Zhao H, Wei Q, Amos CI, Zhang K, Guo Z, et al. XPA polymorphism associated with reduced lung cancer risk and a modulating effect on nucleotide excision repair capacity. *Carcinogenesis* 2003; 24: 505-9.
- [73] Park JY, Park SH, Choi JE, Lee SY, Jeon HS, Cha SI, et al. Polymorphisms of the DNA repair gene xeroderma pigmentosum group A and risk of primary lung cancer. *Cancer Epidemiol Biomarkers Prev* 2002; 11: 993-7.
- [74] Sugimura T, Kumimoto H, Tohnai I, Fukui T, Matsuo K, Tsurusako S, et al. Gene-environment interaction involved in oral carcinogenesis: molecular epidemiological study for metabolic and DNA repair gene polymorphisms. *J Oral Pathol Med* 2006; 35: 11-8.
- [75] Bau DT, Tsai MH, Huang CY, Lee CC, Tseng HC, Lo

- YL, et al. Relationship between polymorphisms of nucleotide excision repair genes and oral cancer risk in Taiwan: evidence for modification of smoking habit. *Chin J Physiol* 2007; 50: 294-300.
- [76] Abbasi R, Ramroth H, Becher H, Dietz A, Schmezer P, Popanda O. Laryngeal cancer risk associated with smoking and alcohol consumption is modified by genetic polymorphisms in ERCC5, ERCC6 and RAD23B but not by polymorphisms in five other nucleotide excision repair genes. *Int J Cancer* 2009; 125: 1431-9.
- [77] Jelonek K, Gdowicz-Klosok A, Pietrowska M, Borkowska M, Korfanty J, Rzeszowska-Wolny J, et al. Association between single-nucleotide polymorphisms of selected genes involved in the response to DNA damage and risk of colon, head and neck, and breast cancers in a Polish population. *J Appl Genet* 2010; 51: 343-52.
- [78] Liu J, Zhang Z, Cao XL, Lei DP, Wang ZQ, Jin T, et al. XPA A23G polymorphism and susceptibility to cancer: a meta-analysis. *Mol Biol Rep* 2012; 39: 6791-9.
- [79] Huang WY, Berndt SI, Kang D, Chatterjee N, Chanock SJ, Yeager M, et al. Nucleotide excision repair gene polymorphisms and risk of advanced colorectal adenoma: XPC polymorphisms modify smoking-related risk. *Cancer Epidemiol Biomarkers Prev* 2006; 15: 306-11.
- [80] Khan SG, Metter EJ, Tarone RE, Bohr VA, Grossman L, Hedayati M, et al. A new xeroderma pigmentosum group C poly(AT) insertion/deletion polymorphism. *Carcinogenesis* 2000; 21: 1821-5.
- [81] Vodicka P, Kumar R, Stetina R, Sanyal S, Soucek P, Haufroid V, et al. Genetic polymorphisms in DNA repair genes and possible links with DNA repair rates, chromosomal aberrations and single-strand breaks in DNA. *Carcinogenesis* 2004; 25: 757-63.
- [82] Shen H, Sturgis EM, Khan SG, Qiao Y, Shahlavi T, Eicher SA, et al. An intronic poly (AT) polymorphism of the DNA repair gene XPC and risk of squamous cell carcinoma of the head and neck: a case-control study. *Cancer Res* 2001; 61: 3321-5.
- [83] An J, Liu Z, Hu Z, Li G, Wang LE, Sturgis EM, et al. Potentially functional single nucleotide polymorphisms in the core nucleotide excision repair genes and risk of squamous cell carcinoma of the head and neck. *Cancer Epidemiol Biomarkers Prev* 2007; 16: 1633-8.
- [84] Yang M, Kang MJ, Choi Y, Kim CS, Lee SM, Park CW, et al. Associations between XPC expression, genotype, and the risk of head and neck cancer. *Environ Mol Mutagen* 2005; 45: 374-9.
- [85] Wood RD. Nucleotide excision repair in mammalian cells. *J Biol Chem* 1997; 272: 23465-8.
- [86] Zhang L, Wang J, Xu L, Zhou J, Guan X, Jiang F, et al. Nucleotide excision repair gene ERCC1 polymorphisms contribute to cancer susceptibility: a meta-analysis. *Mutagenesis* 2012; 27: 67-76.
- [87] Li Y, Gu S, Wu Q, Fu X, Mao Y, Huang Y, et al. No association of ERCC1 C8092A and T19007C polymorphisms to cancer risk: a meta-analysis. *Eur J Hum Genet* 2007; 15: 967-73.
- [88] Yang M, Kim WH, Choi Y, Lee SH, Kim KR, Lee HS, et al. Effects of ERCC1 expression in peripheral blood on the risk of head and neck cancer. *Eur J Cancer Prev* 2006; 15: 269-73.
- [89] Sturgis EM, Dahlstrom KR, Spitz MR, Wei Q. DNA repair gene ERCC1 and ERCC2/XPD polymorphisms and risk of squamous cell carcinoma of the head and neck. *Arch Otolaryngol Head Neck Surg* 2002; 128: 1084-8.
- [90] Yang ZH, Dai Q, Kong XL, Yang WL, Zhang L. Association of ERCC1 polymorphisms and susceptibility to nasopharyngeal carcinoma. *Mol Carcinog* 2009; 48: 196-201.
- [91] Cleaver JE. Common pathways for ultraviolet skin carcinogenesis in the repair and replication defective groups of xeroderma pigmentosum. *J Dermatol Sci* 2000; 23: 1-11.
- [92] Hou SM, Falt S, Angelini S, Yang K, Nyberg F, Lambert B, et al. The XPD variant alleles are associated with increased aromatic DNA adduct level and lung cancer risk. *Carcinogenesis* 2002; 23: 599-603.
- [93] Benhamou S, Sarasin A. ERCC2/XPD gene polymorphisms and cancer risk. *Mutagenesis* 2002; 17: 463-9.
- [94] Majumder M, Sikdar N, Ghosh S, Roy B. Polymorphisms at XPD and XRCC1 DNA repair loci and increased risk of oral leukoplakia and cancer among NAT2 slow acetylators. *Int J Cancer* 2007; 120: 2148-56.
- [95] Harth V, Schafer M, Abel J, Maintz L, Neuhaus T, Besuden M, et al. Head and neck squamous-cell cancer and its association with polymorphic enzymes of xenobiotic metabolism and repair. *J Toxicol Environ Health A* 2008; 71: 887-97.
- [96] Ji YB, Tae K, Lee YS, Lee SH, Kim KR, Park CW, et al. XPD Polymorphisms and Risk of Squamous Cell Carcinoma of the Head and Neck in a Korean Sample. *Clin Exp Otorhinolaryngol* 2010; 3: 42-7.
- [97] Gugatschka M, Dehchamani D, Wascher TC, Friedrich G, Renner W. DNA repair gene ERCC2 polymorphisms and risk of squamous cell carcinoma of the head and neck. *Exp Mol Pathol* 2011; 91: 331-4.
- [98] Hu YY, Yuan H, Jiang GB, Chen N, Wen L, Leng WD, et al. Associations between XPD Asp312Asn polymorphism and risk of head and neck cancer: a meta-analysis based on 7,122 subjects. *PLoS One* 2012; 7: e35220.
- [99] Yuan H, Niu YM, Wang RX, Li HZ, Chen N. Association between XPD Lys751Gln polymorphism and risk of head and neck cancer: a meta-analysis. *Genet Mol Res* 2011; 10: 3356-64.
- [100] Krupa R, Kasznicki J, Gajecka M, Rydzanicz M, Kiwerska K, Kaczmarczyk D, et al. Polymorphisms of the DNA repair genes XRCC1 and ERCC4 are not associated with smoking- and drinking-dependent larynx cancer in a Polish population. *Exp Oncol* 2011; 33: 55-6.
- [101] Shi TY, He J, Qiu LX, Zhu ML, Wang MY, Zhou XY, et al. Association between XPF polymorphisms and cancer risk: a meta-analysis. *PLoS One* 2012; 7: e38606.
- [102] Kiyohara C, Yoshimasu K. Genetic polymorphisms in the nucleotide excision repair pathway and lung cancer

- risk: a meta-analysis. *Int J Med Sci* 2007; 4: 59-71.
- [103] Cui Y, Morgenstern H, Greenland S, Tashkin DP, Mao J, Cao W, et al. Polymorphism of Xeroderma Pigmentosum group G and the risk of lung cancer and squamous cell carcinomas of the oropharynx, larynx and esophagus. *Int J Cancer* 2006; 118: 714-20.
- [104] Manuguerra M, Matullo G, Veglia F, Autrup H, Dunning AM, Garte S, et al. Multi-factor dimensionality reduction applied to a large prospective investigation on gene-gene and gene-environment interactions. *Carcinogenesis* 2007; 28: 414-22.
- [105] Yuan H, Li H, Ma H, Niu Y, Wu Y, Zhang S, et al. Genetic polymorphisms in key DNA repair genes and risk of head and neck cancer in a Chinese population. *Exp Ther Med* 2012; 3: 719-24.
- [106] Ito E, Yanagisawa Y, Iwahashi Y, Suzuki Y, Nagasaki H, Akiyama Y, et al. A core promoter and a frequent single-nucleotide polymorphism of the mismatch repair gene hMLH1. *Biochem Biophys Res Commun* 1999; 256: 488-94.
- [107] Arita M, Zhong X, Min Z, Hemmi H, Shimatake H. Multiple sites required for expression in 5'-flanking region of the hMLH1 gene. *Gene* 2003; 306: 57-65.
- [108] Xu JL, Yin ZQ, Huang MD, Wang XF, Gao W, Liu LX, et al. MLH1 polymorphisms and cancer risk: a meta-analysis based on 33 case-control studies. *Asian Pac J Cancer Prev* 2012; 13: 901-7.
- [109] Pan XM, Yang WZ, Xu GH, Bai P, Qin HJ, Zhang LS, et al. The association between MLH1 -93 G>A polymorphism of DNA mismatch repair and cancer susceptibility: a meta-analysis. *Mutagenesis* 2011; 26: 667-73.
- [110] Kouso H, Yoshino I, Miura N, Takenaka T, Ohba T, Yohena T, et al. Expression of mismatch repair proteins, hMLH1/hMSH2, in non-small cell lung cancer tissues and its clinical significance. *J Surg Oncol* 2008; 98: 377-83.
- [111] Skarda J, Fridman E, Plevova P, Hajdich M, Radova L, Ofek E, et al. Prognostic value of hMLH1 and hMSH2 immunohistochemical expression in non-small cell lung cancer. A tissue microarray study. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2006; 150: 255-9.
- [112] Spaepen M, Vankeirsbilck B, Van Opstal S, Tejpar S, Van Cutsem E, Geboes K, et al. Germline mutations of the hMLH1 and hMSH2 mismatch repair genes in Belgian hereditary nonpolyposis colon cancer (HNPCC) patients. *Fam Cancer* 2006; 5: 179-89.
- [113] Ravnik-Glavac M, Potocnik U, Glavac D. Incidence of germline hMLH1 and hMSH2 mutations (HNPCC patients) among newly diagnosed colorectal cancers in a Slovenian population. *J Med Genet* 2000; 37: 533-6.
- [114] Ellison AR, Lofing J, Bitter GA. Functional analysis of human MLH1 and MSH2 missense variants and hybrid human-yeast MLH1 proteins in *Saccharomyces cerevisiae*. *Hum Mol Genet* 2001; 10: 1889-900.
- [115] Drotschmann K, Clark AB, Kunkel TA. Mutator phenotypes of common polymorphisms and missense mutations in MSH2. *Curr Biol* 1999; 9: 907-10.
- [116] Peltomaki P. Role of DNA mismatch repair defects in the pathogenesis of human cancer. *J Clin Oncol* 2003; 21: 1174-9.
- [117] Nicolaides NC, Papadopoulos N, Liu B, Wei YF, Carter KC, Ruben SM, et al. Mutations of two PMS homologues in hereditary nonpolyposis colon cancer. *Nature* 1994; 371: 75-80.
- [118] Mead LJ, Jenkins MA, Young J, Royce SG, Smith L, St John DJ, et al. Microsatellite instability markers for identifying early-onset colorectal cancers caused by germ-line mutations in DNA mismatch repair genes. *Clin Cancer Res* 2007; 13: 2865-9.
- [119] Benhamou S, Tuimala J, Bouchardy C, Dayer P, Sarasin A, Hirvonen A. DNA repair gene XRCC2 and XRCC3 polymorphisms and susceptibility to cancers of the upper aerodigestive tract. *Int J Cancer* 2004; 112: 901-4.
- [120] Yen CY, Liu SY, Chen CH, Tseng HF, Chuang LY, Yang CH, et al. Combinational polymorphisms of four DNA repair genes XRCC1, XRCC2, XRCC3, and XRCC4 and their association with oral cancer in Taiwan. *J Oral Pathol Med* 2008; 37: 271-7.
- [121] Figueroa JD, Malats N, Rothman N, Real FX, Silverman D, Kogevinas M, et al. Evaluation of genetic variation in the double-strand break repair pathway and bladder cancer risk. *Carcinogenesis* 2007; 28: 1788-93.
- [122] Brenneman MA, Weiss AE, Nickoloff JA, Chen DJ. XRCC3 is required for efficient repair of chromosome breaks by homologous recombination. *Mutat Res* 2000; 459: 89-97.
- [123] Araujo FD, Pierce AJ, Stark JM, Jasin M. Variant XRCC3 implicated in cancer is functional in homology-directed repair of double-strand breaks. *Oncogene* 2002; 21: 4176-80.
- [124] Yin QH, Liu C, Li L, Zu XY, Wang YJ. Association Between the XRCC3 T241M Polymorphism and Head and Neck Cancer Susceptibility: a Meta-analysis of Case-control Studies. *Asian Pac J Cancer Prev* 2012; 13: 5201-5.
- [125] Flores-Obando RE, Gollin SM, Ragin CC. Polymorphisms in DNA damage response genes and head and neck cancer risk. *Biomarkers* 2010; 15: 379-99.
- [126] Thacker J. The RAD51 gene family, genetic instability and cancer. *Cancer Lett* 2005; 219: 125-35.
- [127] Ricks-Santi LJ, Sucheston LE, Yang Y, Freudenheim JL, Isaacs CJ, Schwartz MD, et al. Association of Rad51 polymorphism with DNA repair in BRCA1 mutation carriers and sporadic breast cancer risk. *BMC Cancer* 2011; 11: 278.
- [128] Yu KD, Yang C, Fan L, Chen AX, Shao ZM. RAD51 135G>C does not modify breast cancer risk in non-BRCA1/2 mutation carriers: evidence from a meta-analysis of 12 studies. *Breast Cancer Res Treat* 2011; 126: 365-71.
- [129] Sliwinski T, Walczak A, Przybylowska K, Rusin P, Pietruszewska W, Zielinska-Blizniewska H, et al. Polymorphisms of the XRCC3 C722T and the RAD51 G135C genes and the risk of head and neck cancer in a Polish

- population. *Exp Mol Pathol* 2010; 89: 358-66.
- [130] Werbroutck J, De Ruyck K, Duprez F, Van Eijkeren M, Rietzschel E, Bekaert S, et al. Single-nucleotide polymorphisms in DNA double-strand break repair genes: association with head and neck cancer and interaction with tobacco use and alcohol consumption. *Mutat Res* 2008; 656: 74-81.
- [131] Gresner P, Gromadzinska J, Polanska K, Twardowska E, Jurewicz J, Wasowicz W. Genetic variability of Xrcc3 and Rad51 modulates the risk of head and neck cancer. *Gene* 2012; 504: 166-74.
- [132] Lu J, Wang LE, Xiong P, Sturgis EM, Spitz MR, Wei Q. 172G>T variant in the 5' untranslated region of DNA repair gene RAD51 reduces risk of squamous cell carcinoma of the head and neck and interacts with a P53 codon 72 variant. *Carcinogenesis* 2007; 28: 988-94.
- [133] Tauchi H. Positional cloning and functional analysis of the gene responsible for Nijmegen breakage syndrome, NBS1. *J Radiat Res* 2000; 41: 9-17.
- [134] Zheng J, Zhang C, Jiang L, You Y, Liu Y, Lu J, et al. Functional NBS1 polymorphism is associated with occurrence and advanced disease status of nasopharyngeal carcinoma. *Mol Carcinog* 2011; 50: 689-96.
- [135] Tseng HC, Tsai MH, Chiu CF, Wang CH, Chang NW, Huang CY, et al. Association of XRCC4 codon 247 polymorphism with oral cancer susceptibility in Taiwan. *Anticancer Res* 2008; 28: 1687-91.
- [136] Chiu CF, Tsai MH, Tseng HC, Wang CL, Wang CH, Wu CN, et al. A novel single nucleotide polymorphism in XRCC4 gene is associated with oral cancer susceptibility in Taiwanese patients. *Oral Oncol* 2008; 44: 898-902.
- [137] Pfeiffer P, Goedecke W, Kuhfittig-Kulle S, Obe G. Pathways of DNA double-strand break repair and their impact on the prevention and formation of chromosomal aberrations. *Cytogenet Genome Res* 2004; 104: 7-13.
- [138] Rothkamm K, Kruger I, Thompson LH, Lobrich M. Pathways of DNA double-strand break repair during the mammalian cell cycle. *Mol Cell Biol* 2003; 23: 5706-15.
- [139] West SC. Molecular views of recombination proteins and their control. *Nat Rev Mol Cell Biol* 2003; 4: 435-45.
- [140] Bau DT, Tseng HC, Wang CH, Chiu CF, Hua CH, Wu CN, et al. Oral cancer and genetic polymorphism of DNA double strand break gene Ku70 in Taiwan. *Oral Oncol* 2008; 44: 1047-51.
- [141] Hsu CF, Tseng HC, Chiu CF, Liang SY, Tsai CW, Tsai MH, et al. Association between DNA double strand break gene Ku80 polymorphisms and oral cancer susceptibility. *Oral Oncol* 2009; 45: 789-93.
- [142] Yang H, Lippman SM, Huang M, Jack Lee J, Wang W, Spitz MR, et al. Genetic polymorphisms in double-strand break DNA repair genes associated with risk of oral premalignant lesions. *Eur J Cancer* 2008; 44: 1603-11.
- [143] Ferguson DO, Sekiguchi JM, Chang S, Frank KM, Gao Y, DePinho RA, et al. The nonhomologous end-joining pathway of DNA repair is required for genomic stability and the suppression of translocations. *Proc Natl Acad Sci U S A* 2000; 97: 6630-3.
- [144] Ding Z, Zhang J, Cheng Q, Shen J, Hu Y, Chen H. The LIG4 Ile658Val polymorphism does not affect the risk of cervical carcinoma. *Pathol Res Pract* 2010; 206: 556-9.
- [145] Iwakuma T, Sakumi K, Nakatsuru Y, Kawate H, Igarashi H, Shiraishi A, et al. High incidence of nitrosamine-induced tumorigenesis in mice lacking DNA repair methyltransferase. *Carcinogenesis* 1997; 18: 1631-5.
- [146] Zhang Z, Wang L, Wei S, Liu Z, Wang LE, Sturgis EM, et al. Polymorphisms of the DNA repair gene MGMT and risk and progression of head and neck cancer. *DNA Repair (Amst)* 2010; 9: 558-66.
- [147] Kietthubthew S, Sriplung H, Au WW, Ishida T. Polymorphism in DNA repair genes and oral squamous cell carcinoma in Thailand. *Int J Hyg Environ Health* 2006; 209: 21-9.
- [148] Hall J, Hashibe M, Boffetta P, Gaborieau V, Moullan N, Chabrier A, et al. The association of sequence variants in DNA repair and cell cycle genes with cancers of the upper aerodigestive tract. *Carcinogenesis* 2007; 28: 665-71.
- [149] Hardy J, Singleton A. Genomewide association studies and human disease. *N Engl J Med* 2009; 360: 1759-68.
- [150] McKay JD, Truong T, Gaborieau V, Chabrier A, Chuang SC, Byrnes G, et al. A genome-wide association study of upper aerodigestive tract cancers conducted within the INHANCE consortium. *PLoS Genet* 2011; 7: e1001333.
- [151] Yuan H, Ma H, Lu F, Yuan Z, Wang R, Jiang H, et al. Genetic variants at 4q23 and 12q24 are associated with head and neck cancer risk in China. *Mol Carcinog* 2012.
- [152] Ghousaini M, Song H, Koessler T, Al Olama AA, Kote-Jarai Z, Driver KE, et al. Multiple loci with different cancer specificities within the 8q24 gene desert. *J Natl Cancer Inst* 2008; 100: 962-6.