



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Research paper

## The impact of COVID-19 on social support perception and stress of prehospital care providers



Medine Unal<sup>a</sup>, Atakan Yilmaz<sup>b,\*</sup>, Halis Yilmaz<sup>c</sup>, Gulay Yigitoglu Tasdemir<sup>d</sup>, Mehmet Uluturk<sup>a</sup>, Aykut Kemanci<sup>a</sup>, Hande Senol<sup>e</sup>, Burak Altan<sup>f</sup>, Mert Ozen<sup>b</sup>, Murat Seyit<sup>b</sup>, Alten Oskay<sup>g</sup>, Ibrahim Turkcu<sup>g</sup>

<sup>a</sup> Pamukkale University, Faculty of Medicine, Department of Emergency Medicine, 20070 Denizli, Turkey

<sup>b</sup> Pamukkale University, Medical Faculty, Department of Emergency Medicine, 20070 Denizli, Turkey

<sup>c</sup> Pamukkale University, Health Science Institute, 20160 Denizli, Turkey

<sup>d</sup> Pamukkale University, Faculty of Health Science, Department of Psychiatric Nursing, 20160 Denizli, Turkey

<sup>e</sup> Pamukkale University, Medical Faculty, Department of Biostatistics, 20070 Denizli, Turkey

<sup>f</sup> Health Services Department, Provincial Health Directorate, Denizli, Turkey

<sup>g</sup> Pamukkale University, Medical Faculty, Department of Emergency Medicine, 20070 Denizli, Turkey

## ARTICLE INFO

## Article history:

Received 5 March 2022

Received in revised form 10 April 2022

Accepted 28 April 2022

## Keywords:

Acute stress

Prehospital care providers

COVID-19

Paramedic

Social support

## ABSTRACT

**Background:** This study seeks to explore the impact of COVID-19 outbreak on the social support perception and acute stress disorder of prehospital care providers (PCPs) in the province of Denizli.

**Methods:** This descriptive and cross-sectional study was conducted between December 25, 2020 and January 25, 2021. Out of 510 ambulatory care staff constituting the study population, there were 287 PCPs (56.2%), including 13 physicians, 89 paramedics, 134 emergency medical technicians, and 51 individuals from other occupational groups (nurse, driver, cleaning staff, medical secretary) based at emergency health services. The data collection tools employed in the study include an introductory information form, Multidimensional Scale of Perceived Social Support (MSPSS), and National Stressful Events Survey Acute Stress Disorder Short Scale (NSESSS), which was organized as an online questionnaire.

**Results:** We analyzed the data from 287 PCPs that completed the form and scales. The mean score of the NSESSS was calculated as  $1.53 \pm 0.79$ . The PCPs who experienced health problems ( $1.85 \pm 0.69$ ), suffered from mental problems and received psychotherapy and medication ( $2.57 \pm 0.57$ ), encountered COVID-19 patients ( $1.58 \pm 0.8$ ), provided care for COVID-19 patients ( $1.59 \pm 0.79$ ), and took polymerase chain reaction (PCR) tests ( $1.68 \pm 0.78$ ) had higher acute stress symptom levels. The total mean score of MSPSS was calculated as  $66.28 \pm 17.22$ . Total MSPSS scores of the participants varied significantly in terms of age, marital status, taking a COVID-19 test, suffering from mental problems, status of encountering a COVID-19 patient, and workplace satisfaction ( $p < 0.05$ ).

**Conclusions:** The findings are suggestive of high perceptions of multidimensional social support and low acute stress symptom levels of the PCPs during the COVID-19 pandemic period.

© 2022 College of Emergency Nursing Australasia. Published by Elsevier Ltd. All rights reserved.

## 1. Introduction

The need for service groups, whether public, private or non-governmental organizations, is increasing in the event of disasters and emergencies. The need for healthcare staff carries more weight than that of other service personnel, most notably in the case of

medical disasters, such as epidemics and pandemics. Currently, healthcare staff is fighting at the frontline against the COVID-19 disease in the ongoing pandemic period, as in the epidemics of SARS, MERS, Influenza experienced in previous years. A systematic review indicated that the healthcare staff who were female, directly contacted patients with COVID-19 manifestations or confirmed cases, and had specific personal characteristics turned out to be more susceptible to stress, anxiety, and depression [1]. Prehospital care providers as an emergency medical service (EMS) are in charge of out-of-hospital care for critically ill patients, get in the first contact with infected or suspected individuals, administer on-scene

\* Correspondence to: Department of Emergency Medicine, Pamukkale University Hospital, Pamukkale University of Medical Sciences, Kinikli, 20070 Denizli, Turkey.  
E-mail address: [dr\\_atakanyilmaz@yahoo.com](mailto:dr_atakanyilmaz@yahoo.com) (A. Yilmaz).

treatment, if necessary, and transport these individuals to the relevant healthcare institutions. Prehospital care providers, like other members of the society, are reportedly affected physically, socially and psychologically due to (a) fear of contracting the disease, (b) anxiety of infecting the family and acquaintances, (c) lockdown measures, (d) lack of sufficient knowledge and experience about the disease, (e) shortage of personal protective equipment (PPE), (f) being treated as a potential carrier by the society, (g) increasing working hours, (h) increasing number of patients cared for, and (i) less support from colleagues and administrators by pandemic processes [2–4]. Therefore, the physical, psychological, and social needs of frontline prehospital care providers should be identified in order to provide timely and effective medical care and support to individuals in need of health care during the pandemic.

This study seeks to explore the impact of COVID-19 outbreak on the social support perception and acute stress disorder of pre-hospital care providers in the province of Denizli.

## 2. Methods

### 2.1. Study design

This is a cross-sectional study of prehospital care providers and their social support perception and acute stress disorder. The ethics approval of this study was granted from Non-interventional Studies Ethics Board of Pamukkale University (date: 11.12.2020 and number: 192.168.89.237–35219). The research was carried out within one month (between December 25, 2020 and January 25, 2021) following the approval of the ethics committee. Out of 510 ambulatory healthcare staff based at emergency health services in the province of Denizli, 358 agreed to participate in the study. Of these participants, 71 individuals who did not meet the inclusion criteria were excluded. We ultimately analyzed the data of 287 PCPs (%56.2), including 13 physicians, 89 paramedics, 134 emergency medical technicians, and 51 individuals from other occupational groups (nurse, driver, cleaning staff, medical secretary) based at emergency medical services. The inclusion criteria can be listed as refusal to participate, failure to answer the surveys fully, and being on a leave of absence or taking sick leave during the study period. The exclusion criteria were defined as refusing to participate in the study, leaving questionnaire form incomplete, and being on a leave of absence or taking sick leave while the study was in progress.

### 2.2. Data collection

The research data was collected by means of an introductory information form devised by the study researchers, Multidimensional Scale of Perceived Social Support (MSPSS), and National Stressful Events Survey Acute Stress Disorder Short Scale (NSESSS). The forms and scales were organized by the form of an online survey website (<https://survey.zohopublic.com>) and sent to the prehospital care providers as an email. Before filling out the surveys, the eligible participants were informed about the study through an introductory information form, and those answering the surveys were considered to have given their informed consent to participate in the study. The introductory information form devised by the researchers questions the participants' age, gender, marital status, parenthood status, profession, educational level, years of work experience, work experience at the current department, health status, taking a PCR test and its results, status of encountering, examining or caring for Covid-19 patients, and job satisfaction. The respondents were granted one-month period to complete the surveys.

### 2.3. Main results and measuring instruments

#### 2.3.1. Multidimensional Scale of Perceived Social Support (MSPSS)

MSPSS was developed by Zimet et al. [5] the validity and reliability analysis of MSPSS in Turkey was performed in 2001 by Eker and Arkar, who calculated its Cronbach's alpha coefficient as 0.80–0.95 [6,7]. MSPSS, which is made up of 12 items (four items for each subscale), subjectively evaluates the adequacy of social support received from family, friends, and significant others. The options of the scale are designed in the 7-point Likert type, ranging from "strongly disagree" to "strongly agree". While sub-scale scores are obtained by summing the scores of four items in each sub-scale, the sum of all the sub-scale scores yields the total score of the scale. Accordingly, a higher score indicates a stronger perceived social support [6].

#### 2.3.2. National Stressful Events Survey Acute Stress Disorder Short Scale (NSESSS)

American Psychiatric Association has issued the DSM-5 Severity of Acute Stress Symptom Scale-Adult to assess the severity of acute stress symptoms. NSESSS is a seven-item scale that assesses the severity of acute stress disorder symptoms developing after an extremely stressful event or experience in individuals aged 18 and over. The validity and reliability study of the Turkish version of the DSM-5 NSESSS was conducted by Aşçı İbaşı et al., who calculated its Cronbach's alpha coefficient as 0.95 [8]. Each item in the scale which asks the respondent to assess the severity of the acute stress disorder enduring for the past seven days is graded with scores ranging from "0 =none" to "4 =extremely". Total score ranges from 0 to 28 points, and average total score is calculated by dividing the total raw score by the number of items in the scale [8]. The higher the overall score is, the more severe the symptoms of acute stress disorder are.

### 2.4. Data analysis

All the statistical analyses were performed using SPSS v25.0 (IBM SPSS Statistics 25 software (Armonk, NY: IBM Corp.)). The continuous variables were defined by mean  $\pm$  standard deviation, while categorical variables were presented as number and percent. Normality assumptions were tested with Kolmogorov Smirnov and Shapiro Wilk analyses. For independent group comparisons in which parametric test assumptions were met, we used Independent samples t test for 2 group comparisons and One Way Analysis of Variance (post hoc: Tukey Test) for 2 + group comparisons. When parametric test assumptions were violated, we performed Mann-Whitney U test for 2 group comparisons and Kruskal Wallis Variance Analysis (post hoc: Mann Whitney U test with Bonferroni Correction) for 2 + group comparisons. We also conducted Spearman correlation analysis to investigate the relationship between continuous variables. Statistical significance was set at  $p < 0.05$ .

## 3. Results

In the final analysis, the dataset formed through the information provided by 287 PCPs in total was computerized and then analyzed. The rate of participation in the surveys amounted to 56.2% of the whole emergency medical staff in the city. The mean age of the participants was calculated as  $32.5 \pm 7.4$ . Moreover, 42.9% ( $n = 123$ ) were male, while 57.1% ( $n = 164$ ) were female. In terms of occupational status, the largest participating group was emergency medical technicians with 46.7% ( $n = 134$ ), followed by paramedics with 31% ( $n = 89$ ), other occupational groups with 17.8% ( $n = 51$ ), and physicians with 4.5% ( $n = 13$ ). The descriptive characteristics of the participants are presented in Table 1.

The Cronbach's alpha coefficient was calculated as 0.951 for MSPSS and 0.846 for NSESSS in this study. The mean score of the

**Table 1**  
Descriptive characteristics of the participants (n = 287).

		n	%
<b>Age</b>	18–25	48	16.7
	26–35	148	51.6
	36–45	72	25.1
	46 and over	19	6.6
<b>Gender</b>	Male	123	42.9
	Female	164	57.1
<b>Marital Status</b>	Single	85	29.6
	Married	202	70.4
<b>Number of Children</b>	No child	116	40.4
	1	73	25.4
	2	79	27.5
	3 and more	19	6.6
<b>Profession</b>	Physician	13	4.5
	Paramedic	89	31
	Emergency medical technician	134	46.7
	Other	51	17.8
<b>Educational Status</b>	Primary school	7	2.4
	High School	46	16
	Under-graduate and post-graduate	234	81.5
<b>Working time in the current department</b>	1–5 years	170	59.2
	6–11 years	83	28.9
	12–17 years	28	9.8
	18 and over years	6	2.1
<b>Have you ever encountered a COVID-19 patient?</b>	Yes	254	88.5
	No	33	11.5
<b>Have you ever examined or cared for a COVID-19 patient?</b>	Yes	231	80.5
	No	56	19.5
<b>Have you ever taken a PCR test?</b>	Yes	184	64.1
	No	103	35.9
<b>Mental health problems</b>	I did not experience any mental problems	210	73.2
	I suffered from a mental problem but did not receive any support	49	17.1
	I followed a drug regimen	20	7
	I followed a drug regimen and underwent psychotherapy	8	2.8
<b>Job satisfaction</b>	Satisfied	130	45.3
	Partly satisfied	136	47.4
	Dissatisfied	21	7.3

participants filling in NSESSS was found as  $1.53 \pm 0.79$ , while total mean score of MSPSS was  $66.28 \pm 17.22$ . As for the sub-dimensions of MSPSS, the highest mean scores belonged to 'family' ( $23.24 \pm 5.9$ ), followed by 'significant other' ( $22.24 \pm 7.23$ ) and 'friends' ( $20.8 \pm 6.56$ ).

Table 2 illustrates the breakdown of MSPSS subscale and total score averages of the participating prehospital care providers. Total MSPSS scores of the participants varied significantly in terms of age, marital status, taking a COVID-19 test, suffering from mental problems, status of encountering a COVID-19 patient, and workplace satisfaction ( $p < 0.05$ ).

We did not detect any significant difference in a range of variables, including total MSPSS scores, gender, number of children, occupation (physician, paramedic, emergency medical technician, and others), educational status, caring for a COVID-19 patient before, Covid-19 PCR test result, working experience, working time in the current department, presence of health problems ( $p > 0.05$ ).

NSESSS scores yielded significant differences with regard to age, gender, parenthood status, occupation, status of encountering, examining and/or caring for COVID-19 patients, years of work experience, health problems (coronary artery disease, diabetes mellitus, hypertension, chronic renal failure, etc.), taking a COVID-19 test, suffering from mental problems, and job satisfaction levels ( $p < 0.05$ ). However, no statistical significance was observed in

NSESSS scores in terms of marital and educational status, PCR test results, and work experience at the current department ( $p > 0.05$ ) (Table 3).

NSESSS scores indicated a weak negative correlation with 'family' and 'significant other' subscales, total MSPSS scores, but a moderate negative correlation with 'friends' subscale ( $r = -0.331$ ,  $p < 0.05$ ;  $r = -0.272$ ,  $p < 0.05$ ;  $r = -0.366$ ,  $p < 0.05$ ;  $r = -0.401$ ,  $p < 0.05$ , respectively).

#### 4. Discussion

This study carried out within the first year of the COVID-19 outbreak set out to investigate the impact of this outbreak on the perception of social support and stress in prehospital care providers in the province of Denizli. The resulting picture reveals that acute stress symptoms of this staff are mild, and that perceived multi-dimensional social support is high both as a whole and in the subscales.

Prolonged stress produces adverse mental, emotional, and physiological results. Symptoms, such as burnout, emotional fatigue, or work-related stress, have been reported in almost half of the healthcare professionals during their regular working periods [9]. Increased workload and intense work pressure during the outbreak are exhausting the healthcare staff physically and psychologically, bringing about higher levels of stress [10]. A substantial body of research investigating the psychological impact of the COVID-19 outbreak on the healthcare staff reveals that the frontline personnel is less affected than those working in the background, administrative staff, and society [11–16]. Paradoxically, many lines of evidence indicate the tremendous psychosocial impact of the COVID-19 pandemic on medical staff [2,17–20]. Our study has found mild levels of acute stress symptoms in the frontline prehospital care providers. However, the stress level of those who encountered, examined or treated COVID-19 patients turned out to be significantly higher than their counterparts who did not. Mounting clinical data support the favorable effects of social support against stress [15,21,22]. In line with the literature, our study has found the total social support perception of the prehospital care providers to be at high levels in the subscales of family, friend and significant other. Note that the lower stress levels of the participating prehospital care providers in this study may have resulted from the high levels of perceived social support. In addition, prehospital care providers fight at the frontline during the pandemic and they are the first people to do the first aid to the patients, so they bear a 'savior' role in coping with the heavy emotional burden of the pandemic can be considered as an important underlying factor. Besides, the following situations might have contributed to the lower stress levels of our participants:

- The healthcare staff has been able to gain access to protective equipment since the onset of the pandemic.
- Turkish society has taken a supportive attitude towards healthcare personnel (i.e. applauding on the balconies at 9 p.m.).
- While the number of confirmed cases peaked rapidly in Asian and European countries after China in the early period of the pandemic, there were fewer cases in Turkey due to the implementation of the recommendations issued by the Ministry of Health's Coronavirus Scientific Committee to prevent the transmission of the disease.
- The number of admissions to the emergency health services during the pandemic period has been lower than those in the normal period.
- A separate living space has been arranged for healthcare staff who want to stay away from their beloved ones for fear of infecting the disease.

**Table 2**  
Mean MSPSS scores of the prehospital care providers (n = 287).

	Family		Friends		Significant other		Overall MSPSS	
	X ± SS	p	X ± SS	p	X ± SS	p	X ± SS	p
<b>Age</b>								
18–25	20,9 ± 7,34	0.004 *	19,27 ± 7,86	0.179	20,33 ± 8,1	0.17	60,5 ± 21,02	0.038 *
26–35	23,51 ± 5,4		21,11 ± 6,14		22,13 ± 7,62		66,76 ± 16,36	
36–45	23,38 ± 5,96		20,44 ± 6,7		23,04 ± 5,98		66,86 ± 16,59	
46 and over	26,53 ± 2,76		23,63 ± 4,54		24,84 ± 5,04		75 ± 10,73	
<b>Gender</b>								
Male	22,84 ± 6,04	0.292	20,93 ± 6,69	0.633	22,17 ± 7,24	0.898	65,94 ± 17,46	0.886
Female	23,54 ± 5,79		20,71 ± 6,47		22,29 ± 7,24		66,54 ± 17,09	
<b>Marital Status</b>								
Single	21,35 ± 6,66	< 0.001 **	20,49 ± 6,69	0.61	18,78 ± 8,47	< 0.001 **	60,62 ± 18,81	< 0.001 **
Married	24,03 ± 5,37		20,94 ± 6,51		23,69 ± 6,1		68,66 ± 15,97	
<b>Number of Children</b>								
No child	22,47 ± 6,28	0.039 *	21,1 ± 6,6	0.136	21,09 ± 8	0.143	64,66 ± 17,87	0.115
1	22,34 ± 6,71		19,1 ± 7,2		21,68 ± 7,6		63,12 ± 19,59	
2	24,84 ± 3,89		21,82 ± 5,74		24,22 ± 5,11		70,87 ± 12,55	
3 and more	24,79 ± 5,88		21,32 ± 6,16		23,16 ± 7,08		69,26 ± 17,22	
<b>Profession</b>								
Physician	22,46 ± 6,33	0.604	19,69 ± 8,78	0.394	21,08 ± 8,22	0.718	63,23 ± 22,57	0.49
Paramedic	22,7 ± 6,23		20,57 ± 7,21		21,28 ± 7,85		64,55 ± 18,74	
Emergency medical technician	23,26 ± 5,89		20,46 ± 6,29		22,53 ± 7,08		66,25 ± 16,46	
Other	24,33 ± 5,19		22,41 ± 5,21		23,43 ± 6,11		70,18 ± 14,58	
<b>Educational Status</b>								
Primary school	26,57 ± 2,51	0.232	23,57 ± 5,22	0.454	24,29 ± 5,96	0.607	74,43 ± 12,27	0.327
High School	22,91 ± 6,32		20,65 ± 5,97		21,83 ± 7,38		65,39 ± 16,31	
Under-graduate and post-graduate	23,21 ± 5,87		20,75 ± 6,7		22,26 ± 7,25		66,21 ± 17,51	
<b>Working time in the current department</b>								
1–5 years	22,59 ± 6,22	0.013 *	20,37 ± 6,84	0.526	21,56 ± 7,88	0.13	64,52 ± 18,55	0.14
6–11 years	23,76 ± 5,37		21,19 ± 6,04		23,02 ± 6,41		67,98 ± 15,05	
12–17 years	24,64 ± 5,31		21,71 ± 6,58		22,86 ± 5,36		69,21 ± 14,88	
18 and over years	28 ± 0		23,5 ± 5,05		27,67 ± 0,82		79,17 ± 5,38	
<b>Have you ever encountered a COVID-19 patient?</b>								
Yes	22,86 ± 6,1	0.003 **	20,63 ± 6,58	0.171	21,78 ± 7,37	0.002 **	65,27 ± 17,68	0.014 **
No	26,18 ± 2,63		22,18 ± 6,33		25,73 ± 4,86		74,09 ± 10,42	
<b>Have you ever examined or cared for a COVID-19 patient?</b>								
Yes	22,82 ± 6,07	0.008 **	20,86 ± 6,51	0.804	22,08 ± 7,23	0.403	65,76 ± 17,53	0.408
No	24,98 ± 4,78		20,57 ± 6,79		22,88 ± 7,25		68,43 ± 15,86	
<b>Have you ever taken a PCR test?</b>								
Yes	22,73 ± 6,08	0.047 **	20,01 ± 6,73	0.006 **	21,69 ± 7,50	0.153	64,44 ± 17,61	0.012 **
No	24,14 ± 5,47		22,21 ± 6		23,21 ± 6,63		69,57 ± 16,06	
<b>Mental health problems</b>								
I did not experience any mental problems	23,96 ± 5,62	0.003 *	21,8 ± 6,22	< 0.001 *	23,05 ± 6,78	0.002 *	68,8 ± 16,34	< 0.001 *
I suffered from a mental problem but did not receive any support	21,14 ± 5,84		17,86 ± 7,09		19,55 ± 7,9		58,55 ± 17,63	
I followed a drug regimen	21,15 ± 6,87		19,35 ± 6,62		21,65 ± 7,79		62,15 ± 18,72	
I followed a drug regimen and underwent psychotherapy	22,5 ± 7,6		16,38 ± 4,34		18,88 ± 9,43		57,75 ± 17,97	
<b>Job satisfaction</b>								
Satisfied	25,22 ± 4,61	< 0.001 *	22,99 ± 5,63	< 0.001 *	23,88 ± 6,47	< 0.001 *	72,09 ± 14,71	< 0.001 *
Partly satisfied	22,15 ± 5,99		19,54 ± 6,39		21,17 ± 7,32		62,85 ± 17,06	
Dissatisfied	18,1 ± 7,55		15,48 ± 7,93		18,95 ± 8,91		52,52 ± 19,61	

MSPSS: Multidimensional Scale of Perceived Social Support; \*Kruskal-Wallis Variance Analysis; \*\*Mann-Whitney U test.

Many studies point to the association between being young and increased psychosocial impact. Some line of evidence suggests the relative prevalence of acute stress symptoms in young healthcare providers during the COVID-19 outbreak [23,24]. The rationale behind this finding is that excessive amount of information obtained from rapidly changing, questionable content and sources can cause fear and stress. In a qualitative inquiry carried out in Pakistan, media was portrayed as the major contributor of increased anxiety and stress levels of general public over the course of pandemic, since the accuracy of updates and news concerning the breakout was unable to be verified [25]. Avoiding this information pollution is required to minimize the adverse effects of the pandemic on individuals [26]. Paradoxically, different studies conducted in China during this outbreak have found lower stress levels in young healthcare staff than in other age groups. The justification provided for such a conclusion is that middle- and advanced-aged healthcare workers may have been in service during the SARS epidemic [16,27]. Age is also linked

with greater cognizance of risk, which contributes to greater emotional and psychophysical strain. Accordingly, younger healthcare providers tend to be more eager to launch initiatives, make use of new possibilities, and counter challenges, bringing about a stronger sense of resilience and personal fulfillment [28]. However, a study conducted in Italy reports high stress levels across all age groups in the COVID-19 pandemic [29]. When it comes to our study, as the age of the prehospital care providers increase, the acute stress level decreases, and the perception of total social support and the family subscale is significantly higher. This may result from the fact that, as the age of the individuals' increases, so does their professional experience and the social support they receive from their families, and they can manage the stress they face due to their job more easily.

There has been extensive work on both society at large and healthcare staff suggesting that women experience higher levels of stress than men during the COVID-19 outbreak [30–38]. In line with the aforementioned literature, our data identified higher levels of



**Table 3**  
Mean NSESSS scores of the prehospital care providers (n = 287).

Descriptive characteristics	ASSSS X ± SS	Statistical test
<b>Age band</b>		
18–25	1.67 ± 0.89	<b>0.026*</b>
26–35	1.6 ± 0.75	
36–45	1.37 ± 0.78	
46 and over	1.14 ± 0.75	
<b>Gender</b>		
Male	1.38 ± 0.8	<b>0.006**</b>
Female	1.63 ± 0.77	
<b>Marital Status</b>		
Single	1.46 ± 0.77	0.33
Married	1.56 ± 0.8	
<b>Number of Children</b>		
No child	1.48 ± 0.79	<b>0.003*</b>
1	1.81 ± 0.8	
2	1.39 ± 0.71	
3 and more	1.29 ± 0.86	
<b>Profession</b>		
Physician	1.37 ± 0.71	<b>&lt; 0.001*</b>
Paramedic	1.72 ± 0.83	
Emergency medical technician	1.57 ± 0.73	
Other	1.1 ± 0.74	
<b>Educational Status</b>		
Primary school	1.49 ± 0.51	0.127
High School	1.31 ± 0.7	
Under-graduate and post-graduate	1.57 ± 0.81	
<b>Working experience</b>		
1–5 years	1.59 ± 0.83	<b>0.01*</b>
6–11 years	1.63 ± 0.78	
12–17 years	1.49 ± 0.74	
18 and over years	1.13 ± 0.75	
<b>Working time in the current department</b>		
1–5 years	1.58 ± 0.82	0.515
6–11 years	1.46 ± 0.75	
12–17 years	1.45 ± 0.76	
18 and over years	1.24 ± 0.73	
<b>Have you ever encountered a COVID-19 patient?</b>		
Yes	1.58 ± 0.8	<b>0.003***</b>
No	1.14 ± 0.63	
<b>Have you ever examined or cared for a COVID-19 patient?</b>		
Yes	1.59 ± 0.79	<b>0.006***</b>
No	1.27 ± 0.74	
<b>Have you ever taken a PCR test?</b>		
Yes	1.68 ± 0.78	<b>0.001***</b>
No	1.23 ± 0.71	
<b>What was your PCR test result?</b>		
Positive	1.64 ± 0.67	0.62
Negative	1.7 ± 0.83	
<b>Health problem</b>		
No	1.42 ± 0.79	<b>&lt; 0.001***</b>
Yes	1.85 ± 0.69	
<b>Mental health problems</b>		
I did not experience any mental problems	1.35 ± 0.76	<b>&lt; 0.001*</b>
I suffered from a mental problem but did not receive any support	1.94 ± 0.66	
I followed a drug regimen	1.95 ± 0.63	
I followed a drug regimen and underwent psychotherapy	2.57 ± 0.57	
<b>Job satisfaction</b>		
Satisfied	1.22 ± 0.7	<b>&lt; 0.001****</b>
Partly satisfied	1.68 ± 0.71	
Dissatisfied	2.4 ± 0.9	

National Stressful Events Survey Acute Stress Disorder Short Scale (NSESSS); \*One-Way Anova test; \*\*Mann-Whitney U test; \*\*\*T-test; \*\*\*\* Kruskal-Wallis Variance Analysis

acute stress symptoms in female participants than in their male counterparts. Neff and Karney state that women face more stress factors, and that both women and men display common supportive behaviors, but the former provide better social support than the latter [39]. In contrast to this report, despite more increased stress

levels of women, our study found no significant difference between genders in terms of perceived social support.

Various studies conducted during the COVID-19 and SARS outbreak report stronger psycho-social influence in single individuals than in their married counterparts [2,40,41]. A Chinese study documents more psychiatric symptoms in married caregivers of patients with COVID-19 [12]. On the other hand, no significant difference was evident in acute stress symptom levels with respect to marital status in our study, in which perceived social support was higher in the married respondents, family, and a special person subscales. In the family subscale, the healthcare staff with two or more children reported significantly higher perception of social support and lower acute stress levels than those with no children. The underlying reasons for the lower stress level of our participants with children can be cited as their ability to spare more time for themselves, and the longer time spent together as well as the increased financial and moral support of other family members (grandparents or others) in direct proportion to the number of children.

The presence of a physical and mental illness or a negative mood is likely to enhance the level of stress [2,42–44]. Similarly, a range of studies in the pandemic period document more increased levels of stress in individuals inflicted with health problems [15,45]. Our findings also validate those of other research in that higher stress levels were observed in the participants suffering from chronic diseases (hypertension, diabetes, COPD, immunodeficiency, etc.) and mental problems. Fear of contracting the infection in chronic disease patients and social isolation enforced during the outbreaks add to the stress level of those with existing mental problems. Besides, social support is a key factor in preserving individuals' well-being or coping with the stress caused by any disease [21,46]. In this context, our data analysis reveals decreased levels of social support in those manifesting psychiatric symptoms.

A wide range of studies performed in different parts of the world during the pandemic period have revealed higher levels of stress in medical staff with low educational levels [15,47,48]. In contrast, the relevant research in the Turkish context has reported no marked correlation between educational level and stress [49]. Likewise, our research data did not signal a noteworthy relationship between the educational and stress levels during the pandemic process. This scenario may have resulted from the fact that the COVID-19 Guide has been issued online, that the healthcare staff can easily access the accurate medical information regardless of their educational level, and that they may have felt more prepared for the pandemic.

Various studies on healthcare staff report conflicting results on the correlation between occupation and stress levels during pandemic periods. While some studies indicate more increased levels of stress in nurses, others report higher degrees of stress in doctors, and some others find even higher stress levels in paramedics [37,41,50,51]. In our study, the paramedics manifested by far the highest acute stress level of all the occupational groups. The risk of transmission, fear of getting sick, and infecting their beloved ones may have enhanced the stress levels of our paramedics on the grounds that they did not use PPEs appropriately in the field for faster emergency intervention in patients who did not manifest trauma or COVID-19 symptoms.

Individuals who develop work-related stress may manifest physical, behavioral, emotional and psychological disorders [52]. The more experienced individuals are in their profession and the more satisfied they are with their workplace, the lower stress they end up with [2,53]. Consistent with the above-mentioned findings, our data also confirm the link between job satisfaction of the healthcare staff and their low stress levels and high levels of perceived social support. As the working time increases, factors such as increasing professional experience, improving bilateral relations with co-workers, and sustaining a regular income may produce favorable effects on

stress. It is assumed that working with a fixed job and income in the same workplace for long facilitates getting married and having children, which in turn multiplies the social support tools perceived by the individual from the family. Moreover, many employees have either lost their jobs or closed down their workplaces due to the adverse financial conditions of the pandemic, but the healthcare staff have maintained their job guarantee and income during this period, which may have improved their job satisfaction and thus led to lower stress and high levels of perceived social support.

While stress level was reportedly higher in the medical personnel in Spain who did not take RT-PCR test [54], a Turkish study performed on 939 healthcare staff during the pandemic process demonstrated more increased stress levels in those who took this test [55]. Similar to the latter study, our results also implicate higher levels of stress in the prehospital care providers taking the test. The likely factors for increased stress levels could be the anxiety of being quarantined if the test result is positive, the status of being distanced from working, educational, and social life for a while, and the fear of being stigmatized by the society.

#### 4.1. Limitations

The current research suffers three main limitations. Initially, the evidence garnered from this study is bounded by the specified dates, and the psychosocial approaches of the participants to the pandemic and their psychosocial impact caused by the pandemic may have changed from the onset of the outbreak to the initiation date of the study. Besides, our data may not be generalizable to all prehospital care providers in Turkey and around the world due to the cross-sectional nature of the study. Finally, the questionnaires and the scales that we drew on in this study were based upon the self-reports of the respondents and so reflect their subjective evaluations. These evaluations may have been influenced by other psycho-social events in their daily lives.

#### 5. Conclusion

Performed in the ongoing process of the COVID-19 pandemic, this study reveals that prehospital care providers perceived strong multidimensional social support and manifested low acute stress symptoms. The second major finding is that their acute stress levels and perceived multidimensional social support indicate significant correlations with some of their sociodemographic characteristics. For example, the level of acute stress symptoms correlates negatively with age, experience, and job satisfaction of the healthcare staff but positively with female staff and those who are afflicted with health or mental problems, encounter, examine or care for COVID-19 patients, and take a PCR test. In addition, the level of acute stress symptoms reveals a negative relationship between social support subscales like family, friends, a special person and overall score of multidimensional perceived social support.

#### Funding

There was no funding for this work.

#### Ethics

The study protocol was accepted by the ethical committees of Pamukkale University (date: 08.12.2020 and number: 2020/23).

#### Conflict of interest

The authors have no conflict of interest to declare.

#### Acknowledgements

The authors would like to thank the prehospital care providers who participated in this research.

#### Conflict of interest statement

The authors declare that they have no conflicts of interests.

#### References

- [1] Soto-Cámara R, García-Santa-Basilía N, Onrubia-Baticón H, Cárdena-García RM, Jiménez-Alegre JJ, Reques-Marugán AM, et al. Psychological impact of the COVID-19 pandemic on out-of-hospital health professionals: a living systematic review. *J Clin Med* 2021;10(23):5578. <https://doi.org/10.3390/jcm10235578>
- [2] Elbay RY, Kurtulmuş A, Arpacioğlu S, Karadere E. Depression, anxiety, stress levels of physicians and associated factors in Covid-19 pandemics. *Psychiatry Res* 2020;290:113130. <https://doi.org/10.1016/j.psychres.2020.113130>
- [3] Maguire B, Dean S. Epidemic and bioterrorism preparation among emergency medical services systems. *Prehosp Disaster Med* 2007;22(3):237–42. <https://doi.org/10.1017/s1049023x0000474x>
- [4] Silverman A, Simor A, Loutfy MR. Toronto emergency medical services and SARS. *Emerg Infect Dis* 2004;10(9):1688–9. <https://doi.org/10.3201/eid1009.040170>
- [5] Zimet G, Dahlem N, Zimet S, Farley G. The multidimensional scale of perceived social support. *J Personal Assess* 1988;52:30–41. [https://doi.org/10.1207/s15327752jpa5201\\_2](https://doi.org/10.1207/s15327752jpa5201_2)
- [6] Eker D, Arkar H, Yaldız H. Çok Boyutlu Algılanan Sosyal Destek Ölçeği'nin gözden geçirilmiş formunun faktör yapısı, geçerlik ve güvenilirliği. *Türk Psikiyatr Derg* 2001;12(1):17–25.
- [7] Prasad K, McLoughlin C, Stillman M, Poplau S, Goelz E, Taylor S, et al. Prevalence and correlates of stress and burnout among U.S. healthcare workers during the COVID-19 pandemic: A national cross-sectional survey study. *EClinicalMedicine* 2021;35:100879.
- [8] Aşçıbaşı K, Çökmüş FP, Aydemir Ö. DSM-5 Akut Stres Belirti Şiddeti Ölçeği Türkçe Formu'nun geçerliliği ve güvenilirliği. *Anadolu Psikiyatr Derg* 2017;18:38–44.
- [9] Amerikan Psikiyatri Birliği. Mental bozuklukların tanınması ve sayımsal el kitabı. 5th edition. Koroğlu E, (Ed.). Ankara: Hekimler Yayın Birliği; 2013.
- [10] Zhang C, Yang L, Liu S, Ma S, Wang Y, Cai Z, et al. Survey of insomnia and related social psychological factors among medical staff involved in the 2019 Novel Coronavirus disease outbreak. *Front Psychiatry* 2020;11:306. <https://doi.org/10.3389/fpsy.2020.00306>
- [11] Uyurdağ N, Eskicioğlu G, Aksu S, Soyata AZ. Risk and resilience factors for psychosocial impact in healthcare workers during COVID-19 pandemic: A systematic review. *Anatol Clin* 2021;26(1):122–40. <https://doi.org/10.21673/anadoluklin.783596>
- [12] Li Z, Ge J, Yang M, Feng J, Qiao M, Jiang R, et al. Vicarious traumatization in the general public, members, and non-members of medical teams aiding in COVID-19 control. *Brain Behav Immun* 2020;88:916–9. <https://doi.org/10.1016/j.bbi.2020.03.007>
- [13] Wu Y, Wang J, Luo C, Hu S, Lin X, Anderson AE, et al. A comparison of burnout frequency among oncology physicians and nurses working on the frontline and usual wards during the COVID-19 epidemic in Wuhan, China. *J Pain Symptom Manag* 2020;60(1):e60–5. <https://doi.org/10.1016/j.jpainsymman.2020.04.008>
- [14] Liu CY, Yang YZ, Zhang XM, Xu X, Dou QL, Zhang WW, et al. The prevalence and influencing factors in anxiety in medical workers fighting COVID-19 in China: a cross-sectional survey. *Epidemiol Infect* 2020;148:e98. <https://doi.org/10.1017/S0950268820001107>
- [15] Song X, Fu W, Liu X, Luo Z, Wang R, Zhou N, et al. Mental health status of medical staff in emergency departments during the Coronavirus disease 2019 epidemic in China. *Brain Behav Immun* 2020;88:60–5. <https://doi.org/10.1016/j.bbi.2020.06.002>
- [16] Hou T, Zhang T, Cai W, Song X, Chen A, Deng G, et al. Social support and mental health among health care workers during Coronavirus Disease 2019 outbreak: A moderated mediation model. *PLoS One* 2020;15(5):e0233831. <https://doi.org/10.1371/journal.pone.0233831>
- [17] Lu W, Wang H, Lin Y, Li L. Psychological status of medical workforce during the COVID-19 pandemic: A cross-sectional study. *Psychiatry Res* 2020;288:112936. <https://doi.org/10.1016/j.psychres.2020.112936>
- [18] García-Fernández L, Romero-Ferreiro V, López-Roldán PD, Padilla S, Calero-Sierra I, Monzó-García M, et al. Mental health impact of COVID-19 pandemic on Spanish healthcare workers. *Psychol Med* 2020;1–3. <https://doi.org/10.1017/S0033291720002019>
- [19] Wang Y, Duan Z, Peng K, Li D, Ou J, Wilson A, et al. Acute stress disorder among frontline health professionals during the COVID-19 outbreak: a structural equation modeling investigation. *Psychosom Med* 2021;83(4):373–9. <https://doi.org/10.1097/PSY.0000000000000851>
- [20] Osório FL, Silveira ILM, Pereira-Lima K, Crippa JAS, Hallak JEC, Zuardi AW, et al. Risk and protective factors for the mental health of Brazilian healthcare workers in the frontline of COVID-19 pandemic. *Front Psychiatry* 2021;12:662742. <https://doi.org/10.3389/fpsy.2021.662742>
- [21] Ardahan M. Sosyal destek ve hemşirelik. *Atatürk Üniversitesi Hemşire Yükkokulu Derg* 2006;9(2):68–75.

- [22] Xiao H, Zhang Y, Kong D, Li S, Yang N. The Effects of Social Support on Sleep Quality of Medical Staff Treating Patients with Coronavirus Disease 2019 (COVID-19) in January and February 2020 in China. *Med Sci Monit* 2020;26:e923549 <https://doi.org/10.12659/MSM.923549>
- [23] Madanes SB, Palmer RL, Szuhany KL, Malgaroli M, Jennings EL, Anbarasan D, et al. Acute stress disorder and the COVID-19 pandemic. *Psychiatr Ann* 2020;50(7):295–300. <https://doi.org/10.3928/00485713-20200611-01>
- [24] Salari N, Hosseini-Far A, Jalali R, Vaisi-Raygani A, Rasoulpoor S, Mohammadi M, et al. Prevalence of stress, anxiety, depression among the general population during the COVID-19 pandemic: a systematic review and meta-analysis. *Glob Health* 2020;16(1):57. <https://doi.org/10.1186/s12992-020-00589-w>
- [25] Munawar K, Choudhry FR. Exploring stress coping strategies of frontline emergency health workers dealing Covid-19 in Pakistan: A qualitative inquiry. *Am J Infect Control* 2021;49(3):286–92. <https://doi.org/10.1016/j.ajic.2020.06.214>
- [26] Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 2020;395(10227):912–20. [https://doi.org/10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8)
- [27] Sun D, Yang D, Li Y, Zhou J, Wang W, Wang Q, et al. Psychological impact of 2019 novel coronavirus (2019-nCoV) outbreak in health workers in China. *Epidemiol Infect* 2020;148:e96 <https://doi.org/10.1017/S0950268820001090>
- [28] Vagni M, Giostra V, Maiorano T, Santaniello G, Pajardi D. Personal accomplishment and hardiness in reducing emergency stress and burnout among COVID-19 emergency workers. *Sustainability* 2020;12(21):9071. <https://doi.org/10.3390/su12219071>
- [29] Mazza C, Ricci E, Biondi S, Colasanti M, Ferracuti S, Napoli C, et al. A Nationwide Survey of Psychological Distress among Italian People during the COVID-19 Pandemic: Immediate Psychological Responses and Associated Factors. *Int J Environ Res Public Health* 2020;17(9):3165. <https://doi.org/10.3390/ijerph17093165>
- [30] Huang L, Wang Y, Liu J, Ye P, Cheng B, Xu H, et al. Factors Associated with Resilience Among Medical Staff in Radiology Departments During The Outbreak of 2019 Novel Coronavirus Disease (COVID-19): A Cross-Sectional Study. *Med Sci Monit* 2020;26:e925669 <https://doi.org/10.12659/MSM.925669>
- [31] Wang Y, Di Y, Ye J, Wei W. Study on the public psychological states and its related factors during the outbreak of coronavirus disease 2019 (COVID-19) in some regions of China. *Psychol Health Med* 2021;26(1):13–22. <https://doi.org/10.1080/13548506.2020.1746817>
- [32] Liu D, Ren Y, Yan F, Li Y, Xu X, Yu X, et al. Psychological impact and predisposing factors of the Coronavirus Disease 2019 (COVID-19) pandemic on general public in China. *Lancet Psychiatry* 2020. <https://doi.org/10.2139/ssrn.3551415>
- [33] Zhou SJ, Zhang LG, Wang LL, Guo ZC, Wang JQ, Chen JC, et al. Prevalence and socio-demographic correlates of psychological health problems in Chinese adolescents during the outbreak of COVID-19. *Eur Child Adolesc Psychiatry* 2020;29(6):749–58. <https://doi.org/10.1007/s00787-020-01541-4>
- [34] Moghanibashi-Mansourieh A. Assessing the anxiety level of Iranian general population during COVID-19 outbreak. *Asian J Psychiatr* 2020;51:102076 <https://doi.org/10.1016/j.ajp.2020.102076>
- [35] Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *Int J Environ Res Public Health* 2020 6;17(5):1729. <https://doi.org/10.3390/ijerph17051729>
- [36] Kang L, Ma S, Chen M, Yang J, Wang Y, Li R, et al. Impact on mental health and perceptions of psychological care among medical and nursing staff in Wuhan during the 2019 novel coronavirus disease outbreak: A cross-sectional study. *Brain Behav Immun* 2020;87:11–7. <https://doi.org/10.1016/j.bbi.2020.03.028>
- [37] Tengilimoğlu D, Zekioglu A, Tosun N, Işık O, Tengilimoğlu O. Impacts of COVID-19 pandemic period on depression, anxiety and stress levels of the healthcare employees in Turkey. *Leg Med (Tokyo)* 2021;48:101811 <https://doi.org/10.1016/j.legalmed.2020.101811>
- [38] Usul E, Şan I, Bekgöz B. The Effect of the COVID-19 Pandemic on the Anxiety Level of Emergency Medical Services Professionals. *Psychiatr Danub* 2020;32(3–4):563–9. <https://doi.org/10.24869/psyd.2020.563>
- [39] Neff LA, Karney BR. Gender differences in social support: a question of skill or responsiveness? *J Pers Soc Psychol* 2005;88(1):79–90. <https://doi.org/10.1037/0022-3514.88.1.79>
- [40] Hawkley LC, Cacioppo JT. Loneliness matters: a theoretical and empirical review of consequences and mechanisms. *Ann Behav Med* 2010;40(2):218–27. <https://doi.org/10.1007/s12160-010-9210-8>
- [41] Chan AO, Huak CY. Psychological impact of the 2003 severe acute respiratory syndrome outbreak on health care workers in a medium size regional general hospital in Singapore. *Occup Med (Lond)* 2004;54(3):190–6. <https://doi.org/10.1093/occmed/kqh027>
- [42] Erdal MB. İşletmelerde stres kaynakları, sonuçları ve yönetim teknikleri ve bir uygulama, 2009. [September 1, 2021]. Available from: (<https://tez.yok.gov.tr/UlusalTezMerkezi/tezDetay.jsp?id=16pKQ1F76UFBEuZblQ1nA&no=cShpEf2zam2B930pHHU7y>).
- [43] Baltaş A., Baltaş Z. Stres ve başa çıkma yolları. İstanbul: Remzi Kitabevi; 2002.
- [44] Yue NH, Foley S, Loi R. Work role stressors and turnover intentions: A study of professional clergy in Hong Kong. *Int J Hum Resour Manag* 2005;16(11):2133–46. <https://doi.org/10.1080/09585190500315141>
- [45] Lin D, Friedmann DB, Qiao S, Tam CC, Li X, Li X. Information uncertainty: a correlate for acute stress disorder during the COVID-19 outbreak in China. *BMC Public Health* 2020;20(1):1867. <https://doi.org/10.1186/s12889-020-09952-3>
- [46] Sorias O. Hasta ve sağlıklı öğrencilerde yaşam stresi, sosyal destek ve ruhsal hastalık ilişkisinin incelenmesi. *Ege Üniversitesi Edeb Fakültesi Yayın* 1992;9:33–49.
- [47] Carmassi C, Gesi C, Corsi M, Cremone IM, Bertelloni CA, Massimetti E, et al. Exploring PTSD in emergency operators of a major University Hospital in Italy: a preliminary report on the role of gender, age, and education. *Ann Gen Psychiatry* 2018;17:17. <https://doi.org/10.1186/s12991-018-0184-4>
- [48] Hosseinzadeh-Shanjani Z, Hajimiri K, Rostami B, Ramazani S, Dadashi M. Stress, Anxiety, and Depression Levels Among Healthcare Staff During the COVID-19 Epidemic. *Basic Clin Neurosci* 2020;11(2):163–70. <https://doi.org/10.32598/bcn.11.covid19.651.4>
- [49] Baykal D, Koç Tütüncü S. The perceived stress of paramedics and paramedic students on their stress coping behaviors during COVID. *J Gen Health Sci* 2021;3(2):90–101. <https://doi.org/10.51123/jgehes.2021.20>
- [50] Gouliou P, Mantas C, Dimitroula D, Mantis D, Hyphantis T. General hospital staff worries, perceived sufficiency of information and associated psychological distress during the A/H1N1 influenza pandemic. *BMC Infect Dis* 2010;10:322. <https://doi.org/10.1186/1471-2334-10-322>
- [51] Shahzad F, Du J, Khan I, Fateh A, Shahbaz M, Abbas A, et al. Perceived threat of COVID-19 contagion and frontline paramedics' agonistic behaviour: employing a stressor-strain-outcome perspective. *Int J Environ Res Public Health* 2020;17(14):5102. <https://doi.org/10.3390/ijerph17145102>
- [52] Clegg A. Occupational stress in nursing: a review of the literature. *J Nurs Manag* 2001;9(2):101–6. <https://doi.org/10.1046/j.1365-2834.2001.00216.x>
- [53] Avcı GG, Öztürk G, Azaklı N, Çekinmez ST. Özel bir grup hastanesinde çalışan hemşirelerin işe bağlı gerginlik düzeylerinin ve stresle başa çıkma tarzlarının belirlenmesi. *İzmir Kâtip Çelebi Üniversitesi Sağlık Bilim Fakültesi Derg* 2018;3(1):1–7.
- [54] Erquicia J, Valls L, Barja A, Gil S, Miquel J, Leal-Blanquet J, et al. Emotional impact of the Covid-19 pandemic on healthcare workers in one of the most important infection outbreaks in Europe. *Med Clin* 155. Barç.; 2020. p. 434–40. <https://doi.org/10.1016/j.medcli.2020.07.006>
- [55] Şahin MK, Aker S, Şahin G, Karabekiroğlu A. Prevalence of Depression, Anxiety, Distress and Insomnia and Related Factors in Healthcare Workers During COVID-19 Pandemic in Turkey. *J Community Health* 2020;45(6):1168–77. <https://doi.org/10.1007/s10900-020-00921-w>