



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Prevalence of avian respiratory viruses in broiler flocks in Egypt

Kareem E. Hassan, Salama A. S. Shany, A. Ali, Al-Hussien M. Dahshan, Azza A. El-Sawah, and Magdy F. El-Kady<sup>1</sup>

*Department of Poultry Diseases, Faculty of Veterinary Medicine, Beni-Suef University, Beni-Suef 62511, Egypt*

**ABSTRACT** In this study, respiratory viral pathogens were screened using real-time RT-PCR in 86 broiler chicken flocks suffering from respiratory diseases problems in 4 Egyptian governorates between January 2012 and February 2014. The mortality rates in the investigated flocks ranged from 1 to 47%. Results showed that mixed infection represented 66.3% of the examined flocks. Mixed infectious bronchitis (IBV) and avian influenza (AI)-H9N2 viruses were the most common infection (41.7%). Lack of AI-H9N2 vaccination and high rates of mixed infections in which AI-H9N2 is involved indicate an early AI-H9N2 infection with a potential immunosuppressive effect that predisposes for other viral infections. High pathogenic AI-H5N1 and virulent Newcastle disease virus (vNDV) infections were also detected (26.7% and 8.1%, respectively). Interestingly, co-infection of AI-H9N2 with either AIV-H5N1 or vNDV rarely resulted in high mortality. Partial cell-mediated immunity against similar

internal AI genes, as well as virus interference between AI and vNDV, could be an explanation for this. Highly prevalent IBV and AI-H9N2 were isolated and were molecularly characterized based on S1 gene hypervariable region 3 (HVR3) and hemagglutinin gene (HA) sequences, respectively. IBV strains were related to the variant group of IBV with multiple mutations in HVR3. Though AI-H9N2 viruses showed low rate of evolution in comparison to recent strains, few amino acid substitutions indicative of antibody selection pressure were observed in the HA gene. In conclusion, mixed viral infections, especially with IBV and AI-H9N2 viruses, are the predominant etiology of respiratory disease problems in broiler chickens in Egypt. Further investigations of the role of AI, IBV, and ND viruses' co-infections and interference in terms of altering the severity of clinical signs and lesions and/or generating novel reassortants within each virus are needed.

**Key words:** chicken, Egypt, H9N2, infectious bronchitis, respiratory outbreaks

2016 Poultry Science 95:1271–1280  
<http://dx.doi.org/10.3382/ps/pew068>

## INTRODUCTION

Respiratory disease outbreaks with variable mortality rates and different clinical manifestations have been increasing in Egyptian commercial chicken flocks during the last few years. Respiratory diseases represent a large problem to the poultry industry because of their multifactorial nature (Roussan et al., 2008). Different pathogens such as avian influenza (AI), virulent Newcastle disease virus (vNDV), and infectious bronchitis (IBV) are associated with high mortality rates in broiler chicken flocks (Haghighat-Jahromi et al., 2008). These pathogens are of major significance and have a large economic impact because they are able to induce disease independently or in association with each other (Roussan et al., 2008). The failure of control of poultry diseases in Egypt is a final result of retarded biosecu-

rity, shortage of vaccine coverage, and/or the mismatch between circulating and vaccine strains.

Extensive surveillance and genetic studies revealed that AI-H5N1 viruses became endemic in poultry in many countries including Egypt. Clinically unapparent, infected free-ranging ducks and geese, as well as mixed species backyard holdings, are suspected to play a pivotal role in avian influenza epidemiology in Egypt (Aly et al., 2008). The detection of the AI-H9N2 subtype during 2010 and 2011 in chickens and commercial quail was an additional challenge facing the poultry industry in Egypt (El-Zoghby et al., 2012). As both H5N1 and H9N2 viruses are endemic in Egypt, the genetic characterization of AIVs is necessary for screening the dynamic evolution of each subtype and potential reassortment events.

One of the major components of mixed infections is IBV that produces air sacculitis in chickens, which may result in condemnation of broilers during processing (Hofstad, 1984). In spite of regular vaccinations with Massachusetts (Mass) strains, IBV still has diverse effects on the poultry industry in the country. In Egypt, IBV has been recognized since the 1950s

© 2016 Poultry Science Association Inc.

Received October 26, 2015.

Accepted December 28, 2015.

<sup>1</sup>Corresponding author: [mfelkady@yahoo.com](mailto:mfelkady@yahoo.com)

(Sheble et al., 1986) with the isolation of an IB variant shown by neutralization tests to be closely related to the Dutch variant D3128 and subsequently variants related to Mass (Eid, 1998), D3128, D274, D-08880, 4/91 were characterized. Novel genotypes endemic in Egypt represented by the isolate Egypt/Beni-Suef/01 and nephropathogenic IBV strains Egypt/Beni-Suef/03, which are closely related to Israeli variant 2 and Mass serotype, respectively, were isolated from poultry (Abdel-Moneim et al., 2006). Recently, other novel genotypes were reported in Egypt and classified as Egyptian variant I (e.g., CK/Eg/BSU-1,4,5/2011) and Egyptian variant II (e.g., CK/Eg/BSU-2,3/2011) (Abdel-Moneim et al., 2012).

In spite of strict vaccination regimes, the NDV has been a very serious problem for poultry production in many countries and enormous efforts have been made for controlling the disease. NDV strains from recent outbreaks affecting poultry farms in Egypt between 2011 and 2012 were confirmed to belong to class II, genotype VII, sub genotype d (lineage 5 sub lineage d). Genotype VII is thought to be spreading in Egypt through the trading of poultry and poultry products with Middle Eastern countries and China (Mohamed et al., 2011; Radwan et al., 2013).

For avian respiratory viruses, virus isolation is the standard test for diagnosis although it tends to be costly, labor intensive, and slow (Suarez et al., 2007). Rapid diagnostic tests have been developed for detection of either the viral nucleic acids or viral antigens including rapid chromatographic tests for antigen detection, reverse transcriptase polymerase chain reaction (**RT-PCR**) and real-time RT-PCR (**rRT-PCR**) (Spackman et al., 2002).

In this study the rRT-PCR technique was utilized to investigate the current field situation of avian respiratory pathogens especially AI, vNDV, and IBV viruses in Egypt. Moreover, selected respiratory viruses were molecularly identified to monitor the genetic properties of the circulating viruses.

## MATERIAL AND METHODS

### Field Samples

Tracheal and/or oropharyngeal swabs and tissues were collected from 86 broiler chicken flocks (10 samples per flock) suffering from respiratory disease problems between January 2012 and February 2014. Samples collected varied between seasons and the largest numbers of samples (39 flocks) were collected during winter of 2012 to 2013. The area of investigation included the Beni-Suef, Fayoum, Menia, and Qalubia governorates in Egypt. Tissue samples were ground in phosphate buffer saline pH 7.0 to 7.4 containing gentamycin (50 µg/mL) and mycostatin (1,000 units/mL) in a 1:5 (w/v) dilution, centrifuged, and tissue supernatant was collected (OIE, 2014).

### Viral RNA Extraction and Real-Time RT-PCR (rRT-PCR)

The viral RNA was extracted by Bioflux viral RNA Mini Spin column kit (Bioflux, China) in accordance with manufacturer's instructions. Single step rRT-PCR assays using Verso 1-Step qRT-PCR Kit Plus ROX Vial (Thermo Scientific, US) were conducted using specific oligonucleotide primers and probes for each target virus. The final reaction volume was 25 µL including; 5 µL RNA template, 12.5 µL 2X 1-step PCR ready mix, 1.25 µL RT-enhancer, 0.25 µL Verso enzyme mix, 1 µL of each of the forward and reverse primers, and 0.25 µL probe for AI-H5N1 (Slomka et al., 2007), AI-H9N2 (Ben Shabat et al., 2010), IBV (Callison et al., 2006), and vNDV (Wise et al., 2004), together with 3.75 µL nuclease free water. Thermocycling rRT-PCR conditions were 50°C for 15 min, 95°C for 15 min, followed by 40 cycles at 95°C for 15 sec and 30 sec at 60°C (for AI-H9N2 and IBV) or at 54°C (for AI-H5N1 and NDV) with reading of fluorescence in this step.

### Virus Isolation

Five samples were selected for each of the highly prevalent respiratory viruses (i.e., IBV and AI-H9N2) based on rRT-PCR detection, seasonal and geographical distribution, and mortality rates observed (Table 1). Tracheal swabs and/or tissue suspensions were collected from diseased birds and prepared for inoculation through allantoic route of 10-day-old specific pathogen free embryonated chicken eggs) (**SPF-ECE**) (OIE, 2014). Inoculated eggs were incubated at 37°C for 96 hours and candled daily for embryo viability. In IBV positive rRT-PCR samples, 7 serial blind passages in SPF-ECE were done for isolation. All inoculated egg allantoic fluids from dead and surviving embryos were harvested and tested for hemagglutination using 1% washed chicken red blood cells (OIE, 2014).

### Conventional RT-PCR and Gene Sequencing

The viral RNA was extracted from harvested allantoic fluids by Bioflux viral RNA Mini Spin column kit (Bioflux, China) according to the manufacturer's instructions. RT-PCR used for molecular characterizations of isolates using specific oligonucleotide primers for partial amplification of target genes of AI viruses (hemagglutinin gene (**HA**) gene) (Shany, 2015), IBV (S1 gene) (Selim et al., 2013), and vNDV genes (Wise et al., 2004). Quant One Step RT-PCR Kit (TIANGEN Biotech, China) was used with 25 µL reaction volume including: 7 µL RNA template, 5 µL 5X RT-PCR enhancer, 2.5 µL 10X RT-PCR buffer, 1.5 µL of each of the forward and reverse primers (Table 1), 1.25 µL Hot master Taq-polymerase, 1 µL Super Pure dNTPS,

**Table 1.** Isolated viruses geographic distribution and history.

Virus	Isolate (acc. number)	Governorate	Age (Day)	Vaccination	Mortality	Lesions
<b>IBV</b>	Ck/Eg/BSU-FA-KB23/13 (KR010940)	Fayoum	30	IB Hitchner (D1), H5N1(D9) IB Clone (D15)	10%	Tracheal caseous plug, Enteritis
	Ck/Eg/BSU-FA-KB27/13 (KR010941)	Fayoum	38		4%	
	Ck/Eg/BSU-MN-KB44/13 (KR010942)	Menia	21	IB Hitchner (D7), H5N1(D10)	1%	
	Ck/Eg/BSU-BS-KB60/14 (KR010943)	Beni-Suef	29	IB (4/91) (D1), H9ND (D5), H5N1(D9), NDV Lasota (D17)	30%	
	Ck/Eg/BSU-BS-KB61/14 (KR010944)	Beni-Suef	16	Vaccitec Hatchery, IB (H120) (D1) H9ND (D5), IB (4/91) (D14)	4%	
<b>AIV-H9</b>	A/Ck/Eg/BSU-FA-KB1/12 (KR010954)	Fayoum	27	IB (Ma 5) (D1), IB Hitchner (D7), H5N1 (D9), IB (4/91) (D14)	6%	Caseous tracheitis, caseous plug, Enteritis Congested trachea,
	A/Ck/Eg/BSU-MN-KB15/13 (KR010955)	Menia	34	IB Hitchner (D1), H5N1 (D9), IB-Clone (D15)	6%	
	A/Ck/Eg/BSU-FA-KB55/13 (KR010956)	Fayoum	30	IB Hitchner (D7), IB (Ma 5) (D14), NDV Avinew (D17)	2%	
	A/Ck/Eg/BSU-BS-KB59/13 (KR010957)	Beni-Suef	28	IB (Ma 5) (D1), H5N1 (D10)	20%	
	A/Ck/Eg/BSU-BS-K7T/12 (KF998212)	Beni-Suef	26	IB (Ma 5) (D1), IB Hitchner (D7), H5N1(D9), IB-Clone (D15)	18%	
<b>AIV-H5</b>	A/Ck/Eg/BSU-BS-KF27/12 (KR010950)	Fayoum	31	IB (Ma 5) (D1), IB Hitchner (D7)	16%	General congestion, trachitis Shank Hemorrhage
	A/Ck/Eg/BSU-BS-KF28/12 (KR010951)	Fayoum	42	IB Hitchner (D7), IB-Clone (D15)	15%	
	A/Ck/Eg/BSU-BS-KB56/13 (KR010952)	Beni-Suef	34	IB Primer (D1), H9ND (D5), H5N1(D10), NDV Lasota (D17)	30%	
	A/Ck/Eg/BSU-BS-KB62/14 (KR010953)	Beni-Suef	32	IB Hitchner (D7), NDV Avinew (D17)	23%	
<b>vNDV</b>	VII/Ck/Eg/BSU-BS-KN1/13 (KR010945)	Beni-Suef	34	IB (Ma 5) (D1), IB Hitchner (D6), IB-Clone (D14), NDV Avinew (D20)	13%	Hemorrhage on proventricular glands
	VII/Ck/Eg/BSU-BS-KN2/13 (KR010946)	Beni-Suef	27	IB Hitchner (D1), H5N1(D9) IB Clone (D17)	3%	
	VII/Ck/Eg/BSU-BS-KB57/13 (KR010947)	Beni-Suef	27	IB Hitchner (D7), NDV Lasota (D17)	13%	
	VII/Ck/Eg/BSU-BS-KB58/13 (KR010948)	Beni-Suef	30		8%	

0.25  $\mu$ L RNasin, 0.25  $\mu$ L Quant RTase, and 4.75  $\mu$ L RNase free water. Thermocycling RT-PCR conditions were 50°C for 30 min, 95°C for 2 min, followed by 35 cycles at 95°C for 15 sec and 30 sec at 50 to 52°C according to the oligonucleotide primers, final extension was performed at 65°C for 10 min.

Amplified RT-PCR products were purified using PCR purification Kit (Thermo Scientific) according to manufacturer's instructions and were then sequenced directly using the ABI Prism 3100 automated sequencing machine (Applied Biosystems, Foster City, CA). A BLAST search was conducted for each

**Table 2.** Real-time RT-PCR detection of IBV, H9N2, H5N1, and vNDV viruses from broiler flocks.

	Single viral infection				Mixed viral infection							Total
	IB	H9	H5	VNDV	IB-H9	IB-VND	IB-H5	H9-H5	IB-H9-H5	IB-H5-ND	IB-H5-H9-VND	
Total IBV	16	–	–	–	36	2	7	–	9	1	1	72 (83.7)
Total H9	–	6	–	–	36	–	–	1	9	–	1	53 (61.6%)
Total H5	–	–	4	–	–	–	7	1	9	1	1	23 (26.7%)
Total vNDV	–	–	–	3	–	2	–	–	–	1	1	7 (8.1%)
No. (%)	16 (18.6%)	6 (7%)	4 (4.6%)	3 (3.5%)	36 (41.9%)	2 (2.3%)	7 (8%)	1 (1.2%)	9 (10.5%)	1 (1.2%)	1 (1.2%)	86 (100%)

sequence (<http://www.ncbi.nlm.nih.gov/BLAST>). Sequence comparisons and phylogenetic relationships through a bootstrap of 1,000 trials were determined with the MEGA version 6 program using the Clustal W alignment algorithm (Tamura et al., 2013), also Nucleotide and amino acid identities were determined using Geneious 7.1.3 (Biomatters Ltd, New Zealand).

## RESULTS

### Clinical Finding and Mortality Rates of Examined Flocks

Sixty-eight broiler chicken flocks suffering from respiratory affections from the Beni-Suef, Fayoum, Menia, and Qalubia governorates were examined (supplementary Table 1). Clinical manifestations and postmortem lesions in investigated respiratory disease outbreaks varied according to the infecting virus strain, the vaccination program of the flock, and whether the disease was due to single or multiple infections. The main clinical manifestations were respiratory distress in the form of gasping, rales, and nasal discharges. In some flocks nervous manifestations were observed in the form of head tilting. Tracheal caseation at tracheal bifurcation, tracheitis, and congested viscera were the main pathological features noticed upon postmortem examination. Typical clinical findings of AI-H5N1 and vNDV infected flocks were observed (supplementary Table 1). The incidence of clinical disease was 59.3% in young birds (20–30 days old), which was higher than that of birds aged 31–40 days or those above 40 days old (34.8% and 5.8%, respectively).

Mortality rates in the flocks under investigation ranged from 1 to 47% with the highest mortality rate found in a flock suffering triple IBV, AI-H5N1, AI-H9N2 infection though previous vaccinations against IBV (at 1 and 14 days-old), AI-H5N1 (at 5 days), and AI-H9N2 (at 9 days). Mortality rates in flocks co-infected with IBV and AI-H9N2 were as high as 30%, even in flocks vaccinated against both pathogens. Also AI-H9N2 single infection in some flocks resulted in respiratory signs, tracheal caseation, and 20% mortality rate. Notably, co-infection with AI-H5N1 and H9N2 did not result in very high mortality rates in 11 flocks that were found to be co-infected with both subtypes beside other viruses. Also natural infection with vNDV has resulted in 8 to 13% mortality rates.

### Avian respiratory Diseases Viruses' Detection by rRT-PCR


The results of rRT-PCR revealed that mixed infection is the most common cause of respiratory affection in Egypt and that mixed infection with IB and AIV-H9 viruses was the most common situation in the examined flocks (Table 2). Mixed infection represented 66.3% (57 flocks) while single viral infection of the tested viruses was found in 33.7% (29 flocks). High prevalence of both IBV and AI-H9N2 was observed in investigated flocks (83.7 and 61.6%, respectively). Subsequently, mixed infection with AI-H9N2 with IBV has been found to be the most common mixed infection (36 flocks) representing 41.9% of the total investigated flocks.

### Molecular Identification of Avian Respiratory Viruses

Five isolates of IBV and AI-H9N2 and 4 isolates of AI-H5N1 and vNDV viruses were molecularly identified. The IBV S1 gene, AI virus hemagglutinin gene, and the vNDV F gene sequences were submitted to GenBank and accession numbers were assigned (Table 1).

**Avian IB Viruses.** Sequences encoding S1 spike glycoprotein subunit hyper variable region (HVR 3) of IBV were analyzed. The phylogenetic analysis revealed that four of obtained IBV isolates were closely related to each other and to recent Egyptian isolates belonging to variant 2 Egyptian IBV. Only the isolate Ck/Eg/BSU-FA-KB23/2013 was distant from the other 4 isolates and clustered alone (Figure 1). Deduced amino acid alignments showed that 4 of IBV isolates obtained; Ck/Eg/BSU-FA-KB27/2013, Ck/Eg/BSU-MN-KB44/2013, Ck/Eg/BSU-BS-KB60/2014, and Ck/Eg/BSU-BS-KB61/2014 (accessions KR010941 to 44, respectively) shared 99.2 to 100% and 98.2 to 99.4% S1 nucleotide and amino acid identities, respectively. The isolates were also 98.8 to 100% and 96.3 to 100% similar to the recent Egyptian isolates based on S1 gene nucleotides and amino acid identity, respectively.

The fifth isolate Ck/Eg/BSU-FA-KB23/2013 was genetically distant from the other isolates of the current study and from currently circulating IBV variants in Egypt (83.6 to 86.4% and 82.4 to 82.9% nucleotide and amino acid identities, respectively). This isolate showed 18 amino acid substitutions at the HVR3, of which 3 successive amino



**Figure 1.** Phylogenetic analysis of partial S1 gene sequence of isolated IBV strains (Bold and ◀). Abbreviations: EG, Egypt; CK, chicken). Phylogenetic relationships through a bootstrap trial of 1,000 were determined with the MEGA version 6 using the Clustal W alignment algorithm and neighbor-joining method for tree construction.

acid substitutions at positions 304 to 306 (asparagine to lysine, phenylalanine to tyrosine, and asparagine to aspartic acid, respectively) (Figure 2). It showed relatively higher nucleotides and amino acid identity to the IS/885/00 isolate (88.4 and 85.7% nucleotide and amino acid identities, respectively).

Amino acid substitutions observed in IBV isolates include glutamine to histidine at position 296 (QTA/HTA), while isolate Eg/BSU-FAY-B23/2013 showed glutamine to serine (QTA/STA). In another VN epitope, amino acid substitution arginine to leucine was observed at position 378 (PRG/PRL).

	240	250	260	270	280	290	300	310
JX174185_IBV_Ck/Eg/BSU-2/2011	-----LYPFTNTS	LVKKEKVVYRE	ENSVNITLVL	TNFTFHNESNA	HPNNGGVHT	ISLYQTHTA	QSGYYNFNS	FLSSFV
KR010940_Ck/Eg/BSU-FA-KB23/13	TGNFSDGF	.....I.....R.....	.....I.....N.....R.....S.....S.....S.....FT.F.....S.....A.....KYD.....G.....					
KR010941_Ck/Eg/BSU-FA-KB27/13	TGNFSDG	.....Y.....						
KR010942_CK/Eg/BSU-MN-KB44/13	TGNFSDG	.....Y.....						
KR010943_CK/Eg/BSU-BS-KB60/14	TGNFSDG	.....Y.....						
KR010944_CK/Eg/BSU-BS-KB61/14	TGNFSDG	.....Y.....						
KC197202_IBV_ck/Eg/12vir6109/12	---FSDG	.....Y.....						
AY279533_IBV_IS/885	TGNFSDGF	.....F.....Y.....S.....V.....N.....N.....						.....Y.....L.....G.....
JX174183_IBV_Eg/Beni-Suef/01	-----F.....V.....W.....T.....	.....T.....V.....S.....I.....N.....NI.....Q.....						
X15832_IBV_strain_D274	TGNFSDGF	.....S.....I.....S.....E.....S.....V.....T.....T.....Q.....Q.....S.....L.....I.....						
AY561713_IBV(strain_Ma5)	TGNFSDGF	.....S.....Q.....I.....	.....FT.H.....	.....TG.N.....PS.....QN.QT.....Q.....				
M21970_IBV(strain_H120)	TGNFSDGF	.....S.....Q.....I.....	.....FT.H.....	.....TG.N.....PS.....QN.QT.....Q.....				

	320	330	340	350	360	370	380	▼
JX174185_IBV_Ck/Eg/BSU-2/2011	YKPSDF	MYGSYHPQC	DFRPETINNG	LWFNSLSVSL	AYGPLQGGCK	QSVFNRRATCC	YAYSNGPRL	----
KR010940_Ck/Eg/BSU-FA-KB23/13	.....	.....K.S.....			.....T.....	.....S.K.....		C----
KR010941_Ck/Eg/BSU-FA-KB27/13	.....							CKGVY
KR010942_CK/Eg/BSU-MN-KB44/13	.....		.....L.....					CKGVY
KR010943_CK/Eg/BSU-BS-KB60/14	.....		.....L.I.....					CKGVY
KR010944_CK/Eg/BSU-BS-KB61/14	.....		.....L.I.....					CKGVY
KC197202_IBV_ck/Eg/12vir6109/12	.....							CKGVY
AY279533_IBV_IS/885	.....A.....	.....K.....						C----
JX174183_IBV_Eg/Beni-Suef/01	.....Q.....	.....K.....				.....S.....		CKGVY
X15832_IBV_strain_D274	.....A.Y.....	.....S.K.L.....		.....G.....I.....	.....A.....	.....S.....		CKGVY
AY561713_IBV(strain_Ma5)	.....E.N.....	.....S.N.L.....		.....I.....	.....SG.....	.....G.....L.....		CKGVY
M21970_IBV(strain_H120)	.....E.N.....	.....S.N.L.....		.....I.....	.....SG.....	.....G.....L.....		CKGVY

Dots indicate identical amino acids; arrow heads (▼) denotes the HVR-3. Black underlined letters are: glycosylation sites (NXT/S) except where X = P). Dotted boxes indicate important sites reported to be responsible for virus neutralization epitopes (Moore et al., 1997).

**Figure 2.** Deduced amino acid sequences of S1 gene of IBV strains in comparison to selected variant and classical strains and vaccines. IBV numbering is according to IBV strain H120 Strain (accession M21970).

**AIV-H9N2 Viruses.** The partial HA gene sequences of AI-H9N2 viruses showed that all isolates obtained were related to each other and also related to recent Egyptian and Middle East circulating H9N2 strains, which belong to G1-like lineage (Figure 3). Results also showed that all AI-H9N2 isolates share 99.2% nucleotide and amino acid identities; while they share 97.1 to 97.9% nucleotide and 96.2 to 98.5% amino acid identities with recent Egyptian strain A/Ck/Egypt/S4454E/2011 (accession CY110927) and currently used vaccine strain A/Ck/Egypt/114940v/2011(accession JQ440373).

Deduced amino acid sequence analysis of the HA of AI-H9N2 showed only 4 amino acid substitutions in the signal peptide. Other mutations including serine to threonine in the overlapping area of antigenic epitope at position 127 and glycine to alanine at the receptor binding site at position 128 were observed in the HA of the isolate Ck/Eg/BSU-FA-KB1/2012 (Figure 4). Another isolate Ck/Eg/BSU-BS-KB59/2013 (accession KR010957) also had an amino acid substitution threonine to arginine at the antigenic epitope 2 at position 182 and alanine to valine in the previously reported receptor binding site at position 180 (H9 numbering).


**AIV-H5N1 and vNDV Viruses.** All AI-H5N1 isolates obtained in the current study (Table 1) were closely related to each other and recent Egyptian strains which belong to clade 2.2.1 (class C). They shared 97.2 to 99% nucleotide identities and 99.2 to 100% amino acid identities with the recent Egyptian strains. The vNDV isolates phylogeny revealed that all isolates were classified as genotype VIIId (lineage 5d class II) responsible of vNDV outbreaks in Egypt with no significant genetic changes (data not shown).

## DISCUSSION

The rapid development of the poultry industry in Egypt as well as international trade and movement of poultry has been accompanied with the emergence and spread of many viral diseases (Abdelwhab et al., 2010). Thus ecology and epidemiology of different avian pathogens are now of worldwide concern. In this study, the prevalence of viral pathogens, with special focus on AI, IBV and NDV, was investigated in broiler chickens suffering respiratory disease problems.

There were widespread respiratory disease problems in the winter season where the cold climate is thought to favor virus survival, a high ammonia level, and bad ventilation in different farms, and at least in part, is associated with increased poultry stocking capacity in Egypt (Seififi et al., 2010). The onset of clinical respiratory disease and the highest mortalities in investigated flocks were mainly reported as early as 20 days of age. The lack of vaccination against AI-H9N2 in the majority of the investigated flocks could explain the appearance of disease signs at early ages (supplementary Table 1). Moreover, the high rate of mixed respiratory viral infections observed further indicates a potential immunosuppressive role of AI-H9N2. It was reported that early H9N2 infection causes atrophy and lymphoid depletion of the thymus, and probably also some other lymphoid organs, causing immunosuppression and predispose the chickens to secondary infections (Hadipour et al., 2011).

Data in the current study revealed that IBV alone can cause high mortality (up to 19%) even in vaccinated flocks. Mortalities increased with co-infection of IBV and AIV-H9 ( $\geq 30\%$ ) even in flocks vaccinated against




**Figure 3.** Phylogenetic analysis of partial the HA gene sequence of isolated AIV-H9N2 strains (Bold and ◀). Abbreviations: EG, Egypt; CK, chicken). Phylogenetic relationships through a bootstrap trial of 1,000 were determined with the MEGA version 6 using the Clustal W alignment algorithm and neighbor-joining method for tree construction.

both pathogens. Also AI-H9N2 infection alone resulted in severe respiratory signs, tracheal caseation, and 20% mortality rate. These data highlight the pathogenicity of AI-H9N2 as previously reported (Nili and Asasi, 2002). Mixed infections of AI-H9N2 with other respiratory pathogens (particularly IBV, *Mycoplasma gallisept-*

*ticum*, and *Escherichia coli*) are thought to be responsible for severe clinical disease with subsequent high mortality (Nili and Asasi, 2003; Haghghat-Jahromi et al., 2008). Though not analyzed in this study, the role of secondary agents as bacteria, and environmental factors in the AI-H9N2 pathogenesis, cannot be neglected.





Dots indicate identical amino acids.

H9 antigenic sites (Kaverin *et al.*, 2004a) are underlined [blue (—): epitope I, olive green (—): epitope II, pink (—): overlapping sites].

Dark red (—) underlined letters: antigenic sites reported in other publications.

Black underlined letters are potential N glycosylation sites.

Stars denote the receptor binding site (\*, \*, \*—left side, pocket and right side, respectively).

Gray shaded letters are potential O glycosylation sites.

H9 numbering is indicated and followed by H3 numbering in parenthesis.

□: Cleavage site /: HA1-HA2 connection ---: Partial sequencing

**Figure 4.** Deduced amino acid sequences of the HA protein of Egyptian AIV-H9N2 strains in comparison to selected Egyptian strains.

Interestingly, co-infection with AI-H5N1 and H9N2 infrequently resulted in very high mortality rates. The early exposure to H9N2 viruses possibly provoked cell-mediated immunity against H5N1 due to their similar internal genes. Therefore, partial protection by what is called the “protected window” after H9N2 infection can mask the lethal infection of H5N1 that may go unnoticed while the virus is shed by infected birds. This is a potentially important problem for countries such as Egypt that use H5N1 vaccines and cull flocks only when clinical disease is evident (Khalenkov *et al.*, 2009). Although co-circulation of H5N1 and H9N2 has been suggested to limit the spread and the epizootiologic pattern of the infections for both subtypes (Arafa *et al.*, 2013), their co-circulation in susceptible host populations can increase the likelihood of generating novel reassortant viruses with public health implications (Guan *et al.*, 1999).

Low mortality rates in vNDV infected flocks may be the result of partial protection offered by previous vaccination with live lentogenic NDV vaccines (Hadipour *et al.*, 2011; Munir *et al.*, 2012). Moreover, speculation regarding viral interference exerted by wide and rapid spread of AI-H9N2 LPAI may explain the low mortalities caused by vNDV. Viral interference could be explained by different mechanisms including: competing by attachment interference or blocking of receptor sites for the superinfecting virus; competing intracellularly

for replication host machinery; and virus-induced interferon (Ge *et al.*, 2007).

The virus-specific rRT-PCR showed that mixed infections are the predominant etiology of respiratory diseases, especially co-infection with the IBV and AI-H9N2 viruses. The high prevalence of IBV (Abdel-Moneim *et al.*, 2006) and AI-H9N2 (El-Zoghby *et al.*, 2012) in Egyptian poultry has previously been reported. Higher rates of co-infection with both viruses further indicate a potential role for IBV in increasing the severity of AI-H9N2 infection through HA cleavage activation by providing trypsin-like proteases encoded by coronavirus IBV (Klenk and Garten, 1994; Ng and Liu, 2000; Perk *et al.*, 2004; Haghghat-Jahromi *et al.*, 2008).

It is worth mentioning that rRT-PCR assay alone cannot differentiate between field and vaccine strains of IBV and S1 gene sequencing is thought to be the only method used to discriminate between all IBV strains. Therefore, partial IBV S1 gene phylogenetics were analyzed and it was revealed that 4 isolates were closely related to recent Egyptian IBV isolate Eg/BSU-2/2011 (accession JX174185) that was isolated in northern and middle Egypt and showed a unique genetic signature and subsequently was suggested as a new genotype variant 2 (Abdel-Moneim *et al.*, 2012). One isolate, Ck/Eg/BSU-FA-KB23/2013 (accession KR010940) was distant from the other 4 isolates and it was relatively closer to the nephropathogenic Israeli isolate IS/885/00

(Meir et al., 2004). The strain (IS/885/00) also shared 96.6% S1 similarity to the Egyptian isolate Eg/Beni-Suef/01 (accession JX174183).

Avian IBV serotypic determinants have been identified in the first 395 amino acid region of the S1 subunit, which contains three major hyper variable regions (HVRs). The HVRs are associated with virus-neutralizing antigenic sites and are located between the amino acid residues 38 to 67 (HVR-1), 97 to 141 (HVR-2), and 274 to 387 (HVR-3) (Moore et al., 1997). In this study, amino acid substitutions at virus neutralization (VN) critical antigenic epitopes were observed in IBV isolates. At position 296; QTA/HTA or QTA/STA and position 378; PRG/PRL substitutions were previously documented to cause reduced vaccine protection and generation of IBV escape mutant (Yu et al., 2001). Additionally, the widespread use of various vaccines, including illegally imported heterologous IBVs vaccines, may also exert pressure resulting in the increase of new genetic variants. In addition to serotype changes, the genetic variation may result in changes of the tissue tropism and pathogenicity of the virus (Cavanagh et al., 1992).

Both groups A and B of the G1-like lineage of AI-H9N2 viruses have been circulating extensively in Middle Eastern countries since 1999 (Bashashati et al., 2013). All Egyptian isolates, including those of the current study, are grouped within group B of G1-like lineage according to (Monne et al., 2012). Though the isolated H9N2 viruses showed minor evolution based on genetic analysis of their HA gene, few amino acid substitutions, which are indicative of antibody selection pressure on H9N2 viruses in the field, were determined. Among those substitutions which were observed were that in the overlapping area of antigenic epitope at position 127 previously reported by (Kaverin et al., 2004), receptor binding sites 128 and 180 (H9 numbering) (Wan et al., 2014), and antigenic epitope 2 at position 182 (Figure 4). The later substitution (i.e., position 182) is unique and was reported as indicative of antibody selection pressure on H9N2 viruses in the field (Wan et al., 2014).

To conclude, the co-infection of IBV and AI-H9N2 are the main causes of increased severity and high mortality rates of field outbreaks of respiratory infections in broiler chickens in Egypt. Circulation of variant IBV strains with multiple mutations at the virus neutralizing epitopes in the S1 genes with continuous emergence of new variants were observed. Public health implications may arise due to the co-circulation of AI-H9N2 with H5N1 in susceptible host populations. The AI, IBV, and NDV viruses' interference and co-infections in terms of altering the severity of clinical signs and lesions need further investigation. Identification of factors that influence avian respiratory virus interference will provide new insights in the pathogenesis and subsequently improvement of control programs could be achieved.

## ACKNOWLEDGMENTS

This study was supported by a small project research grant from the Project Support and Financing Unit, Research Center, Beni-Suef University, Egypt.

## SUPPLEMENTARY DATA

**Table S1.** Clinical history, vaccination programs and mortalities of examined chicken flocks.

Supplementary data is available at *PSA Journal* online.

## REFERENCES

- Abdel-Moneim, A. S., M. A. Affi, and M. F. El-Kady. 2012. Emergence of a novel genotype of avian infectious bronchitis virus in Egypt. *Arch. Virol.* 157:2453–2457.
- Abdel-Moneim, A. S., M. F. El-Kady, B. S. Ladman, and J. Gelb, Jr. 2006. S1 gene sequence analysis of a nephropathogenic strain of avian infectious bronchitis virus in Egypt. *Virol J.* 3:78. doi: 10.1186/1743-422X-3-78.
- Abdelwhab el, S. M., A. S. Arafa, A. M. Erfan, M. M. Aly, and H. M. Hafez. 2010. Modified H5 real-time reverse transcriptase-PCR oligonucleotides for detection of divergent avian influenza H5N1 viruses in Egypt. *Avian Dis.* 54:1301–1305.
- Aly, M. M., A. Arafa, and M. K. Hassan. 2008. Epidemiological findings of outbreaks of disease caused by highly pathogenic H5N1 avian influenza virus in poultry in Egypt during 2006. *Avian Dis.* 52:269–277.
- Arafa, A. S., N. M. Hagag, N. Yehia, A. M. Zanaty, M. M. Naguib, and S. A. Nasef. 2013. Effect of cocirculation of highly pathogenic avian influenza H5N1 subtype with low pathogenic H9N2 subtype on the spread of infections. *Avian Dis.* 56:849–857.
- Bashashati, M., M. Vafsi Marandi, and F. Sabouri. 2013. Genetic diversity of early (1998) and recent (2010) avian influenza H9N2 virus strains isolated from poultry in Iran. *Arch. Virol.* 158:2089–2100.
- Ben Shabat, M., R. Meir, R. Haddas, E. Lapin, I. Shkoda, I. Raibstein, S. Perk, and I. Davidson. 2010. Development of a real-time TaqMan RT-PCR assay for the detection of H9N2 avian influenza viruses. *J. Virol. Methods.* 168:72–77.
- Callison, S. A., D. A. Hilt, T. O. Boynton, B. F. Sample, R. Robison, D. E. Swayne, and M. W. Jackwood. 2006. Development and evaluation of a real-time Taqman RT-PCR assay for the detection of infectious bronchitis virus from infected chickens. *J. Virol. Methods.* 138:60–65.
- Cavanagh, D., P. J. Davis, and J. K. A. Cook. 1992. Infectious bronchitis virus: evidence for recombination within the Massachusetts serotype. *Avian Pathol.* 21:401–408.
- Eid, A. M. 1998. Infectious bronchitis virus infection in Egypt. Pages 145–156. *Proc. the International Symposium on infectious bronchitis and pneumovirus infections in Poultry, Rauischholzhausen, Germany.*
- El-Zoghby, E. F., A. S. Arafa, M. K. Hassan, M. M. Aly, A. Selim, W. H. Kilany, U. Selim, S. Nasef, M. G. Aggor, E. M. Abdelwhab, and H. M. Hafez. 2012. Isolation of H9N2 avian influenza virus from bobwhite quail (*Colinus virginianus*) in Egypt. *Arch. Virol.* 157:1167–1172.
- Ge, J., G. Deng, Z. Wen, G. Tian, Y. Wang, J. Shi, X. Wang, Y. Li, S. Hu, Y. Jiang, C. Yang, K. Yu, Z. Bu, and H. Chen. 2007. Newcastle disease virus-based live attenuated vaccine completely protects chickens and mice from lethal challenge of homologous and heterologous H5N1 avian influenza viruses. *J. Virol.* 81:150–158.
- Guan, Y., K. F. Shortridge, S. Krauss, and R. G. Webster. 1999. Molecular characterization of H9N2 influenza viruses: were they the donors of the “internal” genes of H5N1 viruses in Hong Kong? *Proc. Natl. Acad. Sci. USA.* 96:9363–9367.

- Hadipour, M. M., G. H. Habibi, P. Golchin, M. R. Hadipourfard, and N. Shayanpour. 2011. The role of avian influenza, newcastle disease and infectious bronchitis viruses during the respiratory disease outbreak in commercial broiler farms of Iran. *Int. J. Anim. Vet. Adv.* 3:69–72.
- Haghighat-Jahromi, M., K. Asasi, H. Nili, H. Dadras, and A. H. Shooshtari. 2008. Coinfection of avian influenza virus (H9N2 subtype) with infectious bronchitis live vaccine. *Arch. Virol.* 153:651–655.
- Hofstad, M. S. 1984. Avian infectious bronchitis. Pages 429–443 in *Diseases of Poultry*. M. S. Hofstad, H. J. Barnes, B. W. Calnek, W. M. Reid, and H. W. Yoder, eds. Iowa State University Press: Ames, IA.
- Kaverin, N. V., I. A. Rudneva, N. A. Ilyushina, A. S. Lipatov, S. Krauss, and R. G. Webster. 2004. Structural differences among hemagglutinins of influenza A virus subtypes are reflected in their antigenic architecture: analysis of H9 escape mutants. *Viol.* 78:240–249.
- Khalenkov, A., S. Perk, A. Panshin, N. Golender, and R. G. Webster. 2009. Modulation of the severity of highly pathogenic H5N1 influenza in chickens previously inoculated with Israeli H9N2 influenza viruses. *Virology*. 383:32–38.
- Klenk, H. D., and W. Garten. 1994. Host cell proteases controlling virus pathogenicity. *Trends Microbiol.* 2:39–43.
- Meir, R., E. Rosenblut, S. Perl, N. Kass, G. Ayali, S. Perk, and E. Hemsani. 2004. Identification of a novel nephropathogenic infectious bronchitis virus in Israel. *Avian Dis.* 48:635–641.
- Mohamed, M. H., S. Kumar, A. Paldurai, and S. K. Samal. 2011. Sequence analysis of fusion protein gene of Newcastle disease virus isolated from outbreaks in Egypt during 2006. *Viol. J.* 18:237. doi: 10.1186/1743-422X-8-237.
- Monne, I., H. A. Hussein, A. Fusaro, V. Valastro, M. M. Hamoud, R. A. Khalefa, S. N. Dardir, M. I. Radwan, I. Capua, and G. Cattoli. 2012. H9N2 influenza A virus circulates in H5N1 endemically infected poultry population in Egypt. *Influenza Other Respir. Viruses*. 7:240–243.
- Moore, K. M., M. W. Jackwood, and D. A. Hilt. 1997. Identification of amino acids involved in a serotype and neutralization specific epitope within the S1 subunit of avian infectious bronchitis virus. *Arch. Virol. J.* 142:2249–2256.
- Munir, M., M. Abbas, M. T. Khan, S. Zohari, and M. Berg. 2012. Genomic and biological characterization of a velogenic Newcastle disease virus isolated from a healthy backyard poultry flock in 2010. *Viol. J.* 9:01–11.
- Ng, L. F., and D. X. Liu. 2000. Further characterization of the coronavirus infectious bronchitis virus C-like proteinase and determination of a new cleavage site. *Virology*. 272:27–39.
- Nili, H., and K. Asasi. 2002. Natural cases and an experimental study of H9N2 avian influenza in commercial broiler chickens of Iran. *Avian Pathol.* 31:247–252.
- Nili, H., and K. Asasi. 2003. Avian influenza (H9N2) outbreak in Iran. *Avian Dis.* 47:828–831.
- OIE. 2014. *Manual of Diagnostic Tests and Vaccines for Terrestrial Animals*, chapter 2.3.4. Avian influenza. Accessed Nov. 2015. [http://www.oie.int/fileadmin/Home/fr/Health\\_standards/tahm/2.03.04\\_AI.pdf](http://www.oie.int/fileadmin/Home/fr/Health_standards/tahm/2.03.04_AI.pdf).
- Perk, S., S. Pokamunski, N. Elkin, and B. Perelman. 2004. Low pathogenicity avian influenza H9N2 in Israel a threat to the poultry industry. Pages 72–80. *Proc. 5<sup>th</sup> International Symposium on Turkey Diseases*.
- Radwan, M. M., S. F. Darwish, I. M. El-Sabagh, A. A. El-Sanousi, and M. A. Shalaby. 2013. Isolation and molecular characterization of Newcastle disease virus genotypes II and VIId in Egypt between 2011 and 2012. *Virus Genes*. 47:311–316.
- Roussan, D. A., R. Haddad, and G. Khawaldeh. 2008. Molecular survey of avian respiratory pathogens in commercial broiler chicken flocks with respiratory diseases in Jordan. *Poult. Sci.* 87:444–448.
- Seififi, S., K. Asasi, and A. Mohammadi. 2010. Natural co-infection caused by avian influenza H9 subtype and infectious bronchitis viruses in broiler chicken farms. *Veterinarski Arhiv.* 80:269–281.
- Selim, K., A. Arafat, A. Hussein, and A. A. El-Sanousi. 2013. Molecular characterization of infectious bronchitis viruses isolated from broiler and layer chicken farms in Egypt during 2012. *Int. J. Vet. Sci. Med.* 1:102–108.
- Shany, S. A. 2015. *Further Studies on the Current Situation of Avian Influenza in Egypt*. PhD Diss. Beni-Suef University, Egypt.
- Sheble, A., M. Z. Sabry, F. G. Davelaar, A. G. Burger, A. R. Khafagy, M. M. Moustafa, and M. Henna. 1986. Present status of infectious bronchitis in Egypt. *Journal of the Egyptian Veterinary Medical Association*. 46:393–411.
- Slomka, M. J., T. Pavlidis, J. Banks, W. Shell, A. McNally, S. Esen, and I. H. Brown. 2007. Validated H5 Eurasian real-time reverse transcriptase-polymerase chain reaction and its application in H5N1 outbreaks in 2005–2006. *Avian Dis.* 51:373–377.
- Spackman, E., D. A. Senne, T. J. Myers, L. L. Bulaga, L. P. Garber, M. L. Perdue, K. Lohman, L. T. Daum, and D. L. Suarez. 2002. Development of a real-time reverse transcriptase PCR assay for type A influenza virus and the avian H5 and H7 hemagglutinin subtypes. *J. Clin. Microbiol.* 40:3256–3260.
- Suarez, D. L., A. Das, and E. Ellis. 2007. Review of rapid molecular diagnostic tools for avian influenza virus. *Avian Dis.* 51:201–208.
- Tamura, K., G. Stecher, D. Peterson, A. Filipinski, and S. Kumar. 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol. Biol. Evol.* 30:2725–2729.
- Wan, Z., J. Ye, L. Xu, H. Shao, W. Jin, K. Qian, H. Wan, and A. Qin. 2014. Antigenic mapping of the hemagglutinin of an H9N2 avian influenza virus reveals novel critical amino acid positions in antigenic sites. *J. Virol.* 88:3898–3901.
- Wise, M. G., D. L. Suarez, B. S. Seal, J. C. Pedersen, D. A. Senne, D. J. King, D. R. Kapczynski, and E. Spackman. 2004. Development of a real-time reverse-transcription PCR for detection of Newcastle disease virus RNA in clinical samples. *J. Clin. Microbiol.* 42:329–338.
- Yu, L., Y. Jiang, S. Low, Z. Wang, S. J. Nam, W. Liu, and J. Kwangac. 2001. Characterization of three infectious bronchitis virus isolates from China associated with proventriculus in vaccinated chickens. *Avian Dis.* 45:416–424.