

Efficacy, safety, and tolerability of a ready-to-drink bowel preparation in overweight and obese adults: subanalysis by body mass index from a phase III, assessor-blinded study

Lawrence Hookey, Gerald Bertiger^{ID}, Kenneth Lee Johnson II, Mena Boules, Masakazu Ando and David N. Dahdal

Abstract

Background: We performed a *post hoc* secondary analysis for the effect of body mass index (BMI) on the efficacy, tolerability, and safety of ready-to-drink sodium picosulfate, magnesium oxide, and citric acid (SPMC oral solution) bowel preparation.

Methods: A phase III, randomized, assessor-blinded, multicenter, noninferiority study was conducted comparing split-dose, low-volume SPMC oral solution with a powder formulation for oral solution. A *post hoc* secondary analysis assessed efficacy, safety, and tolerability of SPMC oral solution stratified by BMI. BMI was classified by Centers for Disease Control and Prevention definitions (underweight and normal weight: BMI < 25 kg/m²; overweight: BMI 25–29.9 kg/m²; class I obesity: BMI 30–34.9 kg/m²; class II obesity: BMI 35–39.9 kg/m²; class III/severe obesity: BMI ≥ 40 kg/m²). Prespecified primary efficacy endpoint ('responders') was the proportion of participants with 'excellent' or 'good' ratings on a modified Aronchick Scale (AS). Secondary efficacy outcomes were the quality of cleansing of the right colon as assessed by the Boston Bowel Preparation Scale (BBPS); as well as selected findings from the Mayo Clinic Bowel Prep Tolerability Questionnaire. Safety assessments included adverse events (AEs) and laboratory evaluations.

Results: Between 82.8% and 92.5% of participants in any BMI group were responders by AS, and between 91.3% and 100% were responders by BBPS in the right colon. Efficacy was consistent across BMI groups, with no clear trends. Greater than 83% of participants in any BMI group found the preparation 'easy' or 'acceptable' to ingest, and the majority (>58%) rated SPMC oral solution as 'better' than a prior bowel preparation. In all BMI groups, safety data were similar to the overall cohort. Commonly reported, drug-related, treatment-emergent AEs were, by ascending BMI group, nausea (1.1%, 5.3%, 1.0%, 5.7%, and 0%) and headache (1.1%, 4.1%, 1.0%, 5.7%, and 0%).

Conclusions: Ready-to-drink SPMC oral solution had consistent, good quality colon cleansing, and favorable tolerability among participants of all BMI groups.

ClinicalTrials.gov Registration: NCT03017235

Keywords: BMI, bowel preparation, colon cleansing, screening colonoscopy, inadequate bowel preparation, oral solution, obesity, overweight

Received: 8 October 2019; revised manuscript accepted: 5 February 2020.

Ther Adv Gastroenterol

2020, Vol. 13: 1–10

DOI: 10.1177/
1756284820910050

© The Author(s), 2020.
Article reuse guidelines:
sagepub.com/journals-
permissions

Correspondence to:

Gerald Bertiger
Hillmont Gl, 1811
Bethlehem Pike, Building
C-300, Flourtown, PA
19031, USA

gbertiger@hillmontgi.com

Lawrence Hookey
Gastrointestinal
Disease Research Unit,
Department of Medicine,
Queen's University,
Kingston, ON, Canada

Kenneth Lee Johnson II
Vidant Medical Group,
Kinston, NC, USA

Mena Boules
Masakazu Ando
David N. Dahdal
Ferring Pharmaceuticals,
Inc., Parsippany, NJ, USA

Introduction

Regular colorectal cancer (CRC) screening for people over 50 years old has been shown to be an important step to reduce CRC incidence and mortality.¹⁻⁴ Furthermore, certain individuals are at higher risk of developing CRC and should follow their clinicians' recommendations to undergo CRC screening at recommended intervals.⁵⁻⁷

Adults with obesity [body mass index (BMI) ≥ 30 kg/m²] have an approximately 30% greater risk of developing CRC compared with individuals with lower BMI.^{8,9} Adults with obesity have worse disease-free survival and increased risk of cancer recurrence compared with individuals of normal BMI.¹⁰ Likewise, higher BMI is associated with poorer CRC prognosis, including a 14% increased risk of cancer-specific mortality compared with adults of normal BMI.¹¹ Adults with CRC who have low BMI (< 20 kg/m²) also have a 50% increased risk of cancer-specific mortality compared with those with normal BMI.¹⁰

An effective bowel preparation is essential for an optimal CRC screening colonoscopy,¹²⁻¹⁵ but BMI has been shown to affect the quality of bowel preparation. Higher BMI has been cited as an independent risk factor for poorer quality of bowel preparation, with estimates of up to 46% higher risk of inadequate bowel preparation for those with BMI ≥ 30 kg/m².¹⁶⁻¹⁹ However, the rates reported in the literature are not uniform, and a recent prospective, observational study did not find an association between BMI and quality of bowel preparation.²⁰ In real-world settings, the association of high BMI with poorer quality of bowel preparation may be related to nonadherence with dietary restrictions.^{21,22}

Most of the available data on bowel preparation and colonoscopy outcomes by BMI are from retrospective or cohort studies, which may introduce confounding factors into the outcomes.²¹

Results from a phase III, randomized, assessor-blinded, multicenter study of ready-to-drink sodium picosulfate, magnesium oxide, and citric acid (SPMC oral solution) have recently been described.⁶ Here, we performed a *post hoc* secondary analysis for the effect of BMI on the efficacy, tolerability, and safety of SPMC oral solution to determine whether the efficacy trends by BMI in the literature are consistent in the setting of a randomized, controlled trial. This analysis is a

follow-up analysis to the primary analysis, where the superiority of ready-to-drink SPMC oral solution was demonstrated compared with a powder formulation of the same ingredients.

Methods

Study design

A phase III, randomized, assessor-blinded, multicenter, noninferiority study was conducted comparing split-dose, low-volume SPMC oral solution (Clenpiq®, Ferring Pharmaceuticals Inc., Parsippany, NJ) with split-dose, low-volume sodium picosulfate, magnesium oxide, and citric acid powder for oral solution (Prepopik®, Ferring Pharmaceuticals Inc., Parsippany, NJ) [ClinicalTrials.gov identifier: NCT03017235]. Details of the full study have been published previously.²³ The study was conducted in accordance with the principles set forth in the Declaration of Helsinki and in compliance with ICH-GCP standards. The study protocol was approved by Schulman IRB (protocol #000253).

Eligible participants included females and males, 18–80 years of age, who were undergoing elective colonoscopy (screening, surveillance, or diagnostic). (Full inclusion and exclusion criteria have been published previously.²³)

Eligible participants must have had an average of at least three spontaneous bowel movements per week for 1 month prior to the colonoscopy, and been willing, able, and competent to complete the procedure and comply with study instructions. Written informed consent was obtained at screening.

Interventions

The colon-cleansing regimen was a split-dose preparation, with one dose taken the evening before and one dose taken the same day as the colonoscopy, within 5–9 h prior to the procedure.

SPMC oral solution (two 5.4-oz doses) is a ready-to-drink formulation and was consumed as supplied (without mixing, stirring, or dilution), followed by five or more 8-oz glasses of clear liquid within 5 h of the first dose, and four or more 8-oz glasses of clear liquid within 4 h of the second dose.

In both cases, participants were instructed to maintain a diet of clear liquids from 24 h before

the colonoscopy and to stop taking anything by mouth 2h before. Immediately prior to the colonoscopy, participants returned the Mayo Clinic Bowel Prep Tolerability Questionnaire,²⁴ and chemistry and hematology laboratory samples were obtained. Following the colonoscopy, participants returned for visits at 1–2 days, 7 days, and 4 weeks per study protocol to assess laboratory and safety measures.

Endpoints

The primary efficacy outcome was overall quality of colon cleansing as measured by the modified Aronchick Scale (AS) prior to irrigation of the colon, assessed by a treatment-blinded endoscopist. Investigators had an average of 33 years of experience each as practicing gastroenterologists. The prespecified primary efficacy endpoint ('responders') by AS was the proportion of participants with 'excellent' or 'good' ratings.

Secondary efficacy outcomes were the quality of cleansing of the right colon as assessed by the Boston Bowel Preparation Scale (BBPS), as well as the findings of selected questions from the Mayo Clinic Bowel Prep Tolerability Questionnaire. The prespecified key secondary efficacy rate ('responders') by BBPS was the proportion of participants with a segmental score of '3' or '2' in the right colon. The proportion of participants with a BBPS score of ≥ 2 in each of the three colon segments was calculated.

Safety assessments included adverse events (AEs), laboratory evaluations, and electrocardiograms. AEs were classified according to the Medical Dictionary for Regulatory Activities (MedDRA) version 20.1.

The endoscopist noted the number of lesions found during the colonoscopy (recorded as an AE) and removed polyps when possible and appropriate. Lesion biopsies were sent for histological analysis. All malignancies found during the study period, including colonic lesions that were determined to be cancerous, were reported as a serious AE. Polyp and adenoma findings were not a key efficacy endpoint in the study.

Statistical analysis

A *post hoc* secondary analysis was performed to assess efficacy, safety, and tolerability of SPMC

oral solution. BMI was classified according to the Centers for Disease Control and Prevention (CDC) definitions (underweight and normal weight: BMI < 25 kg/m²; overweight: BMI 25–29.9 kg/m²; class I obesity: BMI 30–34.9 kg/m²; class II obesity: BMI 35–39.9 kg/m²; severe obesity: BMI ≥ 40 kg/m²).²⁵

The analysis included all participants who were randomized and received at least one dose of the study drug [modified intent-to-treat (mITT) population]. Baseline and demographic characteristics were descriptively summarized.

The responder rates in the primary, key secondary, and other secondary efficacy endpoints were summarized with exact 95% confidence intervals (CIs), calculated by the Clopper-Pearson method. Tolerability endpoints were descriptively summarized.

Adenoma detection rate (ADR) and polyp detection rate (PDR) were calculated as the proportion of participants who had at least one adenoma or polyp, respectively, in the treatment group.

Results

Of the 448 participants, most (61.2%; 274/448) had a BMI of 25–29.9 or 30–34.9 kg/m² (Table 1). There were few participants with a BMI ≥ 40 kg/m². There were two individuals (2.2%; 2/93) in the BMI < 25 kg/m² group who were 'underweight' by CDC definition (BMI < 18.5 kg/m²). The mean age was consistent across BMI groups (by ascending BMI group: 56.6 years, 57.7 years, 57.7 years, 57.5 years, 54.9 years). Type 2 diabetes prevalence increased by ascending BMI group.

Efficacy

By ascending BMI group, 82.8%, 90.1%, 85.4%, 92.5%, and 89.3% of participants receiving SPMC oral solution were responders (those with 'excellent' or 'good' ratings) for the primary efficacy endpoint, overall colon cleansing by modified AS (Table 2; Figure 1). There was no consistent pattern of responder rate by BMI. Rates of 'inadequate' rating, by ascending BMI group, were 1.1%, 0.6%, 1.9%, 0%, and 0%.

For the key secondary efficacy endpoint, responders by BBPS in the right colon (those with a rating

Table 1. Demographic and baseline characteristics in BMI subgroups, mITT population.

	BMI subgroup					Overall cohort (n = 448)
	<25 (n = 93)	25–29.9 (n = 171)	30–34.9 (n = 103)	35–39.9 (n = 53)	≥40 (n = 28)	
Age (years), mean (SD)	56.6 (11.5)	57.7 (11.6)	57.7 (10.0)	57.5 (9.6)	54.9 (12.5)	57.2 (11.0)
Female, n (%)	59 (63.4)	89 (52.0)	49 (47.6)	38 (71.7)	17 (60.7)	252 (56.3)
Race, n (%)						
White	83 (89.2)	145 (84.8)	85 (82.5)	44 (83.0)	19 (67.9)	376 (83.9)
Black/African American	4 (4.3)	18 (10.5)	11 (10.7)	7 (13.2)	9 (32.1)	49 (10.9)
Asian	6 (6.5)	6 (3.5)	-	1 (1.9)	-	13 (2.9)
Other	-	2 (1.2)	7 (6.8)	1 (1.9)	-	10 (2.2)
BMI (kg/m ²), mean (SD)	22.5 (1.7)	27.5 (1.4)	32.3 (1.3)	36.7 (1.3)	44.5 (4.5)	29.7 (6.1)
Type 2 diabetes mellitus, n (%)	4 (4.3)	15 (8.8)	14 (13.6)	15 (28.3)	11 (39.3)	59 (13.2)

BMI, body mass index; mITT, modified intent-to-treat; SD, standard deviation.

Table 2. Primary efficacy endpoint, overall colon cleansing quality by modified AS, mITT population.

% (n)	BMI subgroup					Overall cohort (n = 448)
	<25 (n = 93)	25–29.9 (n = 171)	30–34.9 (n = 103)	35–39.9 (n = 53)	≥40 (n = 28)	
Excellent	58.1 (54)	55.0 (94)	45.6 (47)	60.4 (32)	50.0 (14)	53.8 (241)
Good	24.7 (23)	35.1 (60)	39.8 (41)	32.1 (17)	39.3 (11)	33.9 (152)
Fair	11.8 (11)	8.2 (14)	11.7 (12)	5.7 (3)	10.7 (3)	9.6 (43)
Inadequate	1.1 (1)	0.6 (1)	1.9 (2)	-	-	0.9 (4)
No Rating	4.3 (4)	1.2 (2)	1.0 (1)	1.9 (1)	-	1.8 (8)
Responders ^a [95% CI for proportion]	82.8 (77) [73.6, 89.8]	90.1 (154) [84.6, 94.1]	85.4 (88) [77.1, 91.6]	92.5 (49) [81.8, 97.9]	89.3 (25) [71.8, 97.7]	87.7 (393) [84.3, 90.6]

AS, Aronchick scale; BMI, body mass index; CI, confidence interval; mITT, modified intent-to-treat.
^aResponders were those rated 'excellent' or 'good' on the modified AS by an endoscopist blinded to the treatment group, and the 95% CI of the responder rate was calculated using the Clopper-Pearson method.

of '3' or '2'), overall 94.2% of participants receiving SPMC oral solution were responders. There was no consistent pattern in responder rate by BBPS in the right colon across BMI groups (by ascending BMI group: 92.5%, 95.3%, 91.3%, 96.2%, 100%) (Table 3). The mean total BBPS score by BMI group was similar to the overall population score of 7.7 (out of a possible 9; Table 3). At least 87% of participants in each BMI

group had a BBPS score of 2 or better in each of the three colon segments (Figure 1; Table 3).

Tolerability

Depending on the BMI group, 96.4–100% of participants were able to complete the majority of the bowel preparation (at least 75% of preparation consumed). At least 83.0% of participants in




Figure 1. In the SPMC oral solution treatment arm, a substantial majority of participants in each BMI group were responders on the modified AS, rated by a treatment-blinded endoscopist. Likewise, at least 87% of participants had a BBPS score of 2 or better in all 3 colon segments. The efficacy ratings were consistent across BMI groups.

AS, Aronchick scale; BBPS, Boston Bowel Preparation Scale, BMI, body mass index; SPMC, sodium picosulfate, magnesium oxide, and citric acid.

each BMI group stated that SPMC oral solution was ‘easy’ or ‘acceptable’ to ingest (Figure 2). Of participants who had experience with a prior

colonoscopy, the majority (>58%) of those in the SPMC oral solution arm rated the preparation as ‘better’ than a prior bowel preparation (Figure 3). There were no clear, consistent trends of tolerability by BMI in these data.

Polyp detection

PDR varied by BMI group, without any observable trend in the data (Figure 4). PDR ranged from 39.3% for those with BMI ≥ 40 kg/m² to 62.3% for those with BMI 35–39.9 kg/m². ADR values were similar across BMI groups, with no consistent trend (by increasing BMI group: 30.1%, 28.7%, 33.0%, 39.6%, and 32.1%).

Safety

Across all BMI groups, there were no deaths, no treatment-emergent adverse events (TEAEs) leading to study discontinuation, and no serious adverse drug reactions (Table 4). Rates of serious TEAEs were $\leq 5.7\%$ in any subgroup, with none reported for those with BMI ≥ 40 kg/m², and rates of severe TEAEs were $\leq 7.5\%$ in any subgroup. Rates of adverse drug reactions across BMI

Table 3. Colon cleansing quality by BBPS, mITT population.

% (n)	BMI subgroup					Overall cohort (n=448)
	<25 (n=93)	25–29.9 (n=171)	30–34.9 (n=103)	35–39.9 (n=53)	≥ 40 (n=28)	
Key secondary efficacy endpoint, right colon cleansing						
3	52.7 (49)	53.2 (91)	44.7 (46)	58.5 (31)	50.0 (14)	51.6 (231)
2	39.8 (37)	42.1 (72)	46.6 (48)	37.7 (20)	50.0 (14)	42.6 (191)
1	3.2 (3)	3.5 (6)	7.8 (8)	1.9 (1)	-	4.0 (18)
0	-	-	-	-	-	-
No rating	4.3 (4)	1.2 (2)	1.0 (1)	1.9 (1)	-	1.8 (8)
Responders ^a [95% CI for proportion]	92.5 (86) [85.1, 96.9]	95.3 (163) [91.0, 98.0]	91.3 (94) [84.1, 95.9]	96.2 (51) [87.0, 99.5]	100 (28) [87.7, 100]	94.2 (422) [91.6, 96.2]
Total BBPS score, mean (SD)	7.7 (1.4)	7.7 (1.4)	7.4 (1.7)	8.0 (1.1)	7.7 (1.5)	7.7 (1.4)
Score ≥ 2 in all 3 segments, % (n)	91.4 (85)	93.6 (160)	87.4 (90)	96.2 (51)	92.9 (26)	92.0 (412)

BBPS, Boston Bowel Preparation Scale; BMI, body mass index; CI, confidence interval; mITT, modified intent-to-treat.
^aResponders were those rated ‘3’ or ‘2’ on the BBPS by an endoscopist blinded to the treatment group, and the 95% CI of the responder rate was calculated using the Clopper-Pearson method.




Figure 2. Participants were asked ‘Was the bowel preparation tolerable?’ on the Mayo Clinic Bowel Prep Tolerability Questionnaire. At least 83% of participants in each BMI group indicated that SPMC oral solution was ‘easy’ or ‘acceptable’ to ingest. BMI, body mass index; SPMC, sodium picosulfate, magnesium oxide, and citric acid.




Figure 3. Participants who had experience with a prior colonoscopy (51–64% of each BMI cohort) were asked to rate the tolerability of the study bowel preparation compared with the previous preparation. The majority of participants rated SPMC oral solution bowel preparation as ‘better’ than a previous preparation. No consistent trend of responses was seen across BMI groups. Participants with no response are not shown on the graph and, therefore, numbers may not add to 100%. BMI, body mass index; SPMC, sodium picosulfate, magnesium oxide, and citric acid.

groups were largely consistent with the overall population rate of 13.2%, except for those who had BMI ≥ 40 kg/m² (3.6%).

Gastrointestinal AEs were the most frequently reported drug-related AE category in the entire study. Here, they occurred in $\leq 7.5\%$ of participants in any BMI group (Table 5). No consistent




Figure 4. The PDR varied by BMI, with no consistent trend. ADR was above the guideline-recommended target for all BMI groups. Any polyps found during the colonoscopy were removed, recorded as AEs, and sent for histological analysis. PDR and ADR were calculated as the percentage of any participants who had at least one polyp or adenoma, respectively. ADR, adenoma detection rate; AE, adverse event; BMI, body mass index; PDR, polyp detection rate.

patterns of drug-related TEAEs by BMI group were evident. By ascending BMI group, rates of nausea were 1.1%, 5.3%, 1.0%, 5.7%, and 0%, and rates of headache were 1.1%, 4.1%, 1.0%, 5.7%, and 0%. Similarly, rates of hypermagnesemia were 1.1%, 2.9%, 1.0%, 3.8%, and 0%.

Discussion

CRC screening in adults with high BMI is essential due to their increased risk of developing CRC and increased risk of CRC-related mortality.^{8–11} A high quality bowel preparation is needed to achieve optimal CRC screening colonoscopy.^{12,15}

Results of this subanalysis from a phase III trial of ready-to-drink SPMC oral solution displayed consistent and high efficacy of colon cleansing across all BMI groups, as measured by responder rates by AS and BBPS. There were very low rates of bowel cleansing rated ‘inadequate’ or ‘unprepared’ (score of ‘0’ by BBPS) in patients across all BMI groups. Previous studies reported an increased risk of inadequate bowel preparation by AS and BBPS for those with higher BMI^{16–18,20,26}; however, this trend was not observed for participants receiving SPMC oral solution.

Lower BMI has also been cited as a factor in incomplete colonoscopy.²⁷ A retrospective review of 2000 colonoscopies revealed that 49% of

Table 4. TEAEs, safety population.

% (n)	BMI subgroup					Overall cohort (n = 448)
	<25 (n = 93)	25–29.9 (n = 171)	30–34.9 (n = 103)	35–39.9 (n = 53)	≥40 (n = 28)	
Any TEAE ^a	80.6 (75)	85.4 (146)	84.5 (87)	92.5 (49)	75.0 (21)	84.4 (378)
Deaths	-	-	-	-	-	-
Serious TEAEs	3.2 (3)	1.2 (2)	1.0 (1)	5.7 (3)	-	2.0 (9)
TEAEs leading to study discontinuation	-	-	-	-	-	-
Severe TEAEs	5.4 (5)	0.6 (1)	-	7.5 (4)	3.6 (1)	2.5 (11)
Adverse drug reaction	12.9 (12)	14.0 (24)	11.7 (12)	18.9 (10)	3.6 (1)	13.2 (59)
Serious adverse drug reaction	-	-	-	-	-	-

AE, adverse event; BMI, body mass index; MedDRA, Medical Dictionary for Regulatory Activities; TEAE, treatment-emergent adverse event.
^aA TEAE was any AE that occurred or a pretreatment AE/medical condition that worsened in intensity after starting the study drug and within 30 days of last exposure to study drug. All endoscopic findings were reported as TEAEs; malignancies were reported as serious TEAEs. AEs were classified according to the MedDRA, version 20.1.

women with BMI < 22 kg/m² had a failed colonoscopy.²⁸ In addition to poor bowel preparation, another factor that may be related to incomplete colonoscopy in individuals with low BMI is a smaller area of adipose tissue, both the quantity and location of which may impact insertion of the colonoscope.^{29,30}

The consistent colon cleansing efficacy of SPMC oral solution across BMI groups may be attributed to preparation factors, such as the SPMC dual mechanism of action of peristaltic stimulation and osmotic agent; dosing factors, such as split dosing and hydration instructions; or due to study design with less risk of confounding factors in the setting of a randomized, controlled trial, and a patient population that has received intense instruction and may be more motivated to adhere to dosing instructions.

Results from a recent meta-analysis suggested that individuals who receive ratings of ‘excellent’, ‘good’, or ‘fair’ by AS have adequate enough colon cleansing to follow regular intervals for CRC screening.³¹ In this study, the corresponding population would be, by ascending BMI group, 94.6%, 98.2%, 97.1%, 98.1%, and 100%, which is the vast majority of each group and largely consistent across BMI. Interestingly, prior studies have reported that patients with total BBPS scores of 6–7 (‘good’ colon cleansing) had higher ADR and PDR than patients with BBPS

scores of 8 or 9 (‘excellent’ colon cleansing).^{32,33} The BBPS allows endoscopists to influence the presence of (aspirate/wash) any residual stool in the colon prior to grading, which may affect endoscopists’ perception of the relative cleanliness of each segment.³³

Higher BMI has previously been associated with increased rates of polyp and adenoma detection.³⁴ Various studies have reported that individuals with high BMI have 1.2–1.8 times higher risk of adenomas than individuals with normal BMI, though the studies did not use a uniform definition of high BMI.^{27,35–37} Other studies have found no association between BMI and the presence of polyps or adenomas.^{38,39} It is unknown if the higher rates of polyps measured in individuals with higher BMI were related to biological factors that increased the absolute number of polyps, or related to earlier missed polyps because of poorer quality of bowel cleansing.⁴⁰

Our subanalysis also showed no trends in polyp detection by BMI. This could be due to high quality and consistent colon cleansing achieved in all BMI groups here, which allowed for similar visualization of polyps in all groups.^{38,39} ADR calculations in this study included average-risk patients undergoing screening colonoscopy and high-risk patients undergoing surveillance colonoscopy; therefore, ADR values may be slightly overestimated, given that some high-risk patients

Table 5. Treatment-emergent, drug-related AEs of interest, safety population.

% (n)	BMI subgroup					Overall cohort (n = 448)
	<25 (n = 93)	25–29.9 (n = 171)	30–34.9 (n = 103)	35–39.9 (n = 53)	≥40 (n = 28)	
GI disorders	3.2 (3)	7.0 (12)	2.9 (3)	7.5 (4)	-	4.9 (22)
Nausea	1.1 (1)	5.3 (9)	1.0 (1)	5.7 (3)	-	3.1 (14)
Vomiting	2.2 (2)	1.2 (2)	-	3.8 (2)	-	1.3 (6)
Abdominal pain	1.1 (1)	0.6 (1)	1.0 (1)	-	-	0.7 (3)
Abdominal distention	1.1 (1)	0.6 (1)	-	-	-	0.4 (2)
Hypermagnesemia	1.1 (1)	2.9 (5)	1.0 (1)	3.8 (2)	-	2.0 (9)
Headache	1.1 (1)	4.1 (7)	1.0 (1)	5.7 (3)	-	2.7 (12)

AE, adverse event; BMI, body mass index; GI, gastrointestinal; MedDRA, Medical Dictionary for Regulatory Activities. AEs were classified according to the MedDRA, version 20.1.

were included in the total population for calculation. As this was a *post hoc* secondary analysis, it was not designed to assess ADR/PDR as the primary outcome or capture confounding factors associated with ADR/PDR.

Participants receiving SPMC oral solution also reported good tolerability of the bowel preparation, with high completion rates and high rates of ‘easy’ or ‘acceptable’ ratings in all BMI groups. The majority of participants with prior colonoscopy experience also rated SPMC oral solution as ‘better’ than a prior bowel preparation. There is limited literature on the tolerability of bowel preparation by BMI, although a single retrospective review in Poland showed increasing tolerability for bowel preparation with increasing BMI.²⁷

The safety data were consistent across all BMI groups, with no new safety signals, indicating that SPMC oral solution is a safe bowel preparation for patients of any BMI.

A limitation of this subanalysis is the small sample size for those with BMI ≥ 40 kg/m², thus conclusions drawn from those data should be made judiciously.

The strength of this subanalysis is the use of data from a randomized, controlled trial with pre-specified efficacy endpoints using validated instruments, and standardized safety reporting methods. Importantly, there are no existing large-scale studies of bowel preparation in individuals

with severe obesity. Data from 2015–2016 show that 7.7% of the adult population in the United States suffers from severe obesity, and this percentage continues to rise.⁴¹

Conclusion

Ready-to-drink SPMC oral solution had consistent and good quality colon cleansing among participants of all BMI groups, as measured by two independent and validated colon cleansing scales. In this heterogeneous study population, ADR was above the ASGE/ACG Taskforce guideline-recommended targets in all BMI groups. Participants in all BMI groups reported favorable tolerability for SPMC oral solution, with most participants preferring SPMC oral solution over a prior bowel preparation. No new safety signals were seen in participants of any BMI group. SPMC oral solution should be considered for patients of any BMI who are undergoing colonoscopy.

Acknowledgments

Medical writing and editorial support was provided by Agnella Izzo Matic (AIM Biomedical, LLC) and was funded by Ferring Pharmaceuticals Inc. The authors would like to thank the investigators, study staff, and participants who were involved in the trial.

Author Contributions

LH, GB, KLJ, MB, MA, and DND analyzed and interpreted the data, drafted and critically revised the article for important intellectual content, and approved the article for publication.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by Ferring Pharmaceuticals Inc., Parsippany, NJ.

Conflict of interest statement

LH has participated in the speaker's bureau for Ferring Pharmaceuticals Inc. GB was an investigator for clinical trials sponsored by Ferring Pharmaceuticals Inc. and has served as a consultant and has participated in the speaker's bureau for Ferring Pharmaceuticals Inc. KLJ is an investigator on clinical trials sponsored by Ferring Pharmaceuticals Inc. MB, MA, and DND are employees of Ferring Pharmaceuticals Inc.

Prior Publication

Portions of the data contained in this manuscript appeared in abstract/poster form at Digestive Disease Week 2019, May 18–21, 2019, Poster Mo1682. This manuscript has not been submitted and is not under consideration for publication anywhere else.

ORCID iD

Gerald Bertiger  <https://orcid.org/0000-0003-4731-2362>

References

1. ASGE Standards of Practice Committee, Saltzman JR, Cash BD, *et al.* Bowel preparation before colonoscopy. *Gastrointest Endosc* 2015; 81: 781–794.
2. Rex DK. Optimal bowel preparation - a practical guide for clinicians. *Nat Rev Gastroenterol Hepatol* 2014; 11: 419–425.
3. Corley DA, Jensen CD, Marks AR, *et al.* Adenoma detection rate and risk of colorectal cancer and death. *N Engl J Med* 2014; 370: 1298–1306.
4. Zauber AG. The impact of screening on colorectal cancer mortality and incidence: has it really made a difference? *Dig Dis Sci* 2015; 60: 681–691.
5. US Preventative Services Task Force, Bibbins-Domingo K, Grossman DC, *et al.* Screening for colorectal cancer: US preventative services task force recommendation. *JAMA* 2016; 315: 2564–2575.
6. Corley DA, Jensen CD, Marks AR, *et al.* Variation of adenoma prevalence by age, sex, race, and colon location in a large population: implications for screening and quality programs. *Clin Gastroenterol Hepatol* 2013; 11: 172–180.
7. Rex DK, Boland CR, Dominitz JA, *et al.* Colorectal cancer screening: recommendations for physicians and patients from the U.S. multi-society task force on colorectal cancer. *Am J Gastroenterol* 2017; 112: 1016–1030.
8. Larsson SC and Wolk A. Obesity and colon and rectal cancer risk: a meta-analysis of prospective studies. *Am J Clin Nutr* 2007; 86: 556–565.
9. Ma Y, Yang Y, Wang F, *et al.* Obesity and risk of colorectal cancer: a systematic review of prospective studies. *PLoS One* 2013; 8: e53916.
10. Doleman B, Mills KT, Lim S, *et al.* Body mass index and colorectal cancer prognosis: a systematic review and meta-analysis. *Tech Coloproctol* 2016; 20: 517–535.
11. Shaukat A, Dostal A, Menk J, *et al.* BMI is a risk factor for colorectal cancer mortality. *Dig Dis Sci* 2017; 62: 2511–2517.
12. Clark BT and Laine L. High-quality bowel preparation is required for detection of sessile serrated polyps. *Clin Gastroenterol Hepatol* 2016; 14: 1155–1162.
13. Chokshi RV, Hovis CE, Hollander T, *et al.* Prevalence of missed adenomas in patients with inadequate bowel preparation on screening colonoscopy. *Gastrointest Endosc* 2012; 75: 1197–1203.
14. Harewood GC, Sharma VK and de Garmo P. Impact of colonoscopy preparation quality on detection of suspected colonic neoplasia. *Gastrointest Endosc* 2003; 58: 76–79.
15. Lebwahl B, Kastrinos F, Glick M, *et al.* The impact of suboptimal preparation on adenoma miss rates and the factors associated with early repeat colonoscopy. *Gastrointest Endosc* 2011; 73: 1207–1214.
16. Borg BB, Gupta NK, Zuckerman GR, *et al.* Impact of obesity on bowel preparation for colonoscopy. *Clin Gastroenterol Hepatol* 2009; 7: 670–675.
17. Hassan C, Fuccio L, Bruno M, *et al.* A predictive model identifies patients most likely to have inadequate bowel preparation for colonoscopy. *Clin Gastroenterol Hepatol* 2012; 10: 501–506.
18. Fayad NF, Kahi CJ, Abd el-jawad KH, *et al.* Association between body-mass index and quality

- of split bowel preparation. *Clin Gastroenterol Hepatol* 2013; 11: 1478–1485.
19. Parra Pérez VF, Watanabe Yamamoto J, Nago A, *et al.* [Factors related to a suboptimal bowel preparation for colonoscopy]. *Acta Gastroenterol Latinoam* 2015; 45: 24–30.
 20. Anklesaria AB, Ivanina EA, Chudy-Onwugaje KO, *et al.* The effect of obesity on the quality of bowel preparation for colonoscopy: results from a large observational study. *J Clin Gastroenterol* 2019; 53: e214–e220.
 21. Sharara AI, Harb AH, Sarkis FS, *et al.* Body mass index and quality of bowel preparation: real life vs. clinical trials. *Arab J Gastroenterol* 2016; 17: 11–16.
 22. Sharara AI, Chalhoub JM, Beydoun M, *et al.* A customized mobile application in colonoscopy preparation: a randomized controlled trial. *Clin Transl Gastroenterol* 2017; 8: e211.
 23. Hookey L, Bertiger G, Lee Johnson K II, *et al.* Efficacy and safety of a ready-to-drink bowel preparation for colonoscopy: a randomized, controlled, non-inferiority trial. *Ther Adv Gastroenterol* 2019; 12: 1–13.
 24. Patel M, Staggs E, Thomas CS, *et al.* Development and validation of the Mayo Clinic Bowel Prep Tolerability Questionnaire. *Dig Liver Dis* 2014; 46: 808–812.
 25. Centers for Disease Control and Prevention. Defining adult overweight and obesity, 2017. <https://www.cdc.gov/obesity/adult/defining.html> (accessed 12 July, 2019).
 26. Cheng C-L, Liu N-J, Tang J-H, *et al.* Predictors of suboptimal bowel preparation using 3-l of polyethylene glycol for an outpatient colonoscopy: a prospective observational study. *Dig Dis Sci* 2016; 62: 345–351.
 27. Kobiela J, Wieszczy P, Reguła J, *et al.* Association of obesity with colonic findings in screening colonoscopy in a large population-based study. *United European Gastroenterol J.* 2018; 6: 1538–1546.
 28. Anderson JC, Gonzalez JD, Messina CR, *et al.* Factors that predict incomplete colonoscopy: thinner is not always better. *Am J Gastroenterol* 2000; 95: 2784–2787.
 29. Nagata N, Sakamoto K, Arai T, *et al.* Predictors for cecal insertion time: the impact of abdominal visceral fat measured by computed tomography. *Dis Colon Rectum* 2014; 57: 1213–1219.
 30. Kim EJ and Kim YJ. Can body mass index predict the difficulty of colonoscopy? *Clin Endosc* 2016; 49: 106–107.
 31. Clark BT, Rustagi T and Laine L. What level of bowel prep quality requires early repeat colonoscopy: systematic review and meta-analysis of the impact of preparation quality on adenoma detection rate. *Am J Gastroenterol* 2014; 109: 1714–1723; quiz 1724.
 32. Calderwood AH, Thompson KD, Schroy PC III, *et al.* Good is better than excellent: bowel preparation quality and adenoma detection rates. *Gastrointest Endosc* 2015; 81: 691–699.
 33. Adike A, Buras MR, Gurudu SR, *et al.* Is the level of cleanliness using segmental Boston Bowel Preparation Scale associated with a higher adenoma detection rate? *Ann Gastroenterol* 2018; 31: 217–223.
 34. Comstock SS, Hortos K, Kovan B, *et al.* Adipokines and obesity are associated with colorectal polyps in adult males: a cross-sectional study. *PLoS One.* 2014; 9: e85939.
 35. Ashktorab H, Paydar M, Yazdi S, *et al.* BMI and the risk of colorectal adenoma in African-Americans. *Obesity* 2014; 22: 1387–1391.
 36. Shapero TF, Chen GI, Devlin T, *et al.* Obesity increases prevalence of colonic adenomas at screening colonoscopy: a Canadian community-based study. *Can J Gastroenterol Hepatol* 2017; 2017: 8750967.
 37. Sato Y, Nozaki R, Yamada K, *et al.* Relation between obesity and adenomatous polyps of the large bowel. *Dig Endosc* 2009; 21: 154–157.
 38. Wallace K, Baron JA, Karagas MR, *et al.* The association of physical activity and body mass index with the risk of large bowel polyps. *Cancer Epidemiol Biomarkers Prev* 2005; 14: 2082–2086.
 39. Terlizzi J, Zheng A, Fuzesi S, *et al.* Polyp detection rates among body mass index categories at first screening colonoscopy. *Am Surg* 2017; 83: 54–57.
 40. Tandon K, Imam M, Ismail BES, *et al.* Body mass index and colon cancer screening: the road ahead. *World J Gastroenterol* 2015; 21: 1371–1376.
 41. Fryar CD, Carroll MD and Ogden CL. Prevalence of overweight, obesity, and severe obesity among adults aged 20 and over: United States, 1960–1962 through 2015–2016. *National Center for Health Statistics* 2018.