

Differential mechanisms underlying responses of soil bacterial and fungal communities to nitrogen and phosphorus inputs in a subtropical forest

Yong Li¹, Dashuan Tian², Jinsong Wang², Shuli Niu², Jing Tian², Denglong Ha³, Yuxi Qu³, Guangwei Jing³, Xiaoming Kang¹ and Bing Song⁴

¹ Beijing Key Laboratory of Wetland Services and Restoration, Institute of Wetland Research, Chinese Academy of Forestry, Beijing, China

² Key Laboratory of Ecosystem Network Observation and Modeling, Institute of Geographic Sciences and Natural Resources Research, Beijing, China

³ Jigongshan Natural Reserve, Xinyang, China

⁴ School of Resources and Environmental Engineering, Ludong University, Yantai, China

ABSTRACT

Atmospheric nitrogen (N) deposition and phosphorus (P) addition both can change soil bacterial and fungal community structure with a consequent impact on ecosystem functions. However, which factor plays an important role in regulating responses of bacterial and fungal community to N and P enrichments remains unclear. We conducted a manipulative experiment to simulate N and P inputs ($10 \text{ g N} \cdot \text{m}^{-2} \cdot \text{yr}^{-1} \text{ NH}_4\text{NO}_3$ or $10 \text{ g P} \cdot \text{m}^{-2} \cdot \text{yr}^{-1} \text{ NaH}_2\text{PO}_4$) and compared their effects on soil bacterial and fungal species richness and community composition. The results showed that the addition of N significantly increased NH_4^+ and Al^{3+} by 99.6% and 57.4%, respectively, and consequently led to a decline in soil pH from 4.18 to 3.75 after a 5-year treatment. P addition increased Al^{3+} and available P by 27.0% and 10-fold, respectively, but had no effect on soil pH. N addition significantly decreased bacterial species richness and Shannon index and resulted in a substantial shift of bacterial community composition, whereas P addition did not. Neither N nor P addition changed fungal species richness, Shannon index, and fungal community composition. A structural equation model showed that the shift in bacterial community composition was related to an increase in soil acid cations. The principal component scores of soil nutrients showed a significantly positive relationship with fungal community composition. Our results suggest that N and P additions affect soil bacterial and fungal communities in different ways in subtropical forest. These findings highlight how the diversity of microbial communities of subtropical forest soil will depend on future scenarios of anthropogenic N deposition and P enrichment, with a particular sensitivity of bacterial community to N addition.

Submitted 12 April 2019
Accepted 6 August 2019
Published 6 September 2019

Corresponding authors
Xiaoming Kang, xmkang@ucas.ac.cn
Bing Song, songbing@ldu.edu.cn

Academic editor
Xavier Le Roux

Additional Information and
Declarations can be found on
page 17

DOI 10.7717/peerj.7631

© Copyright
2019 Li et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Bioinformatics, Ecology, Microbiology, Climate Change Biology, Biogeochemistry
Keywords Nitrogen deposition, Phosphorus addition, Microbial diversity, Community composition, Amplicon sequencing, Subtropical forest, Soil bacteria and fungi

INTRODUCTION

Nitrogen (N) and phosphorus (P) inputs to the ecosystem have greatly increased due to anthropogenic activities, mainly originating from fossil fuel combustion, agricultural fertilization, and dust or ash production (Galloway *et al.*, 2008; Stevens, 2019; Wang *et al.*, 2014). Excessive N and P inputs could result in many adverse impacts, including soil acidification (Guo *et al.*, 2010; Mao *et al.*, 2017; Tian & Niu, 2015) and nutrient imbalance (Peñuelas *et al.*, 2013), although N and P have been considered limiting factors for plant growth (Harpole *et al.*, 2011). Moreover, elevated N availability may aggravate P limitation of ecosystem productivity (Li, Niu & Yu, 2016). A large body of research focusing on aboveground plant community responses have demonstrated multiple effects with nutrient additions, including biodiversity loss (Hooper *et al.*, 2012), species composition, and associated ecosystem services, such as terrestrial carbon dynamics (Isbell *et al.*, 2013; LeBauer & Treseder, 2008). However, the responses of belowground microorganisms to nutrient additions, including general taxonomic traits and related function shifts, remain unclear (Leff *et al.*, 2015; Ma *et al.*, 2016, 2019; Wei *et al.*, 2018). In particular, integrated field experimental investigations of both bacterial and fungal responses to N and P additions are needed to improve our understanding of how the soil microbial community structure shifts in response to nutrient addition and whether the bacterial and fungal responses are consistent.

Soil microbial communities are sensitive to N enrichment (Dai *et al.*, 2018; Ramirez, Craine & Fierer, 2012; Treseder, 2008). On average and based on a global meta-analysis, across all of the studies in agro-ecosystems, the abundances of *Proteobacteria* and *Actinobacteria* significantly increased by 2.2% and 1.1%, respectively, and the abundance of *Acidobacteria* decreased by 2.3% under N addition, which suggest differential sensitivity of bacterial phyla to N inputs (Dai *et al.*, 2018). N addition could lead to elevated N availability (Chen *et al.*, 2019), ammonium toxicity (Van Den Berg *et al.*, 2005), acid cation toxicity (Chen *et al.*, 2016; Tian & Niu, 2015), and loss of base mineral cations (Bowman *et al.*, 2008). Meanwhile, several potential driving factors were proposed to account for shifts in sensitive microbial communities in some ecosystems (Nie *et al.*, 2018). First, soil pH has been reported as a strong predictor of soil bacterial community composition at the continental scale (Lauber *et al.*, 2009; Rousk *et al.*, 2010). A significantly positive correlation between diversity of bacteria and soil pH might be attributable to narrow pH ranges for optimal growth of bacteria (Rousk *et al.*, 2010). However, fungal community composition is less affected by soil pH because fungi generally exhibit wider pH ranges for optimal growth (Rousk *et al.*, 2010). Second, N addition could directly affect soil bacterial community composition through modified ammonium N concentration (Nie *et al.*, 2018; Zeng *et al.*, 2016; Zhou *et al.*, 2017). Soil bacterial community composition is closely related to soil NH_4^+ -N content in tropical forest soil under N enrichment due to a narrow pH decrease in severely acidic soil (pH < 4.5) (Nie *et al.*, 2018). Finally, a positive relationship between plant and fungal beta diversity has been reported under N enrichment, soil properties including soil inorganic N, pH, and associated extractable cations being correlated with compositional changes in plant and fungal communities

(Chen *et al.*, 2018b). Although those mechanisms have previously been investigated, comprehensive studies on different regulatory pathways of N addition on bacterial and fungal community structure in forest ecosystems are limited.

Phosphorus availability plays an important role in affecting microbial growth; however, our understanding of soil microbial community responses to elevated P inputs remains limited (Leff *et al.*, 2015; Ma *et al.*, 2016, 2019). Moreover, the effects of P input on bacterial and fungal community composition can differ (Cassman *et al.*, 2016; Jorquera *et al.*, 2014; Liu *et al.*, 2012b, 2013). Previous studies have demonstrated that P enrichment induces shifts in microbial community composition by increasing P availability and altering soil and plant chemistry (DeForest & Scott, 2010; Lagos *et al.*, 2016; Leff *et al.*, 2015; Liu *et al.*, 2012a, 2013; Teng *et al.*, 2018). Other studies have reported that only N enrichment influences bacterial abundance and community composition, whereas P input did not because soil pH decreases only with N addition (Jorquera *et al.*, 2014; Wang *et al.*, 2018) or because bacterial community was limited by other resources (Ma *et al.*, 2019). The abovementioned pathways have largely been studied separately, and the relative contributions of these pathways to N- and P-induced changes in bacterial and fungal communities, however, have not been investigated using field experiments.

Subtropical forests in southern China undergo extensive N deposition with signs of N saturation, such as soil acidification (Zhu *et al.*, 2015). Previous studies have demonstrated that forest productivity and soil respiration are sensitive to N and P enrichment, but little is known about how P addition affects microbial communities (Li *et al.*, 2018a, 2018b; Yu *et al.*, 2017). To better understand the responses of bacterial and fungal communities to N and P addition and the underlying mechanisms, we conducted a N and P addition experiment in this region. We specifically addressed the following questions: (1) how do bacterial/fungal taxa and community structures respond to N and P enrichment? (2) How do soil biotic and abiotic factors regulate the responses of bacterial/fungal taxa and community structures to N and P enrichment? Finally, (3) what are the mechanisms underlying shifts in bacterial/fungal taxa and community structures to N and P enrichment?

MATERIALS AND METHODS

Site description

This N and P addition experiment was setup at the Jigongshan Nature Reserve, China (31°51'58"N, 114°5'12"E). The site is experiencing northern subtropical to warm temperate climates because it is located within a climate-transitional region. The mean annual surface air temperature and mean annual rainfall at the reserve is 15.2 °C and 1,118 mm, respectively. The soil type is yellow-brown and soil thickness is about 0.3–0.6 m (Yan *et al.*, 2014). The forest type is deciduous oak mixed forest. The predominant tree species in the canopy layer include *Quercus acutissima* and *Q. variabilis* and *Liquidambar formosana*, *Lindera glauca*, and *Viburnum dilatatum* dominates the understory arborous layer. The age of the stand is about 45–50 years and it is a secondary forest due to harvest in the late 1950s (Xu & Liang, 1965). The total N deposition in this region is about

30 kg N ha⁻¹ yr⁻¹ (Zhu *et al.*, 2015). The soil properties at the beginning of the experiment is listed in Table S1.

Experimental design

The detailed information of N and P addition experiment could be found in previous studies (Li *et al.*, 2018a, 2018b). Briefly, this N and P addition experiment was setup in July 2013 using a complete randomized block design. Each block had four treatments that were randomly assigned to 10 × 10 m plots and four replicate blocks were established. Control (CK, without N and P addition), N addition (N, 10 g m⁻² yr⁻¹ NH₄NO₃), P addition (P, 10 g m⁻² yr⁻¹ NaH₂PO₄), and NP addition (NP, 10 g · m⁻² · yr⁻¹ NH₄NO₃ + 10 g · m⁻² · yr⁻¹ NaH₂PO₄) were included. Backpack sprayer were used to spray 50 L of water dissolved additions for each plot onto the forest floor monthly from May to October each year. The control plot received 50 L of water (equivalent to 0.5 mm precipitation) each time.

Soil properties

Six soil cores (0–10 cm depth) were randomly collected from each plot and mixed to obtain one composite sample in October 2017. The samples were passed through a two mm sieve and three parts were divided. One part of the soil was used for the analysis of ammonium (NH₄⁺), nitrate (NO₃⁻), microbial biomass carbon and nitrogen (MBC and MBN), and dissolved organic carbon (DOC). The second part of fresh soil was collected in a 50-mL centrifuge tube, which was stored at -80 °C for soil DNA extraction. The remaining soil was air-dried for the determination of soil pH, total soil organic carbon (SOC), total nitrogen (TN), available phosphorus (AP), and extractable cations including Al³⁺, Ca²⁺, Mg²⁺, and Na⁺. The plant fine roots collected from six soil cores using the two mm-sieving were washed, dried, and then weighted. Soil ammonia and nitrate concentrations were determined by colorimetric analysis on a FIAstar 5000 Analyzer (FIAstar 5000 Analyzer; Foss Tecator, Hillerød, Denmark). Soil DOC was analyzed using a TOC analyzer (multi N/C 3100; Analytik Jena, Jena, Germany). Soil MBC and MBN were determined by a chloroform fumigation extraction method (Brookes *et al.*, 1985). Soil pH was analyzed in a soil water solution (1:2.5 w/v). SOC and TN were analyzed by a C/N analyzer (vario EL III, CHNOS Elemental Analyzer; Elementar, Langensfeld, Germany). Available P was determined following by molybdenum blue colorimetry (Jin *et al.*, 2019). A modified extraction procedure was used to measure extractable cations including Al³⁺, Ca²⁺, Mg²⁺, and Na⁺ (Rauret *et al.*, 2000).

Soil DNA extraction, PCR amplification, and sequencing

Soil DNA was extracted from each sample using a Fast DNA Stool Mini Kit (Tiangen Biotech Beijing Co., Ltd., Beijing, China) according to the manufacturer's instructions. The quality of the purified DNA was assessed based on the 260/280 and 260/230 nm absorbance ratios obtained using a NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies Inc., Wilmington, DE, USA).

Microbial community diversity and composition were assessed by amplification of the *16S rRNA* gene for the bacteria and the internal transcribed spacer (ITS) region of the fungi, as described in [Prober et al. \(2015\)](#). Marker genes in the isolated DNA were PCR-amplified and barcoded in triplicate reactions for both the *16S rRNA* gene (using the 515f/907r primer pair) and the ITS1 region (using the ITS1F/ITS2R primer pair). Sequencing was conducted on an Illumina Miseq platform at Majorbio Biopharm Technology Co., Ltd., Shanghai, China.

Sequencing data accession numbers

Raw sequences were trimmed of reads containing ambiguous bases and long homopolymers and merged using QIIME v 1.7.0 ([Caporaso et al., 2012](#)). All filtered sequences from 16S rRNA and ITS gene amplicons were clustered into Operational Taxonomic Units (OTU) at 97% similarity cutoff using UPARSE (version 7.1, <http://drive5.com/uparse>), followed by chimera filtering using the ribosomal database project (RDP) ([Tian et al., 2018](#)) and UCHIME ([Edgar et al., 2011](#)). The bacterial and fungal OTU sequences were classified using the RDP classifier against Greengenes and UNITE reference database, respectively. All of the raw sequencing data (.fastq files) were submitted to the sequence read archive at the National Center for Biotechnology Information under accession number [PRJNA531787](#).

Statistical analysis

The bacterial and fungal richness was determined by the number of OTUs, and alpha diversity was estimated using the Shannon index. Linear regressions were applied to assess the relationships between bacterial and fungal Shannon index and richness and environmental factors. Stepwise multiple regressions were used to identify the most influential environmental variables on bacterial and fungal Shannon index and richness due to collinearity among environmental factors. Differences in microbial species richness, Shannon index, soil properties and fine root biomass among different treatments were determined by one-way ANOVA with Duncan test. To determine the effect size of N, P and combined NP treatment on the relative abundances of the dominant bacterial and fungal taxa, the response ratio was calculated as $\ln(X_t/X_c)$, where X_t is the mean value of experimental treatment and X_c is the mean value of the control treatment. One-sample *t*-test was used to estimate whether each response ratio was significantly different from zero ([She et al., 2018](#)).

Principal component analysis (PCA) and nonmetric multidimensional scaling (NMDS) were used to determine changes in the bacterial and fungal communities. Analyses, including Anosim, Adonis, and MRPP, were further performed to assess the significant differences in community among different treatments. The partial Mantel test was used to evaluate the linkages between the microbial community structure and the environmental variables. Pairwise taxonomic distance between microbial communities (Bray–Curtis) and Euclidean distance of environmental variables were chosen for partial Mantel. PCA, NMDS, and partial Mantel were conducted using the *vegan* package with R v3.5.0 ([R Core Team, 2018](#)).

Structural equation modeling (SEM) was performed to analyze the hypothetical pathways of N and P addition effects on bacterial and fungal diversity and community composition. Before the SEM analysis, soil N availability including NH_4^+ and NO_3^- , soil acid cations including H^+ and Al^{3+} , soil nutrients including SOC, TN, DOC, MBC and MBN, and microbial community composition (OTUs) were subjected to PCA (Chen et al., 2013). Soil N availability, soil acidity, soil nutrients, and microbial community composition were presented by the first principal components (PC1) in the following SEM analysis. Maximum likelihood estimation method was applied in the process of SEM. The goodness of the models was determined by χ^2 tests, Akaike information criteria (AIC), and root square mean errors of approximation (RMSEA) (Li et al., 2018a).

RESULTS

Fine root biomass and soil properties

Nitrogen, P, and combined NP additions significantly reduced fine root biomass (Table S2). Soil pH decreased with N addition, whereas P and NP additions did not have any effect compared to the control. Soil NH_4^+ was increased by 99.6% under N addition, whereas it did not change after P and NP additions (Table S2). P and NP additions resulted in a significant increase in soil AP, but N addition did not (Table S2). Soil Al^{3+} increased by 57.4%, 27.0%, and 59.1%, respectively, under N, P and NP additions. Soil Ca^{2+} decreased by 33.6% and 25.4%, respectively, under N and NP additions, but did not significantly change under P addition. N, P, and NP additions did not significantly change SOC, total nitrogen (TN), ratio of SOC to TN (C/N ratio), DOC, MBC and MBN, NO_3^- , Mg^{2+} , and Na^+ .

Relative abundance of dominant microbial taxa

A total of 16,573 to 26,745 (average: 21,019) and 46,414–70,354 (average: 64,655) valid sequences were consequently obtained per sample for bacterial and fungi, respectively. These sequences were grouped into 1,666 and 2,487 OTUs at the 97% similarity level for bacterial and fungi, respectively. All of the samples were compared at an equivalent sequencing depth of 16,573 and 46,414 per sample for bacteria and fungi, respectively.

The predominant bacterial phyla across all of the samples were *Proteobacteria*, *Acidobacteria*, and *Actinobacteria* (mean relative abundance > 5%), which accounted for more than 82% of the bacterial sequences on average (Fig. 1A). In addition, *Planctomycetes*, *Chloroflexi*, *Firmicutes*, *Bacteroidetes*, and *Gemmatimonadetes* were also present at relatively low abundance. The dominant fungal phyla across all of the samples were *Ascomycota*, *Basidiomycota*, *Zygomycota*, and *Rozellomycota* (Fig. 1B).

At the phylum level, *Omnitrophica*, *Chlorobi*, and *Nitrospirae* presented significant decrease and *Saccharibacteria* increased in relative abundance under N treatment (Fig. 2A). P addition significantly increased the relative abundances of *Elusimicrobia*, but N addition had no effect compared to the control plots (Fig. 2A). Only NP treatment increased the relative abundance of *Firmicutes* and decreased that of *Chloroflexi* (Fig. 2A). No significant differences were observed in the fungal relative abundance of *Ascomycota*, *Rozellomycota*, and *Chytridiomycota* at the phylum level among different treatments (Fig. 2B). *Basidiomycota* presented significant decrease in their relative abundance under N




Figure 1 Relative abundance of the dominant bacterial (A) and fungal (B) groups at the phylum level under different treatments. [Full-size !\[\]\(fd7fe780e8fd8eece60268c87d0c3e04_img.jpg\) DOI: 10.7717/peerj.7631/fig-1](https://doi.org/10.7717/peerj.7631/fig-1)

and NP additions and only NP addition decreased the relative abundance of *Zygomycota* (Fig. 2B). At the fungal genus level, *Chaetomium* and *Purpureocillium* presented significant decrease in their relative abundance under N and P treatments, and N and NP treatments decreased the relative abundance of *Mycoarthritis* (Fig. 2B). The relative abundance of *Mortierella* significantly decreased only under NP addition (Fig. 2B).

Bacterial and fungal α diversity

Nitrogen addition significantly decreased bacterial α diversity, including phylotype richness (OTU numbers) and Shannon diversity index, but did not change fungal α diversity (Fig. 3). P addition had no effect on both bacterial and fungal α diversity. NP addition only decreased fungal phylotype richness (Fig. 3). N addition decreased the number of unique phylotypes compared to the control (Table S3).




Figure 2 Responses ratio analysis of changes in the relative abundance of dominant bacterial phyla (A) and fungal phyla/genera (B) in response to N, P and NP treatment compared to the control treatment at the 95% confidence interval. Red points indicate significant changes compared with the control treatment. [Full-size !\[\]\(5fd6ef84f97f42d7f8b34275f1b65312_img.jpg\) DOI: 10.7717/peerj.7631/fig-2](https://doi.org/10.7717/peerj.7631/fig-2)

The bacterial richness exhibited the largest correlation with soil pH among all of the environmental factors (Table 1, $r = 0.80$, $P < 0.01$). In addition to DOC ($r = -0.72$, $P < 0.01$), soil pH was the best predictor of bacterial diversity (Table 1, $r = 0.71$, $P < 0.01$). Bacterial richness and diversity also significantly correlated with fine root biomass, SOC, MBC, Al^{3+} , and Ca^{2+} (Table 1). Soil Al^{3+} ($r = -0.62$, $P < 0.05$) presented the largest correlation with fungal richness, followed by soil pH ($r = 0.57$, $P < 0.05$). No significant correlations were noted between fungal diversity and any of the environmental variables (Table 1).

Stepwise regression showed that bacterial richness significantly correlated with pH (64.3%) and DOC (7.1%) (Table S4). Soil pH explained the largest part of the variation in bacterial diversity (50.1%), followed by TN (14.9%), DOC (13.4%), and MBC (9.7%). Soil pH explained 33.0% of the variation in fungal richness, whereas pH (17.0%), SOC (19.1%), and MBC (41.4%) collectively contributed to 65.2% of the variation in fungal diversity (Table S4).

Bacterial and fungal community structure and relationships with environmental factors

Significant differences in bacterial community structure between N and NP additions and the control were observed in NMDS and PCA plots (Fig. 4; Fig. S1). However, only NP addition exhibited significant differences in fungal community structure compared to the control (Fig. 4). The results of anosim, adonis, and MRPP analyses further confirmed the significant differences among treatments (Table S5). This finding was confirmed by a cluster analysis based on a Bray–Curtis distance matrix of soil samples under different treatments (Fig. S2).




Figure 3 Bacterial and fungal phylotype richness (A) and (B) and Shannon diversity index (C) and (D) under different treatments. The error bars represent the SE of the mean ($n = 4$). Different letters above the bars indicate significant difference at $P < 0.05$. [Full-size !\[\]\(fcc3264021d438d9732560e78099f674_img.jpg\) DOI: 10.7717/peerj.7631/fig-3](https://doi.org/10.7717/peerj.7631/fig-3)

A partial Mantel test was performed to reveal the major environmental variables shaping microbial community structure. pH ($r_M = 0.57$, $P = 0.001$), Al^{3+} ($r_M = 0.40$, $P = 0.002$) and fine root biomass ($r_M = 0.36$, $P = 0.005$) were the most important factors that independently contributed to variations in soil bacterial community structure (Table 2). TN, C:N ratio, DOC, and MBC showed weak but significant correlations ($r_M = 0.21$, $r_M = 0.22$, $r_M = 0.27$, $r_M = 0.24$, respectively, $P < 0.05$) (Table 2). pH ($r_M = 0.57$, $P = 0.001$) and MBC ($r_M = 0.40$, $P = 0.004$) were the most important factors that independently contributed to variations in soil fungal community structure (Table 2). C:N ratio, DOC, Al^{3+} , and Mg^{2+} showed weak but significant correlations ($r_M = 0.25$, $r_M = 0.26$, $r_M = 0.21$, $r_M = 0.27$, respectively, $P < 0.05$) (Table 2).

Pathways determining the responses of bacterial and fungal diversity and community composition

The SEM explained 12% of the variations in soil N availability, 83% of the variation in soil acid cations (H^+ and Al^{3+}), 5% of the variation in soil nutrients and 80% of the variation in soil available P (Fig. 5). The total variation in bacterial Shannon index was mainly explained (64%) by soil acid cations and soil available P (Fig. 5A), but the total

Table 1 Pearson correlations between bacterial and fungal richness and diversity and plant and soil characteristics.

<i>r</i>	Bacteria		Fungi	
	Richness	Diversity	Richness	Diversity
Fine root biomass	0.59*	0.56*	0.51*	0.32
pH	0.80**	0.71**	0.57*	0.41
SOC	-0.52*	-0.40	-0.29	-0.47
TN	-0.47	-0.35	-0.16	-0.33
C:N ratio	-0.40	-0.35	-0.42	-0.47
DOC	-0.75**	-0.72**	-0.32	-0.20
NH ₄ ⁺	-0.48	-0.34	-0.23	-0.17
NO ₃ ⁻	-0.40	-0.31	-0.15	-0.02
AP	0.07	0.24	-0.34	-0.42
MBC	-0.64**	-0.49	-0.54*	-0.48
MBN	-0.41	-0.27	-0.40	-0.42
Al ³⁺	-0.54*	-0.58*	-0.62*	-0.33
Ca ²⁺	0.58*	0.54	0.40	0.33
Mg ²⁺	0.24	0.09	0.22	0.13
Na ⁺	-0.06	-0.07	-0.33	-0.17

Notes:* 0.01 < *P* ≤ 0.05.** 0.001 < *P* ≤ 0.01.

SOC, soil organic carbon; TN, total nitrogen content; DOC, dissolved organic carbon; AP, available phosphorus; MBC, microbial biomass carbon; MBN, microbial biomass nitrogen.

variation in fungal Shannon index was mainly explained (39%) by soil available P (marginally significant, *P* = 0.08, Fig. 5B). The total variation in bacterial community composition was mainly explained (79%) by soil acid cations (Fig. 5A), but the total variation in fungal community composition was mainly explained (45%) by soil nutrients (Fig. 5B). Linear regression demonstrated that principal component (PC1) scores of fungal community composition was significantly correlated with PC1 scores of soil nutrients ($R^2 = 0.41$, *P* = 0.008, Fig. S3).

Standardized total effects from SEM demonstrated that N enrichment had stronger effects on bacterial diversity and community composition compared to P addition (Fig. 6). Specifically, acid cations showed the most powerful negative effect on bacterial diversity (Fig. 6A) and had a stronger positive effect on bacterial community composition than soil N availability (Fig. 6B). Both N and P enrichment had negative effects on fungal diversity, and soil nutrients exerted the most powerful positive effect on fungal diversity (Fig. 6C). N enrichment had stronger positive effects on fungal community composition, and soil nutrients exerted strongest negative effect on fungal community composition (Fig. 6D).

DISCUSSION

N but not P addition significantly influences bacterial communities

Our results showed that N addition significantly decreased bacterial OTU richness and Shannon index and altered the bacterial community composition, with *Saccharibacteria*




Figure 4 Non-metric multidimensional scaling (NMDS) ordination of the soil bacterial (A) and fungal (B) community structure under different treatments. The Bray–Curtis distance matrix based on the abundance of OTUs was used to determine the compositional variation.

Full-size  DOI: [10.7717/peerj.7631/fig-4](https://doi.org/10.7717/peerj.7631/fig-4)

increasing, but *Chloroflex* decreasing in terms of their relative abundances (Figs. 2–4; Table S5). These results indicate that N addition tends to favor the copiotrophic phylum (*Saccharibacteria*) and counterselects the oligotrophic phylum (*Chloroflex*) in bacterial communities, which is in line with the copiotrophic hypothesis in previous reports (Fierer, Bradford & Jackson, 2007; Fierer et al., 2012; Leff et al., 2015). However, N addition did not change the relative abundance of dominant phylum of *Proteobacteria*, *Acidobacteria*,

Table 2 Partial mantel test of soil bacterial and fungal community structure with environmental characteristics.

	Bacteria		Fungi	
	<i>r</i>	<i>P</i>	<i>r</i>	<i>P</i>
Fine root biomass	0.36	0.005	0.10	0.18
pH	0.57	0.001	0.40	0.001
SOC	0.16	0.10	0.19	0.07
TN	0.21	0.04	0.12	0.14
C:N ratio	0.22	0.04	0.25	0.02
DOC	0.27	0.02	0.26	0.01
NH ₄ ⁺	0.08	0.21	-0.009	0.49
NO ₃ ⁻	0.07	0.26	0.02	0.45
AP	-0.14	0.95	-0.13	0.92
MBC	0.24	0.03	0.36	0.004
MBN	0.10	0.15	0.17	0.06
Al ³⁺	0.40	0.002	0.21	0.03
Ca ²⁺	0.12	0.18	-0.06	0.66
Mg ²⁺	0.06	0.28	0.27	0.02
Na ⁺	-0.15	0.91	-0.03	0.62

Notes:

The correlation and significance were determined between bacterial and fungal community structure (Bray–Curtis distance) and environmental variables (Euclidean distance) based on 999 permutations. The bold numbers indicate significant correlations.

SOC, soil organic carbon; TN, total nitrogen content; DOC, dissolved organic carbon; AP, available phosphorus; MBC, microbial biomass carbon; MBN, microbial biomass nitrogen.

and *Actinobacteria* (Fig. 2). The results are in contrast with the findings of previous studies, which showed that added N stimulates and decreases the relative abundances of *Proteobacteria* (copiotrophic) and *Acidobacteria* (oligotrophic), respectively (Chen et al., 2019; Fierer et al., 2012; Leff et al., 2015). This discrepancy could be attributable to neutral effects of N enrichment on soil available C and N (including DOC and NO₃⁻, Table S2) (Turlapati et al., 2013). It has been well documented that *Proteobacteria* contains N-fixing and N transforming genera (e.g., *Bradyrhizobium*, *Burkholderia*, *Magnetospirillum*, and *Mesorhizobium*) and thus are closely related to N cycling. However, in our study, added N only increased NH₄⁺ and did not affect NO₃⁻, indicating the N transforming-related bacterial phylum (*Proteobacteria*) may not be affected by the addition of N. Additionally, *Acidobacteria* has been reported to be negatively correlated with soil C availability (Fierer, Bradford & Jackson, 2007). The neutral effect of elevated N on DOC in our study may have contributed to the unaltered relative abundance of *Acidobacteria*, which is favored by low C availability (Fierer, Bradford & Jackson, 2007; Turlapati et al., 2013). Finally, the absence of a decrease in the abundance of *Actinobacteria* may be attributed to its relatively high tolerance to environmental stress such as low pH (Dai et al., 2018).

In contrast to most previous findings (Ling et al., 2017; Tan et al., 2013), neither the bacterial diversity nor the community composition was substantially influenced by




Figure 5 SEM analysis of the effects of N and P enrichment on soil bacterial (A) and fungal (B) community structure. Results of model fitting: (A) $\chi^2 = 19.38$, $df = 13$, $P = 0.11$, RMSEA = 0.18, AIC = 65.38; (B) $\chi^2 = 18.39$, $df = 13$, $P = 0.14$, RMSEA = 0.17, AIC = 64.39. Black and blue arrows represent significant positive and negative pathways, respectively, and dashed arrows indicate non-significant pathways. Numbers at the arrows are standardized path coefficients and arrow width is proportional to the strength of the relationship. R^2 values on top of response variables indicate the proportion of variation explained by relationships with other variables. Prior to SEM analysis, soil N availability (NH_4^+ and NO_3^-), soil acid cations (H^+ and Al^{3+}), soil nutrients (SOC, TN, DOC, MBC and MBN), bacterial and fungal community composition (OTUs) were subject to PCA procedure to reduce the number of variables.

Full-size DOI: 10.7717/peerj.7631/fig-5




Figure 6 SEM-derived standardized total effects of N and P additions, and variables including N availability, acid cations, soil nutrients, and soil available P on bacterial diversity and community composition (A-B) and fungal diversity (C-D). [Full-size !\[\]\(1679558f37f6db0dd8360a2a7e913e90_img.jpg\) DOI: 10.7717/peerj.7631/fig-6](https://doi.org/10.7717/peerj.7631/fig-6)

P addition in our study (Figs. 3 and 4; Table S5), but this finding is consistent with a previous study reporting that most bacterial taxa were not limited by P in Tibetan meadow (Ma *et al.*, 2019). Among all of the bacterial phyla identified, only *Elusimicrobia* were negatively correlated with soil available P (Table S6). There are several potential explanations for the insensitive response to added P. First, it is well established that nutrient addition-induced pH decline is an important mechanism in shifting the bacterial community composition (Chen *et al.*, 2015; Ling *et al.*, 2017; Zeng *et al.*, 2016). However, P addition did not change the soil pH in our study (Table S2). Additionally, P input could increase the P availability, and thus affect the bacterial richness and community structure, especially in P-limited ecosystems (Li *et al.*, 2015; Liu *et al.*, 2013). Our results showed that increased soil P availability did not change microbial biomass, suggesting that microbial growth is not limited to P in this subtropical forest. Finally, Liu *et al.* (2013) pointed out that a decrease in labile SOC is associated with alterations in soil microbial community structure under P addition. In our study, however, neither SOC or DOC was not significantly influenced by P addition (Table S2). Overall, our results suggest that only increased P availability may not be able to change the bacterial community composition in subtropical forest.

Neither N nor P addition significantly influenced fungal community

Our results demonstrated that fungal OTU richness is positively correlated to fine root biomass and soil pH and negatively related to MBC and soil Al^{3+} . However, fungal Shannon index was not significantly associated with soil parameters (Table 1). Fungal community structure was closely related to soil pH and MBC, followed by soil C:N ratio, DOC, Al^{3+} , and Mg^{2+} (Table 2). In contrast to the bacterial community, both N and P additions did not significantly change fungal richness and diversity (Fig. 3), which suggests that the tolerances of fungi to environmental changes are generally stronger than those of bacteria (Fierer *et al.*, 2012; Rousk *et al.*, 2010). The responses of fungal diversity to N enrichment is not always consistent due to ecosystem type and nutrient dose (Chen *et al.*, 2019; Mueller, Balasch & Kuske, 2014). Some studies have demonstrated that experimental N deposition results in increased fungal richness and diversity in a low fertility loblolly pine forest (Mueller, Balasch & Kuske, 2014; Weber, Vilgalys & Kuske, 2013), but other studies showed the opposite response partly due to variations in nutrient availability (Chen *et al.*, 2018a, 2019). Previous studies have demonstrated that N enrichment increases the relative abundance of copiotrophic phylum *Ascomycota* and decreased that of oligotrophic phylum *Basidiomycota* according to the copiotroph-oligotroph concept (Chen *et al.*, 2018b; Fierer, Bradford & Jackson, 2007; She *et al.*, 2018; Yao *et al.*, 2017). Although no significant effect of N enrichment on fungal community composition was observed (Fig. 4; Table S5), regression analysis also showed that the relative abundance of *Basidiomycota* was negatively correlated with soil N availability (NH_4^+ and NO_3^- , Table S6), indicating that N enrichment likely directly affects *Basidiomycota* via increasing N availability. Overall, the limited responses of fungal richness, diversity, and community composition under N enrichment in our study may be due to increased N loss from soil through denitrification and leaching (Niu *et al.*, 2016).

Compared to many studies analyzing the response of soil microorganisms to N addition, relatively little research has been done on the effects of P enrichment on soil fungi in grasslands and forests (Li *et al.*, 2015; Liu *et al.*, 2013), and much fewer studies using high-resolution technology, such as 16S *rRNA* sequencing (Cassman *et al.*, 2016; He *et al.*, 2016). The understanding of fungal community responses to elevated P inputs remains limited (Leff *et al.*, 2015). The effect of P addition on soil fungal community composition depends on ecosystem type, soil properties, and nutrient type and dose (Bao *et al.*, 2013; Beauregard *et al.*, 2010; Li *et al.*, 2015; Liu *et al.*, 2013). Previous studies have demonstrated that experimental P fertilization reduces the species richness of arbuscular mycorrhizal fungi, which could increase soil nutrient capture of their hosts in return for plant C resource (Camenzind *et al.*, 2014; Cheng *et al.*, 2013; Liu *et al.*, 2013). However, elevated P input could alleviate P deficiency of soil microbes and increase the fungal biomass, suggesting P availability is one of the limiting factors for fungal growth in an old tropical forest (Liu *et al.*, 2012a). Another explanation for the not significant change of fungal community composition in response to P enrichment might be the relatively short-term experimental duration of P addition (Cheng *et al.*, 2013), and these detailed parameters on fungal community composition warrant further investigation.

Mechanisms underlying responses of soil bacterial and fungal communities to N and P inputs

Several mechanisms behind the observed shift of bacterial and fungal community composition under N and P additions have been proposed, including increased N and P availability (Liu *et al.*, 2012a, 2013; Nie *et al.*, 2018; Zhou *et al.*, 2017), a decline in soil pH (Chen *et al.*, 2015; Zeng *et al.*, 2016), and increased soil acid cations availability (Chen *et al.*, 2015). Contrary to the N and P availability theory (Liu *et al.*, 2012a; Nie *et al.*, 2018), our results showed that N and P availability (including NH_4^+ , NO_3^- , and AP) had no significant relationship with bacterial and fungal diversity and community structure (Tables 1 and 2). Meanwhile, both partial Mantel test and SEM results have demonstrated that N and P availability had no significant relationship with bacterial and fungal community composition (Table 2; Fig. 5), suggesting that N and P availability could not explain the shifts in bacterial and fungal community composition. The remaining possible reasons for the decline of bacterial diversity and shift of bacterial community composition include increased concentration of H^+ and Al^{3+} (Table S2). Furthermore, SEM results showed that soil acid cations (H^+ and Al^{3+}) induced a significant decrease in bacterial diversity and substantial change of bacterial community composition (Figs. 5 and 6). However, SEM results have demonstrated that soil nutrients are significantly correlated to fungal community composition (Fig. 5). Taken together, in this study, increase of soil acid cations and soil nutrients significantly contributed to the shift in bacterial and fungal community composition, respectively.

CONCLUSIONS

After a 5-year N and P addition in a subtropical forest, our results showed that N addition significantly decreased bacterial species richness and diversity, and resulted in a substantial

shift of bacterial community composition, whereas P addition did not. Neither N nor P addition changed fungal species richness, diversity, and fungal community composition. A structural equation model showed that the shift in bacterial community composition is attributable to an increase in soil acid cations. The principal component scores of soil nutrients showed a significantly positive relationship with fungal community composition. Our results show how the diversity of microbial communities of subtropical forest soil will depend on future scenarios of anthropogenic N deposition and P enrichment, with a particular sensitivity of bacterial community to N addition.

ACKNOWLEDGEMENTS

The authors would like to thank the staff from Jigongshan Natural Reserve for experimental site setup and data collection. Two reviewers and academic editor Xavier Le Roux are appreciated for their constructive comments and valuable input.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This study was financially supported by National Natural Science Foundation of China (Grant Nos. 41403073 and 31600356) and the Thousand Young Talents program to Shuli Niu. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
National Natural Science Foundation of China: 41403073 and 31600356.
Thousand Young Talents program.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Yong Li conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Dashuan Tian contributed reagents/materials/analysis tools.
- Jinsong Wang contributed reagents/materials/analysis tools.
- Shuli Niu conceived and designed the experiments, contributed reagents/materials/analysis tools.
- Jing Tian contributed reagents/materials/analysis tools, prepared figures and/or tables.
- Denglong Ha performed the experiments.
- Yuxi Qu performed the experiments.
- Guangwei Jing performed the experiments.
- Xiaoming Kang authored or reviewed drafts of the paper.
- Bing Song analyzed the data, authored or reviewed drafts of the paper.

Data Availability

The following information was supplied regarding data availability:

The raw sequencing data are available at the Sequence Read Archive (SRA):
[PRJNA531787](https://www.ncbi.nlm.nih.gov/sra/PRJNA531787).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.7631#supplemental-information>.

REFERENCES

- Bao Z, Matsushita Y, Morimoto S, Hoshino YT, Suzuki C, Nagaoka K, Takenaka M, Murakami H, Kuroyanagi Y, Urashima Y, Sekiguchi H, Kushida A, Toyota K, Saito M, Tsushima S. 2013.** Decrease in fungal biodiversity along an available phosphorous gradient in arable Andosol soils in Japan. *Canadian Journal of Microbiology* **59(6)**:368–373 DOI [10.1139/cjm-2012-0612](https://doi.org/10.1139/cjm-2012-0612).
- Beauregard MS, Hamel C, Atul N, St-Arnaud M. 2010.** Long-term phosphorus fertilization impacts soil fungal and bacterial diversity but not AM fungal community in Alfalfa. *Microbial Ecology* **59(2)**:379–389 DOI [10.1007/s00248-009-9583-z](https://doi.org/10.1007/s00248-009-9583-z).
- Bowman WD, Cleveland CC, Halada L, Hresko J, Baron JS. 2008.** Negative impact of nitrogen deposition on soil buffering capacity. *Nature Geoscience* **1(11)**:767–770 DOI [10.1038/ngeo339](https://doi.org/10.1038/ngeo339).
- Brookes PC, Landman A, Pruden G, Jenkinson DS. 1985.** Chloroform fumigation and the release of soil nitrogen: a rapid direct extraction method to measure microbial biomass nitrogen in soil. *Soil Biology and Biochemistry* **17(6)**:837–842 DOI [10.1016/0038-0717\(85\)90144-0](https://doi.org/10.1016/0038-0717(85)90144-0).
- Camenzind T, Hempel S, Homeier J, Horn S, Velescu A, Wilcke W, Rillig MC. 2014.** Nitrogen and phosphorus additions impact arbuscular mycorrhizal abundance and molecular diversity in a tropical montane forest. *Global Change Biology* **20(12)**:3646–3659 DOI [10.1111/gcb.12618](https://doi.org/10.1111/gcb.12618).
- Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM, Betley J, Fraser L, Bauer M, Gormley N, Gilbert JA, Smith G, Knight R. 2012.** Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *ISME Journal* **6(8)**:1621–1624 DOI [10.1038/ismej.2012.8](https://doi.org/10.1038/ismej.2012.8).
- Cassman NA, Leite MFA, Pan Y, De Hollander M, Van Veen JA, Kuramae EE. 2016.** Plant and soil fungal but not soil bacterial communities are linked in long-term fertilized grassland. *Scientific Reports* **6(1)**:23680 DOI [10.1038/srep23680](https://doi.org/10.1038/srep23680).
- Chen D, Lan Z, Bai X, Grace JB, Bai Y. 2013.** Evidence that acidification-induced declines in plant diversity and productivity are mediated by changes in below-ground communities and soil properties in a semi-arid steppe. *Journal of Ecology* **101(5)**:1322–1334 DOI [10.1111/1365-2745.12119](https://doi.org/10.1111/1365-2745.12119).
- Chen D, Lan Z, Hu S, Bai Y. 2015.** Effects of nitrogen enrichment on belowground communities in grassland: relative role of soil nitrogen availability vs. soil acidification. *Soil Biology and Biochemistry* **89**:99–108 DOI [10.1016/j.soilbio.2015.06.028](https://doi.org/10.1016/j.soilbio.2015.06.028).
- Chen D, Li J, Lan Z, Hu S, Bai Y. 2016.** Soil acidification exerts a greater control on soil respiration than soil nitrogen availability in grasslands subjected to long-term nitrogen enrichment. *Functional Ecology* **30(4)**:658–669 DOI [10.1111/1365-2435.12525](https://doi.org/10.1111/1365-2435.12525).
- Chen D, Xing W, Lan Z, Saleem M, Wu Y, Hu S, Bai Y. 2019.** Direct and indirect effects of nitrogen enrichment on soil organisms and carbon and nitrogen mineralization in a semi-arid grassland. *Functional Ecology* **33(1)**:175–187 DOI [10.1111/1365-2435.13226](https://doi.org/10.1111/1365-2435.13226).

- Chen W, Xu R, Chen J, Yuan X, Zhou L, Tan T, Fan J, Zhang Y, Hu T. 2018a. Consistent responses of surface- and subsurface soil fungal diversity to N enrichment are mediated differently by acidification and plant community in a semi-arid grassland. *Soil Biology and Biochemistry* 127:110–119 DOI 10.1016/j.soilbio.2018.09.020.
- Chen W, Xu R, Wu Y, Chen J, Zhang Y, Hu T, Yuan X, Zhou L, Tan T, Fan J. 2018b. Plant diversity is coupled with beta not alpha diversity of soil fungal communities following N enrichment in a semi-arid grassland. *Soil Biology and Biochemistry* 116:388–398 DOI 10.1016/j.soilbio.2017.10.039.
- Cheng Y, Ishimoto K, Kuriyama Y, Osaki M, Ezawa T. 2013. Ninety-year-, but not single, application of phosphorus fertilizer has a major impact on arbuscular mycorrhizal fungal communities. *Plant and Soil* 365(1–2):397–407 DOI 10.1007/s11104-012-1398-x.
- Dai Z, Su W, Chen H, Barberán A, Zhao H, Yu M, Yu L, Brookes PC, Schadt CW, Chang SX, Xu J. 2018. Long-term nitrogen fertilization decreases bacterial diversity and favors the growth of *Actinobacteria* and *Proteobacteria* in agro-ecosystems across the globe. *Global Change Biology* 24(8):3452–3461 DOI 10.1111/gcb.14163.
- DeForest JL, Scott LG. 2010. Available organic soil phosphorus has an important influence on microbial community composition. *Soil Science Society of America Journal* 74(6):2059–2066 DOI 10.2136/sssaj2009.0426.
- Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. 2011. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics (Oxford, England)* 27(16):2194–2200 DOI 10.1093/bioinformatics/btr381.
- Fierer N, Bradford MA, Jackson RB. 2007. Toward an ecological classification of soil bacteria. *Ecology* 88(6):1354–1364 DOI 10.1890/05-1839.
- Fierer N, Lauber CL, Ramirez KS, Zaneveld J, Bradford MA, Knight R. 2012. Comparative metagenomic, phylogenetic and physiological analyses of soil microbial communities across nitrogen gradients. *ISME Journal* 6(5):1007–1017 DOI 10.1038/ismej.2011.159.
- Galloway JN, Townsend AR, Erisman JW, Bekunda M, Cai Z, Freney JR, Martinelli LA, Seitzinger SP, Sutton MA. 2008. Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science* 320(5878):889–892 DOI 10.1126/science.1136674.
- Guo JH, Liu XJ, Zhang Y, Shen JL, Han WX, Zhang WF, Christie P, Goulding KWT, Vitousek PM, Zhang FS. 2010. Significant acidification in major Chinese croplands. *Science* 327(5968):1008–1010 DOI 10.1126/science.1182570.
- Harpole WS, Ngai JT, Cleland EE, Seabloom EW, Borer ET, Bracken MES, Elser JJ, Gruner DS, Hillebrand H, Shurin JB, Smith JE. 2011. Nutrient co-limitation of primary producer communities. *Ecology Letters* 14(9):852–862 DOI 10.1111/j.1461-0248.2011.01651.x.
- He D, Xiang X, He J-S, Wang C, Cao G, Adams J, Chu H. 2016. Composition of the soil fungal community is more sensitive to phosphorus than nitrogen addition in the alpine meadow on the Qinghai-Tibetan Plateau. *Biology and Fertility of Soils* 52(8):1059–1072 DOI 10.1007/s00374-016-1142-4.
- Hooper DU, Adair EC, Cardinale BJ, Byrnes JEK, Hungate BA, Matulich KL, Gonzalez A, Duffy JE, Gamfeldt L, O'Connor MI. 2012. A global synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature* 486(7401):105–108 DOI 10.1038/nature11118.
- Isbell F, Reich PB, Tilman D, Hobbie SE, Polasky S, Binder S. 2013. Nutrient enrichment, biodiversity loss, and consequent declines in ecosystem productivity. *Proceedings of the National Academy of Sciences of the United States of America* 110(29):11911–11916 DOI 10.1073/pnas.1310880110.

- Jin Z, Chen C, Chen X, Hopkins I, Zhang X, Han Z, Jiang F, Billy G. 2019. The crucial factors of soil fertility and rapeseed yield—a five year field trial with biochar addition in upland red soil, China. *Science of the Total Environment* **649**:1467–1480 DOI [10.1016/j.scitotenv.2018.08.412](https://doi.org/10.1016/j.scitotenv.2018.08.412).
- Jorquera MA, Martínez OA, Marileo LG, Acuña JJ, Saggat S, Mora ML. 2014. Effect of nitrogen and phosphorus fertilization on the composition of rhizobacterial communities of two Chilean Andisol pastures. *World Journal of Microbiology and Biotechnology* **30**(1):99–107 DOI [10.1007/s11274-013-1427-9](https://doi.org/10.1007/s11274-013-1427-9).
- Lagos LM, Acuña JJ, Maruyama F, Ogram A, De la Luz Mora M, Jorquera MA. 2016. Effect of phosphorus addition on total and alkaline phosphomonoesterase-harboring bacterial populations in ryegrass rhizosphere microsites. *Biology and Fertility of Soils* **52**(7):1007–1019 DOI [10.1007/s00374-016-1137-1](https://doi.org/10.1007/s00374-016-1137-1).
- Lauber CL, Hamady M, Knight R, Fierer N. 2009. Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Applied and Environmental Microbiology* **75**(15):5111–5120 DOI [10.1128/AEM.00335-09](https://doi.org/10.1128/AEM.00335-09).
- LeBauer DS, Treseder KK. 2008. Nitrogen limitation of net primary productivity in terrestrial ecosystems in globally distributed. *Ecology* **89**(2):371–379 DOI [10.1890/06-2057.1](https://doi.org/10.1890/06-2057.1).
- Leff JW, Jones SE, Prober SM, Barberan A, Borer ET, Firn JL, Harpole WS, Hobbie SE, Hofmockel KS, Knops JMH, McCulley RL, La Pierre K, Risch AC, Seabloom EW, Schuetz M, Steenbock C, Stevens CJ, Fierer N. 2015. Consistent responses of soil microbial communities to elevated nutrient inputs in grasslands across the globe. *Proceedings of the National Academy of Sciences of the United States of America* **112**(35):10967–10972 DOI [10.1073/pnas.1508382112](https://doi.org/10.1073/pnas.1508382112).
- Li J, Li Z, Wang F, Zou B, Chen Y, Zhao J, Mo Q, Li Y, Li X, Xia H. 2015. Effects of nitrogen and phosphorus addition on soil microbial community in a secondary tropical forest of China. *Biology and Fertility of Soils* **51**(2):207–215 DOI [10.1007/s00374-014-0964-1](https://doi.org/10.1007/s00374-014-0964-1).
- Li Y, Niu S, Yu G. 2016. Aggravated phosphorus limitation on biomass production under increasing nitrogen loading: a meta-analysis. *Global Change Biology* **22**(2):934–943 DOI [10.1111/gcb.13125](https://doi.org/10.1111/gcb.13125).
- Li Y, Sun J, Tian D, Wang J, Ha D, Qu Y, Jing G, Niu S. 2018a. Soil acid cations induced reduction in soil respiration under nitrogen enrichment and soil acidification. *Science of the Total Environment* **615**:1535–1546 DOI [10.1016/j.scitotenv.2017.09.131](https://doi.org/10.1016/j.scitotenv.2017.09.131).
- Li Y, Tian D, Yang H, Niu S. 2018b. Size-dependent nutrient limitation of tree growth from subtropical to cold temperate forests. *Functional Ecology* **32**(1):95–105 DOI [10.1111/1365-2435.12975](https://doi.org/10.1111/1365-2435.12975).
- Ling N, Chen D, Guo H, Wei J, Bai Y, Shen Q, Hu S. 2017. Differential responses of soil bacterial communities to long-term N and P inputs in a semi-arid steppe. *Geoderma* **292**:25–33 DOI [10.1016/j.geoderma.2017.01.013](https://doi.org/10.1016/j.geoderma.2017.01.013).
- Liu L, Gundersen P, Zhang T, Mo J. 2012a. Effects of phosphorus addition on soil microbial biomass and community composition in three forest types in tropical China. *Soil Biology and Biochemistry* **44**(1):31–38 DOI [10.1016/j.soilbio.2011.08.017](https://doi.org/10.1016/j.soilbio.2011.08.017).
- Liu Y, Shi G, Mao L, Cheng G, Jiang S, Ma X, An L, Du G, Johnson NC, Feng H. 2012b. Direct and indirect influences of 8 yr of nitrogen and phosphorus fertilization on *Glomeromycota* in an alpine meadow ecosystem. *New Phytologist* **194**(2):523–535 DOI [10.1111/j.1469-8137.2012.04050.x](https://doi.org/10.1111/j.1469-8137.2012.04050.x).
- Liu L, Zhang T, Gilliam FS, Gundersen P, Zhang W, Chen H, Mo JM. 2013. Interactive effects of nitrogen and phosphorus on soil microbial communities in a tropical forest. *PLOS ONE* **8**(4):e61188 DOI [10.1371/journal.pone.0061188](https://doi.org/10.1371/journal.pone.0061188).

- Ma W, Jiang S, Assemien F, Qin M, Ma B, Xie Z, Liu Y, Feng H, Du G, Ma X, Le Roux X. 2016. Response of microbial functional groups involved in soil N cycle to N, P and NP fertilization in Tibetan alpine meadows. *Soil Biology and Biochemistry* **101**:195–206 DOI [10.1016/j.soilbio.2016.07.023](https://doi.org/10.1016/j.soilbio.2016.07.023).
- Ma B, Zhou X, Zhang Q, Qin M, Hu L, Yang K, Xie Z, Ma W, Chen B, Feng H, Liu Y, Du G, Ma X, Le Roux X. 2019. How do soil micro-organisms respond to N, P and NP additions? Application of the ecological framework of (co-)limitation by multiple resources. *Journal of Ecology* **107**:2329–2345 DOI [10.1111/1365-2745.13179](https://doi.org/10.1111/1365-2745.13179).
- Mao Q, Lu X, Zhou K, Chen H, Zhu X, Mori T, Mo J. 2017. Effects of long-term nitrogen and phosphorus additions on soil acidification in an N-rich tropical forest. *Geoderma* **285**:57–63 DOI [10.1016/j.geoderma.2016.09.017](https://doi.org/10.1016/j.geoderma.2016.09.017).
- Mueller RC, Balasch MM, Kuske CR. 2014. Contrasting soil fungal community responses to experimental nitrogen addition using the large subunit rRNA taxonomic marker and cellobiohydrolase I functional marker. *Molecular Ecology* **23**(17):4406–4417 DOI [10.1111/mec.12858](https://doi.org/10.1111/mec.12858).
- Nie Y, Wang M, Zhang W, Ni Z, Hashidoko Y, Shen W. 2018. Ammonium nitrogen content is a dominant predictor of bacterial community composition in an acidic forest soil with exogenous nitrogen enrichment. *Science of the Total Environment* **624**:407–415 DOI [10.1016/j.scitotenv.2017.12.142](https://doi.org/10.1016/j.scitotenv.2017.12.142).
- Niu S, Classen AT, Dukes JS, Kardol P, Liu L, Luo Y, Rustad L, Sun J, Tang J, Templer PH, Thomas RQ, Tian D, Vicca S, Wang Y-P, Xia J, Zaehle S. 2016. Global patterns and substrate-based mechanisms of the terrestrial nitrogen cycle. *Ecology Letters* **19**(6):697–709 DOI [10.1111/ele.12591](https://doi.org/10.1111/ele.12591).
- Peñuelas J, Poulter B, Sardans J, Ciais P, Van der Velde M, Bopp L, Boucher O, Godderis Y, Hinsinger P, Llusia J, Nardin E, Vicca S, Obersteiner M, Janssens IA. 2013. Human-induced nitrogen–phosphorus imbalances alter natural and managed ecosystems across the globe. *Nature Communications* **4**(1):2934 DOI [10.1038/ncomms3934](https://doi.org/10.1038/ncomms3934).
- Prober SM, Leff JW, Bates ST, Borer ET, Firn J, Harpole WS, Lind EM, Seabloom EW, Adler PB, Bakker JD, Cleland EE, DeCrappeo NM, DeLorenze E, Hagenah N, Hautier Y, Hofmockel KS, Kirkman KP, Knops JMH, La Pierre KJ, MacDougall AS, McCulley RL, Mitchell CE, Risch AC, Schuetz M, Stevens CJ, Williams RJ, Fierer N. 2015. Plant diversity predicts beta but not alpha diversity of soil microbes across grasslands worldwide. *Ecology Letters* **18**(1):85–95 DOI [10.1111/ele.12381](https://doi.org/10.1111/ele.12381).
- R Core Team. 2018. *R: a language and environment for statistical computing*. Vienna: The R Foundation for Statistical Computing. Available at <http://www.R-project.org/>.
- Ramirez KS, Craine JM, Fierer N. 2012. Consistent effects of nitrogen amendments on soil microbial communities and processes across biomes. *Global Change Biology* **18**(6):1918–1927 DOI [10.1111/j.1365-2486.2012.02639.x](https://doi.org/10.1111/j.1365-2486.2012.02639.x).
- Rauret G, Lopez-Sanchez JF, Sahuquillo A, Barahona E, Lachica M, Ure AM, Davidson CM, Gomez A, Luck D, Bacon J, Yli-Halla M, Muntau H, Quevauviller P. 2000. Application of a modified BCR sequential extraction (three-step) procedure for the determination of extractable trace metal contents in a sewage sludge amended soil reference material (CRM 483), complemented by a three-year stability study of acetic acid and EDTA extractable metal content. *Journal of Environmental Monitoring* **2**(3):228–233 DOI [10.1039/b001496f](https://doi.org/10.1039/b001496f).
- Rousk J, Bååth E, Brookes PC, Lauber CL, Lozupone C, Caporaso JG, Knight R, Fierer N. 2010. Soil bacterial and fungal communities across a pH gradient in an arable soil. *ISME Journal* **4**(10):1340–1351 DOI [10.1038/ismej.2010.58](https://doi.org/10.1038/ismej.2010.58).

- She W, Bai Y, Zhang Y, Qin S, Feng W, Sun Y, Zheng J, Wu B. 2018. Resource availability drives responses of soil microbial communities to short-term precipitation and nitrogen addition in a desert shrubland. *Frontiers in Microbiology* 9:186 DOI 10.3389/fmicb.2018.00186.
- Stevens CJ. 2019. Nitrogen in the environment. *Science* 363(6427):578–580 DOI 10.1126/science.aav8215.
- Tan H, Barret M, Mooij MJ, Rice O, Morrissey JP, Dobson A, Griffiths B, O’Gara F. 2013. Long-term phosphorus fertilisation increased the diversity of the total bacterial community and the phoD phosphorus mineraliser group in pasture soils. *Biology and Fertility of Soils* 49:661–672 DOI 10.1007/s00374-012-0755-5.
- Teng Z, Cui J, Wang J, Fu X, Xu X. 2018. Effect of exogenous nitrogen and phosphorus inputs on the microbe-soil interaction in the secondary *Castanopsis sclerophylla* forest in east China. *iForest—Biogeosciences and Forestry* 11(6):794–801 DOI 10.3832/ifor2673-011.
- Tian J, He N, Hale L, Niu S, Yu G, Liu Y, Blagodatskaya E, Kuzyakov Y, Gao Q, Zhou J. 2018. Soil organic matter availability and climate drive latitudinal patterns in bacterial diversity from tropical to cold temperate forests. *Functional Ecology* 32(1):61–70 DOI 10.1111/1365-2435.12952.
- Tian D, Niu S. 2015. A global analysis of soil acidification caused by nitrogen addition. *Environmental Research Letters* 10(2):24019 DOI 10.1088/1748-9326/10/2/024019.
- Treseder KK. 2008. Nitrogen additions and microbial biomass: a meta-analysis of ecosystem studies. *Ecology Letters* 11(10):1111–1120 DOI 10.1111/j.1461-0248.2008.01230.x.
- Turlapati SA, Minocha R, Bhiravarasa PS, Tisa LS, Thomas WK, Minocha SC. 2013. Chronic N-amended soils exhibit an altered bacterial community structure in Harvard Forest, MA, USA. *FEMS Microbiology Ecology* 83(2):478–493 DOI 10.1111/1574-6941.12009.
- Van Den Berg LJJ, Dorland E, Vergeer P, Hart MAC, Bobbink R, Roelofs JGM. 2005. Decline of acid-sensitive plant species in heathland can be attributed to ammonium toxicity in combination with low pH. *New Phytologist* 166(2):551–564 DOI 10.1111/j.1469-8137.2005.01338.x.
- Wang R, Balkanski Y, Boucher O, Ciais P, Peñuelas J, Tao S. 2014. Significant contribution of combustion-related emissions to the atmospheric phosphorus budget. *Nature Geoscience* 8(1):48–54 DOI 10.1038/ngeo2324.
- Wang Q, Wang C, Yu W, Turak A, Chen D, Huang Y, Ao J, Jiang Y, Huang Z. 2018. Effects of nitrogen and phosphorus inputs on soil bacterial abundance, diversity, and community composition in Chinese fir plantations. *Frontiers in Microbiology* 9:1543 DOI 10.3389/fmicb.2018.01543.
- Weber C, Vilgalys R, Kuske C. 2013. Changes in fungal community composition in response to elevated atmospheric CO₂ and nitrogen fertilization varies with soil Horizon. *Frontiers in Microbiology* 4:78 DOI 10.3389/fmicb.2013.00078.
- Wei H, Peng C, Yang B, Song H, Li Q, Jiang L, Wei G, Wang K, Wang H, Liu S, Liu X, Chen D, Li Y, Wang M. 2018. Contrasting soil bacterial community, diversity, and function in two forests in China. *Frontiers in Microbiology* 9:1693 DOI 10.3389/fmicb.2018.01693.
- Xu G, Liang C. 1965. Division and transformation of secondary forest in Jigongshan Forest Farm. *Practical Forestry Technology* (6):8–10 DOI 10.13456/j.cnki.lykt.1965.06.005 [in Chinese].
- Yan J, Zhang W, Wang K, Qin F, Wang W, Dai H, Li P. 2014. Responses of CO₂, N₂O and CH₄ fluxes between atmosphere and forest soil to changes in multiple environmental conditions. *Global Change Biology* 20(1):300–312 DOI 10.1111/gcb.12327.
- Yao F, Yang S, Wang Z, Wang X, Ye J, Wang X, DeBruyn JM, Feng X, Jiang Y, Li H. 2017. Microbial taxa distribution is associated with ecological trophic cascades along an elevation gradient. *Frontiers in Microbiology* 8:2071 DOI 10.3389/fmicb.2017.02071.

- Yu L, Wang Y, Zhang X, Dörsch P, Mulder J. 2017.** Phosphorus addition mitigates N₂O and CH₄ emissions in N-saturated subtropical forest, SW China. *Biogeosciences* **14**(12):3097–3109 DOI [10.5194/bg-14-3097-2017](https://doi.org/10.5194/bg-14-3097-2017).
- Zeng J, Liu X, Song L, Lin X, Zhang H, Shen C, Chu H. 2016.** Nitrogen fertilization directly affects soil bacterial diversity and indirectly affects bacterial community composition. *Soil Biology and Biochemistry* **92**:41–49 DOI [10.1016/j.soilbio.2015.09.018](https://doi.org/10.1016/j.soilbio.2015.09.018).
- Zhou Z, Wang C, Zheng M, Jiang L, Luo Y. 2017.** Patterns and mechanisms of responses by soil microbial communities to nitrogen addition. *Soil Biology and Biochemistry* **115**:433–441 DOI [10.1016/j.soilbio.2017.09.015](https://doi.org/10.1016/j.soilbio.2017.09.015).
- Zhu J, He N, Wang Q, Yuan G, Wen D, Yu G, Jia Y. 2015.** The composition, spatial patterns, and influencing factors of atmospheric wet nitrogen deposition in Chinese terrestrial ecosystems. *Science of the Total Environment* **511**:777–785 DOI [10.1016/j.scitotenv.2014.12.038](https://doi.org/10.1016/j.scitotenv.2014.12.038).