

highest metastatic rate, however G4 γ showed the worst survival. **CONCLUSION:** We identified molecular subgroups and subtypes of MBs based on gene expression and DNA methylation profile in children in our cohort series. The results may contribute to the establishment of nation-wide correlated optimal diagnosis and treatment strategies for MBs in infant and children.

MBRS-16. MYC REGULATED LONG NONCODING RNA LNC-HLX-2-7 IS A PUTATIVE MOLECULAR MARKER AND A THERAPEUTIC TARGET FOR GROUP 3 MEDULLOBLASTOMAS IN CHILDREN

Keisuke Katsushima¹, Piyush Joshi², Stacie Stapleton², Alexandra Garancher³, Rajeev Vibhakar⁴, Eric Raabe⁵, Charles Eberhart⁵, Robert Wechsler-Reya³, George Jallo², and **Ranjan Perera**¹; ¹Johns Hopkins University, St. Petersburg, FL, USA, ²JHACH, St. Petersburg, FL, USA, ³Sanford Burnham, La-Jolla, CA, USA, ⁴University of Colorado, Boulder, CO, USA, ⁵Johns Hopkins University, Baltimore, MD, USA

Medulloblastoma (MB), a central nervous system tumor that predominantly affects children, requires aggressive therapy. Recent advances in the noncoding RNA genome could contribute to the sub-classification of medulloblastoma. The focus of this study is to identify novel long noncoding RNAs (lncRNAs) as molecular markers and potential therapeutic targets within each subgroup of MBs, in particular within Group 3. We analyzed publicly available 175 RNA-seq datasets to identify a group of putative lncRNA signatures that may be able to differentiate medulloblastoma subgroups accurately. Among those, lncRNA *lnc-HLX-2-7* was highly upregulated in Group 3 MB cell lines, patient-derived xenografts, FFPE samples compared to other groups. CRISPR/Cas9 deletion of the *lnc-HLX-2-7* followed by the fluorescence-activated sorting and generating monoclonal Group 3 MB cells significantly reduced the cell growth and 3-D colony formation together with the induction of apoptosis. Intracranial injection to mouse cerebellum using *lnc-HLX-2-7* deleted cells resulted in reduced tumor growth compared to parental cells, and tumors were further characterized by single-cell sequencing. We identified that oncogene MYC regulates *lnc-HLX-2-7* and its expression can be controlled by the small molecule JQ1, a BET-bromodomain (BRD4) inhibitor that disrupts interactions with MYC. RNA-FISH analysis using FFPE, PDX, and tissue microarrays revealed that *lnc-HLX-2-7* expression is specific to Group 3 MB compared to other groups. We present supporting evidence that *lnc-HLX-2-7* is a novel molecular marker and a potential therapeutic target for Group 3 MBs in children.

MBRS-17. EXAMINING THE ROLE OF LHX9 IN GROUP 3 MEDULLOBLASTOMA

Sarah Injac^{1,2}, L. Frank Huang³, Stephen Mack^{1,2}, Frank Braun^{2,1}, Yuchen Du⁴, Mari Kogiso^{1,2}, Holly Lindsay^{1,2}, D. William Parsons^{1,2}, Hong Zhao⁵, Ching C. Lau⁶, Stephen T.C. Wong³, and Xiao-nan Li^{1,4}; ¹Baylor College of Medicine, Houston, TX, USA, ²Texas Children's Cancer and Hematology Centers, Houston, TX, USA, ³Experimental Hematology and Cancer Biology, Brain Tumor Center, Cincinnati Children's Hospital, Cincinnati, OH, USA, ⁴Department of Pediatrics, Northwestern University, Chicago, IL, USA, ⁵Department of Systems Medicine and Bioengineering, Houston Methodist Research Institute and Cancer Center, Weill Cornell School of Medicine, Houston, TX, USA, ⁶Connecticut Children's Medical Center, The University of Connecticut School of Medicine, Hartford, CT, USA

Medulloblastoma (MB) is the most common malignant brain tumor of childhood. Despite major advances in our understanding of the biology of MB, novel treatments remain urgently needed. Using a chemical-genomics driven drug repositioning strategy, we identified the cardiac glycoside family of compounds as potential treatments for Group 3 MB. We subsequently demonstrated that single-agent treatment with digoxin prolongs survival in a patient-derived xenograft model (PDOX) of Group 3 MB to a degree comparable to radiation therapy, a mainstay in the treatment of MB. Finally, we examined the mechanism of digoxin-mediated cell killing using RNA-seq. This work identified LHX9, a member of the LIM homeobox family of transcription factors, as the gene most significantly down-regulated following treatment (Huang and Injac et al, *Sci Trans Medicine*, 2018). Homologs of LHX9 play key roles in cerebellar development via spatially and temporally restricted expression and LHX9 has been proposed as a core transcription factor (TF) in the regulatory circuitry of Group 3 tumors. Loss of function of other core TFs has been shown to impact MB growth. The role of LHX9 in MB, however, has not been previously experimentally evaluated. We now report that knockdown of LHX9 in MB-derived cell lines results in marked growth inhibition raising the possibility that loss of LHX9 plays a major role in digoxin-mediated cell killing and that LHX9 represents a key dependency required for the growth of Group 3 MB. Clinical targeting of core TFs would represent a novel approach to targeting this devastating disease.

MBRS-18. TUMOR SUPPRESSOR P53 DEFINES THE THERAPEUTIC RESPONSES IN TREATMENT OF MEDULLOBLASTOMA

Avinash L. Mohan¹, Anubhav G. Amin¹, Michael E. Tobias¹, Mohan K. Das¹, Raphael SS de Medeiros², Nelci Zanon², Chirag D. Gandhi¹, Sidnei Epelman², and **Meena Jhanwar-Unival**¹; ¹New York Medical College/Westchester Medical Center, Valhalla, NY, USA, ²Hospital Santa Marcelina, São Paulo, Brazil

Medulloblastoma (MB) is the most common primary pediatric malignant brain tumor. Current molecular analysis classifies MB into 4 groups, classic (WNT), sonic hedgehog (Shh), group 3, and group 4. Furthermore, atypical p53 signaling is associated with disease progression and confers poor prognosis. This study investigated the correlation of mutational status of p53 and iSO17q with disease progression and metastatic potential. In addition, we used small molecule inhibitors of PI3K (Buparlisib; BKM120) and HDAC (LBH-589) on a p53-mutant MB cell line to find novel therapeutic targets. Efficacy of these drugs were assessed using functional assays (cell proliferation, migration, cell cycle and drug resistance). MB tumors (n=53) were evaluated for G1I-1, GAB-1, NPR, KV1, YAP expression and mutant p53 via immunohistochemistry and correlated to patient outcomes. Results demonstrated that: 1) high expression of GAB-1 and YAP were found in the Shh group, while KV1 expression was present in all subtypes; 2) mutant p53 expression was present in various subsets of MB with no apparent correlation with metastasis or disease progression; 3) patients displaying iSO17q (determined by fluorescence in situ hybridization (FISH) technique) exhibited metastatic disease; 4) LBH-589 and BKM120 caused both time and dose-dependent inhibition of MB cell proliferation and migration; 5) combined treatment of BKM120 and LBH-589 had a synergistic effect; 6) MB cells demonstrated drug-resistance to BKM120. In conclusion, these findings underscore use of Buparlisib and LBH-589 in treatment of MB. Further, the role of mutant p53 in disease progression remains elusive, whereas presence of iSO17q defines metastatic potential.

MBRS-19. SYNERGISM OF HDAC AND PARP INHIBITORS IN MYC-DRIVEN GROUP 3 MEDULLOBLASTOMA CELLS

Johanna Vollmer^{1,2}, Jonas Ecker^{1,2}, Thomas Hielscher³, Gintivė Valinciuė^{1,2}, Sina Oppermann^{1,2}, Heike Peterziel^{1,2}, Ina Oehme^{1,2}, Olaf Witt^{1,2}, and **Till Milde**¹; ¹Hopp Children's Cancer Center Heidelberg (KITZ), Heidelberg, Germany, ²Clinical Cooperation Unit Pediatric Oncology, German Cancer Research Center (DKFZ), Heidelberg, Germany, ³Division of Biostatistics, German Cancer Research Center (DKFZ), Heidelberg, Germany

Patients with MYC-driven Group 3 medulloblastoma (MB) show particularly poor outcome. It was previously shown that MYC-driven MBs are highly sensitive to class I histone deacetylase inhibition (HDACi). We studied the molecular effects of the class I HDACi entinostat in MYC-driven MB cells to identify potentially synergistic drug combinations, prioritizing drug clinical availability to enable clinical translation. Gene expression profiles of the MYC-amplified group 3 MB cell line HD-MB03 treated with entinostat were analyzed using bioinformatic approaches, identifying 29 altered biomechanisms. Overlay with a translational drug library of n=76 compounds resulted in 44 compounds targeting 9 biomechanisms. Filtering for publications supporting each drug's role in MYC-driven entities, or functional interaction with HDACs, without publication of this combination in MBs, resulted in 5 compounds (olaparib, idasanutlin, ribociclib, selinexor, vinblastine). Synergism testing identified olaparib as the drug with the strongest synergism. Validation of the combination olaparib and entinostat by p.H2AX and PI staining as well as trypan blue exclusion showed increased double strand breaks (DSBs), increased cell death, loss of viability and cell numbers. Selectivity of MYC-amplified MB cells was shown by comparison to MYC-non amplified cell lines, which showed higher IC50s, and reacted with cell cycle arrest as opposed to cell death to the combination treatment. The role of HDACi in DNA damage repair was confirmed by increased DSBs when entinostat was added to the combination of olaparib with doxorubicin. Our study identified olaparib as a potential combination partner with entinostat for the treatment of MYC-driven Group 3 MB.

MBRS-20. CSF-DERIVED CIRCULATING TUMOR DNA AS A BIOMARKER FOR DISEASE PROGRESSION AND TUMOR EVOLUTION IN MEDULLOBLASTOMA

Anthony Pak-Yin Liu¹, Rahul Kumar^{2,3}, Smith Kyle², Leena Paul², Murali Chintagumpala⁴, Eric Bouffter⁵, Michael J Fisher⁶, Tim Hassall⁷, Sri Gururangan⁸, Jordan Hansford⁹, Richard Cohn¹⁰, David W. Ellison¹¹, Richard J. Gilbertson¹², Amar Gajjar¹, Giles W. Robinson¹, and Paul A. Northcott²; ¹Division of Neuro-Oncology, Department of Oncology, St. Jude Children's Research Hospital, Memphis, TN, USA, ²Division of Brain Tumor Research, Department of Developmental Neurobiology, St. Jude Children's Research Hospital, Memphis, TN, USA, ³St. Jude Graduate School of Biomedical Sciences, Memphis, TN, USA, ⁴Department of Pediatrics, Texas Children's Cancer Center, Baylor College