Open access Editorial

General Psychiatry

Special issue for psychosomatic medicine

Wenhao Jiang, ¹ Zheng Lu, ² Yonggui Yuan ¹

To cite: Jiang W, Lu Z, Yuan Y. Special issue for psychosomatic medicine. *General Psychiatry* 2022;**35**:e100961. doi:10.1136/gpsych-2022-100961

Received 18 October 2022 Accepted 01 November 2022 Psychosomatic medicine is a branch of medicine that studies psychosomatic interactions. It focuses on the aetiology, pathology, diagnosis, treatment and prevention of psychosomatic diseases. Increasing effort is being made to improve the understanding of psychosomatic medicine in patients, their families, general hospitals and society. In addition, the development of psychosomatic medicine calls for wide collaboration with psychiatry, psychology, behavioural science, sociology, biology, neurochemistry and immunology. We initiated this special issue to encourage interdisciplinary studies in psychosomatic medicine.

This issue has generated a very informative collection of articles. We cover current psychosomatic development, cross-cultural issues, sexual studies, gastroenterology, cardiology, cancer, stress, anxiety, depression, schizophrenia and others. In the first article, Dr Lee, the president of the International College of Psychosomatic Medicine, comprehensively introduces the history of psychosomatic societies and their global presence. The author noted the rapid growth of psychosomatic communities in Asian countries, especially China. In addition, Dr Lee called for friendship and collegiality to foster a better future for our rapidly developing psychosomatic society.

Cultural diversity is highly represented in this issue. Khaustova et al shared their experiences providing proactive psychiatry support for people suffering from chronic disease in the Ukrainian population. They highlighted that such support could reduce subclinical anxiety and somatisation symptoms and improve quality of life. Shah et al introduced and compared the mental health laws and coronavirus disease 2019 (COVID-19) mental health policies in Pakistan and China. They concluded that mental health issues could be reduced by adequate mental health laws and appropriate implementation. Amone-P'Olak et ali explored the prevalence and risks of post-traumatic stress disorder among war-affected youth in Uganda. They detected background and post-war environmental risk

factors among them and highlighted the disastrous effects of war on young people's mental health. Liang *et al*⁸ translated the Central Sensitisation Inventory into the Chinese language, and they proved excellent reliability and validity. The authors also recommended the usefulness of this tool in Chinese populations.

Specific groups with psychosomatic problems are represented in this issue. Joshi et at focused on women with Dhat syndrome and studied its phenomenology, disability and sexual functioning in these patients. The authors concluded that women with non-pathological vaginal discharge similar to men with Dhat syndrome, having a deficit understanding of pathology; they recommend comprehensive assessments and management to decrease the suffering of women with the disorder experience. Sun et al^{10} studied the relationship between faecal microbiota and long-term Tibetan meditation in monks. The enriched microbiotas in monks were associated with a lower risk of anxiety, depression, cardiovascular disease and better immune function.

Wang et al¹¹ described the psychological relationships among cancer survivors. They noted that cancer survivors commonly experienced self-concept impairments, body image disturbance, sexual troubles and social relationship crises. The caregivers exhibited critical burdens and health condition issues. In addition, physicians in the related departments were experiencing challenges in patient-doctor communication and burnout. Zhang et al¹² focused on patients with gastrointestinal diseases. They discussed the necessity of generating a conventional clinical model for both functional and organic gastrointestinal illness. They reviewed the challenges, limitations and concept development. The authors emphasised the significance of a novel systematic psychosomatic model.

Depression is the most prevalent and disturbing mental disorder in modern society, and its interaction with the body is a major focus of psychiatry. Liu *et al*¹³ studied the brain's connectivity in major



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Psychosomatics and Psychiatry, Southeast University Zhongda Hospital, Nanjing, Jiangsu, China ²Department of Psychiatry, Tongji Hospital of Tongji University, School of Medicine, Shanghai, China

Correspondence to

Dr Yonggui Yuan; yygylh2000@sina.com

1

depressive episodes with mixed features. They stated that the abnormal connectivity within the default mode network's subsystem might be a critical mechanism for the pathology of major depressive episodes with mixed features. In addition, Liu et al¹⁴ explored interleukin 6, high-sensitivity C reactive protein and tumour necrosis factor-α in melancholic in atypical and anxious depression. They suggested these dynamic immunoinflammatory components could be used as potential biomarkers in depression. Simultaneously, a novel treatment study of depression was also presented in this issue. Wang et al¹⁵ focused on ketamine and its rapid antidepressant effect. The authors noted that increased activity of AMPA receptors, enhanced BDNF production and release, BDNFmTORC1 signal pathways, disinhibition, interaction with 5-HT and dopamine system and other mechanisms might be behind ketamine's rapid antidepressant effect. The relationship between depression and other clinical situations was explored. Li et al¹⁶ explored the relationship between depression and handgrip strength (HGS) and suggested that increased left HGS was associated with lower depression risk.

Anxiety and stress were other critical subjects addressed in this issue. Jin *et al*¹⁷ summarised the neuro-humoral-immune regulatory network in psychiatric and psychosomatic disorders. The authors discussed potential mechanisms of stress-induced neuroinflammation, neuron damage and apoptosis. Cai *et al*¹⁸ demonstrated the causal effects between blood pressure and anxiety, depressive symptoms, neuroticism and well-being. The authors also suggested that proper blood pressure management lowers the risk of mood disorders and cardiovascular diseases.

Finally, Chen *et al*¹⁹ studied the effect of obesity on schizophrenia's mortality risk and presented the results from a large cohort with body mass index records. They found that underweight status was associated with higher mortality in this disorder group; however, this was potentially affected by older ages.

We foresee that the articles included in this special issue will expand the knowledge of psychosomatic medicine and interdisciplinary research. Ongoing collaborative studies from psychosomatic societies will further reveal cross-cultural issues and under-represented populations throughout the world. Finally, additional research on biopsychosocial mechanisms would contribute to the clinical practice of psychosomatic disorders.

Contributors WJ, ZL and YY edited the topic. WJ and YY wrote the editorial. All authors approved the submitted version.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID ID

Yonggui Yuan http://orcid.org/0000-0001-6496-3998

REFERENCES

- 1 Fava GA, Cosci F, Sonino N. Current psychosomatic practice. Psychother Psychosom 2017;86:13–30.
- 2 Fava GA, Sonino N. Psychosomatic medicine: emerging trends and perspectives. *Psychother Psychosom* 2000;69:184–97.
- 3 Lee HB, Oldham M. The biopsychosocial model and consultationliaison psychiatry: legacy of George Engel and John Romano. J Acad Consult Liaison Psychiatry 2022;63:187–8.
- 4 Lee HB. Psychosomatic medicine and consultation-liaison psychiatry around the world: finding unity in biopsychosocial model. *Gen Psychiatr*. In press 2022.
- 5 Khaustova OO, Markova MV, Driuchenko MO, et al. Proactive psychological and psychiatric support of patients with chronic non-communicable diseases in a randomised trial: a Ukrainian experience. Gen Psychiatr 2022;35:e100881.
- 6 Shah SM, Sun T, Xu W, et al. The mental health of China and Pakistan, mental health laws and COVID-19 mental health policies: a comparative review. Gen Psychiatr 2022:e100885.
- 7 Amone-P'Olak K, Omech B, Kakinda AI, et al. Echoes of the past: prevalence and correlates of PTSD among formerly abducted youths in northern Uganda: findings from the ways study. Gen Psychiatr 2022;35:e100840.
- 8 Liang D, Yu X, Guo X. Adaptation and validation of the Chinese version of the Central Sensitisation Inventory in patients with chronic pain. Gen Psychiatr. In press 2022.
- 9 Joshi S, Tripathi A, Agarwal S, et al. Phenomenology, disability and sexual functioning in female Dhat syndrome: a study of tertiary care gynaecology outpatients. Gen Psychiatr 2022;35:e100863.
- 10 Sun Y, Ju P, Xue T, et al. The alteration of faecal microbiota balance related to long-term deep meditation. Gen Psychiatr. In Press 2022.
- 11 Wang Y, Feng W. Cancer-related psychosocial challenges. Gen Psychiatr 2022;35:e100871.
- 12 Zhang Q, Ding L, Cao J. Evolution and significance of the psychosomatic model in gastroenterology. *Gen Psychiatr* 2022;35:e100856.
- 13 Liu R, Qi H, Guan L, et al. Functional connectivity of the default mode network subsystems in patients with major depressive episodes with mixed features. Gen Psychiatr. In Press 2022.
- 14 Liu H, Wu X, Wang Y, et al. TNF-α, IL-6 and hsCRP in patients with melancholic, atypical and anxious depression: an antibody array analysis related to somatic symptoms. Gen Psychiatr 2022;35:e100844.
- 15 Wang S, Tang S, Huang J, et al. Rapid-Acting antidepressants targeting modulation of glutamatergic system: clinical and preclinical evidence and mechanisms. *Gen Psychiatr*. In Press 2022.
- 16 Li N, Zhou R, Zhang B. Handgrip strength and the risk of major depressive disorder: a two-sample Mendelian randomisation study. *Gen Psychiatr* 2022;35:e100807.
- 17 Jin R, Zhao H, Hu J. Disturbed sensitive equilibrium in psychiatry: led by stress induced inflammation. *Gen Psychiatr*. In press 2022.
- 18 Cai L, Liu Y, He L. Investigating genetic causal relationships between blood pressure and anxiety, depressive symptoms, neuroticism and subjective well-being. Gen Psychiatr. In press 2022.
- 19 Chen J, Perera G, Shetty H, et al. Body mass index and mortality in patients with schizophrenia spectrum disorders: a cohort study in a South London catchment area. Gen Psychiatr 2022;35:e100819.





Wenhao Jiang, MD/PhD/PhD. He received a doctoral degree in Neurology from the Southeast University in Nanjing, China and a doctoral degree in psychology from Georgia State University in the USA. His main research interests include imaging genetic studies in schizophrenia, bipolar disorder, and ADHD. He participated in the data analysis and coordinating phase of the ENIGMA schizophrenia project. He also participated in the remission of ADHD adults study held by the Center for Translational Research in Neuroimaging and Data Science in China.