




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Editorial

One year on: Are we ready for COVID? ☆

Y un año después, ¿estamos preparados para la COVID?



On February 15, 2020, we warned of the imminent pandemic expansion of the 2019-nCoV virus.¹ Just one year later, we have a better understanding of the antigenic characteristics, replicative cycle and genome of SARS-CoV-2, and candidate therapeutic targets have been identified. The high transmissibility of this virus via aerosols has also been confirmed, a factor that we know increases the risk of contagion in indoor environments, even if safe distances are maintained. This means that more stringent preventive measures are needed, especially among health professionals, given the risk to which they are exposed in their clinical practice.^{2,3} The rate of Covid-19 infection among health professionals has been as high as 21.4%. It is essential that we understand the reasons for and consequences of this high infection rate, as not only are health workers at risk from these transmission mechanisms, their close and continuous contact with people affected by Covid-19 has also made them vectors of transmission.

During this year, changes made in procedures and interventions as knowledge developed have generated confusion among professionals and given them the impression that the correct measures were being introduced late. The shortage of personal protective equipment and support systems for the most serious patients led the authorities to roll out plans for prioritizing and rationalizing the use of supplies, while material that offered better protection was reserved for professionals exposed to aerosol-generating procedures.⁴

The SANICOVI^{®5} study concludes by recommending measures that include the appropriate management of preventive, protective, educational, and organizational resources, both human and material. This not only prevents infections among healthcare professionals, but also ensures safer patient care. Workloads must be adjusted according to the care setting, and prevention procedures must be reinforced, training improved, and protocols disseminated and implemented to provide tailored protective measures.

Covid-19 diagnosis has increased with the incorporation of rapid tests that detect viral antigens or agglutinating antibodies, thus improving detection in the acute or convalescent phases. The sensitivity and specificity of each test depend on the stage of infection, so each one has a maximum effective period beyond which its results may be erroneous.⁶ Each test, therefore, must be selected according to time and circumstance.⁷

The course of Covid-19, characterized by an initial viral phase followed by a hyperinflammatory phase and a prothrombotic phase, is better characterized. All 3 phases impact on the various manifestations of the disease.⁷ Several prognostic factors have also been identified that have significantly influenced the management of patients during the months of the pandemic,⁸ while the early, unproven use of lopinavir/ritonavir, hydroxychloroquine, and azithromycin has been stopped after they were shown to be ineffective in trials.

The current therapeutic gold standard is oxygen therapy; high-flow oxygen therapy has emerged as an essential modality, and the need to develop intermediate respiratory care units is clear. In these units, the need for orotracheal intubation (OTI), a procedure associated with complications and mortality (OTI 54% vs. non-OTI 7%), is significantly lower thanks to non-invasive mechanical ventilation and high-dependence care (e.g. proning).⁹ In terms of medical treatment, we continue to use remdesivir as an antiviral, dexamethasone or other steroids as anti-inflammatories,¹⁰ and anticoagulation to prevent or treat thromboembolic disease, despite their obvious limitations. Tocilizumab, an IL-6 receptor blocker, continues to be used, despite conflicting results in clinical trials.

Promising new data are being obtained with convalescent plasma, hyperimmune gammaglobulin, and monoclonal antibodies that neutralize viral epitopes, such as bamlanivimab or casirivimab plus imdevinab or etesevimab.¹¹ The FDA has authorized bamlanivimab alone or in combination with etesevimab for mild-to-moderate non-hospitalized patients. Pending further evidence, the antiparasitic ivermectin, the JAK inhibitor baricitinib, and IL-1 inhibitors are not yet recommended due to lack of data.¹² Two trials have shown that colchicine shortened recovery time and decreased hospitalizations and complications. In preclinical studies, plitidepsin, approved for multiple myeloma, has recently shown 27.5 times higher anti-SARS-CoV-2 activity than remdesivir.

Fortunately, in contrast to the scant progress made in therapy, several vaccines have been authorized in just 10 months, including novel mRNA vaccines (Pfizer, Moderna), vector-based vaccines, similar to those used for Ebola (Astra-Zeneca), and other vaccines in the pipeline, such as mRNA (CureVac) and vector-based products (Janssen and the Russian Gamaleya) and adjuvanted subunit vaccines (Novavax, Sanofi-GSK).¹³ All of them target the spike (S) protein.

The very high viral replication has led to the appearance of mutant variants, including spike mutations, which demand close surveillance. Asian countries prioritized traditional techniques

☆ Please cite this article as: Peña-Otero D, Bello Dronca S, Díaz-Pérez D, de la Rosa Carrillo D, Y un año después, ¿estamos preparados para la COVID? Archivos de Bronconeumología, 2021;57:517–518.

(inactivated virus) that are less potent but offer more viral targets. Current vaccines are safe and very effective, although the duration of immunity is not known. Nor is it known whether they will avert disease or infection by preventing transmission from vaccinated individuals. Some interesting projects are investigating apparently powerful mucosal vaccines that offer sterilizing immunity. All the evidence suggests that we will need to receive more than one vaccine.

Covid-19 causes persistent symptoms in most patients (fatigue, muscle weakness, difficulty sleeping, anxiety, or depression).¹⁴ The long-term complications remain unknown. In the acute phase, this infection has put a stop to humanized healthcare and forced hospitals to restrict, or even ban, patient visiting. Despite efforts to improve the response capacity of care homes, health centers, and hospitals, the pandemic is generating delays in care¹⁵ and limiting active user participation. Wherever possible, the switch to telemedicine has accelerated, but are we prepared?

After one year, we can say that activities associated with Covid-19 biosecurity and treatment, if they are to be effective, must be backed up by high-quality science and evidence. “Evidence-based policy”¹⁶ must count from the outset on the collaboration of all actors involved in current and future challenges. In addition to the impact on the general population, the effects on health professionals are beginning to be made themselves felt, given the generalized accumulation of working hours, infections, and, not least, moral distress.

References

- Peña-Otero D, Díaz-Pérez D, de-la-Rosa-Carrillo D, Bello-Dronda S. Are we ready for the new coronavirus? *Arch Bronconeumol.* 2020;56:195–6.
- Tang S, Mao Y, Jones RM, Tan Q, Ji JS, Li N, et al. Aerosol transmission of SARS-CoV-2? Evidence, prevention and control. *Environ Int.* 2020;144:106039.
- Peña Otero D, Eguillor Mutiloa M. Medidas de protección individual y colectiva en la Covid-19. *Rev Patol Respir.* 2020;23 Supl. 3:S268–71.
- Aranaz Andrés JM, Gea Velázquez de Castro MT, Vicente-Guijarro J, Beltrán Peribáñez J, García Haro M, Valencia-Martín JL, et al. Mascarillas como equipo de protección individual durante la pandemia de COVID-19: cómo, cuándo y cuáles deben utilizarse. *J Healthc Qual Res.* 2020;35:245–52.
- Grupo S. Factores relacionados con el contagio por SARS-CoV-2 en profesionales de la salud en España. *Proyecto SANICOVI. Enferm Clin.* 2020;30:360–70.
- Sethuraman N, Jeremiah SS, Ryo A. Interpreting diagnostic tests for SARS-CoV-2. *JAMA.* 2020;323:2249–51.
- Lupu L, Palmer A, Huber-Lang M. Inflammation, thrombosis, and destruction: the three-headed cerberus of trauma- and SARS-CoV-2-induced ARDS. *Front Immunol.* 2020;11:584514.
- Mudatsir M, Fajar JK, Wulandari L, Soegiarto G, Ilmawan M, Purnamasari Y, et al. Predictors of COVID-19 severity: a systematic review and meta-analysis. *F1000Research.* 2020;9:1107.
- Hernandez-Rubio JC, Abad MS, Leon AY, Higuera GD, Rodriguez LG, Torrejon CG, et al. Outcomes of an intermediate respiratory care unit in the COVID-19 pandemic. *PLoS One.* 2020;15:e0243968.
- Horby P, Lim WS, Emberson JR, Mafham M, Bell JL, Linsell L, et al. Dexamethasone in hospitalized patients with Covid-19 – preliminary Report. *N Engl J Med.* 2020;384:693–704.
- Weinreich DM, Sivapalasingam S, Norton T, Ali S, Gao H, Bhole R, et al. REGN-COV2, a neutralizing antibody cocktail, in outpatients with Covid-19. *N Engl J Med.* 2021;384:238–51.
- COVID-19 Treatment Guidelines Panel [Accessed 11 February 2021]. Available from: <https://www.covid19treatmentguidelines.nih.gov/>, 2021.
- Izda V, Jeffries MA, Sawalha AH. COVID-19: a review of therapeutic strategies and vaccine candidates. *Clin Immunol.* 2021;222:108634.
- Huang C, Huang L, Wang Y, Li X, Ren L, Gu X, et al. 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study. *Lancet.* 2021;397:220–32.
- Hortal-Carmona J, Padilla-Bernáldez J, Melguizo-Jiménez M, Ausín T, Cruz-Piqueras M, López de la Vieja MT, et al. Efficiency is not enough. Ethical analysis and recommendations for the allocation of scarce resources in a pandemic situation. *Gac Sanit.* 2021;30192–8, <http://dx.doi.org/10.1016/j.gaceta.2020.07.006>, 192–8 (S0213-9111(20)).
- Kemp L, Aldridge DC, Booy O, Bower H, Browne D, Burgmann M, et al. 80 questions for UK biological security. *PLoS One.* 2021;16:e0241190.

David Peña-Otero,^a Salvador Bello Dronda,^{b,*} David Díaz-Pérez,^c
David de la Rosa Carrillo^d

^a *Enfermería, Hospital Sierrallana, Subdirección de Cuidados del Servicio Cántabro de Salud, Miembro IDIVAL e IISGM, Área de Enfermería Respiratoria-SEPAR, Torrelavega, Cantabria, Spain*

^b *Servicio de Neumología, Hospital Universitario Miguel Servet, Zaragoza, Spain*

^c *Enfermería, Hospital Universitario Nuestra Señora de Candelaria, Servicio Canario de Salud, Área de Enfermería Respiratoria-SEPAR, Santa Cruz de Tenerife, Spain*

^d *Servicio de Neumología, Hospital de la Santa Creu i Sant Pau, Área TIR-SEPAR, Barcelona, Spain*

* Corresponding author.

E-mail address: sbello@salud.aragon.es
(S. Bello Dronda).