

POSTER PRESENTATION

Open Access

# Expansion and characterization of tumor-infiltrating lymphocytes from human sarcoma

MacLean Hall<sup>1\*</sup>, John Mullinax<sup>1</sup>, Erica Royster<sup>1</sup>, Angela Reagan<sup>1</sup>, Ricardo Gonzalez<sup>1</sup>, Jose Pimiento<sup>1</sup>, Shari Pilon-Thomas<sup>2</sup>, Amod Sarnaik<sup>1</sup>

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

## Background

Adoptive Cell Transfer (ACT) using Tumor Infiltrating Lymphocytes (TIL) has previously been shown at our institution and others to be an effective treatment for metastatic melanoma, resulting in a 38% response rate [1]. We applied this therapy to other solid tumors that have demonstrated a positive correlation between immune infiltrates and patient outcome [2]. Specifically in sarcoma, intra-tumoral CD4+ and CD8+ T cells have been detected, but their potential for *ex vivo* expansion is still relatively unexplored [3]. In this study, we investigated the feasibility of expanding TIL from surgically resected sarcoma specimens and analyzed the phenotype of these lymphocytes.

## Methods

Four different subtypes of sarcoma were surgically resected from patients accrued under an IRB approved research protocol (MCC50064). A portion of the tumor specimens was digested and immediately phenotyped by flow cytometry. The remaining tumor was minced and plated as fragments for the isolation of TIL, which were expanded *in vitro* for six weeks using high dose IL-2. Eight separate TIL cultures were established and phenotyped by flow cytometry.

## Results

Analysis of enzymatically digested human sarcoma specimens showed that 64% of lymphocytic infiltrates were CD3+ cells. TIL were isolated from fragments of each of the four sarcoma specimens as eight individual cultures and propagated *in vitro*, with TIL observed in 59 out of 84 (70%) fragments. Of the expanded CD3+ TIL, on average 39% were CD8+ T cells that expressed both the

co-stimulatory molecule 4-1BB (33%) and inhibitory PD-1 (41%) by flow cytometry.

## Conclusions

Human sarcoma specimens yield CD3+ CD8+ TIL which can be expanded *in vitro*, supporting further investigation into the feasibility of adoptive cell transfer as a therapy for these patients. Efforts are currently focused on the scalability of this process and the functional capacity of these TIL. Additionally, the expression of 4-1BB and PD-1 on a substantial of CD3+ CD8+ TIL demonstrates the opportunity to modulate these pathways to improve both yield and function, another endeavor we are presently investigating.

## Authors' details

<sup>1</sup>H. Lee Moffitt Cancer Center, Tampa, FL, USA. <sup>2</sup>H. Lee Moffitt Cancer Center & Research Institute, Tampa, FL, USA.

Published: 4 November 2015

## References

1. Pilon-Thomas S, Kuhn L, Ellwanger S, Janssen W, Royster E, Marzban S, et al: Efficacy of adoptive cell transfer of tumor-infiltrating lymphocytes after lymphopenia induction for metastatic melanoma. *J Immunother* 35(8):615-620.
2. Turcotte S, Gros A, Hogan K, Tran K, Hinrichs CS, Wunderlich JR, et al: Phenotype and function of T cells infiltrating visceral metastases from gastrointestinal cancers and melanoma: implications for adoptive cell transfer therapy. *J Immunol* 191(5):2217-2225.
3. Tseng W, Malu S, Zhang M, Chen J, Sim GC, Wei W, et al: Analysis of the Intratumoral Adaptive Immune Response in Well Differentiated and Dedifferentiated Retroperitoneal Liposarcoma. *Sarcoma* 2015:547460.

doi:10.1186/2051-1426-3-S2-P19

Cite this article as: Hall et al.: Expansion and characterization of tumor-infiltrating lymphocytes from human sarcoma. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):P19.

<sup>1</sup>H. Lee Moffitt Cancer Center, Tampa, FL, USA

Full list of author information is available at the end of the article