



Dyslipidemia is associated with risk for rotator cuff repair failure: a systematic review and meta-analysis



Andrew P. Gatto, BA^{a,*}, Daniel A. Hu, BS^b, Brian T. Feeley, MD^c, Drew Lansdown, MD^c

^aCollege of Osteopathic Medicine, Touro University California, Vallejo, CA, USA

^bFeinberg School of Medicine, Northwestern University, Chicago, IL, USA

^cDepartment of Orthopedic Surgery, University of California San Francisco, San Francisco, CA, USA

ARTICLE INFO

Keywords:

Shoulder
Rotator cuff
Repair
Risk
Retear
Function
Dyslipidemia
Hyperlipidemia

Level of evidence: Level IV; Systematic Review

Background: Lipid deposition secondary to dyslipidemia (DLD) is shown to have a significant impact on tendon pathology, including tendon elasticity, fatty infiltration, and healing properties. Rotator cuff repair is a common procedure, susceptible to influence from many tear-related and patient-related characteristics. The purpose of this study was to determine the relationship between DLD and rotator cuff repair outcomes with analysis of re-tear risk and function.

Methods: PubMed, Embase, and SPORTDiscus were searched for all English-language, peer-reviewed studies between 2000 and the present, which analyzed relationships between patient-related factors and outcomes of rotator cuff repair. Studies that explicitly examined the effect of DLD on rotator cuff repair outcomes were chosen for inclusion. Included studies were assessed for methodological quality, and data were extracted for meta-analysis.

Results: Of the 3087 titles, 424 were screened by abstract, and 67 were reviewed in full. Inclusion criteria were met by 11 studies. Of these studies, 5 studies assessed re-tear, 2 studies measured function, 3 studies reported both re-tear and function, and 1 study evaluated the risk of re-tear necessitating a revision surgery. The studies report no significant difference in functional outcomes. Meta-analysis revealed that DLD patients had a significantly higher risk of re-tear after primary rotator cuff repair (odds ratio 1.32, 95% confidence interval 1.06-1.64).

Conclusion: DLD leads to an increased risk of re-tear after rotator cuff repair, although function appears to be unimpaired. DLD should be considered among other risk factors when counseling patients regarding expected rotator cuff repair outcomes.

© 2022 The Authors. Published by Elsevier Inc. on behalf of American Shoulder & Elbow Surgeons. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Rotator cuff tears, most commonly due to avulsion of the tendon from its insertion, are one of the leading causes of shoulder pain and dysfunction.⁵⁸ Rotator cuff repair reliably improves function and quality of life,^{15,22,43} although repair failure is not uncommon. The odds of re-tear are multifactorial, and occurrence ranges from 20% to 60% after rotator cuff repair surgery.¹³

Extensive evidence has identified tear-related and patient-related characteristics, which may adversely affect rotator cuff repair outcomes. Increasing age is the most predictive characteristic for rotator cuff repair failures.^{24,29,35,37} Although clear mechanisms for this increase in failure rate remain incompletely defined. Fatty infiltration, atrophy, delamination, and increasing preoperative tear size are specific factors that increase the odds of

re-tear.^{9,24,30,35,37} Systemic conditions, such as diabetes mellitus, obesity, and tobacco, have also been implicated.^{14,25,29,62} Dyslipidemia (DLD) is among these patient characteristics that potentially impact rotator cuff repair outcomes. More than 90 million US adults have a total cholesterol (TC) level >200 mg/dL, and nearly 43 million adults have a high-density lipoprotein (HDL) level <40 mg/dL.^{12,54} As the population ages, the intersection between these diseases is expected to grow.⁴⁴

DLD encompasses disorders of lipoprotein metabolism, including overproduction of TC (ie, hyperlipidemia), low-density lipoprotein (LDL), and triglycerides (TGs) and/or underproduction of HDL. Current evidence suggests that lipids can accumulate in the extracellular matrix of tendons, leading to the formation of lipid deposits altering mechanical properties and increasing local inflammation.^{26,59} Hyperlipidemia animal models display reduced tendon elasticity, greater fatty infiltration, and poor tendon-to-bone healing.^{6,7,14} DLD is shown to be an independent risk factor for Achilles, patellar, and rotator cuff tendon pathology.^{1,21,41,47,52}

Institutional review board approval was not required for this systematic review.

*Corresponding author: Andrew P. Gatto, BA, College of Osteopathic Medicine, Touro University California, 1310 Club Drive, Vallejo, CA 94592, USA.

E-mail address: agatto2@student.touro.edu (A.P. Gatto).

<https://doi.org/10.1016/j.xrrt.2022.02.003>

2666-6391/© 2022 The Authors. Published by Elsevier Inc. on behalf of American Shoulder & Elbow Surgeons. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1
Strategy for PubMed search.

PubMed			
1.	Rotator cuff	7.	lipoprotein*
2.	Dyslipidemia	8.	lipid*
3.	Hyperlipidemia	9.	comorbid*
4.	Hypercholesterolemia	10.	factor*
5.	Hyperlipoproteinemia	11.	predict*
6.	Cholesterol	12.	failure*

Search Term: 1 AND (2 OR 3 OR 4 OR 5 OR 6 OR 7 OR 8 OR 9 OR 10 OR 11 OR 12).

Limits: Date 2000–2021, Full text, English.

*Search function, represents a root term.

Rotator cuff disease in the presence of DLD has been associated with tendinopathy, increased risk of tear, and increased risk of postoperative retear, although the evidence is debated.⁶⁰

The purpose of our systematic review and meta-analysis was to evaluate the association between DLD and outcomes of rotator cuff repair, notably retear and functional outcomes. We hypothesized that DLD patients would have a greater risk of rotator cuff retear and poorer functional outcomes.

Materials and methods

A systematic review was conducted using PubMed, Embase, and SPORTDiscus in January 2022. The study protocol was registered with the PROSPERO International Prospective Register of Systematic Reviews (registration number: CRD42021264487). Our purpose was to identify all English-language, peer-reviewed studies between 2000 and the present, which analyzed the relationships between patient-related factors and outcomes to rotator cuff tear. The search was completed using keywords: “rotator cuff,” “dyslipidemia,” “hyperlipidemia,” “hypercholesterolemia,” “hyperlipoproteinemia,” “cholesterol,” “lipoprotein*,” “lipid*,” “comorbid*,” “factor*,” “predict*,” and “failure*.” The search strategy for PubMed is presented in [Table 1](#).

Criteria

The following inclusion criteria were applied to study selection: (1) Population included adult DLD patients (defined as elevated TC, LDL, TG, or low HDL), confirmed by health record or laboratory test. (2) All subjects underwent surgical repair for primary rotator cuff tear; (3) Available comparison of non-DLD patients undergone primary rotator cuff repair; (4) Measures of outcome, including retear, revision, quantitative range of motion, and/or satisfaction/pain/function score; (5) Study design to be primary research of randomized control, cohort, cross-sectional, case–control, or case series design, with available full text.

Exclusion criteria included (1) Population failed to include surgically treated primary rotator cuff repair, DLD patients; (2) DLD measure was poorly defined; (3) Primary cuff repair was 1 year beyond diagnosis; (4) Lack of comparison to non-DLD, rotator cuff repair patients; (5) Significant difference (ie, $P \leq .05$) in patient characteristics between study groups, without control; (6) Outcomes not specific to rotator cuff repair or DLD; (7) Studies published before 2000; (8) Full text unavailable; and (9) Review articles and case studies.

Literature screening and data extraction

Article screening was done independently by 2 authors (A.G. and D.H.); discrepancies were settled by discussion. All titles were screened and removed if they were obviously irrelevant. Abstracts were reviewed for relevance, determined by clear naming of a DLD

measure, the mention or implication of factors associated with rotator cuff repair outcome, or focus on comorbidities. Those relevant progressed to full text review for inclusion. Data were extracted by 2 authors (A.G. and D.H.) and included basic article characteristics, population characteristics, populations size and source, dropout rate, intervention and control groups, follow-up period, imaging modality, DLD definition and guidelines, preoperative and postoperative outcome scores, and retear totals. All statistical outcomes that aligned with a DLD measure or comparison group (ie, measures of central tendency, relative ratios, confidence intervals, and regression models) were also extracted.

Quality assessment

Two reviewers (A.G. and D.H.) independently assessed the methodological quality and reliability of the included studies using the Newcastle-Ottawa Scale.⁵⁷ Studies were assessed for potential sources of biases, generalizability, and control of confounding factors. For rating comparability, preoperative tear size on magnetic resonance imaging was selected as the most important factor for a study’s analysis to control for, as this seems to be the most reliable predictor for retear.^{9,24,38} A 6-month follow-up was determined adequate.^{32,50} Reviewers resolved discrepancies by consensus. Studies were included if they were, at least, “satisfactory”. The risk of bias across studies to assess cumulative evidence was determined independently by 2 authors and settled by consensus.

Statistical analysis

RevMan 5.4.1 (Review Manager, The Cochrane Collaboration, 2020, London, UK) was used for meta-analysis. Cancienne et al¹¹ and O’Donnell et al⁴⁶ had significant overlap, as both collected their cohort from the Humana database in PearlDiver. O’Donnell et al⁴⁶ had larger population and more recent data, thus it was included in the meta-analysis and Cancienne et al¹¹ were excluded. Meta-analysis was performed for retear, although heterogenous outcome measures and analysis methods of the selected studies prevented us from performing a meta-analysis on these other variables. Odds ratio (OR) was selected for measure of effect size, presented with 95% confidence interval. Analysis was carried out with a random effects model and Mantel-Haenszel estimate. Heterogeneity was evaluated by Cochrane’s Q test and the Higgins I^2 test; heterogeneity thresholds (eg, substantial $I^2 = 50\%$ – 90%) were according to the Cochrane Reviews handbook.²⁷

Results

The search strategy yielded 5106 titles, 1872 of which were removed as duplicates, leaving 3234 unique titles reviewed. Of this, 425 abstracts were screened, and 67 full texts were reviewed. All authors came to a consensus for the 11 studies^{4,8,11,23,25,30,31,38,46,49,61} that met criteria inclusion ([Figure 1](#)).

Table II
Summary of studies.

Title	Study details: author, year, design, level	Population	Design details	Dyslipidemia measure	Statin use	Outcome statistics	Impact of DLD
Factors affecting rotator cuff integrity after arthroscopic repair for medium-sized or larger cuff tears: a retrospective cohort study ³¹	Kim et al, 2017, retrospective cohort III	180, age: 60.4 ± 7.4, M/F: 84/96 single institution	36 DLD 144 non-DLD	TC >240 mg/dL	Not described	Retear: OR 52.814 (95% CI 3.229-808.643), <i>P</i> = .004*	Retear
Factors predictive of healing in large rotator cuff tears: is it possible to predict retear preoperatively? ³⁰	Jeong et al, 2018, case –control III	102 (112 S), age: 65.6 ± 6.6, M/F: 45/67, single institution	21 DLD; 91 non-DLD	Hyperlipidemia diagnosed in record	Not described	Retear: <i>P</i> = .628	No retear
Hypo-high-density lipoproteinemia is associated with preoperative tear size and with postoperative retear in large to massive rotator cuff tears ⁴⁹	Park et al, 2020, case series IV	195, age: 60.5 ± 7.5 M/F: 100/95, single institution	167 DLD: HyTC 58; HyTg 51; HyLDL 142; Hypo-HDL 66; Hy-non-HDL 109 28; non-DLD	HyTC: TC ≥200 mg/dL; HyTG: TG ≥ 150 mg/dL; HyLDL: LDL ≥ 100 mg/dL HypoHDL: HDL <40 mg/dL Male and <50 mg/dL Female Hy-non-HDL: non-HDL ≥130 mg/dL	No statin use	Hypo-HDL: L/Ma tears: OR 3.0 (95% CI, 1.1-8.8), <i>P</i> = .04*	Retear with Hypo-HDL
Perioperative serum lipid status and statin use affect revision surgery rate after arthroscopic rotator cuff repair ^{11,*}	Cancienne et al, 2017, retrospective cohort III	30,638 TC 13,164; LDL 12,337 TG 13,230 Age: 40-84 M/F: TC 6775/6389; LDL 6337/6000; TG 6819/6411 PearlDiver Database	TC: 5001 (mod 3,535, hi 1466) LDL: 6459 (mod 5,518, hi 941) TG: 5536 (mod 2,850, hi 2686) normal TC: 8163 normal LDL: 5878 normal TG: 7694	(Norm<Mod≤Hi) TC: < 200 ≤ 240 mg/dL LDL: <100 ≤ 160 mg/dL TG: <150 ≤ 200 mg/dL	Statin/no statin Norm TC: 4824/3339 Mod TC: 1585/1950 High TC: 765/701 Norm LDL:3812/2066 Mod LDL: 2465/3053 High LDL: 494/447	Retear TC mod: OR 1.20 (95% CI, 1.03-1.40), <i>P</i> = .022* TC hi: OR 1.36 (95% CI, 1.14-1.64), <i>P</i> = .001* LDL mod: OR 1.24 (95% CI, 1.10-1.41), <i>P</i> = .001* LDL hi: OR 1.46 (95% CI, 1.08-1.99), <i>P</i> = .014* TG mod: OR 1.13 (95% CI, 0.94-1.34), <i>P</i> = .190 TG hi: OR 1.09 (95% CI, 0.95-1.26), <i>P</i> = .222 Retear: OR 1.09 (95% CI, 1.01-1.18), <i>P</i> = .32*	Retear with elevated TC and LDL
The effect of patient characteristics and comorbidities on the rate of revision rotator cuff repair ⁴⁶	O'Donnell et al, 2020 Case series IV PearlDiver database	41,467 Age: 58.3% 60-74 M/F: 21,853/19,615	22,068 DLD 19,399 non-DLD	Hyperlipidemia diagnosed in record	Not described	Retear: OR 1.09 (95% CI, 1.01-1.18), <i>P</i> = .32*	Retear
Combination of risk factors affecting retear after arthroscopic rotator cuff repair: a decision tree analysis ²⁵	Harada et al, 2020 Retrospective cohort III	286 Age: 64.9 ± 7.1 M/F: 160/126 Single institution	61 DLD 225 non-DLD	Hyperlipidemia diagnosis from PCP in record; TC determined by record blood test and treated as a continuous variable	Not described	Retear: <i>P</i> = .0178† Functional improvement: JOA: <i>P</i> = .1121 UCLA: <i>P</i> = .1114	Retear; no functional
Does statin-treated hyperlipidemia affect rotator cuff healing or muscle fatty infiltration after rotator cuff repair? ⁴	Amit et al, 2021 Prospective cohort II	77 Age: Statin: 61.7 ± 6.6; Control: 60.2 ± 7.4 M/F: 42/35 Single institution	38 DLD 39 non-DLD	Hyperlipidemia diagnosis from PCP in record	All DLD patients on statins	Retear: OR 0.65 (95% CI 0.24-1.80), <i>P</i> = .41 WORC: <i>P</i> = .087 ASES: <i>P</i> = .84 CSS: <i>P</i> = .76 DASH: <i>P</i> = .064	No retear or function
Hyperlipidemia increases the risk of retear after arthroscopic rotator cuff repair ²³	Garcia et al, 2017 Retrospective cohort III	85 Age: 62.1 (45.3-74.3) M/F: 54/32 Single institution	33 DLD 52 non-DLD	TC, HDL, LDL, TG from PCP in record	All DLD patients on statins	Retear: OR 6.5 (<i>P</i> < .001)† Retear by DLD measure: HDL: <i>P</i> = .23; LDL <i>P</i> = .30; TC <i>P</i> = .25, TG <i>P</i> = .35 Functional (ASES and VAS pain): <i>P</i> = .14 ASES: <i>P</i> = .352 SST: <i>P</i> = .238 SANE: <i>P</i> = .240 VAS function: <i>P</i> = .792 VAS pain: <i>P</i> = .147 FE: <i>P</i> = .034‡ ER: .269	Retear; no functional
Comorbidity effect on speed of recovery after arthroscopic rotator cuff repair ⁸	Berglund et al, 2018 Retrospective cohort III	627 Age: 62.1 (29-87) M/F: 382/245 Single institution	132 DLD 495 non-DLD	Hyperlipidemia diagnosed in record	Not described	ASES: <i>P</i> = .352 SST: <i>P</i> = .238 SANE: <i>P</i> = .240 VAS function: <i>P</i> = .792 VAS pain: <i>P</i> = .147 FE: <i>P</i> = .034‡ ER: .269	No function, except FE

Dyslipidemia with perioperative statin usage is not associated with poorer 24-month functional outcomes after arthroscopic rotator cuff surgery ⁶¹	Zheng et al. 2020 Retrospective cohort Single institution	266 Age: 61.3 ± 9.8 M/F: 117/149	134 DLD 132 non-DLD	Hyperlipidemia diagnosed in record	115 DLD patients on statins	No function
Factors related to symptomatic failed rotator cuff repair leading to revision surgeries after primary arthroscopic surgery ³⁸	Lee et al. 2020 Case-control Single institution	98 Age: I: 63.2 ± 7.4; II: 61.6 ± 7.4 M/F: 44/54	Group I (44): retear that required revision surgery Group II (54): retear that did not require revision surgery	Preoperative lab test: TC and LDL treated as continuous variables	Not described	Revision surgery in retear patients

S, shoulders; M, male; F, female; TC, total cholesterol; LDL, low-density lipoprotein; TG, triglycerides; DLD, dyslipidemia; HyTC, hypertriglyceridemia; HyLDL, hyper-LDLemia; Hypo-HDL, hypo-HDLemia; Hy-nonHDL, hyper-non-HDLemia; PCP, primary care physician; Norm, normal; Mod, moderate; OR, odds ratio. DeOrio and Coifield classification⁴⁹—Sm, small; Me, medium; L, large; Ma, massive.

*Significance by multiple logistics regression.
[†]Significance by chi-squared.
[‡]Significance by t-test.
[§]Significance by univariate logistic regression.

Five studies^{11,30,31,46,49} assessed retear, 2 studies^{8,61} measured function, and 3 studies^{4,23,25} reported both retear and function outcomes. In addition, 1 study³⁸ examined outcomes by evaluating factors associated with a retear necessitating a revision surgery. In 7 studies^{4,8,11,23,25,31,61} data were obtained via cohort design, 6^{4,8,11,23,25,31,61} of which were retrospective, 2 studies^{30,38} were case–control studies, and 2 studies^{46,49} were case series.

Dyslipidemia

DLD was defined differently among the studies (Table II). Six studies^{4,8,23,30,46,61} used a “hyperlipidemia” diagnosis in patient records, 2^{4,23} of which noted corroboration with the patient’s primary care physician. Four studies^{11,31,38,49} accessed preoperative laboratory values, including TC, LDL, HDL, and TG. One study²⁵ used both a recorded “hyperlipidemia” diagnosis from the primary care physician and preoperative laboratory values. Statin use was described in 5 studies; 2 studies^{4,23} had all patients on statins, 1 study⁴⁹ had no patients on statins, and 2 studies^{11,61} detailed a mix of statin use; 6 studies^{8,25,30,31,38,46} did not provide details on statin use.

Retear

DLD was significantly associated with retear in 6 of 8 studies^{11,23,25,31,46,49} that explored retear rates after rotator cuff repair (Table II). Park et al⁴⁹ and Cancienne et al¹¹ isolated hypo-HDL and elevated TC and LDL as significant risk factors for retear, respectively. Harada et al,²⁵ Kim et al,³¹ and O’Donnell et al⁴⁶ identified hyperlipidemia as an independent risk factor among other patient-related and tear-related factors. Harada et al²⁵ went on to determine that a combination of anteroposterior tear size (ie, ≥40 mm), hyperlipidemia, and critical shoulder angle (ie, ≥37°) generated the greatest risk of retear compared with other combinations. Garcia et al²³ found the risk of DLD independent from cholesterol-lowering medication dosage. Amit et al⁴ and Cancienne et al¹¹ found that DLD was not a risk for retear when controlled by statin therapy.

Meta-analysis was performed with results from 7 studies^{4,23,25,30,31,46,49} reporting retear among DLD patients; Cancienne et al¹¹ were excluded because of patient overlap with O’Donnell et al.⁴⁶ In total, 42,402 subjects were included, 52.8% of which had DLD. Patient age ranged from 45.3 to 84 years, with 54.0% males. There were 41,467 subjects drawn from the PearlDiver Database⁴⁶ and 935 patients drawn from their respective institutions.^{4,23,25,30,31,49} The results showed that DLD was associated with retear as DLD patients were significantly more likely to experience retear after cuff repair (OR 1.54, 95% CI 1.01-2.35; Figure II). The combination of studies displayed substantial heterogeneity²⁷ ($\chi^2 = 13.79, P = .03; I^2 = 56%$), with a significant overall effect ($Z = 1.99, P = .05$).

Function

In 5 of 5 studies^{4,8,23,38,61} recording functional outcomes, there was no significant difference in DLD patients versus non-DLD patients across multiple outcome measures and patient-reported assessments (Table II). Function was recorded postoperatively at 3, 6²⁵ 12^{4,8}, and 24 months.⁶¹ The evaluations used included the American Shoulder and Elbow Surgeons score (3 studies^{4,8,23}), UCLA Shoulder Rating Scale (2^{25,61}), Constant Shoulder Score (2^{4,61}), Visual Analog Scale for Pain (3^{8,23,61}) and Function (1⁸), Disability of Arm, Shoulder, and Hand (1⁴), Japanese Orthopedic Association (1²⁵), Simple Shoulder Test (1⁸), Single Assessment Numeric Evaluation (1⁸), and Western Ontario Rotator Cuff index

Table III
Newcastle-Ottawa risk of bias assessment.

Authors	Study design	Level of evidence	Selection	Comparability	Outcome/exposure	Quality [*]
Amit et al, 2021 ⁴	Prospective cohort	II	★★★★	★	★★★	Excellent
Berglund et al, 2018 ⁸	Retrospective cohort	III	★★★★		★	Satisfactory
Cancienne et al, 2017 ¹¹	Retrospective cohort	III	★★★★	★	★★★	Excellent
Garcia et al, 2017 ²³	Retrospective cohort	III	★★★★		★★	Good
Harada et al, 2021 ²⁵	Retrospective cohort	III	★★★★		★★★	Good
Kim et al, 2017 ³¹	Retrospective cohort	III	★★★★	★★	★★	Excellent
Zheng et al, 2020 ⁶¹	Retrospective cohort	III	★★★★		★★	Good
Jeong et al, 2018 ³⁰	Case–control	III	★★★★	★	★	Good
Lee et al, 2020 ³⁸	Case–control	III	★★★	★	★★	Good
O'Donnell et al, 2020 ⁴⁶	Case series	IV	★★★★	★	★★★	Excellent
Park et al, 2020 ⁴⁹	Case series	IV	★★★★	★★	★★	Excellent

Cohort studies: Selection (4 stars) graded on representativeness of exposed cohort, selection of nonexposed cohort, exposure ascertainment, evidence that the outcome is not present at study onset; Comparability (2 stars) graded on control for preoperative tear size and/or additional factors; Outcome (3 stars) graded on mode of assessment, follow-up length of 6 months, and dropout rate. Case–control and Case series studies: Selection (4 stars) graded on case definition, representativeness of cases, selection of controls, definition of controls; Comparability (2 stars) graded on control for preoperative tear size and/or additional factors; Exposure/outcome (3 stars) graded on ascertainment of exposure, continuity in exposure ascertainment between groups, and nonresponse rate.

^{*}Quality: excellent 8-9 stars, good 6-7 stars, satisfactory 4-5 stars, unacceptable 1-3 stars.




Figure 1 Flowchart for the identification of included studies.

(⁴). Berglund et al⁸ quantitatively measured forward elevation and external rotation, revealing reduced forward elevation for hyperlipidemia patients 1 year after rotator cuff repair ($P = .034$). Three of these 5 studies^{4,23,61} had patients on stain therapy.

Rearer requiring revision

Lee et al³⁸ evaluated factors that increased the odds of a rotator cuff rearer necessitating a revision. Patients whose rearer required repair had significantly elevated TC (OR 1.01, 95% CI 1.00-1.03, $P = .015$) and LDL (OR 1.02, 95% CI 1.01-1.04, $P = .11$) when compared with rearer patients who did not require repair; TG did not have a significant impact ($P = .230$). Of note, revision patients had persistent pain and functional impairment for at least 3 months after primary surgery.

Risk of bias

Using the Newcastle-Ottawa Scale,⁵⁷ 5 studies^{4,11,31,46,49} were deemed excellent quality, 5 studies^{23,25,30,38,61} were good quality, 1 study⁸ was satisfactory quality, and no papers were of unsatisfactory quality (Table III). Assessing comparability, 2 studies^{31,48}

controlled for preoperative rearer, which was previously mentioned as the most important factor to control for. For outcomes, all 7 studies^{4,8,11,23,25,31,61} of cohort design had a minimum of 6-month follow-up.

Discussion

This systematic review elucidates the adverse association between DLD and rotator cuff repair outcomes. There was a significant increase in the likelihood of rotator cuff repair rearer for patients with DLD. In the evaluated studies, functional outcomes after rotator cuff repair do not appear to be significantly affected for patients with DLD compared with those without DLD.

Growing evidence suggests that DLD is a risk factor for the development of rotator cuff disease, among other tendon pathology. In a meta-analysis by Lai et al,³⁴ DLD patients had a 2.17 (95% CI, 1.46-3.23) greater odds of rotator cuff disease than non-DLD patients. Lin et al⁴¹ followed hyperlipidemia patients over an 11-year period and found an elevated risk with a hazard ratio of 1.48 (95% CI 1.42-1.55, multivariate analysis). Abboud and Kim³ revealed that patients with rotator cuff tears had a higher incidence of elevated TC, ($\bar{x} = 237$ mg/dL, $P = .03$), LDL ($\bar{x} \approx 155$, $P = .02$), and TG ($\bar{x} \approx 185$, $P = .03$); although lower, HLD was statistically insignificant ($\bar{x} \approx 35$, $P = .10$). Animal studies show reduced tendon elasticity and increased fatty infiltration in hyperlipidemic mice and rabbit rotator cuff tendons,^{6,16} two factors believed to be associated with rearer.^{4,21,25,28} These results suggest rotator cuff tendons in a DLD environment are compromised preoperatively, sustaining damage that may increase the size of primary tear or alter effective healing postoperatively.⁴⁹ Contradictory studies exist, however, making the association between DLD and rotator cuff disease unclear.⁶⁰ A previous meta-analysis by Zhao et al⁶² found borderline contrary findings in evaluating the rearer risk for DLD patients (OR 1.50, 95% CI 0.99-2.26), although this analysis only included 4 studies that were also used in our analysis (Garcia et al,²³ Harada et al,²⁵ Jeong et al,³⁰ and Kim et al³¹). Our study uses rearer data from an additional 4 studies^{4,11,46,49} to offer stronger statistical power, adding to the existing knowledge regarding risk factors for rotator cuff repair failure.

Multiple competing mechanisms likely contribute to the increased odds of rearer imposed by DLD. Circulation-derived lipid deposits in tendon extracellular matrix have been widely observed in nonfamilial hypercholesterolemia patients. The present research suggests that these deposits cause oxidative damage, persistent inflammatory cytokine production, obstructed tissue vascularity,




Figure 2 Meta-analysis forest plot synthesizing the effect of DLD on rotator cuff retear odds. Mantel-Haenszel, random effects model.

and reduced cholesterol efflux and matrix turnover by macrophages and tenocytes.^{2,5,11,55} In a recent study by Li et al,³⁹ high cholesterol inhibited gene expression in tendon-derived stem cells, a newly identified resident cell population theorized to be responsible for tendon maintenance and repair. A subsequent study by Li et al⁴⁰ revealed that cholesterol also induces apoptosis and autophagy of these stem cells. This pathophysiology may underly the stiffness, poor collagen organization, and fatty infiltration observed in healing DLD rotator cuff repair models.^{7,16} In addition, Chung et al¹⁶ found that hyperlipidemia had a deleterious effect on fatty infiltration in a murine rotator cuff model, both pre- and post-repair. Although no studies observe this association in humans, 3 studies in this review investigated DLD concurrently with fatty infiltration as predictors for retear: Harada et al²⁵ and Kim et al³¹ demonstrated both as risk factors in a patient population, whereas Jeong et al³⁰ reported that DLD was not a risk factor. These coinciding results allow for the postulation that DLD patients may experience greater fatty infiltration, contributing to retear risk. Fatty infiltration is the subject of substantial research for its role in rotator cuff tear pathology.^{42,53,54} The stem cells responsible for fatty infiltration harbor potentially restorative value, which can be potentiated via pharmacological intervention.^{20,36,56} This potential could prove therapeutic and DLD patients may uniquely benefit from such intervention, in conjunction with lipid control via statins.

Statin therapy appears to effectively suppress or eliminate the detrimental effects of DLD. In murine studies, simvastatin administration reduced fatty infiltration and improved tendon-to-bone healing, compared with hyperlipidemia mice not on a statin.¹⁶ Amit et al⁴ showed that patients on statin therapy had fatty infiltration progression, retear rate, and function comparable to non-DLD patients. Similar results for functional outcomes are found in Zheng et al.⁶¹ Cancienne et al¹¹ offer perhaps the most convincing evidence that statins mitigate the risk of retear, given the significant size of study and comparison within groups. In this study, DLD patients had a significantly greater risk of retear without statins, although DLD patients on statins had risk equivalent to non-DLD patients. Garcia et al²³ found that statins were ineffective against retear, although this study lacks control of confounding variables statin therapy on tendon healing is not well defined and is still the subject of upcoming research.^{10,17-19}

DLD is a condition that encompasses any disorder of lipoproteins. Therefore, concluding that DLD results in an increased risk of retear is relatively nonspecific. Most of the articles in this review used a recorded “hyperlipidemia” diagnosis or elevated TC to identify subjects. Cancienne et al¹¹ recorded TC, LDL, and TG and found that moderate and high TC and LDL (Table II) were independently associated with retear. Park et al⁴⁹ additionally dissect

DLD into low HDL and high non-HDL (Table II), revealing that low HDL (HDL <40 mg/dL for males and HDL <50 mg/dL for females) alone is associated with retear. They also report that low HDL increased preoperative tear size, which may explain the increased retear risk. Low HDL has also been reported as an independent risk for primary tear.^{2,48} HDL, also known as “good” cholesterol, is responsible for collecting excess cholesterol in the body for disposal or recycling, reducing systemic deposition, and serving a vasculo-protective role. Additional beneficial properties include anti-inflammatory, antioxidant, anti-thrombotic, and angiogenesis regulation.⁵¹ Reduced HDL has been linked to tendon xanthoma formation.⁴⁵ Overall, little is known about the roles of specific lipoproteins in tendon pathology, although specified research may help clearly characterize the detrimental effect of DLD.

Although some risk factors are more apparent than others, it appears reliably evident that rotator cuff repair failure does not rely on one factor.^{9,14,24,25,30,35,37,62} In addition, risk factors rarely appear exclusively in isolation in clinical settings. Harada et al²⁵ developed a novel decision tree analysis to predict the probability of retear with coexisting risk factors. In addition to hyperlipidemia, this analysis included anteroposterior tear size, fatty infiltration, and critical shoulder angle. Kwon et al³³ created a 15-point numerical scoring system based on retraction, fatty infiltration, anteroposterior tear size, age, bone mineral density, and work activity. Further development of evaluation systems such as these is a worthwhile endeavor to provide an accurate and reliable prediction of rotator cuff repair outcome. Although our study suggests that DLD is a risk factor for retear, we cannot determine its true effect, and thus, we recommend it be considered among other coexisting risk factors.

This study has several limitations. Given the design of the systematic review, variability between studies can have unrecognized implications, including variation in study design, subject pools, statin therapy regimen, surgery and rehabilitation protocol, DLD measure, and statistical methods. Statin use varied across studies and is considered a major confounder, as it seems to impact retear odds. In addition, a large portion of the subjects came from a database study, which comes with inherent limitations, such as coding errors and attrition due to insurance. Overall, our meta-analysis had a significant effect (Z = 2.48, P = .01) but was rated as having substantial heterogeneity by the Higgins I² (56%). Second, in several studies, DLD as a risk factor was measured concurrently with other risk factors, likely producing a confounding effect.^{8,25,30,31,46,49} Although we could not separate these factors for meta-analysis, great effort was put into scrutinizing the statistical control of confounding variables in our risk of bias assessment. Although the majority of studies controlled for confounding

variables, only 2 studies adequately controlled for preoperative tear size.^{17,49} Third, retear was not well defined by all studies (ie, biologic vs. human error cause of retear). In addition, there was variability in the criteria for retear. Finally, the adequacy of lipid control over time and length of statin therapy likely play a significant role in the overall effect of DLD. No study in this review recorded such data, although variability likely exists between subjects with differing effects on tendon pathology.

Conclusion

Rotator cuff repair outcomes, particularly retear risk, are associated with DLD. Functional outcomes appear unaffected, with possible exception to range of motion. DLD should be considered among other risk factors when assessing patients' rotator cuff repair outcomes.

Disclaimers:

Funding: No funding was disclosed by the authors.

Conflicts of interest: Brian T Feeley: Editorial Board: Journal of Shoulder and Elbow Surgery – Associate Editor, Current reviews in musculoskeletal medicine – Editor in Chief; Financial Col stocks in Bioniks Inc and Kaliber.ai; Research support from National Institute of Health, Veterans Affairs, Orthofix, and Stryker Corporation.

Drew A Lansdown: Consultant: Vericel, Inc. Educational support: Evolution Surgical/Arthrex. Committees: American Orthopedic Society for Sports Medicine, Arthroscopy Association of North America, and Orthopedic Research Society. Research support: American Orthopedic Society for Sports Medicine, Arthroscopy Association of North America, and Arthritis Foundation.

The other authors, their immediate families, and any research foundation with which they are affiliated have not received any financial payments or other benefits from any commercial entity related to the subject of this article.

References

- Abate M, Schiavone C, Di Carlo L, Salini V. Prevalence of and risk factors for asymptomatic rotator cuff tears in postmenopausal women. *Menopause* 2014;21:275–80. <https://doi.org/10.1097/GME.0b013e31829638e3>.
- Abate M, Schiavone C, Salini V, Andia I. Occurrence of tendon pathologies in metabolic disorders. *Rheumatology (Oxford)* 2013;52:599–608. <https://doi.org/10.1093/rheumatology/kes395>.
- Abboud JA, Kim JS. The effect of hypercholesterolemia on rotator cuff disease. *Clin Orthop Relat Res* 2010;468:1493–7. <https://doi.org/10.1007/s11999-009-1151-9>.
- Amit P, Kuiper JH, James S, Snow M. Does statin-treated hyperlipidemia affect rotator cuff healing or muscle fatty infiltration after rotator cuff repair? *J Shoulder Elbow Surg* 2021;30:2465–74. <https://doi.org/10.1016/j.jse.2021.05.014>.
- Artieda M, Cenaarro A, Junquera C, Lasierra P, Martínez-Lorenzo MJ, Pocióvi M, et al. Tendon xanthomas in familial hypercholesterolemia are associated with a differential inflammatory response of macrophages to oxidized LDL. *FEBS Lett* 2005;579:4503–12. <https://doi.org/10.1016/j.febslet.2005.06.087>.
- Barth J, Andrieu K, Fotiadis E, Hannink G, Barthelemy R, Saffarini M. Critical period and risk factors for retear following arthroscopic repair of the rotator cuff. *Knee Surg Sports Traumatol Arthrosc* 2017;25:2196–204. <https://doi.org/10.1007/s00167-016-4276-x>.
- Beason DP, Abboud JA, Kuntz AF, Bassora R, Soslowsky LJ. Cumulative effects of hypercholesterolemia on tendon biomechanics in a mouse model. *J Orthop Res* 2011;29:380–3. <https://doi.org/10.1002/jor.21255>.
- Beason DP, Hsu JE, Marshall SM, McDaniel AL, Temel RE, Abboud JA, et al. Hypercholesterolemia increases supraspinatus tendon stiffness and elastic modulus across multiple species. *J Shoulder Elbow Surg* 2013;22:681–6. <https://doi.org/10.1016/j.jse.2012.07.008>.
- Berglund DD, Kurowicki J, Giveans MR, Horn B, Levy JC. Comorbidity effect on speed of recovery after arthroscopic rotator cuff repair. *JSES Open Access* 2018;2:60–8. <https://doi.org/10.1016/j.jses.2017.12.003>.
- Beri A, Dwamena FC, Dwamena BA. Association between statin therapy and tendon rupture: a case-control study. *J Cardiovasc Pharmacol* 2009;53:401–4. <https://doi.org/10.1097/FJC.0b013e3181a0ce8b>.
- Cancienne JM, Brockmeier SF, Rodeo SA, Werner BC. Perioperative Serum Lipid Status and Statin Use Affect the Revision Surgery Rate After Arthroscopic Rotator Cuff Repair. *Am J Sports Med* 2017;45:2948–54. <https://doi.org/10.1177/0363546517717686>.
- Carroll M, Fryar C, Nguyen D. HDL, National Health and Nutrition Examination Survey: Total and High-density Lipoprotein Cholesterol in Adults: United States, 2015–2016. NCHS data brief, no. 290. Hyattsville, MD: National Center for Health Statistics; 2017.
- Chona DV, Lakomkin N, Lott A, Workman AD, Henry AC, Kuntz AF, et al. The timing of retears after arthroscopic rotator cuff repair. *J Shoulder Elbow Surg* 2017;26:2054–9. <https://doi.org/10.1016/j.jse.2017.07.015>.
- Chung SW, Kim JY, Kim MH, Kim SH, Oh JH. Arthroscopic repair of massive rotator cuff tears: outcome and analysis of factors associated with healing failure or poor postoperative function. *Am J Sports Med* 2013;41:1674–83. <https://doi.org/10.1177/0363546513485719>.
- Chung SW, Park JS, Kim SH, Shin SH, Oh JH. Quality of life after arthroscopic rotator cuff repair: evaluation using SF-36 and an analysis of affecting clinical factors. *Am J Sports Med* 2012;40:631–9. <https://doi.org/10.1177/0363546511430309>.
- Chung SW, Park H, Kwon J, Choe GY, Kim SH, Oh JH. Effect of Hypercholesterolemia on Fatty Infiltration and Quality of Tendon-to-Bone Healing in a Rabbit Model of a Chronic Rotator Cuff Tear: Electrophysiological, Biomechanical, and Histological Analyses. *Am J Sports Med* 2016;44:1153–64. <https://doi.org/10.1177/0363546515627816>.
- de Oliveira LP, Vieira CP, Da Ré Guerra F, de Almeida Mdos S, Pimentel ER. Statins induce biochemical changes in the Achilles tendon after chronic treatment. *Toxicology* 2013;311:162–8. <https://doi.org/10.1016/j.tox.2013.06.010>.
- Dolkart O, Liron T, Chechik O, Somjen D, Broch T, Maman E, et al. Statins enhance rotator cuff healing by stimulating the COX2/PGE2/EP4 pathway: an in vivo and in vitro study. *Am J Sports Med* 2014;42:2869–76. <https://doi.org/10.1177/0363546514545856>.
- Esenkaya I, Sakarya B, Unay K, Elmali N, Aydin NE. The influence of atorvastatin on tendon healing: an experimental study on rabbits. *Orthopedics* 2010;33:398. <https://doi.org/10.3928/01477447-20100429-06>.
- Feeley BT, Liu M, Ma CB, Agha O, Aung M, Carlin L, et al. Human Rotator Cuff Tears Have an Endogenous, Inducible Stem Cell Source Capable of Improving Muscle Quality and Function After Rotator Cuff Repair. *Am J Sports Med* 2020;48:2660–8. <https://doi.org/10.1177/0363546520935855>.
- Gaida JE, Alfredson L, Kiss ZS, Wilson AM, Alfredson H, Cook JL. Dyslipidemia in Achilles tendinopathy is characteristic of insulin resistance. *Med Sci Sports Exerc* 2009;41:1194–7. <https://doi.org/10.1249/MSS.0b013e31819794c3>.
- Galatz LM, Ball CM, Teefey SA, Middleton WD, Yamaguchi K. The outcome and repair integrity of completely arthroscopically repaired large and massive rotator cuff tears. *J Bone Joint Surg Am* 2004;86:219–24. <https://doi.org/10.2106/0004623-200402000-00002>.
- Garcia GH, Liu JN, Wong A, Cordasco F, Dines DM, Dines JS, et al. Hyperlipidemia increases the risk of retear after arthroscopic rotator cuff repair. *J Shoulder Elbow Surg* 2017;26:2086–90. <https://doi.org/10.1016/j.jse.2017.05.009>.
- Gwark JY, Sung CM, Na JB, Park HB. Outcomes of Arthroscopic Rotator Cuff Repair in Patients Who Are 70 Years of Age or Older Versus Under 70 Years of Age: A Sex- and Tear Size-Matched Case-Control Study. *Arthroscopy* 2018;34:2045–53. <https://doi.org/10.1016/j.arthro.2018.02.047>.
- Harada N, Gotoh M, Ishitani E, Kakuma T, Yano Y, Tataru D, et al. Combination of risk factors affecting retear after arthroscopic rotator cuff repair: a decision tree analysis. *J Shoulder Elbow Surg* 2021;30:9–15. <https://doi.org/10.1016/j.jse.2020.05.006>.
- Hast MW, Abboud JA, Soslowsky LJ. Exploring the role of hypercholesterolemia in tendon health and repair. *Muscles Ligaments Tendons J* 2014;4:275–9.
- Higgins JPT, Green S, editors. *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0; 2011. The Cochrane Collaboration. Available at: www.handbook.cochrane.org. Accessed September 11, 2021.
- Itoigawa Y, Wada T, Kawasaki T, Morikawa D, Maruyama Y, Kaneko K. Supraspinatus Muscle and Tendon Stiffness Changes After Arthroscopic Rotator Cuff Repair: A Shear Wave Elastography Assessment. *J Orthop Res* 2020;38:219–27. <https://doi.org/10.1002/jor.24469>.
- Jenssen KK, Lundgreen K, Madsen JE, Kvakestad R, Dimmen S. Prognostic Factors for Functional Outcome After Rotator Cuff Repair: A Prospective Cohort Study With 2-Year Follow-up. *Am J Sports Med* 2018;46:3463–70. <https://doi.org/10.1177/0363546518803331>.
- Jeong HY, Kim HJ, Jeon YS, Rhee YG. Factors Predictive of Healing in Large Rotator Cuff Tears: Is It Possible to Predict Retear Preoperatively? *Am J Sports Med* 2018;46:1693–700. <https://doi.org/10.1177/0363546518762386>.
- Kim YK, Jung KH, Kim JW, Kim US, Hwang DH. Factors affecting rotator cuff integrity after arthroscopic repair for medium-sized or larger cuff tears: a retrospective cohort study. *J Shoulder Elbow Surg* 2018;27:1012–20. <https://doi.org/10.1016/j.jse.2017.11.016>.
- Kurowicki J, Berglund DD, Momoh E, Disla S, Horn B, Giveans R, et al. Speed of recovery after arthroscopic rotator cuff repair. *J Shoulder Elbow Surg* 2017;26:1271–7. <https://doi.org/10.1016/j.jse.2016.11.002>.
- Kwon J, Kim SH, Lee YH, Kim TI, Oh JH. The Rotator Cuff Healing Index: A New Scoring System to Predict Rotator Cuff Healing After Surgical Repair. *Am J Sports Med* 2019;47:173–80. <https://doi.org/10.1177/0363546518810763>.

34. Lai J, Gagnier JJ. The Effect of Lipid Disorders on the Risk of Rotator Cuff Disease: A Systematic Review and Meta-Analysis. *JB JS Open Access* 2018;3:e0018. <https://doi.org/10.2106/JBJS.OA.18.00018>.
35. Le BT, Wu XL, Lam PH, Murrell GA. Factors predicting rotator cuff retears: an analysis of 1000 consecutive rotator cuff repairs. *Am J Sports Med* 2014;42:1134–42. <https://doi.org/10.1177/0363546514525336>.
36. Lee C, Liu M, Agha O, Kim HT, Feeley BT, Liu X. Beige FAPs Transplantation Improves Muscle Quality and Shoulder Function After Massive Rotator Cuff Tears. *J Orthop Res* 2020;38:1159–66. <https://doi.org/10.1002/jor.24558>.
37. Lee YS, Jeong JY, Park CD, Kang SG, Yoo JC. Evaluation of the Risk Factors for a Rotator Cuff Retear After Repair Surgery. *Am J Sports Med* 2017;45:1755–61. <https://doi.org/10.1177/0363546517695234>.
38. Lee S, Park I, Lee HA, Shin SJ. Factors Related to Symptomatic Failed Rotator Cuff Repair Leading to Revision Surgeries After Primary Arthroscopic Surgery. *Arthroscopy* 2020;36:2080–8. <https://doi.org/10.1016/j.arthro.2020.04.016>.
39. Li K, Deng G, Deng Y, Chen S, Wu H, Cheng C, et al. High cholesterol inhibits tendon-related gene expressions in tendon-derived stem cells through reactive oxygen species-activated nuclear factor- κ B signaling. *J Cell Physiol* 2019;234:18017–28. <https://doi.org/10.1002/jcp.28433>.
40. Li K, Deng Y, Deng G, Chen P, Wang Y, Wu H, et al. High cholesterol induces apoptosis and autophagy through the ROS-activated AKT/FOXO1 pathway in tendon-derived stem cells. *Stem Cell Res Ther* 2020;11:131. <https://doi.org/10.1186/s13287-020-01643-5>.
41. Lin TT, Lin CH, Chang CL, Chi CH, Chang ST, Sheu WH. The effect of diabetes, hyperlipidemia, and statins on the development of rotator cuff disease: a nationwide, 11-year, longitudinal, population-based follow-up study. *Am J Sports Med* 2015;43:2126–32. <https://doi.org/10.1177/0363546515588173>.
42. Liu X, Ning AY, Chang NC, Kim H, Nissenson R, Wang L, et al. Investigating the cellular origin of rotator cuff muscle fatty infiltration and fibrosis after injury. *Muscles Ligaments Tendons J* 2016;6:6–15. <https://doi.org/10.11138/mltj/2016.6.1.006>.
43. Makhni EC, Swart E, Steinhaus ME, Mather RC 3rd, Levine WN, Bach BR Jr, et al. Cost-Effectiveness of Reverse Total Shoulder Arthroplasty Versus Arthroscopic Rotator Cuff Repair for Symptomatic Large and Massive Rotator Cuff Tears. *Arthroscopy* 2016;32:1771–80. <https://doi.org/10.1016/j.arthro.2016.01.063>.
44. Mather RC 3rd, Koenig L, Acevedo D, Dall TM, Gallo P, Romeo A, et al. The societal and economic value of rotator cuff repair. *J Bone Joint Surg Am* 2013;95:1993–2000. <https://doi.org/10.2106/JBJS.L.01495>.
45. Matsuura F, Hirano K, Koseki M, Ohama T, Matsuyama A, Tsuiji K, et al. Familial massive tendon xanthomatosis with decreased high-density lipoprotein-mediated cholesterol efflux. *Metabolism* 2005;54:1095–101. <https://doi.org/10.1016/j.metabol.2005.03.014>.
46. O'Donnell EA, Fu MC, White AE, Taylor SA, Dines JS, Dines DM, et al. The Effect of Patient Characteristics and Comorbidities on the Rate of Revision Rotator Cuff Repair. *Arthroscopy* 2020;36:2380–8. <https://doi.org/10.1016/j.arthro.2020.05.022harada>.
47. Ozgurtas T, Yildiz C, Serdar M, Atesalp S, Kutluay T. Is high concentration of serum lipids a risk factor for Achilles tendon rupture? *Clin Chim Acta* 2003;331:25–8. [https://doi.org/10.1016/s0009-8981\(03\)00075-5](https://doi.org/10.1016/s0009-8981(03)00075-5).
48. Park HB, Gwark JY, Im JH, Jung J, Na JB, Yoon CH. Factors Associated with Atraumatic Posterosuperior Rotator Cuff Tears. *J Bone Joint Surg Am* 2018;100:1397–405. <https://doi.org/10.2106/JBJS.16.01592>.
49. Park HB, Gwark JY, Kwack BH, Jung J. Hypo-High-Density Lipoproteinemia Is Associated With Preoperative Tear Size and With Postoperative Retear in Large to Massive Rotator Cuff Tears. *Arthroscopy* 2020;36:2071–9. <https://doi.org/10.1016/j.arthro.2020.04.043>.
50. Rossi LA, Chahla J, Verma NN, Millett PJ, Ranalletta M. Rotator Cuff Retears. *JBJS Rev* 2020;8:e0039. <https://doi.org/10.2106/JBJS.RVW.19.00039>.
51. Tan JT, Ng MK, Bursill CA. The role of high-density lipoproteins in the regulation of angiogenesis. *Cardiovasc Res* 2015;106:184–93. <https://doi.org/10.1093/cvr/cvv104>.
52. Torgutalp ŞŞ, Babayeva N, Taş S, Dönmez G, Korkusuz F. Effects of hyperlipidemia on patellar tendon stiffness: A shear wave elastography study. *Clin Biomech (Bristol, Avon)* 2020;75:104998. <https://doi.org/10.1016/j.clinbiomech.2020.104998>.
53. Uezumi A, Ito T, Morikawa D, Shimizu N, Yoneda T, Segawa M, et al. Fibrosis and adipogenesis originate from a common mesenchymal progenitor in skeletal muscle. *J Cell Sci* 2011;124:3654–64. <https://doi.org/10.1242/jcs.086629>.
54. Virani SS, Alonso A, Benjamin EJ, Bittencourt MS, Callaway CW, Carson AP, et al. Heart disease and stroke statistics—2020 update: a report from the American Heart Associationexternal icon. *Circulation* 2020;141:e139–596. <https://doi.org/10.1161/CIR.0000000000000757>.
55. von Bahr S, Movin T, Papadogiannakis N, Pikuleva I, Rönnow P, Diczfalusy U, et al. Mechanism of accumulation of cholesterol and cholestanol in tendons and the role of sterol 27-hydroxylase (CYP27A1). *Arterioscler Thromb Vasc Biol* 2002;22:1129–35. <https://doi.org/10.1161/01.atv.0000022600.61391.a5>.
56. Wang Z, Liu X, Liu M, Jiang K, Kajimura S, Kim H, et al. β_3 -Adrenergic receptor agonist treats rotator cuff fatty infiltration by activating beige fat in mice. *J Shoulder Elbow Surg* 2021;30:373–86. <https://doi.org/10.1016/j.jse.2020.06.006>.
57. Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, Tugwell P. The Newcastle-Ottawa Scale (NOS) for assessing the quality if nonrandomized studies in meta-analyses. Ottawa Hospital Research Institute. Available at: http://www.ohri.ca/programs/clinical_epidemiology/oxford.htm. Accessed September 11, 2021.
58. Yamaguchi K, Ditsios K, Middleton WD, Hildebolt CF, Galatz LM, Teefey SA. The demographic and morphological features of rotator cuff disease. A comparison of asymptomatic and symptomatic shoulders. *J Bone Joint Surg Am* 2006;88:1699–704. <https://doi.org/10.2106/JBJS.E.00835>.
59. Yang Y, Lu H, Qu J. Tendon pathology in hypercholesterolaemia patients: Epidemiology, pathogenesis and management. *J Orthop Translat* 2018;16:14–22. <https://doi.org/10.1016/j.jot.2018.07.003>.
60. Yang Y, Qu J. The effects of hyperlipidemia on rotator cuff diseases: a systematic review. *J Orthop Surg Res* 2018;13:204. <https://doi.org/10.1186/s13018-018-0912-0>.
61. Zeng GJS, Lee MJH, Chen JY, Ang BFH, Hao Y, Lie DTT. Dyslipidemia With Perioperative Statin Usage Is Not Associated With Poorer 24-Month Functional Outcomes After Arthroscopic Rotator Cuff Surgery. *Am J Sports Med* 2020;48:2518–24. <https://doi.org/10.1177/0363546520937266>.
62. Zhao J, Luo M, Pan J, Liang G, Feng W, Zeng L, et al. Risk factors affecting rotator cuff retear after arthroscopic repair: a meta-analysis and systematic review [published online ahead of print, 2021 Jun 2]. *J Shoulder Elbow Surg* 2021;30:2660–70. <https://doi.org/10.1016/j.jse.2021.05.010>.